

## Mediterranean Marine Science

Vol 25, No 1 (2024)

VOL 25, No 1 (2024)



**Changes in organic carbon properties during intense plankton blooms and macroaggregate formation in the coastal Adriatic Sea, Croatia (case studies in 2020-2022)**

*NIKI SIMONOVIC, MARIJA MARGUS, PAOLO PALIAGA, ANDREA BUDIŠA, IRENA CIGLENEČKI*

doi: [10.12681/mms.35082](https://doi.org/10.12681/mms.35082)


### To cite this article:

SIMONOVIC, N., MARGUS, M., PALIAGA, P., BUDIŠA, A., & CIGLENEČKI, I. (2024). Changes in organic carbon properties during intense plankton blooms and macroaggregate formation in the coastal Adriatic Sea, Croatia (case studies in 2020-2022). *Mediterranean Marine Science*, 25(1), 160–178. <https://doi.org/10.12681/mms.35082>


## Changes in organic carbon properties during intense plankton blooms and macroaggregate formation in the coastal Adriatic Sea, Croatia (case studies in 2020–2022)

Niki SIMONOVIĆ, Marija MARGUŠ, Paolo PALIAGA, Andrea BUDIŠA and Irena CIGLENEČKI


*Mediterranean Marine Science*, 25 (1) 2024



**Fig. S1:** Calibration plot (linear scale) of  $\Delta i$  versus Triton-X-100 in 0.55 M NaCl. ( $\Delta i$  - adsorption effect measured as the difference between capacity current for base electrolyte and aqueous solutions of surfactant (A).



**Fig. S2:** A.c. voltammograms of the water column sample of Rogoznica Lake sampled at 5m depth, where intense biological activity was recorded. Voltammetric curves revealed the typical desorption wave at around -1.6 V vs. Ag/AgCl in the filtered fractions smaller than 0.45 and 0.22  $\mu\text{m}$ . In a fraction smaller than 0.22  $\mu\text{m}$ , due to the presence of colloidal, highly surface-active OM, the desorption wave was even more expressed and for about 50 mV more negatively positioned. Experimental conditions: accumulation time,  $t_a = 120$  s at -0.6 V vs. Ag|AgCl.



**Fig. S3:** Physico-chemical properties of the Rogoznica Lake water column on October, 20 2020 (A) and June 28th, 2021 (B). NSA parameter (SAS/DOC ratio) was determined up to the chemocline layer, considering that SAS cannot be measured in an anoxic sample due to reduced sulphur species interference (Simonović *et al.*, 2023).

**Table S1.** Organic carbon properties (DOC, SAS, NSA) and plankton abundance in surface samples during: the red tide caused by dinoflagellate *Noctiluca scintillans* (A) and zooplankton bloom of invasive ctenophore *Mnemiopsis leidyi* (B).

A		Coordinates	DOC (mg L <sup>-1</sup> )	SAS (mg L <sup>-1</sup> )	NSA	Abundance (cells L <sup>-1</sup> )
Sampling date	Sampling location					
31.3.2021.	Port of Rovinj-Rovigno*	45.08097°N, 13.63473°E	106.9	13.5	0.127	12008
31.3.2021.	Old Town*	45.08299°N, 13.63195°E	105.8	8.37	0.079	33453
31.3.2021.	Valdibora Bay*	45.09021°N, 13.63339°E	318.1	14.3	0.045	72010
15.4.2021.	Škaraba Bay*	45.06511°N, 13.63845°E	236.3	15.4	0.065	78302
16.4.2021.	Lone Bay*	45.06988°N, 13.63295°E	11.39	2.88	0.253	46800
23.4.2021.	ACI Marina Rovinj-Rovigno*	45.07566°N, 13.63502°E	13.65	2.58	0.211	48696

\*measured in dilution.

B		Coordinates	DOC (mg L <sup>-1</sup> )	SAS (mg L <sup>-1</sup> )	NSA	Abundance (ind m <sup>-3</sup> )
Sampling date	Sampling location					
24.8.2020.			0.959	0.147	0.153	20
25.8.2020.			1.602	0.162	0.101	20
27.8.2020.			1.494	0.203	0.136	0.1
27.8.2020.	Valdibora Bay	45.09021°N, 13.63339°E	2.222	0.334	0.150	60
27.8.2020.			6.768	1.338	0.198	80
27.8.2020.			8.581	1.453	0.169	100
7.9.2020.			2.817	0.611	0.217	400

No.ind- number of individuals.

**Table S2.** Organic carbon properties (DOC, SAS, NSA) of RL (from 2020 to 2022) surface oxic layer (A), chemocline (B) and samples with maximal oxygen saturation (C). Display C also represent samples where specific desorption peak (represented in Fig. 3.A.e) was measured by a.c. voltammetry method.

<i>A</i>	<i>Rogoznica Lake</i> (43.53091°N, 15.95876°E)		DOC (mg L <sup>-1</sup> )	SAS (mg L <sup>-1</sup> )	NSA		
Sampling date							
16.12.2020.			2.378	0.242	0.102		
27.1.2021.			2.125	0.190	0.089		
24.3.2021.			2.089	0.177	0.085		
28.6.2021.	Surface layer [0-2]m		1.994	0.193	0.097		
6.9.2021.			1.742	0.262	0.150		
12.10.2021.			1.178	0.246	0.138		
23.11.2021.			1.984	0.247	0.125		
16.2.2022.			1.691	0.211	0.130		
6.4.2022.			1.577	0.209	0.133		
12.5.2022.			1.576	0.198	0.126		
18.7.2022.			2.214	0.179	0.081		
<i>B</i>			<i>Rogoznica Lake</i> (43.53091°N, 15.95876°E)		DOC (mg L <sup>-1</sup> )	SAS (mg L <sup>-1</sup> )	NSA
Sampling date							
27.1.2021.	Chemocline		2.253	0.202	0.089		
28.6.2021.			2.592	0.198	0.073		
16.9.2021.			1.742	0.243	0.139		
12.10.2021.			1.714	0.235	0.137		
23.11.2021.			1.955	0.214	0.109		
12.10.2021.			1.365	0.163	0.119		
23.11.2021.			1.159	0.163	0.102		
18.7.2022.			2.388	0.150	0.063		
<i>C</i>	<i>Rogoznica Lake</i> (43.53091°N, 15.95876°E)	Depth (m)	DOC (mg L <sup>-1</sup> )	SAS (mg L <sup>-1</sup> )	NSA		
Sampling date							
16.2.2022.	Oxygen saturation maximum ( $\geq 100\%$ )	0	2.018	0.278	0.138		
6.4.2022.		0	1.638	0.235	0.144		
		2	1.515	0.183	0.121		
12.5.2022.		5	2.044	0.333	0.163		
18.7.2022.		0	2.823	0.229	0.081		
		6	1.809	0.200	0.111		
10.11.2022.		0	4.175	0.258	0.062		