



Mediterranean Marine Science

Vol 24, No 3 (2023)

VOL 24, No 3 (2023)



New kid in town: Pinna rudis spreads in the eastern Mediterranean

MARIA ZOTOU, ORESTIS PAPADAKIS, GAETANO CATANESE, YOLANDA STRANGA, MICHAIL RAGKOUSIS, THODOROS E. KAMPOURIS, ROXANI NAASAN AGA - SPYRIDOPOULOU, EVANGELOS PAPADIMITRIOU, DROSOS KOUTSOUBAS, STELIOS KATSANEVAKIS

doi: <u>10.12681/mms.35343</u>

To cite this article:

ZOTOU, M., PAPADAKIS, O., CATANESE, G., STRANGA, Y., RAGKOUSIS, M., KAMPOURIS E., T., NAASAN AGA -SPYRIDOPOULOU, R., PAPADIMITRIOU, E., KOUTSOUBAS, D., & KATSANEVAKIS, S. (2023). New kid in town: Pinna rudis spreads in the eastern Mediterranean. *Mediterranean Marine Science*, *24*(3), 709–721. https://doi.org/10.12681/mms.35343 Mediterranean Marine Science Indexed in WoS (Web of Science, ISI Thomson) and SCOPUS The journal is available on line at http://www.medit-mar-sc.net www.hcmr.gr DOI: http://doi.org/10.12681/mms.35343

New kid in town: Pinna rudis spreads in the eastern Mediterranean

Maria ZOTOU¹, Orestis PAPADAKIS¹, Gaetano CATANESE^{2,3}, Yolanda STRANGA¹, Michail RAGKOUSIS¹, Thodoros E. KAMPOURIS¹, Roxani NAASAN AGA-SPYRIDOPOULOU⁴, Evangelos PAPADIMITRIOU¹, Drosos KOUTSOUBAS¹ and Stelios KATSANEVAKIS¹

 ¹ Department of Marine Sciences, University of the Aegean, Lofos Panepistimiou, 81100 Mytilene, Greece
 ² IRFAP-LIMIA, Institut de Recerca i Formació Agroalimentària i Pesquera de les Illes Balears, Laboratori d'Investigacions Marines i Aqüicultura, Govern de les Illes Balears, 07157 Port d'Andratx, Spain

³ INAGEA-UIB, Agro-Environmental and Water Economics Research Institute, University of the Balearic Islands, 07122 Palma, Spain ⁴ iSea, Environmental Organisation for the Preservation of the Aquatic Ecosystems, Kritis 12, 54645 Thessaloniki, Greece

Corresponding author: Maria ZOTOU; m.maryzotou@gmail.com

Contributing Editor: Fabio CROCETTA

Received: 04 September 2023; Accepted: 30 November 2023; Published online: 19 December 2023

Abstract

In the Mediterranean Sea, the genus *Pinna* encompasses two large fan-shaped bivalve species, *Pinna nobilis* and *Pinna rudis*. Historically, both species coexisted in the western Mediterranean until a mass mortality event (MME) brought *P. nobilis* to the brink of extinction. Notably, *P. rudis* remained unaffected by the MME, and its recent successful recruitment and further spread have been hypothesized to be linked to the local extinction of *P. nobilis*. Although *P. rudis* has been sparsely recorded in the eastern Mediterranean Sea (with some of these records being doubtful), reports emerging in the summer of 2023 from researchers and citizens have unveiled its sudden spread in the region. This study documents the recent presence of *P. rudis* within Greek waters through a dedicated survey and molecularly confirmation through two distinct molecular methods. Information derived from a citizen science initiative, following photo-identification of the species was also included. Furthermore, an updated review of the distribution of *P. rudis* in the Mediterranean Sea, integrating data from literature and online repositories is provided. This research confirms the recent spread of *P. rudis* in Crete, the Greek Ionian Sea, and Cyprus (first verified records of the species in the regions), in marine areas where *P. nobilis* has become extinct, further strengthening the hypothesis that *P. rudis* has benefited from the collapse of *P. nobilis* populations.

Keywords: Pinnidae; Pinna rudis; Distribution; Mediterranean Sea; Molecular identification.

Introduction

The family Pinnidae comprises large bivalve mollusks with a distinctive fan-like shape, anchored in the substrate by byssus threads (Gvozdenović et al., 2019; Deidun et al., 2022) and consists of two genera, Atrina and Pinna (Lemer et al., 2014). The genus Pinna encompasses the two largest bivalve species of the Mediterranean Sea, Pinna nobilis (Linnaeus, 1758) and Pinna rudis (Linnaeus, 1758) (Zavodnik et al., 1991; Galinou-Mitsoudi et al., 2006). These congeneric species, while sharing certain characteristics, also exhibit notable differences. Pinna nobilis can attain an impressive shell length of up to 120 cm (Huber, 2010), and its maximum recorded lifespan exceeds 45 years (Rouanet et al., 2015). In contrast, P. rudis reaches a maximum shell length of only 50 cm (García-March & Kersting, 2006) and a maximum age of 31 years (Nebot-Colomer et al., 2016).

Although both species inhabit the Mediterranean Sea,

Medit. Mar. Sci., 24/3 2023, 709-721

P. nobilis is confined solely to this region while *P. rudis* has a broader distribution, occurring also along the African Atlantic coasts and Macaronesian islands (Barreiros et al., 2016; Gvozdenović et al., 2019; Kersting & Ballesteros, 2021). The habitat preferences and depth distributions of the two species differ: Pinna nobilis is closely associated with Posidonia oceanica meadows and soft substrates (Addis et al., 2009; Vázquez-Luis et al., 2014; Deudero et al., 2015) and is found at depths ranging from 0.5 m to 60 m (Garcia-March & Kersting, 2006; Katsanevakis, 2007; Vázquez-Luis et al., 2017). Pinna rudis can also inhabit similar habitats (such as seagrass meadows, maerl, detritus and boulder beds, as described in Kersting & García-March, 2017), but shows a preference for gravel bottoms and hard substrates (Poppe & Goto, 1993) to depths down to 40 m (Sempere et al., 2006). Their populations overlap in areas where both species coexist, particularly in the western Mediterranean (Kersting & García-March, 2017). This overlap occasionally

leads to the misidentification of P. rudis as P. nobilis (e.g., Gvozdenović et al., 2019; Kersting & Ballesteros, 2021). Morphologically, the most distinct feature of P. rudis lies in its shell, which appears more robust and triangular, with fewer but larger protruding scales compared to that of P. nobilis (Cosentino & Giacobbe, 2006). The external color of *P. rudis* valves is reddish brown, and valves possess a symmetrical, triangular shape with transparent ends, notably delicate, featuring 5-10 radiating ribs, neatly arranged in rows. Adult P. nobilis tends to be larger and lacks the protuberances present on the surface of the shell in P. rudis. Additionally, P. rudis typically displays a white and iridescent mantle border, while P. nobilis exhibits a pink one. Both species have similar-sized nacre lobes (García-March & Vicente, 2006; Gvozdenović et al., 2019).

Pinna spp. play a unique and crucial role in the ecosystem. Pinna nobilis acts as a host to many marine species, including invasive species, on and within its shell, providing a hard substrate amidst the soft bottom surrounding, functioning as an ecosystem engineer (Giacobbe, 2002; Rabaoui et al., 2009; Katsanevakis et al., 2016; Kersting & García-March, 2017; Çinar et al., 2021), as does P. rudis shells (Cosentino & Giacobbe, 2007; Lopes et al., 2020). Furthermore, Pinna spp., being filter-feeding organisms, can decrease water turbidity and retain organic material (Trigos et al., 2014; Basso et al., 2015). From a cultural ecosystem services perspective, they are considered attractions for SCUBA divers (Morrocco et al., 2018; Deidun et al., 2022) since the species are emblematic, adding complexity and dimension to the marine environment. Both species are considered endangered due to various anthropogenic activities threatening their populations, such as poaching, anchoring, trawling, and habitat degradation (Katsanevakis, 2007; Barea-Azcón et al., 2008; Deudero et al., 2015; Kersting et al., 2019). Consequently, they have been included in Annex II of the Bern and the Barcelona Conventions.

Since 2016, P. nobilis populations have experienced a substantial decline due to mass mortality events (MMEs) that emerged across the Mediterranean Basin (Vázquez-Luis et al., 2017; Kersting et al., 2019; Zotou et al., 2020; Özalp & Kersting, 2020; Katsanevakis et al., 2021). The MMEs have been linked to the presence of the parasite Haplosporidium pinnae (Catanese et al., 2018; Grau et al., 2021), although Mycobacterium sp. and Vibrio mediterranei were also detected in dead or moribund individuals (Prado et al., 2020; Carella et al., 2020; 2023). The mortality rates reached 100% in most Mediterranean locations (Katsanevakis et al., 2021). As a result, in 2019, the species' conservation status was evaluated as "critically endangered" on the IUCN Red List (Kersting et al., 2019), and the attention of marine scientists focused on actions to prevent the global extinction of P. nobilis. Remarkably, P. rudis remained unaffected by the pathogen responsible for the high mortality observed in its congeneric species (Catanese et al., 2018). Recently, in regions where the two species co-existed, an increase in P. rudis recruitment has been reported (Kersting & Ballesteros, 2021) and small populations of P. rudis have been observed in areas where the species had not been recorded previously (Donato *et al.*, 2021; present study).

Unfortunately, *P. rudis* is understudied compared to *P. nobilis*, with scarce publications assessing the population dynamics or the distribution of the species in the Mediterranean (Nebot-Colomer *et al.*, 2016; Kersting & García-March, 2017; Gvozdenović *et al.*, 2019). The first mapping of *P. rudis* occurrences in the Mediterranean, based on the existing literature, was published in 2019 (Gvozdenović *et al.*, 2019). Most of these observations were concentrated in the western Mediterranean, with the majority of studies being conducted in Spain. Notably, in Greece, only two research articles have mentioned the presence of *P. rudis* in the Aegean and the Ionian Sea (Salomidi *et al.*, 2016; Galinou-Mitsoudi *et al.*, 2023).

Citizen science represents a collaborative and inclusive approach that engages both professional scientists and citizens (Kelly *et al.*, 2021). The integration of citizen science into monitoring programs offers valuable information while reducing the time, effort, and costs associated with scientific fieldwork. Simultaneously, this integration is viewed as a pivotal process for nurturing a new generation of citizens with a profound comprehension and appreciation of the natural environment (Mokos *et al.*, 2020). In recent years, an increasing number of research papers have incorporated citizen science (Cabanellas-Reboredo *et al.*, 2019; Deidun *et al.*, 2022; Kampouris *et al.*, 2022), and networks of environmentally conscious citizens willingly share their valuable observations with scientists.

The primary aim of this study was to assess if the distribution of P. rudis is changing in the Mediterranean Sea after the MME of P. nobilis. To this aim, an assessment of the distribution of P. rudis in Greek waters, particularly southern Crete, combined with opportunistic records from other areas, was conducted. To describe the changes in the range of the species, the study also includes a comprehensive review of its past distribution in the Mediterranean Sea. The key objectives of this research were: i) to document the recent expansion of P. rudis in the eastern Mediterranean, with a focus in Crete; ii) to provide molecular confirmation of the correct identification of the species; iii) to evaluate information arising from our citizen science network about Pinna populations in Greece and Cyprus; and iv) to map the present distribution of P. rudis in the Mediterranean Sea utilizing existing literature and including supplementary data sourced from online databases (GBIF, WoRMS).

Materials and Methods

Main study area

Crete, the largest island in Greece and the fifth largest in the Mediterranean Sea, boasts a coastline that spans more than 1046 km (Alexandrakis, 2014). Positioned approximately 160 km south of the Greek mainland, it occupies a strategic location within the eastern portion of the Mediterranean basin. The island has an elongated shape, stretching approximately 260 km from its eastern to western extents, with a width fluctuating between 12 and 60 km from north to south. The southern Cretan coastal waters are relatively unaffected by significant river flows; consequently, this region stands out as one of the most oligotrophic regions within the Mediterranean. This is manifested by notably low levels of primary productivity and chlorophyll concentrations (Karakasis & Eleftheriou 1998; Psarra et al., 2000; Bosc et al., 2004). Sea surface temperature along the southern Cretan coastal waters spans from 16.5°C during winter to a peak of 27.5°C in summer (2010-2023 E.U. Copernicus Marine Service Data). Salinity levels within this region span from 39.2 to 39.5 psu (2021-2023 E.U. Copernicus Marine Service Data). The circulation patterns in the region are influenced by multiple factors, including wind patterns, water temperature, and the larger-scale Mediterranean circulation. It is a complex oceanographic circulation system where water masses from the Western Mediterranean, the Ionian, and the Levantine Seas interact (Zodiatis 1993; Akpinar et al., 2016). The local circulation structures of south Crete exhibit a series of permanent and recurring gyres and jets, including the Cretan Cyclone, the Mid Mediterranean Jet, and the Ierapetra Anticyclone (Robinson et al., 1992; Fusco et al., 2003).

Sampling and documentation

Samples for molecular identification were collected through SCUBA diving at a site with high *Pinna* population densities, pinpointed by a citizen scientist (Site #6, Kalypso, S. Crete; Table 1) in July 2023. The sampling methodology adhered to the non-lethal technique proposed by Sanna *et al.* (2013), to reduce any damage to the shell and soft tissues of the specimens (Scarpa *et al.*, 2020). The collected tissue samples were stored in 10ml tubes and preserved in 99.8% ethanol. In total, eight mantle samples were acquired from living individuals. All sampled individuals were photographed *in situ* with a digital camera (Sony RX100 III).

Furthermore, tens of individuals were documented at several other surveyed sites in southern Crete during the same period. To visually depict the distribution of the documented populations, a map was generated using QGIS software (version 3.24.1). Additional observations contributed by both researchers and citizen scientists from other locations in Greek waters, confirmed through photographic evidence, were also mapped.

Molecular confirmation

Total genomic DNA was extracted from the eight mantle biopsies using the DNA NucleoSpin® Tissue extraction kit following the instructions provided by the manufacturer. Two distinct methods were carried out for species identification. A first analysis was performed following the method described by Catanese et al. (2022a) but applied directly to the extracted DNAs from the bivalve samples (approximately 30 ng). The method is based on a multiplex-PCR which selectively amplifies amplicons of different lengths for each P. nobilis (a fragment of 175 bp using the primers Pnob12SF/Pnob12SR) and P. rudis species (a fragment of 123 bp using the primer pair Prud16SF/ Prud16SR). The method employs species-specific primers designed on the mitochondrial genes 12S rDNA and 16S rDNA, which have been previously characterized for their inter-specific differences (Catanese et al., 2022b). In addition, a 28S rDNA fragment (205 bp using the primers Pinna28SF/ Pinna28SR) is included in the multiplex-PCR as a PCR positive control of amplification for Pinna sp. and Atrina sp.

For the second method, DNA samples underwent PCR amplification and sequencing targeting a segment of 1150 base pairs (bp) within the 28S rDNA nuclear gene. Two distinct regions were amplified using the primers 28S_Pinna_F/28S_Pinna_R to target a first portion (fragments A and B in Vázquez-Luis *et al.*, 2021) and the primers 28Sa/28srd5b (fragment C in Vázquez-Luis *et al.*, 2021) to target a second portion. The PCR reactions were conducted in a total volume of 20 μ L. The reaction mixture consisted of 10 μ L of KAPA Taq Ready Mix PCR kit, 0.4 μ L of each primer (stock 20 μ M), 1 μ L of DNA (approximately 200 ng/ μ L), and the remaining volume was made up with water.

Table 1. Locations with coordinates in decimal degrees indicating the start of each dive (and maximum distance of approximately 200 m from reported individuals), and dates when each location was visited. Mantle samples for molecular confirmation were collected from Kalypso (Station #6) and are marked with an asterisk.

Station	Location	Latitude (N)	Longitude (E)	Sampling date
1	Ag. Antonios	35.245514°	23.836447°	5/7/2023
2	Domata	35.229138°	23.905206°	5/7/2023
3	Glika nera	35.200247°	24.106286°	6/7/2023
4	Plakias	35.191408°	24.373022°	8/7/2023
5	Myrtos	34.996291°	25.572163°	10/7/2023
6	Kalypso*	35.172628°	24.398562°	8/7/2023

The temperature profile for the PCR reaction was as follows: an initial denaturation step at 94°C for 2 minutes, succeeded by 40 cycles of denaturation at 94°C for 30 seconds, annealing at 60°C for 20 seconds, and extension at 72°C for 90 seconds. The PCR products were separated on 2% agarose gels in TAE 1x buffer (w/v). The gels were stained with GelRed and included a High-Ranger DNA ladder size standard 50 bp (for multiplex PCR) or 1kb (for nuclear PCR amplification). The amplified DNA fragments were visualized using a UV transilluminator and excised from the agarose gel. Subsequently, they were purified using the Metabion International mi-Gel Extraction Kit, following the manufacturer's instructions. The two obtained PCR fragments underwent bi-directional sequencing using the Applied Biosystems 3130xl DNA automated sequencer. The sequences were edited and aligned using the BioEdit version 7.2.5 software (Hall, 1999) and MEGA X (Kumar et al., 2018).

The citizen science network

For several years, a network of dedicated SCUBA divers has actively collaborated with the 'Marine Biodiversity and Ecosystems Management' lab of the University of the Aegean, sharing valuable information regarding marine biodiversity. Additionally, within the framework of another research project with a specific focus on sharing information regarding the endangered P. nobilis populations in Greece, the pre-existing citizen-science Facebook group Red Fish Project (https://www.facebook.com/groups/481084575908558/), that monitors threatened aquatic species was utilized, following the protocols and principles described by Kampouris et al. (2022). Presence data collected from the environmental organization iSea, through their Facebook group (https:// www.facebook.com/groups/104915386661854) and their field actions were also incorporated in the dataset. As of June 2023, there have been notable citizen observations of live Pinna individuals in locations where P. nobilis has faced local extinction due to recent MMEs; however, upon careful examination of the photographs submitted alongside the reports, it became evident that while the genus was accurately identified, the species in question was not P. nobilis, which is typically found in Greece, but rather P. rudis. To investigate this further and validate the observations, we communicated with the citizens who had made the "uncommon" pinnid observations, and subsequently, a survey was organized in southern Crete to collect samples for molecular species identification.

Review of the past distribution of **Pinna rudis** *in the Mediterranean*

A systematic literature review was undertaken to retrieve information on the past distribution of *P. rudis* in the Mediterranean, employing Scopus and Google Scholar as the primary search platforms. For the Scopus and Google Scholar search, the keywords "*Pinna rudis*" and "Mediterranean Sea" were used to query titles and abstracts, together and apart. The Google Scholar search was restricted to the initial 100 documents (sorted by relevance), as subsequent results did not yield any additional relevant information. Studies conducted beyond the Mediterranean Sea were omitted from the analysis. To augment the dataset, additional records were obtained by backward searching a systematic review article by Gvozdenović *et al.* (2019). Additionally, *P. rudis* georeferenced data from GBIF and WoRMS were included in the review. Subsequent to the elimination of duplicates, all available records from literature and online databases until 31 July 2023 were assimilated. The spatial data was visualized on a map using the QGIS software (version 3.24.1).

Results

Documentation of the population in south Crete

During the visual observations conducted in July 2023 in southern Crete (Fig. 1), *Pinna rudis* individuals were found at six locations (Table 1, Table 2). A total tally of 31 individuals was recorded, of which 28 were alive whilst three were dead. Of the total population in southern Crete, 24 individuals were classified as adults, constituting 77.4% of all documented *P. rudis* individuals, while seven individuals were characterized as presumed juveniles, based on their small size (less than 6 cm shell height; Fig. 2). Notably, all individuals were found on rocky or pebbly substrates at depths ranging between 5 and 20 meters.

Among these locations, Kalypso (Site #6) hosted the largest population, with 19 individuals. This site was the only sampling site where dead (presumed juvenile) individuals were encountered; however, the reason for their mortality was unclear. Samples for molecular analyses were also collected from Site #6. In the three locations (Sites #2, #3, #5) where fewer individuals were observed (1 or 2 individuals/site), the presence of presumed juvenile individuals was consistently noted. These findings provide evidence for the establishment of a *P. rudis* population in southern Crete, characterized by the successful recruitment of juveniles and the presence of adult individuals.

Additional locations

During the summer months of 2023, *P. rudis* individuals were occasionally recorded across various locations in Greece. A total of 29 individuals was documented at 17 sites from three locations (Crete and Antikythera, Ionian Sea, Cyprus) (Fig. 3). Specifically, one individual was observed and photographed by one of the authors (SK) on the island of Antikythera at 5 m depth on a rocky substrate in August 2023. In October 2023, another individual was reported by a researcher (D. Poursanidis, pers, comm,) on the northern coast of Crete at 15 m depth. An additional ten individuals were recorded by citizen



Fig. 1: Photographs taken during sampling in southern Crete, Greece: a, d: View from above of a *Pinna rudis* individual with visible mantle. b, c: *P. rudis* individuals with distinctive shell ornamentation. e: Non-lethal mantle sampling for molecular confirmation of the species.

Station	Location	Living individuals	Dead individuals	Juveniles	Adults
1	Ag. Antonios	3	-	1	2
2	Domata	2	-	1	1
3	Glika nera	1	-	1	-
4	Plakias	4	-	-	4
5	Myrtos	2	-	1	1
6	Kalypso	16	3	3	16

Table 2. Number of individuals found in sampled locations, categorized by age (adult/presumed juvenile) and status (alive/dead).



Pinna rudis population in South Crete

Fig. 2: Population characteristics (adult/presumed juvenile individuals) and their status (alive/dead) in visited locations of South Crete.



Fig. 3: Photograph identification of opportunistic records made by researchers or citizen scientists. a: Print screen of a citizen scientist's post on Red Fish Project with a living individual found in southern Crete. b: Photograph taken by a citizen scientist at Parga (Ionian Sea). c: Photograph of a living specimen taken by S. Katsanevakis at Antikythera Island.

scientists participating in the Red Fish Project in south and east Crete and western Cyprus during the summer of 2023. These individuals were found across five dive sites at depths between 4 and 35 m. The environmental organization iSea documented seven individuals in three islands of the Ionian Sea (Paxos, Corfu, and Ithaca).

Furthermore, reports from citizens, supported by photo-identification of the individuals, indicated the presence of *P. rudis* in other areas of the Ionian Sea. Near Parga, ten individuals were spotted across three dive sites, between 2 and 17 m deep. There was also information on *P. rudis* presence at Kefalonia Island (Ionian Sea), but these individuals were not photographed; thus the information was not included in the distribution map of the species' presence in the Greek seas (Fig. 4). Data about locations and the recorded individuals by site can be found in Table 3.

Molecular confirmation

All 205 bp fragments from the 28S rDNA gene of the examined samples were successfully amplified during the multiplex-PCR tests, serving as a positive control for the two *Pinna* species and *Atrina* spp. Moreover, only the 123 bp PCR product, specific for the 16S rDNA gene of *P. rudis*, was obtained from the eight samples. No cross or simultaneous amplification of the two species-specific fragments were observed in any PCR.

In the PCR targeting the 28S rDNA nuclear gene, two amplicons of differing lengths, approximately 900 bp and 350 bp, were consistently generated in all samples. Upon sequence analysis, the same haplotype was identified across both fragments for all examined individuals. BLAST analysis revealed that the obtained sequences exhibited a 100% identity similarity with a sequence previously deposited in GenBank under accession number LC634517, obtained from *P. rudis*. Analyzing the chromatograms of the 28S rDNA sequences, the nine diagnostic nucleotide sites for the identification of *Pinna* species or putative hybrids as described by Vázquez-Luis *et al.* (2021), showed the typical pattern of *P. rudis* species. Additionally, no insertion/deletion events were detected in the obtained sequences.

Review of the past distribution of Pinna rudis *in the Mediterranean*

A comprehensive total of 399 records containing georeferenced information on *P. rudis* distribution in the Mediterranean were compiled. Among these records, 118 (30%) were derived from the literature review and forward searches of scientific articles, while most data (281 records, 70%) were acquired from online databases (Fig. 5).

Among the contributing countries, Spain stood out with 226 records, followed by France (75), Italy (49), and Algeria (24). Conversely, countries with limited records were Croatia (1 record), Lebanon (1), Libya (1), Montenegro (2), Tunisia (2), Morocco (2), Greece (6), and Malta (10) (Fig. 6).

In terms of ecoregions (as classified in Spalding *et al.*, 2007), the Western Mediterranean Sea exhibited the most substantial presence of *P. rudis* records, encompassing 70.4% of all records. The Alborán Sea accounted for 18.6% of the records, while the remaining ecoregions collectively contributed 11.0% (Ionian Sea 7.3%, Adriat-



Fig. 4: The map displays the herein documented records of *Pinna rudis* in Greece and Cyprus, with circle size representing the number of individuals found in each location. The green color represents the documented population of southern Crete by our dedicated survey, while the pink color indicates opportunistic records made by researchers or citizen scientists. The opacity of symbols is set to 60% to highlight overlapping observations.

Location	Latitude (N)	Longitude (E)	Depth range	Number of indi- viduals
Antikythera Island	35.883338°	23.292293°	5 m	1
Parga (Ionian Sea)	39.278193°	20.386764°	2-17 m	4
Parga (Ionian Sea)	39.282090°	20.404204°	2-17 m	3
Parga (Ionian Sea)	39.279557°	20.400047°	2-17 m	3
Ierapetra (S. Crete)	34.994694°	25.750889°	25 m	1
Ierapetra (S. Crete)	34.999222°	25.798389°	4-11m	3
Sitia (S. Crete)	35.011583°	26.154472°	10 m	1
Heraklion (N. Crete)	35.401219°	25.03381°	15 m	1
Paksimadia Isles (S. Crete)	35.003915°	24.601525°	12-35 m	4
Pegia (W. Cyprus)	34.856784°	32.348314°	10 m	1
Peristeres Island (North Straits Corfu)	39.792189°	19.95975°	4 m	1
Paxos Island	39.225982°	20.164533°	2 m	1
SW. Corfu Island	39.417611°	19.903028°	6-15 m	1
SW. Corfu Island	39.393917°	19.989389°	6-15 m	1
SW. Corfu Island	39.427598°	19.9374°	2.5-5 m	1
SW. Corfu Island	39.410789°	19.912179°	2.5-5 m	1
Ithaca Island (Kioni port)	38.44959°	20.690681°	3 m	1

Table 3. Additional locations where opportunistic records of *Pinna rudis* individuals occurred. Locations with coordinates in decimal degrees indicating the start of each dive (and maximum distance of approximately 200 m from reported individuals), depth range, and number of individuals found in each site.

ic Sea 1.8%, Aegean Sea 1.3%, Tunisian Plateau/Gulf of Sidra 0.5%, Levantine Sea 0.3%; Fig. 7).

Overall, the presence of *P. rudis* in the western Mediterranean Sea has been thoroughly documented and studied, while information about the eastern part of the basin is scarce. This highlights the current knowledge that the distribution range of the species is primarily concentrated in the western Mediterranean Basin (Fig. 8). Regarding the six past records within the Greek seas, five records corresponded to a single article (Salomidi Data contribution per source



Fig. 5: Contribution of sources used for the distribution review of *Pinna rudis* in the Mediterranean. The green color represents records retrieved from online databases (GBIF, WoRMS), while the blue color corresponds to spatial information obtained from the literature.

et al., 2016) and were situated within Santorini's islands complex (South Aegean Sea), and one record occurred from a specimen recently caught on fish farming nets in Thesprotia region (Ionian Sea) (Galinou-Mitsoudi *et al.*, 2023). There was one additional record retrieved from the GBIF database, originating from a citizen scientist's observation in Lesvos Island (North Aegean Sea), but it was a clear misidentification of *P. nobilis*, as judged by observing the accompanying photos, and was discarded.

Discussion

The findings of the present study have illuminated a broader distribution range for *P. rudis* in the Mediterranean Sea than previously known. The integration of data sourced from online databases has significantly augmented the records indicating the species' presence. However, the unexpected discovery of a verified population in southern Crete, alongside reported occurrences from other locations in Greece and Cyprus, suggests that the species either has been present but undocumented or that its distribution range has recently expanded into the eastern Mediterranean. While the former appears implausible given that Crete and the regions of the Ionian Sea (islands and mainland) have been subject to thorough investigations and *P. rudis* had not been previously identified there, the latter hypothesis gains traction. The Marine Biodiversity and Ecosystems Management lab (Dept. Marine Sciences) diving team has extensively surveyed several sites in southern Crete and the Ionian Sea during the past five years, and P. rudis had never been detected. This underscores the novelty of the species' presence in the region. A previous study that focused on mapping endangered species in the Aegean Sea had reported the presence of P. nobilis in southern Crete (Sini et al., 2017). However, the subsequent outbreak of MMEs led to the collapse of all known P. nobilis populations on the island (Zotou et al., 2020). The emergence of P. rudis populations in areas affected by P. nobilis' MMEs has been increasingly reported throughout the Mediterranean by scientist and citizen observations uploaded to citizen science platforms and social media. In locations where the two species coexisted, the successful recruitment of P. rudis has been linked to the decline of P. nobilis (Kersting & Ballesteros, 2021). It is also possible that in locations where mixed populations occurred, misidentifications might have taken place (not only in juvenile individuals but also in adults), thereby overlooking the presence of P. rudis (D. Kersting, pers. comm.). After the MME, P. rudis individuals present in those mixed populations have become more "visible" and easier to correctly identify. Notably, the findings of the present study mirror similar observations made by Donato et al. (2021), who documented P. rudis populations in locations where the species had not previously been observed in Italy. This growing body of evidence underscores the potential interaction between the two species and the ecological dynamics at play in the wake of significant disturbances, as initially hypothesized by Kersting & Ballesteros (2021).

The review of the past distribution in the Mediterranean Sea coincides with the results of Gvozdenović *et al.* (2019), considering that the majority of sites with confirmed presence of *P. rudis* are concentrated in the western Mediterranean Sea, while data originated from the eastern, central, and northern Mediterranean Sea are scarce. The low population density, the cryptic habitat (Gvozdenović *et al.*, 2019), and the potential misidentification of *P. rudis* for *P. nobilis* may have impeded monitoring efforts and underestimated the distribution range of the species.

The distribution range of marine species depends on a complex interplay of factors, including temperature,



Fig. 6: Number of records per country collated for the distribution review of *Pinna rudis* in the Mediterranean.



Fig. 7: Number of records per ecoregion collated for the distribution review of *Pinna rudis* in the Mediterranean.



Fig. 8: Distribution map of *Pinna rudis* in the Mediterranean Sea. Different colored circles correspond to different data sources. The higher opacity of symbols indicates the occurrence of multiple records. The Mediterranean Basin is divided into ecoregions based on the classification proposed by Spalding *et al.* (2007).

currents, interspecific interactions, and habitat availability (Dunstan & Bax, 2007). Species with pelagic larval stages generally exhibit higher dispersal potential, facilitated by ocean currents (Donelson et al., 2019). The recent colonization success and range shift of *P. rudis* in the Mediterranean has yet to be fully understood, with several hypotheses proposed to explain it. One hypothesis posits that interspecific competition between the two Pinna species, when coexisting, may have constrained P. rudis. Another hypothesis proposes the consumption of P. rudis larvae by the larger-sized P. nobilis as the main reason for restriction of the population density of P. rudis. With the subsequent collapse of most P. nobilis populations, the survival and establishment of P. rudis larvae may have been enhanced (Kersing & Ballesteros, 2021). Another possible explanation lies in the thermophilic nature of P. rudis and the ongoing warming trends in the Mediterranean (Kersing & Ballesteros, 2021; Donato et al., 2021). Climate change has triggered local extinctions, substantial changes in food webs, and shifts in species distribution in other groups (Bianchi, 2007; Nikolaou & Katsanevakis, 2023), which may have been beneficial for the thermophilic P. rudis to flourish (Kersting & Ballesteros, 2021; Donato et al., 2021).

Although previous research has hinted at the presence of *P. rudis* in the Aegean Sea (Salomidi *et al.*, 2016) in a study about gathering ecological information for the establishment of a potential Marine Protected Area on Santorini Island, no image documentation was provided. Furthermore, none of the experienced scientific divers of our group, with a collective log of more than 4,000 dives in Greek waters and > 25 publications on *Pinna*, has previously observed *P. rudis* in the region. In a more recent article (Galinou-Mitsoudi *et al.*, 2023), a single juvenile *P. rudis* individual was collected from fish farming nets in the Thesprotia region of the Ionian Sea in February 2022. This individual was subsequently photographed on three separate occasions over a nine-month period. This rarity of documented sightings underscores the significance of our study's findings in shedding light on the noteworthy shift in the species distribution within the Greek Seas.

The integration of data from online repositories (GBIF, WoRMS) was instrumental in enriching the spatial information available for the distribution mapping of P. rudis in the Mediterranea. However, the reliability of such data can be compromised by misidentifications and inadequate documentation. A good example of this pertains to a record from GBIF, which indicated the presence of five dead P. rudis individuals in the Kalloni Gulf, Lesvos Island (Greece), as reported by a citizen. The accompanying photographs clearly depicted P. nobilis individuals. Considering that the area held a large, well-documented population of P. nobilis (Katsanevakis et al., 2019; Zotou et al., 2020), the specific unverified citizen science records from GBIF are a clear case of species misidentification, underscoring the need for cautious consideration of unverified citizen science records.

The studied individuals observed during the field survey exhibited the distinct shell morphology (large protuberances on the surface of their shells) and the iridescent mantle border of *P. rudis* and did not show any indications of the putative hybrids that might manifest traits of both *P. nobilis* and *P. rudis*. Likewise, molecular analyses confirmed the absence of cross-species or introgression. In fact, the molecular results undoubtedly identified all the analyzed individuals to be *P. rudis*, as confirmed both at the mitochondrial (multiplex-PCR) and nuclear level (ribosomal gene sequences).

The contribution of local ecological knowledge was pivotal in documenting the P. rudis population in southern Crete. Similarly, citizen scientists across various regions of the country are actively sharing more information about areas of interest and live pinnids through social media platforms. Networks of environmentally aware citizens, guided by members of the scientific community to validate and oversee shared information, not only promote ocean literacy but also strive to maximize data reliability by training the observers. It is worth noting that since resilient P. nobilis populations still survive in a few areas in Greece (Papadakis et al., 2022; 2023) and putative P. rudis-P. nobilis hybrids have been documented in sites where both species coexisted (Vázquez-Luis et al., 2021; Kersting & Ballesteros, 2021), accurate species identification becomes challenging without molecular confirmation.

Following the mass mortality of P. nobilis in the eastern Mediterranean Sea, the ecological niche it once occupied has remained vacant, and its ecological role has not been replaced; it is premature to assert that P. rudis is fulfilling this ecological role, especially considering the difference in densities of Pinna spp. before and after the MME. The knowledge about the ecology and biology of P. rudis remains limited (Nebot-Colomer et al., 2016; Gvozdenović et al., 2019; Lopes et al., 2020; Kersting & Ballesteros, 2021), let alone about putative hybrids. Hence, further research is essential to gain a deeper understanding of the life cycle, reproduction, recruitment, growth, population dynamics, and ecosystem functioning of *P. rudis*. Extensive monitoring in regions affected by MMEs might reveal further unknown cases of the species' establishment.

In conclusion, given the ecological similarities between *P. rudis* and *P. nobilis*, coupled with the threats both species face from human activities, it is crucial for Greek national and international legislation to encompass *P. rudis* within existing protective laws established for *P. nobilis*. Rigorous enforcement of these protection measures is crucial to prevent poaching and exploitation of *P. rudis*, particularly in the wake of the dramatic decline of its congeneric species, which has historically suffered from illegal activities in the Greek Seas (Katsanevakis, 2007; 2016).

Acknowledgements

The present study was conducted in the framework of the European Union's LIFE PINNARCA project: "Protection and restoration of *Pinna nobilis* populations as a response to the catastrophic pandemic started in 2016" (Reference: LIFE20 NAT/ES/001265), which is focusing on the recovery of the fan mussel in the Mediterranean Sea and has received funding from the "LIFE Programme" of the European Union, and the ERDF Project 'Management Actions for the conservation and protection of *Pinna nobilis* in the Protected Areas GR1220002 & GR1270010 of Thermaikos Gulf' funded by the Natural Environment & Climate Change Agency (NECCA). The sea surface temperatures (SSTs) and salinity data were retrieved from the E.U. Copernicus Marine Service (https://www.copernicus.eu/, accessed on 03 August 2023). We are grateful for the valuable suggestions and feedback from the reviewers. We thank Dimitris Poursanidis, Jérôme Smeets from Kalypso Dive Center, Nikos Chatzopoulos and Tasos Papanikolaou from Parga Diving Center, SCUBA divers David Stich Taylor, Paris Sofos, and Isabel Kass (via Red Fish Project) for contributing to this study by sharing information and photos.

Data Availability Statement: The Sea Surface Temperature (SST) and salinity data are openly available by the E.U. Copernicus Marine Service (https://www.copernicus.eu/, accessed on 03 August 2023).

References

- Addis, P., Secci, M., Brundu, G., Manunza, A., Corrias, S. et al., 2009. Density, size structure, shell orientation and epibiotic colonization of the fan mussel *Pinna nobilis* L. 1758 (Mollusca: Bivalvia) in three contrasting habitats in an estuarine area of Sardinia (W Mediterranean). *Scientia Marina*, 73 (1), 143-152.
- Akpinar, A., Yilmaz, E., Fach, B., Salihoglu, B., 2016. Physical oceanography of the Eastern Mediterranean Sea. p. 1-14.
 In: *The Turkish Part of the Mediterranean Sea*. Turan, K., Salihoglu, B., Ozbek, E.O., Ozturk, B. (Eds). Turkish Marine Research Foundation, Turkey.
- Alexandrakis, G., 2014. Estimation of the climatic change impact to beach tourism using joined vulnerability analysis and econometric modelling. In: ADAPTto- CLIMATE Conference, Nicosia, 27-28 March 2014. Cyprus.
- Barea-Azcón, J.M., Ballesteros-Duperón, E., Moreno, D., 2008. Libro rojo de los invertebrados de Andalucía. Consejería de Medio Ambiente, Junta de Andalucía, Sevilla, 4.
- Barreiros, J.P., Pacheco, R.J.S., Gonçalves, S.C., 2016. A New Insight on the Symbiotic Association between the Fan Mussel *Pinna Rudis* and the Shrimp *Pontonia pinnophylax* in the Azores (NE Atlantic). *Global Journal of Zoology*, 1 (1), 001-002.
- Basso, L., Vázquez-Luis, M., García-March, J.R., Deudero, S., Alvarez, E. *et al.*, 2015. The Pen Shell, *Pinna nobilis*: A review of population status and recommended research priorities in the Mediterranean Sea. *Advances in Marine Biology*, 71, 109-160.
- Bianchi, C.N., 2007. Biodiversity issues for the forthcoming tropical Mediterranean Sea. *Hydrobiologia*, 580, 7–21.
- Bosc, E., Bricaud, A., Antoine, D., 2004. Seasonal and interannual variability in algal biomass and primary production in the Mediterranean Sea as derived from 4 years of SeaWiFS observations. *Global Biogeochem Cycles*, 18 (1), GB1005.
- Cabanellas-Reboredo, M., Vázquez-Luis, M., Mourre, B., Álvarez, E., Deudero, S. *et al.*, 2019. Tracking a mass mortality outbreak of pen shell *Pinna nobilis* populations: A collaborative effort of scientists and citizens. *Scientific reports*, 9 (1), 1-11.
- Carella, F., Antuofermo, E., Farina, S., Salati, F., Mandas, D. et al., 2020. In the wake of the ongoing mass mortality events: Co-occurrence of Mycobacterium, Haplosporidium and

other pathogens in *Pinna nobilis* collected in Italy and Spain (Mediterranean Sea). *Frontiers in Marine Science*, 7, 48.

- Carella, F., Palić, D., Šarić, T., Župan, I., Gorgoglione, B. *et al.*, 2023. Multipathogen infections and multifactorial pathogenesis involved in noble pen shell (*Pinna nobilis*) mass mortality events: Background and current pathologic approaches. *Veterinary Pathology*, 60 (5), 560-577.
- Catanese, G., Coupé, S., Bunet, R., 2022b. Mitogenome sequence comparison in the endangered congeneric *Pinna nobilis* and *Pinna rudis* bivalves. *Molecular Biology Reports*, 49, 3627-3635.
- Catanese, G., Grau, A., Valencia, J.M., Garcia-March, J.R., Vázquez-Luis, M. *et al.*, 2018. *Haplosporidium pinnae* sp. nov., a haplosporidian parasite associated with mass mortalities of the fan mussel, *Pinna nobilis* in the Western Mediterranean Sea. *Journal of Invertebrate Pathology*, 157, 9-24.
- Catanese, G., Tena-Medialdea, J., Bel Dajković, M.A., Mičić, M., García-March, J.R., 2022a. An incubation water eDNA method for a non-destructive rapid molecular identification of the two congeneric *Pinna nobilis* and *Pinna rudis* bivalve juveniles. *MethodX*, 9, 101708.
- Çinar, M.E., Bilecenoğlu, M., Yokeş, M.B., Güçlüsoy, H., 2021. *Pinna nobilis* in the south Marmara Islands (Sea of Marmara); it still remains uninfected by the epidemic and acts as egg laying substratum for an alien invader. *Mediterranean Marine Science*, 22 (1), 161-168.
- Cosentino, A., Giacobbe, S., 2006. Shell ornament in *Pinna nobilis* and *Pinna rudis* (Bivalvia: Pteriomorpha). *Journal of Conchology*, 39 (2), 1-6.
- Cosentino, A., Giacobbe, S., 2007. Aspects of epizoobiontic mollusc assemblages on *Pinna* shells. I. Composition and structure. *Cahiers de Biologie Marine*, 48, 187-197.
- Deidun, A., Previati, M., Marrone, A., Gauci, A., Zammit, A. et al., 2022. Ocean literacy and scientific data acquisition through citizen science campaigns: a mixed approach in the Maltese Islands to collect information on *Pinna nobilis* and *Pinna rudis*. *Mediterranean Marine Science*, 23 (2), 357-365.
- Deudero, S., Vázquez-Luis, M., Alvarez, E., 2015. Human stressors are driving coastal benthic long-lived sessile fan mussel *Pinna nobilis* population structure more than environmental stressors. *PLOS ONE*, 10 (7), e0134530.
- Donato, G., Vázquez-Luis, M., Nebot-Colomer, E., Lunetta, A., Giaccobe, S., 2021. Nobel fan-shell, *Pinna nobilis*, in Lake Faro (Sicily, Italy): ineluctable decline or extreme opportunity? *Estuarine, Coastal and Shelf Science*, 261, 107536.
- Donelson, J.M., Sunday, J.M., Figueira W.F., Gaitán-Espitia, J.D., Hobday, A.J. *et al.*, 2019. Understanding interactions between plasticity, adaptation and range shifts in response to marine environmental change. *Philosophical Transactions of the Royal Society B*, 374: 20180186.
- Dunstan, P.K., Bax, N.J., 2007. How far can marine species go? Influence of population biology and larval movement on future range limits. *Marine Ecology Progress Series*, 344, 15-28.
- Fusco, G., Manzella, G.M.R., Cruzado, A., Gacic, M., Gasparini, G.P. *et al.*, 2003. Variability of mesoscale features in the Mediterranean Sea from XBT data analysis. *Annales Geophysicae*, 21, 21-32.

- Galinou-Mitsoudi, S., Vlahavas, G., Papoutsi, O., 2006. Population study of the protected bivalve *Pinna nobilis* (Linnaeus, 1758) in Thermaikos Gulf (north Aegean Sea). *Journal of Biological Research*, 5, 47-53.
- Galinou-Mitsoudi, S., Zaminos, G., Manousis, T., Kontadakis, C., Stasinos, S. *et al.*, 2023. Mediterranean and Hellenic Seas: New records and distributional status of marine molluscs (by December 2022). *Xenophora Taxonomy*, 39, 57-67.
- García-March, J.R., Kersting, D.K., 2006. Preliminary data on the distribution and density of *Pinna nobilis* and *Pinna rudis* in the Columbretes Islands Marine Reserve (Western Mediterranean, Spain). *Organisms, Diversity & Evolution*, 6, 6-16.
- García-March, J.R., Vicente, N., 2006. Protocol to study and monitor *Pinna nobilis* populations within marine protected areas. Spain, Malta Environment and Planning Authority. (available at: https://www.researchgate.net/ publication/235963792_Protocol_to_study_and_monitor_Pinna_nobilis_populations_within_marine_protected areas).
- Giacobbe, S., 2002. Epibiontic mollusc communities on *Pinna* nobilis L. (Bivalvia, Mollusca). Journal of Natural History, 36, 1385-1396.
- Grau, A., Villalba, A., Navas, J.I., Hansjosten, B., Valencia, J.M. *et al.*, 2022. Wide-geographic and long-term analysis of the role of pathogens in the decline of *Pinna nobilis* to critically endangered species. *Frontiers in Marine Science* 9, 666640.
- Gvozdenović, S., Mačić, V., Pešić, V., Nikolić, M., Peraš, I. et al., 2019. Review on Pinna rudis (Linnaeus, 1758) (Bivalvia: Pinnidae) presence in the Mediterranean. Agriculture & Forestry, 65 (4), 115-126.
- Hall, T.A., 1999. BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/ NT. *Nucleic Acids Symposium Series*, 41, 95-98.
- Huber, M., 2010. *Compendium of Bivalves*. Heckenheim, ConchBooks Press.
- Kampouris, T.E., Koutsoubas, D., Kanelopoulou, K., Zannaki, C., Batjakas, I.E., 2022. Informing the general public on the threat status of the European spiny lobster, *Palinurus elephas* (Fabricius, 1787) through Citizen-Science and social media platforms. A case study from the Aegean Sea. *Mediterranean Marine Science*, 23 (2), 366-373.
- Karakasis, I., Eleftheriou, A., 1998. The Continental Shelf of Crete: The Benthic Environment. *Marine Ecology*, 19 (4), 263-277.
- Katsanevakis, S., 2007. Growth and mortality rates of the fan mussel *Pinna nobilis* in Lake Vouliagmeni (Korinthiakos Gulf, Greece): a generalized additive modelling approach. *Marine Biology*, 152 (6), 1319-1331.
- Katsanevakis, S., 2016. Transplantation as a conservation action to protect the Mediterranean fan mussel *Pinna nobilis*. *Marine Ecology Progress Series*, 546, 113-122.
- Katsanevakis, S., Tsirintanis, K., Tsaparis, D., Doukas, D., Sini, M. et al., 2019. The cryptogenic parasite Haplosporidium pinnae invades the Aegean Sea and causes the collapse of Pinna nobilis populations. Aquatic Invasions, 14 (2), 150-164.
- Katsanevakis, S., Carella, F., Çinar, M.E., Čižmek, H., Jimenez,

C. *et al.*, 2021. The fan mussel *Pinna nobilis* on the brink of extinction in the Mediterranean. *Imperiled: The Encyclope- dia of Conservation*.

- Kelly, R., Evans K., Karen, A., Bettiol, S., Corney, S. *et al.*, 2021. Connecting to the oceans: supporting ocean literacy and public engagement. *Reviews in Fish Biology and Fisheries*, 32, 123-143.
- Kersting, D.K., Ballesteros, E., 2021. Is the local extinction of *Pinna nobilis* facilitating *Pinna rudis* recruitment? *Mediterranean Marine Science*, 22 (3), 623-626.
- Kersting, D.K., García-March, J.R., 2017. Long-term assessment of recruitment, early stages and population dynamics of the endangered Mediterranean fan mussel *Pinna nobilis* in the Columbretes Islands (NW Mediterranean). *Marine Environmental Research*, 130, 282-292.
- Kersting, D.K., Benabdi, M., Cizmek, H., Grau, A., Jimenez, C. et al., 2019. Pinna nobilis. The IUCN Red List of Threatened Species 2019: e.T160075998A160081499. https://dx.doi.org/10.2305/IUCN.UK.2019-3.RLTS. T160075998A160081499.en. Accessed on 26 July 2023.
- Kumar, S., Stecher, G., Li, M., Knyaz, C., Tamura, K., 2018. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution*, 35, 1547-1549.
- Lemer, S., Buge, B., Bemis, A., Giribet, G., 2014. First molecular phylogeny of the circumtropical bivalve family Pinnidae (Mollusca, Bivalvia): Evidence for high levels of cryptic species diversity. *Molecular Phylogenetics and Evolution*, 75, 11-23.
- Lopes, E.P., Monteiro, N., Santos, A.M., 2020. Epibiotic assemblages on the pen shell *Pinna rudis* (Bivalvia, Pinnidae) at Matiota Beach, São Vicente Island, Cabo Verde. *African Journal of Marine Science*, 42 (1), 13-21.
- Mokos, M., Cheimonopoulou, M.T., Koulouri, P., Previati, M., Realdon, G. *et al.*, 2020. Mediterranean Sea Literacy: When Ocean Literacy becomes region-specific. *Mediterranean Marine Science*, 21 (3), 592-598.
- Morrocco, V., Sicuro, A., Zangaro, F., Pinna, M., 2018. First record of the protected species *Pinna nobilis* (Linnaeus, 1758) in the Aquatina Lagoon (NATURA 2000 site IT9150003, South-East Italian coastline). *Nature Conservation*, 28, 51.
- Nebot-Colomer, E., Vázquez-Luis, M., García-March, J.R., Deudero, S., 2016. Population structure and growth of the threatened pen shell, *Pinna rudis* (Linnaeus, 1758) in a Western Mediterranean marine protected area. *Mediterranean Marine Science*, 17 (3), 785-793.
- Nikolaou, A., Katsanevakis, S., 2023. Marine extinctions and their drivers. *Regional environmental change*, 23 (88).
- Özalp, H.B., Kersting, D.K., 2020. A pan-Mediterranean extinction? *Pinna nobilis* mass mortality has reached the Turkish straits system. *Marine Biodiversity*, 50 (81).
- Papadakis, O., Papadimitriou, E., Katsanevakis, S., 2022. Pinna nobilis, in search for the surviving populations in Greece. In: 15th ICZEGAR Congress, Mytilene, Greece, 12-15 October 2022 (DOI: 10.13140/RG.2.2.11353.60007).
- Papadakis, O., Mamoutos, I., Ramfos, A., Catanese, G., Papadimitriou, E. *et al.*, 2023. Status, distribution, and threats of the last surviving fan mussel populations in Greece. *Mediterranean Marine Science*, 24 (3), 679-708.

Poppe, G.T., Goto, Y., 1993. European Seashells. Vol. 2. Hack-

enheim, ConchBooks Press, 221 pp.

- Prado, P., Carrasco N., Catanese, G., Grau A., Cabanes P. et al., 2020. Presence of Vibrio mediterranei associated to major mortality in stabled individuals of Pinna nobilis L. Aquaculture, 519, 734899.
- Psarra, S., Tselepides, A., Ignatiades, L., 2000. Primary productivity in the oligotrophic Cretan Sea (NE Mediterranean): seasonal and interannual variability. *Progress in Oceanography*, 46, 187e204.
- Rabaoui, L., Tlig-Zouari, S., Cosentino, A., Ben Hassine, O.K., 2009. Associated fauna of the fan shell *Pinna nobilis* (Mollusca: Bivalvia) in the northern and eastern Tunisian coasts. *Scientia Marina*, 73, 129-141.
- Robinson, A.R., Malanotte-Rizzoli, P., Hecht, A., Michelato, A., Roether, W. *et al.*, 1992. General circulation of the Eastern Mediterranean. *Earth-science reviews*, 32 (4), 285-309.
- Rouanet, E., Trigos, S., Vicente, N., 2015. From youth to death of old age: the 50-year story of a *Pinna nobilis* fan mussel population at Port-Cros Island (Port-Cros National Park, Provence, Mediterranean Sea). *Scientific Reports of the Port-Cros National Park*, 29, 209-222.
- Salomidi, M., Giacoumi, S., Gerakaris, V., Issaris, Y., Sini, M. et al., 2016. Setting an ecological baseline prior to the bottom-up establishment of a marine protected area in Santorini Island, Aegean Sea. Mediterranean Marine Science, 17 (3), 720-737.
- Sanna, D., Cossu, P., Dedola, G.L., Scarpa, F., Maltagliati, F. et al., 2013. Mitochondrial DNA Reveals Genetic Structuring of *Pinna nobilis* across the Mediterranean Sea. *PLOS ONE*, 8 (6): e67372.
- Scarpa, F., Sanna, D., Azzena, I., Mugetti, D., Cerruti, F. *et al.*, 2020. Multiple Non-Species-Specific Pathogens Possibly Triggered the Mass Mortality in *Pinna nobilis*. *Life*, 10 (238), 1-18.
- Sempere, J.B., Jerez, P.S., Luis, M.V., Martínez, C.O., 2006. Evaluación de las especies marinas de interés ecológico del LIC de las Islas Columbretes. Alicante, Spain: Departamento de Ciencias del Mar y Biología Aplicada, Universidad de Alicante.
- Sini, M., Katsanevakis, S., Koukourouvli, N., Gerovasileioy, V., Dailianis, T. *et al.*, 2017. Assembling Ecological Pieces to Reconstruct the Conservation Puzzle of the Aegean Sea. *Frontiers in Marine Science*, 4, 347.
- Spalding, M.D., Fox, H.E., Allen, G.R., Davidson, N., Ferdaña, Z. et al., 2007. Marine Ecoregions of the World: A Bioregionalization of Coastal and Shelf Areas. *BioScience*, 57 (7), 573-583.
- Trigos, S., García-March, J.R., Vicente, N., Tena, J., Torres, J., 2014. Utilization of muddy detritus as organic matter source by the fan mussel *Pinna nobilis*. *Mediterranean Marine Science*, 13 (3), 667-674.
- Vázquez-Luis, M., March, D., Alvarez, E., Alvarez-Berastegui, D., Deudero, S., 2014. Spatial distribution modelling of the endangered bivalve *Pinna nobilis* in a Marine Protected Area. *Mediterranean Marine Science*, 15 (3), 626-634.
- Vázquez-Luis, M., Álvarez, E., Barrajon, A., Garcia-March, J.R., Grau, A. *et al.*, 2017. SOS *Pinna nobilis*: a mass mortality event in western Mediterranean Sea. *Frontiers in Marine Science*, 4, 220.
- Vázquez-Luis, M., Nebot-Colomer, E., Deudero, S., Planes, S.,

Boissin, E., 2021. Natural hybridization between pen shell species: *Pinna rudis* and the critically endangered *Pinna nobilis* may explain parasite resistance in *P. nobilis*. *Molecular Biology Reports*, 48, 997-1004.

- Zavodnik, D., Hrs-Brenko, M., Legac, M., 1991. Synopsis on the fan shell *Pinna nobilis* L. in the eastern Adriatic Sea. *Les espèces marines à protéger en Méditerranée*, 169-178.
- Zodiatis, G., 1993. Circulation of the Cretan Sea water masses (Eastern Mediterranean Sea). *Oceanologica Acta*, 16 (2), 107-114.
- Zotou, M., Gkrantounis, P., Karadimou, E., Tsirintanis, K., Sini, M. et al., 2020. Pinna nobilis in the Greek seas (NE Mediterranean): on the brink of extinction? Mediterranean Marine Science, 21 (3), 575-591.

Copernicus data

Mediterranean Sea - High Resolution L4 Sea Surface Temperature Reprocessed. E.U. Copernicus Marine Service Information (CMEMS). Marine Data Store (MDS). DOI: 10.48670/moi-00173 (Accessed on 03-08-2023). Mediterranean Sea Physics Reanalysis. E.U. Copernicus Marine Service Information (CMEMS). Marine Data Store (MDS). DOI: 10.25423/CMCC/MEDSEA_MULTIYEAR_PHY_006_004_E3R1 (Accessed on 03-08-2023).

Supplementary Data

The following supplementary information is available online for the article: Zotou_Pinna_rudis_Supplementary_File. xlsx.