

Mediterranean Marine Science

Vol 25, No 1 (2024)

VOL 25, No 1 (2024)



An Atlantic origin for the introduced species *Halimeda incrassata* (Bryopsidales, Chlorophyta)

HEROEN VERBRUGGEN, ENRIC BALLESTEROS

doi: [10.12681/mms.35993](https://doi.org/10.12681/mms.35993)

To cite this article:

VERBRUGGEN, H., & BALLESTEROS, E. (2024). An Atlantic origin for the introduced species *Halimeda incrassata* (Bryopsidales, Chlorophyta). *Mediterranean Marine Science*, 25(1), 55–57. <https://doi.org/10.12681/mms.35993>

An Atlantic origin for the introduced species *Halimeda incrassata* (Bryopsidales, Chlorophyta)

Heroen VERBRUGGEN¹ and Enric BALLESTEROS²

¹ School of BioSciences, University of Melbourne, Victoria 3010, Australia

² Centre d'Estudis Avançats de Blanes (CEAB-CSIC), Carrer Accés Cala Sant Francesc 14, 17300 Blanes, Girona, Spain

Corresponding author: Heroen VERBRUGGEN; heroen@unimelb.edu.au

Contributing Editor: Vasilis GEROVASILEIOU

Received: 01 December 2023; Accepted: 12 February 2024; Published online: 15 March 2024

The tropical calcified green alga *Halimeda incrassata* (J.Ellis) J.V.Lamouroux has been found as an introduced species in the Mediterranean Sea with the first observations in Mallorca dating back to 2011 (Alós *et al.*, 2016). The species spread rapidly at this location, but had not been reported from outside this region until recently. A report published in *Mediterranean Marine Science* provides additional records of this species from Rhodes (Kondylatos *et al.*, 2023). Even though the only observations from that location are from the gut content of silver-cheeked toadfish, *Lagocephalus sceleratus* (Gmelin, 1789), the evidence is very convincing, including a combination of morphological and DNA evidence.

In addition to the possibility of an accidental release from a private aquarium, two alternative colonisation pathways for the arrival of *Halimeda incrassata* in the Eastern Mediterranean Sea based on shipping were described by Kondylatos *et al.* (2023). The first pathway is from the western Mediterranean or the Atlantic via the Strait of Gibraltar; the second from the Red Sea or Indian Ocean via the Suez Canal. While we cannot exclude the possibility of an accidental release, in this paper we argue that of the two shipping scenarios, only the first is realistic. Our argument centres on (1) accurate interpretations of global *Halimeda incrassata* records based on current taxonomic knowledge, (2) historical biogeographic interpretations of *Halimeda* phylogenies, (3) the common nature of west-to-east migration of introduced species in the Mediterranean.

The name *H. incrassata* has been applied across all tropical regions, including the Caribbean region, the Indian Ocean and the Pacific. However, molecular data has shown that Indo-Pacific records are not in fact *H. incrassata* but two look-alike species, the Pacific deep-water species *Halimeda kanaloana* Vroom and the widespread Indo-Pacific species *Halimeda heteromorpha* N'Yeurt (Verbruggen *et al.*, 2006). Other Indo-Pacific species that could possibly be mistaken for *H. incrassata* in terms of overall morphology are *H. borneensis* W.R.Taylor, *H. cylindracea* Decaisne and *H. stuposa* W.R.Taylor, although these species can be distinguished with anatomical characters or DNA-based methods. In Figure 1, we

compile published distribution records of these *Halimeda* species, which were either verified with molecular data or based on morphological identification by expert taxonomists with knowledge of the latest molecular taxonomic insights (Verbruggen *et al.*, 2009; Dijoux *et al.*, 2012; Marcelino, 2012; Pongparadon *et al.*, 2015; Alós *et al.*, 2016; Ximenes *et al.*, 2017; Kondylatos *et al.*, 2023). These results clearly show that *H. incrassata* is limited to the Atlantic Ocean plus the recent records in the Mediterranean Sea. All records from the Red Sea and the Indo-West Pacific are of other species.

Historical biogeographical insights have shown a strong divide of the main *Halimeda* lineages (the taxonomic sections of the genus; Verbruggen & Kooistra, 2004) into Atlantic and Indo-Pacific subclades (Kooistra *et al.*, 2002; Verbruggen *et al.*, 2005a; Verbruggen *et al.*, 2009). The sequences of the introduced Mediterranean species clearly correspond to *H. incrassata*, which is closely related to *H. simulans* M.Howie and *H. monile* (J.Ellis & Solander) J.V.Lamouroux, two species that are also restricted to the Atlantic Ocean based on molecular evidence. It is important to note here that these species too, have misidentified records in the Indo-Pacific in public databases, but molecular data shows such records should be placed under *H. borneensis* for *H. simulans* records or under either *H. cylindracea* or *H. stuposa* for *H. monile* records (Verbruggen *et al.*, 2005b). The fact that the species *H. incrassata* is nested in a lineage of Atlantic endemics reinforces the notion that this region represents the origin of the introduced species found in the Mediterranean Sea.

We acknowledge that a large number of the introduced species in Rhodes, and the eastern Mediterranean more broadly, are Lessepsian migrants originating from the Red Sea and Indian Ocean, and the dispersion of these species in the Mediterranean always proceeds east to west and south to north (it cannot be otherwise). However, there are many examples of species that were introduced or first colonized the western part of the Mediterranean and made their way to the east. These include several fish species, like the bastard grunt *Pomadasys incisus* Bowdich, 1825 (Bodilis *et al.*, 2013) and the Ma-




Fig. 1: Distribution of *Halimeda incrassata* and several Indo-Pacific species that it may be confused with morphologically. The species introduced in the Mediterranean Sea is the Atlantic species and the only realistic migration pathway would have been from the Atlantic Sea through the Strait of Gibraltar.

deira scorpionfish *Scorpaena maderensis* Valenciennes, 1833 (Astruch *et al.*, 2021). Two examples amongst the tunicates are those of *Microcosmus squamiger* Michaelsen, 1927 (Turon *et al.*, 2007; Önen, 2020) and *Didemnum vexillum* Kott, 2002 (Çinar & Özgül, 2023). Amongst the seaweeds we can highlight the well-known invasives *Womersleyella setacea* (Hollenberg) R.E.Norris, *Phyllymenia gibbesii* (Harvey) S.M.Lin, Rodríguez-Prieto, De Clerck, Guiry (Rodríguez-Prieto *et al.*, 2021) and *Codium fragile* (Suringar) Hariot, species that were first reported in the western Mediterranean (Verlaque *et al.*, 2015) and progressed to the east to reach the Levantine and Aegean Seas, respectively (Taskin *et al.*, 2013; Tsiamis *et al.*, 2010). Particularly fast was the expansion of the Atlantic sea urchin crab, *Percnon gibbesi*, that in only 12 years spread from the southwestern Mediterranean to the Aegean and Levantine Seas (Katsanevakis *et al.*, 2011).

Although a release from a private aquarium cannot be ruled out as the introduction pathway of *Halimeda incrassata* in Greece, and *Halimeda* can be easily obtained from online shops, we also consider shipping to be a plausible source of the expansion of *H. incrassata* in the Mediterranean Sea, either directly from the western Atlantic or from the much closer populations in the Balearic Islands. The Caribbean Sea, the Balearic Islands and the Aegean are hot spots for recreational shipping and a lot of boats overwintering in the Caribbean come to the Mediterranean in summer (Gui & Russo, 2011; Rodrigue & Notteboom, 2012) and even more move from the Balearic to the Greek Islands. Ballast water or fouling do not appear likely for *H. incrassata*, but the species can probably survive for several days in the humid conditions inside anchor lockers, as has been suggested for *Caulerpa taxifolia* (Sant *et al.*, 1996), and we argue that a similar mechanism could account for the spreading of *H.*

incrassata in the Mediterranean. The species occurs on sandy bottoms where recreational vessels commonly anchor. Re-attachment and growth of thallus fragments is a common mechanism of asexual propagation in *Halimeda* (Walters *et al.* 2002) and is thought to occur extensively in *H. incrassata*, specifically (Littler *et al.*, 2007). In summary, we argue that eastwards recreational shipping is a plausible explanation for the presence of *H. incrassata* in Rhodes.

References

- Alós, J., Tomas, F., Terrados, J., Verbruggen, H., Ballesteros, E., 2016. Fast-spreading green beds of recently introduced *Halimeda incrassata* invade Mallorca island (NW Mediterranean Sea). *Marine Ecology Progress Series*, 558, 153-158.
- Astruch, P., Belloni, B., Rouanet, E., Schohn, T., Harmelin-Vivien, M. *et al.*, 2022. Dominance of *Scorpaena maderensis* among scorpaenids of littoral rocky reefs in Corsica (NW Mediterranean): Further evidence of Mediterranean Sea warming. *Journal of Fish Biology*, 100, 601-604.
- Bodilis, P., Crocetta, F., Langeneck, J., Francour, P., 2013. The spread of an Atlantic fish species, *Pomadoury inciscus* (Bodwich, 1825) (Osteichthyes: Haemulidae), within the Mediterranean Sea with new additional records from the French Mediterranean coast. *Italian Journal of Zoology*, 80, 273-278.
- Çinar, M.E., Özgül, A., 2023. Clogging nets-*Didemnum vexillum* (Tunicata: Ascidiacea) is in action in the eastern Mediterranean. *Journal of the Marine Biological Association of the United Kingdom*, 103, e89.
- Dijoux, L., Verbruggen, H., Mattio, L., Duong, N., Payri, C., 2012. Diversity of *Halimeda* (Bryopsidales, Chlorophyta) in New Caledonia: a combined morphological and molecu-

- lar study. *Journal of Phycology*, 48, 1465-1481.
- Gui, L., Russo, A.P., 2011. Cruise ports: a strategic nexus between regions and global lines - evidence from the Mediterranean. *Maritime Policy and Management*, 38, 129-150.
- Katsanevakis, S., Poursanidis, D., Yokes, M.B., Mačić, V., Beqiraj, S. *et al.*, 2011. Twelve years after the first record of the crab *Percnon gibbesi* (H. Milne Edwards, 1853) in the Mediterranean: current distribution and invasion rates. *Journal of Biological Research-Thessaloniki*, 16, 224-236.
- Kondylatos, G., Kalaentzis, K., Gratsia, E., Mavrouleas, D., Kasapidis, P. *et al.*, 2023. *Halimeda incrassata* (Bryopsidales, Chlorophyta) in Rhodes, Greece, Eastern Mediterranean. *Mediterranean Marine Science*, 24, 633-638.
- Kooistra, W.H.C.F., Coppejans, E.G.G., Payri, C., 2002. Molecular systematics, historical ecology, and phylogeography of *Halimeda* (Bryopsidales). *Molecular Phylogenetics and Evolution*, 24, 121-138.
- Littler, M.M., Littler, D.S., Brooks, B.L., Lapointe, B.E., 2007. Unusual linear arrays of the coral reef macrophyte *Halimeda incrassata* in the Bahamas. *Coral Reefs*, 26, 817-818.
- Marcelino, V.R., 2012. Dynamics and conservatism of ecological niches in *Halimeda*. MSc thesis. Ghent University, Belgium, 74pp.
- Önen, S.A., 2020. First occurrence of fouling ascidian species *Microcosmus squamiger* Michaelsen, 1927 and *Didemnum ahu* Monniot C. & Monniot F., 1987 in Izmir Bay (Eastern Aegean Sea). *Journal of Natural History*, 54, 1897-1912.
- Pongparadon, S., Zuccarello, G.C., Phang, S.-M., Kawai, H., Hanyuda, T. *et al.*, 2015. Diversity of *Halimeda* (Chlorophyta) from the Thai-Malay Peninsula. *Phycologia*, 54, 349-366.
- Rodrigue, J.P., Notteboom, T., 2012. The geography of cruise shipping: Caribbean and Mediterranean itineraries, capacity deployment and ports of call. *ALRT 2012 Conference*, Vancouver. 20 pp.
- Rodríguez-Prieto, C., Shabaka, S.H., El-Din, N.S., De Clerck, O., 2021. Morphological and molecular assessment of *Grateloupia* (Halymeniales, Rhodophyta) from Egypt revealed a new introduced species in the Mediterranean Sea, *Grateloupia gibbesii*. *European Journal of Phycology*, 60, 83-95.
- Sant, N., Delgado, O., Rodríguez-Prieto, C., Ballesteros, E., 1996. The spreading of the introduced seaweed *Caulerpa taxifolia* (Vahl) C. Agardh in the Mediterranean Sea: testing the boat transportation hypothesis. *Botanica Marina*, 39, 427-430.
- Taskin, E., Öztürk, M., Kurt, O., Ulcay, S., 2013. Benthic marine algae in Northern Cyprus (Eastern Mediterranean Sea). *Journal of Black Sea/Mediterranean Environment*, 19, 143-161.
- Tsiamis, K., Montesanto, B., Panayotidis, P., Katsaros, C., Verlaque, M., 2010. Updated records and range expansion of alien marine macrophytes in Greece (2009). *Mediterranean Marine Science*, 11 (1), 61-79.
- Turon, X., Nishikawa, T., Rius, M., 2007. Spread of *Microcosmus squamiger* (Ascidiacea: Pyuridae) in the Mediterranean Sea and adjacent waters. *Journal of Experimental Marine Biology and Ecology*, 342, 185-188.
- Verbruggen, H., Kooistra, W.H.C.F., 2004. Morphological characterization of lineages within the calcified tropical seaweed genus *Halimeda* (Bryopsidales, Chlorophyta). *European Journal of Phycology*, 39, 213-228.
- Verbruggen, H., De Clerck, O., Schils, T., Kooistra, W.H.C.F., Coppejans, E., 2005a. Evolution and phylogeography of *Halimeda* section *Halimeda*. *Molecular Phylogenetics and Evolution*, 37, 789-803.
- Verbruggen, H., De Clerck, O., Kooistra, W.H.C.F., Coppejans, E., 2005b. Molecular and morphometric data pinpoint species boundaries in *Halimeda* section *Rhipsalis* (Bryopsidales, Chlorophyta). *Journal of Phycology*, 41, 606-621.
- Verbruggen, H., De Clerck, O., N'Yeurt, A.D.R., Spalding, H., Vroom, P. S., 2006. Phylogeny and taxonomy of *Halimeda incrassata*, including descriptions of *H. kanaloana* and *H. heteromorpha* spp. nov. (Bryopsidales, Chlorophyta). *European Journal of Phycology*, 41, 337-362.
- Verbruggen, H., Tyberghein, L., Pauly, K., Vlaeminck, C., Van Nieuwenhuyze, K. *et al.*, 2009. Macroecology meets macroevolution: evolutionary niche dynamics in the seaweed *Halimeda*. *Global Ecology and Biogeography*, 18, 393-405.
- Verlaque, M., Ruitton, S., Mineur, F., Boudouresque, C.F., 2015. *CIESM Atlas of exotic species in the Mediterranean*. CIESM Publishers, Monaco. 362 pp.
- Walters, L.J., Smith, C.M., Coyer, J.A., Hunter, C.L., Beach, K.S. *et al.*, 2002. Asexual propagation in the coral reef macroalga *Halimeda* (Chlorophyta, Bryopsidales): production, dispersal and attachment of small fragments. *Journal of Experimental Marine Biology and Ecology*, 278, 47-65.
- Ximenes, C.F., Cassano, V., Oliveira-Carvalho, M.F., Bandeira-Pedrosa, M.E., Gurgel, C.F.D. *et al.*, 2017. Systematics of the genus *Halimeda* (Bryopsidales, Chlorophyta) in Brazil including the description of *Halimeda jolyana* sp. nov. *Phycologia*, 56, 369-381.