

Mediterranean Marine Science

Vol 25, No 2 (2024)

Mediterranean Marine Science



First record of the West Atlantic *Ariopsis felis* (Linnaeus, 1766) (Actinopterygii: Siluriformes: Ariidae) in the Mediterranean Sea

JEHAD Y. SALAH, MOHAMMED A. ABUTAIR, SARA A.A. AL MABRUK; BRUNO ZAVA; MARIA CORSINI-FOKA, ALAN DEIDUN, ANTONIO DI NATALE; VALENTINA CROBE; FAUSTO TINTI

doi: [10.12681/mms.36177](https://doi.org/10.12681/mms.36177)

To cite this article:

SALAH, J. Y., ABUTAIR, M. A., AL MABRUK, S. A., ZAVA, B., CORSINI-FOKA, M., DEIDUN, A., DI NATALE, A., CROBE, V., & TINTI, F. (2024). First record of the West Atlantic *Ariopsis felis* (Linnaeus, 1766) (Actinopterygii: Siluriformes: Ariidae) in the Mediterranean Sea. *Mediterranean Marine Science*, 25(2), 263–271. <https://doi.org/10.12681/mms.36177>

First record of the West Atlantic *Ariopsis felis* (Linnaeus, 1766) (Actinopterygii: Siluriformes: Ariidae) in the Mediterranean Sea

Jehad Y. SALAH¹, Mohammed A. ABUTAIR², Sara A.A. Al MABRUK^{3,4}, Bruno ZAVA^{5,6}, Maria CORSINI-FOKA⁷, Alan DEIDUN⁸, Antonio Di NATALE⁹, Valentina CROBE¹⁰ and Fausto TINTI¹⁰

¹Department of Fisheries, General Directorate of Fisheries Gaza (DoF), Al Rasheed St. 191/1000, Gaza, Palestine

²Department of Marine Environment, General Directorate of Fisheries Gaza (DoF), Al manar St. 3/49, Abasan alkabira, Khanyunis, Palestine

³Department of General Nursing Technology, Higher institute of Science and Technology, Cyrene, Libya

⁴Marine Biology in Libya Society, El Bayda, Libya

⁵Wilderness studi ambientali, via Cruillas 27, 90146 Palermo, Italy

⁶Museo Civico di Storia Naturale di Comiso, Via degli Studi 9, 97013 Comiso (RG), Italy

⁷Institute of Oceanography, Hellenic Centre for Marine Research. Hydrobiological Station of Rhodes, Cos Street, 85100 Rhodes, Greece

⁸Department of Geosciences, University of Malta, Msida MSD 2080, Malta

⁹Aquastudio Research Institute, Via Trapani 6, 98121 Messina, Italy

¹⁰Department of Biological, Geological and Environmental Sciences, University of Bologna, Via S. Alberto 163, 48123 Ravenna, Italy

Corresponding author: Maria Corsini-Foka; mcorsini@hcmr.gr

Contributing Editor: Paraskevi K. KARACHLE

Received: 12 December 2023; Accepted: 02 May 2024; Published online: 12 June 2024

Abstract

A single individual of Ariidae was captured in 2022 in the Palestinian waters off the Gaza Strip, eastern Mediterranean. A comprehensive description of the morphological and meristic characters of the specimen was conducted. Molecular identification was performed through sequence analysis of the Cytochrome Oxidase subunit I (COI). The specimen was identified as *Ariopsis felis*, both morphologically and genetically. The natural range of this species is the western Atlantic and, to the best of our knowledge, the species has never been reported before from Mediterranean waters. The singular occurrence of this fish in the basin prompts further discussion, raising questions about the possible vectors that may be responsible for this new introduction in the Mediterranean Sea.

Keywords: Rare fish; introduced species; cryptogenic species; Mediterranean Sea; Gaza Sea; DNA barcoding.

Introduction

The family Ariidae (Siluriformes) consists of 156 species belonging to 30 genera, widely distributed worldwide, particularly in tropical and subtropical areas, prevalently in marine waters and occasionally in brackish waters of estuaries and coastal lagoons (Daget *et al.*, 1986; Simier *et al.*, 2021; Froese & Pauly 2023). Within the Mediterranean Sea ichthyofauna, only one species of this family is documented, the Guinean sea catfish *Carlarius parkii* (Günther, 1864) (Kovačić *et al.*, 2021). But this record is based on a single specimen, originally reported as *Arius parkii* Günther, 1864, collected in the Mediterranean from the shallow waters of Haifa Bay, Israel, in 1983 (Golani & Ben-Tuvia, 1986; Golani & Sonin, 1996; Golani, 1996, 2021). The natural distribution range of this

species consists of the eastern Atlantic waters extending from Cape Blanc to Angola, sporadically to the Atlantic coasts of Morocco (Acero & Betancur-R., 2016).

In the present study, the occurrence of the hardhead sea catfish *Ariopsis felis* (Linnaeus, 1766) is documented for the first time in the Mediterranean Sea, after the collection of a single specimen in the coastal waters off the Gaza Strip, Palestine. The natural range of this species is based in the western Atlantic. The study provides a detailed account of morphological characters and presents results from genetic analysis, supporting the assigned species identification. The unexpected occurrence of this exotic fish in the eastern part of the Mediterranean Sea is briefly discussed, raising questions into the possible vectors of introduction that may have facilitated this unprecedented range extension.

Material and Methods

On 17 October 2022, a fish unfamiliar to fishermen was found among the catch of a bottom trawler operating in the fishing area off the Gaza Strip, approximately 2 miles from the shore (Coordinates: 31.43167°N, 34.28642°E), at a depth of 25 m, on a muddy bottom (Fig. 1). Bottom trawlers are commonly employed in the region for the catch of demersal and benthic fish (Abudaya *et al.*, 2013). The fishing areas along the Gaza Strip coast have been described by Abudaya *et al.* (2013) and Hussein *et al.* (2022). The mouth of the Gaza Valley river (Ubeid, 2011; Abualhin, 2016) is located 3-5 km from the capture site. Since the fisherman did not recognize the species, he contacted the Department of Fishery (DoF) of the General Directorate of Fisheries of Gaza and brought the sample to its research team. The specimen is now deposited in the collection of the laboratory of DoF, preserved in ethanol under the registration code NS-felis-01, 17-10-2022.

The main measurements were taken with a caliper to the nearest 0.1 cm, meristic characters were counted under a stereoscope and X-rays analysis was performed.

Identification keys of Ariidae from seas all over the world were consulted, among which Fischer & Bianchi (1984), Kailola & Bussing (1995), Kailola (1999), Acero (2002), Daget (2003), Marceniuk (2005), Marceniuk & Menezes (2007), Acero & Betancur-R. (2016) and Marceniuk *et al.* (2017a, 2024).

Genetic analysis

Total genomic DNA (gDNA) was extracted from approximately 20 mg of tissue preserved in 99% ethanol using the Wizard® SV Genomic DNA Purification System by Promega, according to the manufacturer's instructions. The quality of the extracted gDNA was subsequently assessed on a 1% agarose gel electrophoresis.

The Cytochrome Oxidase subunit I (COI) gene was amplified with the M13-tailed primer cocktail (COI-3; C_FishF1t1-C_FishR1t1) described by Ivanova *et al.* (2007). The PCR reaction was performed in 25 µL total volume containing 3 µL gDNA template, 5 µL 1x PCR buffer, 2.5 µL of MgCl₂ (25 mM), 0.5 µL dNTPs (25 mM), 0.25 µL of each primer cocktail (10 µM) and 0.25 U GoTaq G2 Flexi DNA polymerase (Promega). Amplification was performed in a T-gradient thermocycler (Biometra) with an initial denaturation of 2 min at 94°C, followed by 35 cycles of 30 s at 94°C, 30 s at 52°C and 60 s at 72°C and a final extension step for 10 min at 72°C. The PCR product was evaluated on 2% agarose gel. Purification and sequencing were performed at MacroGen Europe (Milan, Italy), using the M13F and M13R primers (Messing *et al.*, 1983; Ivanova *et al.*, 2007).

Trace files were edited with the software MEGA 11 (Tamura *et al.*, 2021) and the sequence was compared against published sequences through GenBank BLAST (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>, accessed on 20 October 2023) and the Barcode of Life Data System (BOLD, <http://www.boldsystems.org>, accessed on 20 October 2023) Identification Engine. Available COI se-



Fig. 1: Map of the eastern Mediterranean showing the sampling locations of Ariidae records: *Ariopsis felis*, Gaza Sea, Palestine, present study (red star); *Carlarius parkii*, Haifa Bay, Israel (Golani & Ben-Tuvia, 1986; Golani & Sonin, 1996) (yellow triangle).

quences were retrieved for 12 species of Ariidae from both BOLDsystems v.4 (Ratnasingham & Hebert 2007; <http://www.boldsystems.org>) and NCBI databases (<https://www.ncbi.nlm.nih.gov>). A Neighbour-joining (NJ) tree (Saitou & Nei, 1987) was computed with MEGA using p-distance (Collins & Cruickshank, 2013) with pairwise deletion. Bootstrap analysis with 1000 replicates (Felsenstein, 1985) was performed to assess statistical support to nodes of the NJ tree topologies.

Results

Morphological characterization

The fish, 224 mm in total length, 230 g in weight, showed the following characteristics (Fig. 2A, B): body elongated and cylindrical; head rounded and slightly flat; front and rear nostrils close together on each side of snout, the posterior pair partly covered by a flap of skin; humeral process pointed, triangular (Fig. 2B); rugose bony shield on head, with supraoccipital process extending posteromedially to meet the semilunar separate bone at base of dorsal-fin spine (named nuchal plate, pre-dorsal plate) (Fig. 3A); a keel on supraoccipital process; a groove in median depression of head, extending forward to opposite rear halves of eyes and preceded by a fontanel; mouth large and inferior; three pairs of barbels (1

maxillary and 2 mental) around mouth, maxillary barbels the longest, reaching pectoral-fin bases (Fig. 2A, 3A); on the roof of the mouth, premaxillary teeth conical forming a plate slightly curved; conical palatal teeth arranged in 2 patches on each side (2 small irregular median patches associated with vomere, the right smaller than the left, and 2 larger oval shaped lateral patches), the median adjacent to the lateral in each side, the two lateral well separated one from the other (Fig. 3B). A sharp spine, frontally and posteriorly serrated, in the dorsal fin, preceded by a very short one (Fig. 3A); a sharp serrated spine in pectoral fin; adipose fin moderately developed opposite to anal fin; caudal fin forked with pointed lobes, the dorsal lobe longer than the ventral lobe. Dorsal fin rays: I+I+7, Pectoral fin rays: I+9; Anal fin rays: 15; Ventral fin rays: 6. Anterior gill rakers on first arch 13 (including 2 damaged), on second arch 11. A number of at least 46 vertebrae were distinguishable after the X-rays were applied (Fig. 2C). Colour: grey-brown dorsally, silvery whitish ventrally, adipose fin dark, other fins reddish-brown with dark tips. Main morphometric measurements and their proportions as % of standard length are presented in Table 1.

Molecular identification

The COI marker was successfully amplified and the resulting 652 bp sequence has been submitted to GenBank




Fig. 2: The specimen of *Ariopsis felis* from Gaza Strip waters, Palestine (A: freshly caught; B: preserved in liquid, scale bar= 20 mm; C: X-rays).




Fig. 3: *Ariopsis felis* from Gaza Sea, Palestine: dorsal view of bony shield on the head (A) and premaxillary and palatal teeth (B).

Table 1. Main morphometric measurements (mm), expressed also in percent of standard length (SL), of the *Ariopsis felis* specimen collected in the Gaza Sea, Palestine, in October 2022.

Measurements	mm	% SL
Total length	224	
Fork length	193	
Standard length	181	
Head length	40.7	22.5
Snout length	13	7.2
Eye diameter	9	5.0
Interorbital distance	21.4	11.8
Body depth	40	22.1
Snout to dorsal fin	57.3	31.7
Snou to ventral fin	93	51.4
Snout to adipose fin	134	74.0
Snout to anal fin	130	71.8
Distance between dorsal fin end and adipose fin origin	55.7	30.8
Caudal peduncle depth	13.3	7.3
Dorsal fin spine length	30.0	16.6
Pectoral fin spine length	28.5	15.7

under Accession Number OR923365. Molecular identification through BLAST and BOLD yielded high percent identity matches for *A. felis* at 99.81% and 99.85%, respectively, confirming the morphological identification. The NJ tree (Fig. 4) was obtained by comparing our sequence with 12 available sequences from *A. felis* and 9

from the sister species *Ariopsis assimilis* (Günther, 1864) retrieved from BOLD. Additionally, one sequence from each of 10 other species within the Ariidae family was included in the analysis. The generated sequence clustered within the *A. felis* clade with a fully-robust bootstrap support (BP: 100%).




Fig. 4: Neighbor-Joining tree based on COI genetic p-distances of 12 Ariidae taxa. The sequences of *Ariopsis felis* and *Ariopsis assimilis* used in the tree reconstruction were originally recorded on BOLD and GeneBank as *Sciades felis* and *Sciades assimilis*, respectively; nevertheless, based on information from the World Register of Marine Species (WoRMS Editorial Board, 2023) and FishBase (Froese & Pauly, 2023), the correct genus designation for these sequences is recognized as *Ariopsis*. Numbers near nodes indicate bootstrap values ($\geq 70\%$). A distance scale bar is given.

Discussion

The morphological and morphometric characters observed in the specimen from the waters of the Gaza Strip led us to identify it as *A. felis*, with a primary reliance on the criteria outlined by Acero (2002) and Marceniuk *et al.* (2017a). Subsequent genetic analysis confirmed the accuracy of this assignment. The molecular identification provided crucial support to the morphological assignment, particularly due to the highly unusual and out-of-

native-range collection of the specimen. The family Ariidae benefits from numerous publicly available sequences (e.g., 655 public records on BOLD in 54 species), which contributed to a more precise specimen identification within the family. All sequences comprising the well-supported clade of *A. felis*, which includes the sequence of the Gaza sample, are sourced from published papers (Valdez-Moreno *et al.*, 2010; Deeds *et al.*, 2014) supporting the robustness of the clade (Antil *et al.*, 2023).

The pattern of palatal teeth in particular is one of the

distinctive characters used for the identification of species in the family Ariidae. The palatal teeth in our sample were arranged in two pairs of irregularly rounded patches, well-separated by a space smaller than their diameter and the vomere-associated teeth were absent. These features allowed us to separate our sample from *C. parkii* described from the nearby Israel waters, in which the palatal teeth are in single pair of rounded patches (cf. Günther, 1864; Golani & Sonin, 1996; Daget, 2003; Marceniuk & Menezes 2007; Acero & Betancur-R., 2016). The disposition of the palatal teeth of the Gaza specimen showed some similarity to that described for the Indo-west Pacific species *Hexanematachthys sagor* (Hamilton, 1822) (Fischer & Bianchi, 1984; Kailola, 1999; Froese & Pauly, 2023) and to those of some Ariidae of the west-central Pacific ocean, such as *Arius armiger* De Vis, 1884, *A. leptaspis* (Bleeker, 1862), *A. (Hemiarius) stormii* (Bleeker, 1858), *A. utarus* Kailola, 1990 and other species (Kailola, 1999), as well as from Ariidae of the eastern-central Pacific (Kailola & Bussing, 1995). Palatal teeth of the Ariidae *Netuma thalassina* (Rüppell, 1837) and *Plicofollis dussumieri* (Valenciennes, 1840) reported from the Red Sea (Golani & Fricke 2018) differ clearly from the teeth of the Gaza specimen (Fischer & Bianchi, 1984; Marceniuk *et al.*, 2017b). Conversely, the palatal teeth of the sample studied in the present work were especially similar to those of the west Atlantic *A. felis* (Acero, 2002; Marceniuk *et al.*, 2017a).

Meristic counts taken agreed with those cited by Marceniuk *et al.* (2017a), apart from the anal fin rays count (15 vs 17-19) and the number of gill rakers in the second arch (11 vs 13-17).

Proportions of main morphometric measurements of our specimen fell within the ranges given for *A. felis* by Marceniuk *et al.* (2017a), except that obtained for body depth (22.1% of standard length) that exceeded the maximum value of the range of 16.4-20.3% of standard length reported by the same authors.

The subtropical marine species *A. felis* is common, and sometimes locally abundant, on muddy bottoms of shallow coastal waters, brackish waters of estuaries and coastal lagoons, only occasionally entering fresh waters; it is distributed from Cape Cod, Massachusetts, to Florida, USA, and Gulf of Mexico, including Louisiana, USA, to Yucatan (Acero, 2002; Pensinger *et al.*, 2021).

This benthic fish is caught mainly with bottom trawls, seines, and through hook-and-line; it reaches a maximum total length of 70 cm, common 25 cm, and it feeds prevalently on decapods, gastropods and fish (Acero, 2002; Osowski *et al.*, 2023). The hardhead sea catfish constitutes part of the diet of bullsharks and alligators in the western Gulf of Mexico (Marsaly *et al.*, 2023); occasionally, it is source of food for bottlenose dolphins in the northern Gulf of Mexico (Ronje *et al.*, 2017). Generally, *A. felis* is known to show a longevity extending over 20 years, exhibiting large oocytes and low fecundity and males which incubate eggs and juveniles in their mouths (Acero, 2002; Pensinger *et al.*, 2021). The sharp dorsal and pectoral fin spines can inflict painful wounds (Gordon *et al.*, 2021). The hardhead sea catfishes are not com-

mercially or recreationally targeted fishes in their natural range (Pensinger *et al.*, 2021), and are difficult to keep in ordinary aquaria (<https://saltwater.aqua-fish.net/?hard-head-sea-catfish>).

The record of *A. felis* described in the present study from the eastern Levant waters of Palestine is the first for the whole Mediterranean Sea and constitutes the second member of Ariidae found in the basin, after *C. parkii*, the capture of which was only at about 170 km of distance from the Gaza sample location (Fig. 1) (Golani & Ben-Tuvia 1986; Golani & Sonin, 1996).

The Guinean sea catfish *C. parkii* is widely distributed in the east Atlantic waters of Africa, entering estuaries and freshwater of rivers (Froese & Pauly, 2023). It was initially considered a species expanding its range into the Mediterranean through the Strait of Gibraltar, with no relation to human activity (Golani *et al.*, 2002, 2015; Psomadakis *et al.*, 2012). Nevertheless, the absence, for decades, of intermediate records for the species from the basin, have led some authors, including Evans *et al.* (2020) and Zenetos *et al.* (2022), to suggest that the Guinean sea catfish is a cryptogenic species of the basin, that is, a species not demonstrably native or introduced, sensu Carlton (1996).

Explaining the singular occurrence of *A. felis* in the Gaza Sea presents a formidable challenge, given that its native distribution spans from the northwestern Atlantic to the Gulf of Mexico. As a result, assigning a natural range expansion of *A. felis* from the Atlantic into the waters off Gaza is not a plausible hypothesis. In addition, scattered observations in Mauritania, Brazil and Australia have been reported in GBIF (iNaturalist contributors, 2023a, b, c), adding complexity to the understanding of its current geographic distribution.

Other western Atlantic fish species have been detected in the Mediterranean. For instance, the holocentriform *Holocentrus adscensionis* (Osbeck, 1765), identified in Malta waters (Vella *et al.*, 2016), is considered cryptogenic according to Evans *et al.* (2020). The acanthuri-form *Acanthurus coeruleus*, Bloch and Schneider, 1801, a species frequently recorded in the basin, is speculated to have entered the Mediterranean either naturally through the Strait of Gibraltar or by shipping or as a result of aquarium escapes (Golani *et al.*, 2015, 2021). Similarly, the pomacentrid *Stegastes variabilis* (Castelnau, 1855), reported twice, is thought to have reached the Mediterranean via ballast waters or as an aquarium escapee (Vella *et al.*, 2015; Golani *et al.*, 2023).

Human activity, particularly the possibility of introduction via ballast waters, seems more plausible. The site of capture off Gaza lies between and in close proximity to major shipping ports, including Port Said, Tel Aviv, Haifa and Beirut. This complex interplay between human-related factors and environmental dynamics emphasizes the need for further investigation to unravel the mechanisms behind the unexpected presence of *A. felis* in the eastern Mediterranean.

In the wider region, there are rivers in Israel and in the Gaza Strip, emptying into the Mediterranean Sea; also, the brackish waters of Bardawil Lagoon along the north

coast of Sinai and the Nile Delta and its channels in Egypt, are located not far from the Mediterranean sites where the two single records of Ariidae, *C. parkii* (Israel) and *A. felis* (Gaza Sea, Palestine), were made (Fig. 1). While these sites might offer suitable environmental conditions for supporting small populations of marine Ariidae, there is currently no information regarding the potential establishment of both species in the broader region.

Therefore, in our opinion, it is premature to suggest an eventual pathway for the introduction of *A. felis* into the southeastern Levantine waters since the current understanding we have of the species is based solely on our recent singular discovery. This underscores the need for further monitoring of exotic/alien fish species' distributions within the eastern Mediterranean, so as to provide conclusive insights into the potential pathways and mechanisms of introduction for the same species in the region.

The present record significantly contributes valuable information to the interesting but poorly-known ichthyofaunal diversity of the Gaza coast. The Palestinian marine region is characterized by more than 128 bony fish species reported in fishery activities, including Lessepsian migrants and newcomers originating from the Atlantic (Abuamrah, 2018; Abd Rabou, 2019; Bariche *et al.*, 2019; Abd Rabou *et al.*, 2020; Hussein *et al.*, 2022; Azzurro, 2023; Department of Fisheries Gaza, Palestine, unpublished data). The ongoing exploration of this marine environment enhances our comprehension of the intricate dynamics underlying the ichthyofaunal composition of this area.

Acknowledgements

The authors are grateful to Captain Iyad Al Hissi of the fishing vessel Haj rajab (License Number: P-049-4) for promptly providing the specimen described in this study as well as information on its capture. They also warmly thank Dr Danilo Scannella, Institute for Marine Biological Resources and Biotechnology-IRBIM, CNR, Mazara del Vallo (TP). The authors are also indebted to anonymous reviewers for providing useful and constructive comments and suggestions on the first version of the manuscript.

References

- Abd Rabou, A.F.N., 2019. On the occurrence and health risks of the Silver-cheeked Toadfish (*Lagocephalus sceleratus* Gmelin, 1789) in the marine ecosystem of the Gaza Strip, Palestine. *Biodiversitas*, 20 (9), 2620-2627.
- Abd Rabou, A.F.N., Al-Agha, M.R., Abd Rabou, M.A., Abu-altayef, M.T., Elnabris, K.J. *et al.*, 2020. On the Ocean Sunfishes (*Mola mola*, Linnaeus 1758) by-caught off the Mediterranean coast of the Gaza Strip, Palestine. *Biomedical Journal of Scientific and Technical Research*, 29 (4), 22624-22631.
- Abualhin, K., 2016. Mapping of underwater seabed morphology of the Gaza Strip coastal zone using remote sensing technique. *Earth Sciences Research Journal*, 20 (2), C1-C7.
- Abuamrah, H., 2018. A survey of marine Bony Fishes of the Gaza Strip, Palestine. Lap Lambert Academic Publishing, 132 pp.
- Abudaya, M., Harper, S., Ulman, A., Zeller, D., 2013. Correcting mis- and under-reported marine fisheries catches for the Gaza Strip: 1950-2010. *Acta Adriatica*, 54 (2), 241-252.
- Aceró, A., 2002. Order Siluriformes: Ariidae. Sea catfishes. p. 831-852. In: *FAO species identification guide for fishery purposes. The living marine resources of the western central Atlantic. Volume 2 Bony fishes part 1 (Acipenseridae to Grammatidae)*. Carpenter, K.E. (Ed). FAO, Rome.
- Aceró, A., Betancur-R., R., 2016. Siluriformes: Ariidae. p. 1742-1750. In: *FAO species identification guide for fishery purposes. The living marine resources of the eastern Central Atlantic, Vol. 3 bony fishes part 1 (Elopiformes to Scorpaeniformes)*. Carpenter, K.E., De Angelis, N. (Eds). FAO, Rome.
- Antil, S., Abraham, J.S., Sripoorna, S., Maurya, S., Dagar, J. *et al.*, 2023. DNA barcoding, an effective tool for species identification: a review. *Molecular Biology Reports*, 50, 761-775.
- Azzurro, E. (Ed.), 2023. *ORMEF*. World Wide Web electronic publication. www.ormef.eu (02/2023)
- Bariche, M., Sayar, N., Balistreri, P., 2019. Records of two non-indigenous fish species *Synanceia verrucosa* Bloch and Schneider, 1801 and *Acanthurus sohal* (Forsskål, 1775) from the Gaza strip (eastern Mediterranean Sea). *BioInvasions Records*, 8 (3), 699-705.
- Carlton, J.T., 1996. Biological invasions and cryptogenic species. *Ecology*, 77 (6), 1653-1655.
- Collins, R.A., Cruickshank, R.H., 2013. The seven deadly sins of DNA barcoding. *Molecular Ecology Resources*, 13, 969-975.
- Daget, J., 2003. Ariidae. p. 269-276. In: *Faune des poissons d'eaux douces et saumâtres de l'Afrique de l'Ouest, Tome 2*. Paugy, D., Lévêque, C., Teugels, G.G. (Eds). Musée Royal de l'Afrique Centrale, Tervuren, Belgique, Museum National d'Histoire Naturelle, Paris, France and Institut de Recherche pour le Développement, Paris, France. Collection Faune et Flore tropicales 40.
- Daget, J., Gosse, J-P., Thys van den Audenaerde, D.F.E., 1986. *Check-list of the freshwater fishes of Africa. Cloffa*. MRAC Tervuren, Belgium and ORSTOM, Paris, France, 535 pp.
- Deeds, J.R., Handy, S.M., Fry, F. Jr, Granade, H., Williams, J.T. *et al.*, 2014. Protocol for Building a Reference Standard Sequence Library for DNA-Based Seafood Identification. *Journal of AOAC International*, 97 (6), 1626-1633.
- Evans, J., Arndt, E., Schembri, P.J., 2020. Atlantic fishes in the Mediterranean: using biological traits to assess the origin of newcomer fishes. *Marine Ecology Progress Series*, 643, 133-143.
- Felsenstein, J., 1985. Confidence limits on phylogenies: An Approach Using the Bootstrap. *Evolution*, 39, 783-791.
- Fischer, W., Bianchi, G. (Eds), 1984. Ariidae. *FAO species identification sheets for fishery purposes. Western Indian Ocean fishing area 51. Vol. 1. Introductory material, Bony Fishes Families: Acanthuridae to Clupeidae*. FAO, Rome, pages variable.

- Froese, R., Pauly, D. (Eds), 2023. *FishBase*. World Wide Web electronic publication. www.fishbase.org (10/2023) (Accessed on 13-11-2023).
- Golani, D., 1996. The Marine Ichthyofauna of the Eastern Levant-History, Inventory, and Characterization. *Israel Journal of Zoology*, 42, 15-55.
- Golani, D., 2021. An updated Checklist of the Mediterranean fishes of Israel, with illustrations of recently recorded species and delineation of Lessepsian migrants. *Zootaxa*, 4956 (1), 1-108.
- Golani, D., Ben-Tuvia, A., 1986. New records of fishes from the Mediterranean coast of Israel including Red Sea immigrants. *Cybium*, 10, 285-291.
- Golani, D., Fricke, R., 2018. Checklist of the Red Sea Fishes with delineation of the Gulf of Suez, Gulf of Aqaba, endemism and Lessepsian migrants. *Zootaxa*, 4509, 1-215.
- Golani, D., Sonin, O., 1996. The occurrence of the tropical West African marine fishes *Acanthurus monroviae* (Acanthuridae) and *Arius parkii* (Ariidae) in the Levant. *Aqua-Journal of ichthyology and Aquatic Biology*, 2, 1-3.
- Golani, D., Orsi-Relini L., Massutí E., Quignard J.-P., 2002. *CIESM atlas of exotic species in the Mediterranean. Volume 1. Fishes*. Briand, F. (Ed). CIESM Publishers, Monaco, 254 pp.
- Golani, D., Askarov, G., Appelbaum-Golani, B., 2015. First confirmed record of the Blue Tang, *Acanthurus coeruleus* (Actinopterygii: Perciformes: Acanthuridae) in the Mediterranean. *Acta Ichthyologica et Piscatoria*, 45 (3), 311-313.
- Golani, D., Azzurro, E., Dulčić, J., Massutí, E., Orsi-Relini, L., 2021. *Atlas of Exotic Fishes in the Mediterranean Sea, 2nd Edn*. Briand F. (Ed). CIESM Publishers, Paris, 365 pp.
- Golani, D., Einbinder, S., Shemesh, E., Tchernov, D., 2023. First record of the Western Atlantic cocoa damselfish *Stegastes variabilis* from the Levant (Perciformes: Pomacentridae). *Journal of the Marine Biological Association of the United Kingdom*, 103, e22, 1-2.
- Gordon, H., Cousins, B.J., Newberry, M.A., Farcy, D.A., 2021. Clinical considerations in initial evaluation and treatment of Hardhead catfish spine puncture wounds. *Case Reports in Emergency Medicine*, 8841234, 6 pp.
- Günther, A., 1864. *Catalogue of the fishes in the British Museum, Volume 5. Catalogue of the Physostomi, containing the families Siluridae, Characinae, Haplochromidae, Sternopygidae, Scopelidae, Stomiidae in the collection of the British Museum*. Trustees, London, 455 pp.
- Hussein, D.M.A., Lucchetti, M.C., Zaqoot, A.A., Penca, J., Hussein, M.A., 2022. Status of Fisheries in Gaza Strip: Past Trends and Challenges. *International Journal of Euro-Mediterranean Studies*, 15 (2), 179-216.
- iNaturalist contributors, iNaturalist, 2023a. iNaturalist Research-grade Observations. iNaturalist.org. Occurrence dataset <https://doi.org/10.15468/ab3s5x> accessed via GBIF.org on 2023-11-14.
- iNaturalist contributors, iNaturalist, 2023b. iNaturalist Research-grade Observations. iNaturalist.org. Occurrence dataset <https://doi.org/10.15468/ab3s5x> accessed via GBIF.org on 2023-11-14.
- iNaturalist contributors, iNaturalist, 2023c. iNaturalist Research-grade Observations. iNaturalist.org. Occurrence dataset <https://doi.org/10.15468/ab3s5x> accessed via GBIF.org on 2023-11-14.
- Ivanova, N.V., Zemlak, T.S., Hanner, R.H., Hebert, P.D., 2007. Universal primer cocktails for fish DNA barcoding. *Molecular Ecology Notes*, 7(4), 544-548.
- Kailola, P.J., 1999. Order Siluriformes Ariidae (= Tachysuridae). Sea catfishes (fork-tailed catfishes). p. 1827-1879. In: *FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Volume 3. Batoid fishes, chimaeras and bony fishes part 1 (Elopidae to Linophrynidae)*. Carpenter, K.E., Niem, V.H. (Eds). FAO, Rome.
- Kailola, P.J., Bussing, W.A., 1995. Ariidae. Bagres marinos. p. 860-886. In: *Guía FAO para la identificación de especies para los fines de la pesca: Pacífico centro-oriental. Volume 2, Vertebrados, Parte 1*. Fischer, W., Krupp, F., Schneider, W., Sommer, C., Carpenter, K.E., Niem, V.H. (Eds). FAO, Rome.
- Kovačić, M., Lipej, L., Dulčić, J., Iglesias, S.P., Goren, M., 2021. Evidence-based checklist of the Mediterranean Sea fishes. *Zootaxa*, 4998 (1), 1-115.
- Marceniuk, A.P., 2005. Key for identification of the sea catfishes species (Siluriformes, Ariidae) of the Brazilian coast. Chave para identificação das espécies de bagres marinhos (Siluriformes, Ariidae) da costa brasileira. *Boletim Do Instituto De Pesca*, 31(2), 89-101. (In Portuguese, Brazil).
- Marceniuk, A.P., Menezes, N.A., 2007. Systematics of the family Ariidae (Ostariophysi, Siluriformes), with a redefinition of the genera. *Zootaxa*, 1416, 1-126.
- Marceniuk, A.P., Acero, P.A., Cooke, R., Betancur-R, R., 2017a. Taxonomic revision of the New World genus *Ariopsis* Gill (Siluriformes: Ariidae), with description of two new species. *Zootaxa*, 4290 (1), 1-042.
- Marceniuk, A.P., Bogorodsky, S.V., Mal, A.O., Alpermann, T.J., 2017b. Redescription of the blacktip sea catfish *Plicofollis dussumieri* (Valenciennes) (Siluriformes: Ariidae), with a new record from the Red Sea and notes on the diversity and distribution of *Plicofollis* spp. *Marine Biodiversity*, 47, 1239-1250.
- Marceniuk, A.P., Oliveira, C., Ferraris, C.J. Jr., 2024. A new classification of the family Ariidae (Osteichthyes: Ostariophysi: Siluriformes) based on combined analyses of morphological and molecular data. *Zoological Journal of the Linnean Society*, 200(2), 426-476.
- Marsaly, B., Daugherty, D., Shipley, O.N., Gelpi, C., Boyd, N. et al., 2023. Contrasting ecological roles and flexible trophic interactions of two estuarine apex predators in the western Gulf of Mexico. *Marine Ecology Progress Series*, 709, 55-76.
- Messing, J., 1983. New M13 vectors for cloning. *Methods in Enzymology*, 101, 20-78.
- Oowski, A.R., Jargowsky, M.B., Cooper, P.T., Powers, S.P., Drymon, J.M., 2023. Dietary Habits of Hardhead (*Ariopsis felis*) and Gafftopsail (*Bagre marinus*) Catfish revealed through DNA Barcoding of stomach contents. *Fishes*, 8, 539.
- Pensinger, L.G., Brown-Peterson, N.J., Green, C.C., Midway, S.R., 2021. Reproductive biology of hardhead catfish *Ariopsis felis*: evidence for overwintering oocytes. *Journal of Fish Biology*, 99, 308-320.
- Psomadakis, P.N., Giustino, S., Vacchi, M., 2012. Mediterranean

- nean fish biodiversity: an updated inventory with focus on the Ligurian and Tyrrhenian seas. *Zootaxa*, 3263, 1-46.
- Ratnasingham, S., Hebert, P.D.N., 2007. Bold: The barcode of life data system ([Http://www.barcodinglife.org](http://www.barcodinglife.org)). *Molecular Ecology Notes*, 7, 355-364.
- Ronje, E.I., Barry, K.P., Sinclair, C., Grace, M.A., Barros, N. *et al.*, 2017. A common bottlenose dolphin (*Tursiops truncatus*) prey handling technique for marine catfish (Ariidae) in the northern Gulf of Mexico. *PLoS ONE*, 12(7), e0181179.
- Saitou, N., Nei, M., 1987. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, 4(4), 406-25.
- Simier, M., Osse, O.J.F., Sadio, O., Ecoutin, J-M., 2021. Biology and ecology of sea catfish (Ariidae) of estuarine, lagoon and coastal ecosystems in West Africa. *Journal of Fish Biology*, 99, 629-643.
- Tamura, K., Stecher, G., Kumar, S., 2021. MEGA11: Molecular Evolutionary Genetics Analysis version 11. *Molecular Biology and Evolution*, 38, 3022-3027.
- Ubeid, K.F., 2011. Sand characteristics and beach profiles of the coast of Gaza Strip, Palestine. *Serie Correlación Geológica*, 27 (2), 125-136.
- Valdez-Moreno, M., Ivanova, N.V., Elias-Gutierrez, M., Pedersen, S.L., Bessonov, K., *et al.*, 2019. Using eDNA to bio-monitor the fish community in a tropical oligotrophic lake. *Plos One*, 14 (4), e0215505.
- Vella, A., Agius Darmanin, S., Vella, N., 2015. Morphological and genetic barcoding study confirming the first *Stegastes variabilis* (Castelnau, 1855) report in the Mediterranean. *Mediterranean Marine Science*, 16, 606-612.
- Vella, A., Vella, N., Agius Darmanin, S., 2016. The first record of the longjaw squirrelfish, *Holocentrus adscensionis* (Osbeck, 1765) (Holocentriformes: Holocentridae), in the Mediterranean Sea. *International Journal of Natural and Engineering Sciences*, 1 (3), 78-85.
- WoRMS Editorial Board, 2023. *World Register of Marine Species*. Available from <https://www.marinespecies.org> at VLIZ. Accessed 12-12-2023.
- Zenetos, A., Albano, P.G., López Garcia, E., Stern, N., Tsiamis, K. *et al.*, 2022. Established non-indigenous species increased by 40% in 11 years in the Mediterranean Sea. *Mediterranean Marine Science*, 23 (1), 196-212.