

Mediterranean Marine Science

Vol 26, No 1 (2025)

Mediterranean Marine Science



Complementing underwater visual surveys with eDNA metabarcoding to detect Mediterranean non-indigenous fishes

PANAGIOTA XANTHOPOULOU, STELIOS KATSANEVAKIS, MICHALIS RAGKOUSIS, ORESTIS PAPADAKIS, MARIA ZOTOU, NIKOLAOS KAMIDIS, OWEN S. WANGENSTEEN, VASILLIS PAPATHANASIOU, DIMITRIOS KARAMPETSIS, ANTONIOS D. MAZARIS, CHRYSOULA GUBILI

doi: [10.12681/mms.37756](https://doi.org/10.12681/mms.37756)

To cite this article:

XANTHOPOULOU, P., KATSANEVAKIS, S., RAGKOUSIS, M., PAPADAKIS, O., ZOTOU, M., KAMIDIS, N., WANGENSTEEN, O. S., PAPATHANASIOU, V., KARAMPETSIS, D., MAZARIS, A. D., & GUBILI, C. (2025). Complementing underwater visual surveys with eDNA metabarcoding to detect Mediterranean non-indigenous fishes. *Mediterranean Marine Science*, 26(1), 216–229. <https://doi.org/10.12681/mms.37756>

Complementing underwater visual surveys with eDNA metabarcoding to detect Mediterranean non-indigenous fishes

Panagiota XANTHOPOULOU^{1,2}, Stelios KATSANEVAKIS¹, Michalis RAGKOUSIS¹, Orestis PAPADAKIS¹, Maria ZOTOU¹, Nikolaos KAMIDIS², Owen S. WANGENSTEEN³, Vasillis PAPATHANASIOU², Dimitrios KARAMPETSIS^{2,4}, Antonios D. MAZARIS⁴, and Chrysoula GUBILI^{2,5}

¹ Department of Marine Sciences, University of the Aegean, 81100 Mytilene, Greece

² Hellenic Agricultural Organisation-DIMITRA, Fisheries Research Institute, 64007 Nea Peramos, Kavala, Greece

³ Department of Evolutionary Biology, Ecology and Environmental Sciences and Biodiversity Research Institute (IRBio), University of Barcelona, 08028 Barcelona, Catalonia, Spain

⁴ Department of Ecology, School of Biology, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece

⁵ School of Science, Engineering, and Environment, University of Salford, M5 4WT Salford, UK

Corresponding author: Panagiota XANTHOPOULOU; xanthopop@elgo.gr

Contributing Editor: Costas S. TSIGENOPOULOS

Received: 14 May 2024; Accepted: 05 February 2025; Published online: 27 February 2025

Abstract

Non-indigenous species (NIS) are among the main drivers of native biodiversity loss, habitat alteration, and degradation of ecosystem services, with some NIS posing risks to human health. Efficient monitoring strategies are necessary to assess the distribution and impacts of NIS. In this study, we compared the performance of environmental DNA (eDNA) metabarcoding and visual surveys with SCUBA diving to detect marine fish NIS in Greek waters. We collected water samples from 20 coastal sites across the Aegean, Ionian, and Levantine Seas in both warm and cold periods, targeting the 12S rRNA region. A reference 12S Mediterranean NIS sequence database was created to improve regional monitoring. Underwater visual surveys were performed at the same sites to visually detect fish NIS. Overall, 15 non-indigenous fishes were detected, five with both eDNA and visual surveys, seven exclusively by eDNA, and three by visual surveys alone. The southern stations yielded more NIS detections than the northern stations in both periods. Our findings demonstrate that eDNA can provide a rapid, low-cost, and effective tool, advocating for its integration into systematic NIS monitoring in the eastern Mediterranean Sea. A comprehensive barcode reference database is essential in enhancing the effectiveness of eDNA approaches. Thus, the combination of eDNA metabarcoding and traditional underwater visual surveys is recommended for comprehensive monitoring of NIS in marine environments.

Keywords: eDNA metabarcoding; coastal; biological invasions; eastern Mediterranean Sea; 12S rRNA; visual survey.

Introduction

Biological invasions are complex, multistage processes, and their success is influenced by several factors at each stage (Blackburn *et al.*, 2011; Briski *et al.*, 2018). Ecological and socioeconomic impacts of these invasions are escalating globally, yet mitigation measures are often delayed, insufficient, or non-existent (Ahmed *et al.*, 2022). Consequently, post-invasion management costs exceed pre-invasion expenses (for prevention) by more than 25-fold (Cuthbert *et al.*, 2022). Given the exponential increase in the cost of inaction against non-indigenous species (NIS), it is imperative to implement management actions during early invasion stages (Giakoumi *et al.*, 2019; Katsanevakis *et al.*, 2023). Effective decision making in this context requires detailed information about

NIS introduction pathways, documented occurrence, distribution ranges, and population sizes (Briski *et al.*, 2018; Galanidi & Zenetos, 2022). Standardised protocols that yield consistent data and time series are crucial to enhance the quality of current state assessments and future projections (Ziegler, 2013; Borja *et al.*, 2016). Such data can be collected through various approaches, such as targeted scientific surveys, fisheries records, and citizen science initiatives (e.g., Katsanevakis *et al.*, 2012a; Zenetos *et al.*, 2013).

In the Mediterranean Sea, a biodiversity hotspot of over 17,000 marine species (Coll *et al.*, 2010), a total of 993 NIS has been reported, with 751 species confirmed as established by 2021 (Zenetos *et al.*, 2022). NIS establishment rate has increased by 40% from 2010 to 2021 (Zenetos *et al.*, 2022). The steady rise of sea surface tem-

perature during the past decades (Pastor *et al.*, 2020) has facilitated the establishment and spread of thermophilic Lessepsian species in the eastern and central Mediterranean Sea (Raitsos *et al.*, 2010; Katsanevakis *et al.*, 2014; Karachle *et al.*, 2022). However, it has been debated whether the recent increase in NIS records represents an actual increase in NIS introductions or reflects intensified scientific efforts and citizen science initiatives (Bailey *et al.*, 2020; Galanidi & Zenetos, 2022; Zenetos *et al.*, 2022). There is a growing number of NIS and cryptogenic species in the eastern Mediterranean Sea (Zenetos *et al.*, 2018; Ragkousis *et al.*, 2023), with at least 221 species recorded in the Greek Seas, including 17 classified as invasive, and 148 with established populations, while the establishment status of the remaining species remains uncertain (Zenetos *et al.*, 2018).

Fish is a major group of NIS with 173 established or casual species (42 in the western, 65 in the central, and 139 in the eastern Mediterranean Sea, respectively; Zenetos *et al.*, 2022; Galanidi *et al.*, 2023). At least 40 have been detected in Greece, with five being characterised as invasive (Korakaki *et al.*, 2021). The three most reported invasive fishes are *Siganus luridus*, *Siganus rivulatus*, and *Pterois miles* (Ragkousis *et al.*, 2023). *Siganus* grazers tend to overgraze on coastal rocky reefs, causing the degradation of local habitats and communities, whereas *P. miles* is a voracious opportunistic predator feeding on important local biodiversity, even commercial fishes, causing disruption to local food webs and posing threats to human health with its venomous spines (Tsirintanis *et al.*, 2022, 2023). NIS richness is higher in southern Greece (South Aegean and Levantine Seas), due to temperature regimes favourable to thermophilic Lessepsian species (Ragkousis *et al.*, 2023; Evangelopoulos *et al.*, 2024). Moreover, a positive correlation has been observed between the number of NIS introduced as transport stowaways and proximity to major ports (Zenetos *et al.*, 2018; Ragkousis *et al.*, 2023).

NIS early detection is critical for effective management (Katsanevakis *et al.*, 2023). Traditional methods such as fishing nets, scuba diving, underwater cameras, and settlement panels are widely used to detect and monitor NIS species (Tamburini *et al.*, 2021; Evangelopoulos *et al.*, 2024). Each method has significant strengths and can be deployed separately or combined in NIS assessments (Bessell *et al.*, 2023; Robinson *et al.*, 2023; West *et al.*, 2024). They have been included to support decision making and management (Lehtiniemi *et al.*, 2015; Katsanevakis *et al.*, 2023), however, they tend to be costly and could underestimate species occurrence, richness and population state variables, while they might be highly intrusive to local ecosystems (Katsanevakis *et al.*, 2012a). Environmental DNA (eDNA) analysis has evolved as a promising alternative to overcome most limitations of traditional sampling methods (Taberlet *et al.*, 2012; Borrell *et al.*, 2017; Fediajevaite *et al.*, 2021). It relies on the retrieval of the genetic material released into the aquatic environment (e.g., water) and on the taxonomic assign-

ment using DNA metabarcoding (Taberlet *et al.*, 2012). In marine research, eDNA has emerged as a groundbreaking tool for monitoring the diversity and the distribution of marine organisms, ranging from plankton to top predators (Stat *et al.*, 2017; Zhang *et al.*, 2020b). By principal, eDNA requires less sampling effort (e.g., fewer sampling trips), less taxonomic expertise, and is cost effective and less invasive than traditional sampling methods, as there is no direct interaction with the target organisms and the habitat (Deiner *et al.*, 2017). One of the major advantages of eDNA though, is that it enables the detection of low abundance species (Jerde *et al.*, 2011), which is crucial for monitoring cryptic, rare, endangered, and invasive species at the early stages of invasion (Kelly *et al.*, 2014; Aglieri *et al.*, 2023).

The potential of eDNA for marine NIS detection and its integration in management strategies is increasingly discussed (Sepulveda *et al.*, 2020b; Morissette *et al.*, 2021; Zangaro *et al.*, 2021). Nevertheless, eDNA based community assessments and protocols are still to be standardized to be officially included in management recommendations and policy (Sepulveda *et al.*, 2020b; Morissette *et al.*, 2021). Moreover, incomplete reference databases limit the method's applicability (Ruppert *et al.*, 2019; Doble *et al.*, 2020; Duarte *et al.*, 2021). For example, among the 173 NIS fish established or casually encountered across the Mediterranean Sea (Zenetos *et al.*, 2022), 52 are currently lacking an available reference barcode for the fish-specific 12S rRNA primer MiFish (Miya *et al.*, 2015), hampering species assignment and detection of taxa. Here, we aim to investigate the efficacy of eDNA sampling in detecting NIS fishes across Greek coastal zones and to compare the results from eDNA to those from standardized underwater visual surveys.

Material and Methods

Sample collection and processing

Seawater samples for eDNA analyses were collected from 20 sites across Greek territorial waters between February 2021 and March 2022 (Fig. 1). Two sampling expeditions per site were carried out, one during the warm (May-October) and one during the cold period (November-April). In total, 120 water samples were collected using a 2-L Niskin water sampler at approximately 10 m depth (three replicate 1 L samples at each sampling station). Each 1 L sample was filtered through a 0.22 µm Sterivex filter capsule (Merck Millipore) and promptly stored at -20 °C for subsequent processing (one filter was discarded during filtration due to possible contamination by contact). All equipment was thoroughly cleaned with 10% bleach and distilled water before use, and sterilized single-use syringes were employed for filtration. To investigate possible contamination, three *in situ* negative controls and one PCR blank were included, each involving the filtration of bottled commercial water.




Fig. 1: Approximate environmental DNA sampling and underwater visual survey (SCUBA) sites for non-indigenous fish species detection across the four main regions of Greek waters.

eDNA extraction, amplification, and sequencing

Filters were retrieved from Sterivex capsules and DNA was extracted using the Qiagen DNeasy Blood and Tissue Kit sensu Cowart *et al.* (2018) with minor modifications. Firstly, we extracted DNA from the entire filter; secondly, we did not use whitebeads and incubated the filter with 720 µl ATL and 80 µl Proteinase K overnight. The addition of RNase was omitted, DNA was eluted twice with 40 µl AE and stored at -20 °C. A ~170 bp barcoding region of the 12S rRNA was amplified using the MiFish primer set, targeting fish detection (Miya *et al.*, 2015). Each DNA extract was amplified in methodological PCR triplicates, which were pooled to maximize the detection of rare taxa. The PCR mix included 10 µl AmpliTaq gold 360Master mix (Applied Biosystems), 0.16 µl Bovine Serum Albumin (20 µg/µl, Thermo Scientific), 4.84 µl H₂O, 1 µl of the forward and 1 µl of the reverse tagged primer (5 µM, including 2–4 leading Ns and 8-bp sample tags) and 3 µl of the extracted eDNA. The amplification conditions included denaturation at 95 °C for 10 min, 40 cycles at 95 °C for 30 s, 60 °C for 30 s and, 72 °C for 30 s and a final extension at 72 °C for 5 min. Upon PCR amplification, all samples per plate were pooled and concentrated using a MinElute PCR Purification Kit (Qiagen), following the manufacturer's guidelines. The final DNA concentration was measured using a Qubit 4 Fluorometer (Invitrogen). Library preparation was performed through the TruSeq DNA PCR-Free High Throughput Library Prep Kit (Illumina) with 2 µg of DNA and included a)

end repairing, b) MinElute concentration, c) adenylation, d) ligation, and e) clean-up. Library concentration was quantified to avoid overclustering the Illumina flow-cell using the NEBNext Library Quant Kit for Illumina (New England Biolabs), following the kit's quick-start protocol. A pair-end sequencing (2 x 150 bp) was carried out on an Illumina NovaSeq 6000 sequencer (Illumina, USA) at Novogene facilities (Cambridge, UK).

Bioinformatic analyses

Sequenced data in multiplexed fastq files were processed through the MJOLNIR pipeline (<https://github.com/uit-metabarcoding/MJOLNIR>), primarily utilizing tools from the OBITools package (Boyer *et al.*, 2016). Paired-end reads were aligned using *illuminapairedend*, retaining only those with an alignment quality score above 40. Demultiplexing and primer-sequence removal were performed by *ngsfilter*, discarding sequences with mismatched primer tags. Length filtering and dereplication of sequences were performed with *obigrep* and *obiuniq*, retaining sequences of 140-190 bp. Singleton sequences and chimeric amplicons were eliminated using the *uchime denovo* algorithm from VSEARCH (Rognes *et al.*, 2016). Molecular operational taxonomic units (MOTUs) delimitation based on linkage networks was implemented using the SWARM procedure (Mahé *et al.*, 2015).

Taxonomic assignment was performed using the

DNA Universal-databank for Fisheries and Aquaculture reference database (DUFA; last updated on 2022-01-06) for the 12S MiFish fragment (<https://github.com/uit-metabarcoding/DUFA>). Subsequent to the taxonomic assignment, putative pseudogenes were removed using LULU (Frøslev *et al.*, 2017). DUFA NIS availability was cross-referenced with an updated Mediterranean Sea NIS list (Zenetos *et al.*, 2022). Additionally, we manually added 75 published 12S sequences absent in the DUFA 12S database (excluding unverified entries; Table S1). These sequences were retrieved from the National Center for Biotechnology Information (NCBI) Taxonomy database (<https://www.ncbi.nlm.nih.gov/taxonomy>). We also generated 12S sequences for four species that were not included in the NCBI database (*Torquigener flavimaculosus*, *Stephanolepis diaspros*, *Upeneus pori*, and *Pteragogus trispilus*; Accession Numbers: PQ638932-35, respectively), using the NucleoSpin Tissue kit (Macherey-Nagel, Germany) for DNA extraction and the MiFish primer pair (F: 5'-GTCGGTAAACTCGTGCCAGC-3'; R: 5'-CATAGTGGGGTATCTAATCCCAGTTTG-3') for amplification (Miya *et al.*, 2015). The PCR protocol included an initial denaturation at 94 °C for 3 min, followed by 30 cycles at 94 °C for 50 s, 50 °C for 50 s, 72 °C for 50 s, and a final extension at 72 °C for 5 min. PCR products were commercially sequenced (Macrogen, the Netherlands), and sequences were manually checked and edited using ProSeq 3.0 (Filatov, 2002).

Environmental variables

During eDNA sampling expeditions, water temperature, salinity, and dissolved oxygen were measured *in situ* using an Orion multi-parametric device. Turbidity was measured at each station with an Orion AQUAfast turbidity-meter. Additional water samples were collected from all sites to analyze five nutrient salts (PO_4^{3-} , NO_2^- , NO_3^- , SiO_2 , and NH_4^+), total suspended solids (TSS), and chlorophyll-a concentrations at approximately 10 m depth using a Niskin Water Sampler. Approximately 500 mL of sea water were filtrated under vacuum pressure through 0.45 μm pre-weighted nitrocellulose filters and kept frozen at -20 °C for nutrient analyses according to Parsons *et al.* (1984). The filters were also used to estimate TSS according to APHA (1998). For chlorophyll-a analysis, 1 L of water samples were filtrated through 47 mm diameter GF/F glass fiber filters. Filters were subsequently diluted in 10 mL of 90% acetone, stored overnight at 4 °C in the dark and analyzed following APHA (1998). All analyses were performed on a HITACHI U-2001 spectrophotometer. Analytical precision was tested by triplicate analysis of 40% of all samples.

Underwater visual surveys

Separate field expeditions were conducted for underwater visual surveys. To assess the presence, population density, and biomass of alien marine fish species, SCU-

BA diving surveys were conducted by two divers. Strip transects and line transect distance sampling were conducted along three consecutive replicate transects (25 m in length and 5 m in width) at two depth zones (5 m and 15 m), according to Thanopoulou *et al.* (2018). The first diver conducted strip transects targeting mobile species, estimating the total length (TL) of each identified fish. The second diver conducted distance sampling (Buckland *et al.*, 2001) targeting less mobile and cryptic species, recording the vertical distance of each fish from the line transect (Buckland *et al.*, 2001) and estimating its TL. The dominant substrate type (soft or hard) was recorded, and taxon identification was made *in situ* for most species, or by analyzing photographic and video material in cases of uncertainty.

Statistical analysis

For community analysis, environmental triplicate samples were merged and treated as a single sample. A minimum of 97% identity was used for species level identification (Miya *et al.*, 2015). Additionally, MOTUs with relative read abundance per sample below 0.005 or less than five reads were set to zero to minimize tag-switching bias (Antich *et al.*, 2023) and reduce the likelihood of false positives from potential contamination. All datasets were treated as qualitative presence-absence data on each MOTU per sample and all statistical analyses were performed in R 4.1 (<https://cran.r-project.org>). Non-metric multidimensional scaling (NMDS) ordinations, using the *metaMDS* function in the “vegan” package version 2.6.2 (Oksanen *et al.*, 2022), provided a reduced-space graphical representation of the species composition per site. Distance matrices were calculated using the Jaccard coefficient (function *vegdist*). Permutational analysis of variance (PERMANOVA) was performed to assess the influence of substrate (soft or hard), period (warm or cold), and sampling region (North Aegean, Central Aegean, Ionian, and Levantine Seas) on the composition of non-indigenous fish communities based on eDNA. PERMANOVA was conducted using the *adonis* function in the “vegan” package and using presence/absence data with 10,000 permutations. The wrapper function *pairwise.adonis2* from the package “pairwiseAdonis” 0.4.1 (Martinez Arbizu, 2020) was used for multilevel pairwise comparisons. NMDS plots were generated in R, heatmaps in Flourish (<https://flourish.studio/>), maps in QGIS 3.14 (<https://www.qgis.org>), and images were modified using Inkscape 1.1 (<https://inkscape.org/>).

Generalized Additive Models (GAMs; Hastie & Tibshirani, 1990) were used to investigate the effects of environmental variables on the NIS eDNA detectability on sites of species known occurrence, based on our visual surveys, literature, and online databases such as EASIN (Katsanevakis *et al.*, 2012b) and ELNAIS (Zenetos *et al.*, 2015). We applied the *gam* function in the “mgcv” package 1.8.34 in R (Wood *et al.*, 2016) and used the binomial distribution family. Collinearity among environmental variables was assessed using the Pearson correla-

tion coefficient in the “PerformanceAnalytics” package 2.0.4 (Peterson *et al.*, 2020). Predictor variables included water temperature, salinity, turbidity, total suspended solids (TSS), and total dissolved nitrogen (TDN) calculated as the sum of NO_2^- , NO_3^- , and NH_4^+ . Additionally, NIS was set as a factor to control any species-specific detection bias. Candidate models (Venables & Ripley, 2002) are listed in Table S2. The Akaike Information Criterion (AIC; Akaike, 1973; Burnham & Anderson, 2002) was used for model selection.

Results

Approximately, 12.4 million paired-end reads were generated (103,766 on average per sample), 11,410,677 of which were retained for taxonomic assignment after filtering and MOTU delimitation, and 9,286,019 were subsequently assigned to a taxonomic group by the MJOL-NIR pipeline (Fig. S1). Only Actinopterygii reads were retained for downstream analyses; 72,696 reads (~0.59%) belonged to the classes Mammalia and Aves. The PCR blank and one negative control provided no reads for any MOTU. Between the other two negative controls, both

had few reads of a MOTU assigned to *P. miles* (five and three), whereas the second had three additional reads of a *F. commersonii* MOTU. Due to the low number of reads in negative controls, we maintained all MOTUs for downstream analyses. We detected 250 unique MOTUs of fish, representing 38 orders and 72 families. For this study, only fish NIS were considered, comprising 4.41% of the total filtered Actinopterygii reads ($n = 403,816$). In total, 13 MOTUs belonged to 12 established NIS in the Mediterranean Sea (*Atherinomorus forskalii*, *Fistularia commersonii*, *Lagocephalus sceleratus*, *Lagocephalus suezensis*, *Parexocoetus mento*, *Pterois miles*, *Sargocentron rubrum*, *Siganus luridus*, *Siganus rivulatus*, *Stephanolepis diaspros*, *Torquigener flavimaculosus*, and *Upenus pori*). NIS richness was higher in the Levantine and Ionian Seas, with less records in the North Aegean (Fig. 2 and S2). Four species (*L. suezensis*, *S. rubrum*, *T. flavimaculosus*, and *U. pori*) were exclusive to the Levantine Sea, whilst *P. mento* was solely recorded in the central/eastern Aegean Sea (Kalymnos Island; Fig. 2). Five species were recorded only once through eDNA (*U. pori*, *T. flavimaculosus*, *S. rubrum*, *P. mento*, and *L. suezensis*). Three species were exclusively found in the southern Aegean and the Levantine Seas during the cold period (*L.*




Fig. 2: Non-indigenous fish species detected in Greece using environmental DNA analysis, colours indicate species detection during warm (May-October), cold (November-April), and both periods; black masks indicate the detection with *in situ* visual observation by SCUBA diving and pins indicate previously documented occurrence in literature (references in Table S6).

suezensis, *P. mento*, and *U. pori*), whereas *S. rubrum* and *S. diaspros* were recorded in Crete and North/Central Aegean, respectively, during warm months.

Environmental variables

Eleven environmental variables were recorded at each station (Fig. S3). The highest temperatures and lowest salinity levels were documented in the North Aegean Sea (in the warm period). The lowest temperature was recorded at Skroponeria in the cold period (13 °C) and the highest at Thasos Island in the warm period (26.15 °C). Salinity ranged from 35.30 psu (Thasos Island, cold period) to 39.35 psu (Kastelorizo Island, warm period), with values below 38 occurring in the North Aegean sampling sites (Table S3). Chlorophyll-a ranged from 0.058 (Skroponeria, warm period) to 1.065 µg/L (Thasos Island, cold period). North Aegean stations (Thassos, Samothraki, Chalkidiki, Skyros, and Lesbos) demonstrated higher phosphate, nitrate, and silicate concentrations for both periods; Ionian stations showed maximum ammonium, turbidity and TSS values. The Central Aegean stations exhibited the highest nitrite concentrations during both periods. Additionally, collinearity among the environmental variables was checked (Fig. S4). Temperature was negatively correlated with dissolved oxygen and chlorophyll-a, but positively correlated with PO_4^{3-} and NO_3^- . Salinity was negatively correlated with chlorophyll-a, NO_3^- , and SiO_2 . NO_3^- and SiO_2 were significantly correlated with the highest number of environmental variables, whereas TSS did not show any significant correlation with other variables.

The NMDS ordinations (stress: 0.097) indicated that NIS composition was not affected by substrate, sampling period, or geographic area (Fig. S5), which was corroborated by PERMANOVA results ($p > 0.05$; Table S4). However, PERMANOVA showed a statistically significant interaction between period and area ($p = 0.031$). Pairwise comparisons revealed significant differences between the Ionian Sea and both the North Aegean ($R^2 = 0.178$; $p = 0.022$) and Levantine Seas ($R^2 = 0.227$; $p = 0.008$), and a weak clustering was depicted in the corresponding NMDS analysis (Fig. S5a).

Visual survey vs eDNA

Visual surveys identified eight non-indigenous fishes (*Alepes djedaba*, *Cheilodipterus novemstriatus*, *Parupeneus forsskali*, *P. miles*, *S. rubrum*, *S. luridus*, *S. rivulatus*, and *T. flavimaculosus*; Fig. S6). Most fishes were recorded in the Levantine and Ionian Seas, whereas *A. djedaba* was exclusively detected at Skyros Island (North Aegean Sea, Fig. S6). Five species were identified by both visual surveys and eDNA metabarcoding, however, seven were detected only through eDNA analysis. Three species, *A. djedaba*, *C. novemstriatus*, and *P. forsskali* were not detected in eDNA samples (Figs 2, 3, and S6).

GAMs

Five variables with possible implications on eDNA degradation were used for GAMs: temperature, salinity, turbidity, TSS, and TDN. Among the 12 candidate models (Table S2) the highest percentage of deviance explained and adjusted R^2 were 17.80% and 0.087 (m1), respectively. The best GAM, indicated by the lowest AIC value, included water temperature and TDN (m10), with none of them being statistically significant (Tables S2 and S5; Fig. S7); this model explained only 4.71% of the deviance.

Discussion

The Mediterranean ecosystems are under dramatic pressure due to biological invasions and climate change, e.g., sea-surface temperature increase, algal blooms, and heatwaves driving mass mortalities, (Boudouresque *et al.*, 2017; Garrabou *et al.*, 2022; Chatzimentor *et al.*, 2023). The rapid establishment of NIS in the eastern Mediterranean Sea, particularly those with a direct and significant impact on local biodiversity or ecosystem services, highlights the need for proper detection and management of their populations (Galil *et al.*, 2018; Roy *et al.*, 2019). Here, we applied eDNA metabarcoding aiming to detect marine fish NIS, and we were able to report almost twice the species that were detected by visual surveys. These findings support the potential of eDNA as an effective monitoring tool for detecting marine fish NIS in the region. Our study further demonstrated that it is important to expand existing monitoring schemes considering different sources of information.

Invasive fish NIS have more severe negative cumulative impact on coastal hard substrate habitats, being more intense in the South Aegean Sea where NIS are more abundant compared to the North (Tsirintanis *et al.*, 2023). Most of the NIS reported in Greece prefer shallow reef-associated areas, with a few exceptions such as *Tylerius spinosissimus*, that is caught in depths greater than 90 m (Froese & Pauly, 2023). Our eDNA samples were collected from coastal waters, hence only species inhabiting shallow waters of the continental shelf were detected. Fish NIS detected in this study have been recorded in depths down to 132 m (*F. commersonii*), however most are restricted in shallow waters, usually <60 m (Froese & Pauly, 2023). Most of them were detected in lower latitude stations; however, our results suggest broader NIS fish distributions than previously reported or documented by the visual surveys of this study (Fig. 2). In some cases, eDNA revealed NIS presence in northern sites where no publicly available records were available, such as *F. commersonii* and *P. miles*; their presence at these sites had not been documented previously, although specimens have been found in adjacent areas (e.g., Katsanevakis *et al.*, 2020b). For example, *A. forsskali* is one of the first and most abundant lessepsian migrants in the eastern Mediterranean Sea (Tsirintanis *et al.*, 2022 and references herein), however it has few records in Greece (Fig. 2).

Our results revealed a wider distribution range than previously reported, with Lefkada Island (Ionian Sea) being currently the most westerly site of occurrence in the Mediterranean Sea (unpubl. data). Such findings are crucial for species such as *L. scleratus* which was detected at two sites for the first time and is known for posing direct threats to local ecosystems, fisheries, and human health (Katsanevakis *et al.*, 2018; Tsirintanis *et al.*, 2023).

Potential environmental predictors (e.g., salinity, nitrogen content as TDN, turbidity, and TSS) and the substrate type were tested using GAMs and PERMANOVA, respectively. Our analysis did not reveal any potential impact of environmental variation on NIS eDNA detection, likely due to the small sample size (Tables S2 and S5). Additionally, substrate type categories incorporated a variety of ecological niches and interactions, potentially masking any impact. Despite fish NIS occurrence reported at various study sites (Table S6), temporal NIS distribution patterns might be affecting our results; most NIS detected in our study are thermophilic and are anticipated to be present in areas with higher water temperatures (Rilov & Galil, 2009). However, no seasonal variation has been observed to affect abundance and biomass of most NIS species (Mavruk *et al.*, 2017), corroborating our results. Furthermore, no studies have attempted to outline the influence of other variables, such as NO_2^- , NH_4^+ , and or turbidity for lessepsian fish NIS range expansion to the best of our knowledge. Nevertheless, our results might align better with eDNA physical properties and fish NIS life cycle (Harrison *et al.*, 2019 and references herein). Accelerated DNA degradation caused by temperature increase has been reported in other studies (Strickler *et al.*, 2015; Mauvisseau *et al.*, 2022). Similarly, elevated acidity and chlorophyll-a concentration, associated to raised bacterial abundance, negatively affect eDNA quality (Seymour *et al.*, 2018; McKnight *et al.*, 2024). Conversely, salinity itself or related factors, such as diminished microbe abundance, have been as-

sociated with slower eDNA degradation (Collins *et al.*, 2018; Saito & Doi, 2021), whereas TDN has not been linked to significant effects on eDNA decay (Seymour *et al.*, 2018; McKnight *et al.*, 2024). On the other hand, the rate of shedding DNA increases with higher temperature and fish biomass (Jo *et al.*, 2019). Hence, in the warmer regions of the study area, where many thermophilic fishes are more abundant, these two contradicting mechanisms may either increase or decrease detectability through eDNA under specific conditions in comparison to colder regions/periods. Higher detectability during spawning periods may also affect the effectiveness of eDNA (Tsuji & Shibata, 2021; Ostberg & Chase, 2022).

Traditional methods have been used to identify and monitor NIS before the advent of eDNA method, assisting NIS management (Zaiko *et al.*, 2018; Fediajevaite *et al.*, 2021). Here, we demonstrated that eDNA outperforms conventional monitoring methods in fish NIS detection. The higher efficiency of eDNA is corroborated in various studies (Valentini *et al.*, 2016; Stat *et al.*, 2017; Robinson *et al.*, 2023). Marine NIS detection using SCUBA-based visual surveys has few inherent limitations, which eDNA sampling can address. For example, SCUBA surveys might overlook NIS that are cryptic, small, or blend into their surroundings (Marchini *et al.*, 2015). Divers are also limited by depth, time, and the breadth of areas they can survey, making visual surveys challenging in habitats with high complexity (Friedlander & Parrish, 1998). Additionally, some fish NIS might be nocturnal or elusive, escaping daytime SCUBA surveys, yet their DNA fragments can still be detected in water samples (Kelly *et al.*, 2014). Nevertheless, three species were recorded by visual surveys and were not detected by eDNA analyses (Fig. 3). Therefore, eDNA metabarcoding has the potential to become a powerful tool in NIS biomonitoring assessments, complementing but not replacing traditional methods. The complementary use of both techniques in fish biomonitoring is consistently highlighted in recent




Fig. 3: Comparison of non-indigenous marine fish species detected in Greece using eDNA analysis and visual observation by SCUBA diving.

studies, with an apparent overlap in species detection and observed clear advantages of each method (Valdivia-Carrillo *et al.*, 2021; Robinson *et al.*, 2023).

The choice of molecular markers for meta-barcoding undeniably influences the outcome of eDNA studies (Zangaro *et al.*, 2021; Zhang *et al.*, 2020a). Mediterranean fish NIS, as a taxonomic group, have a considerably extensive barcode reference database, mostly referring to the cytochrome c oxidase I (COI) gene (Zangaro *et al.*, 2021), however, there are still species that are absent from public databases; particularly for other barcoding regions, e.g., 12S rRNA (Table S1). In this study, the MiFish primers pair was selected due to its higher fish specificity and its overall improved performance in fish detection to other primers. For example, it produces greater number and proportion of usable fish reads to COI primers despite the more complete reference database of the latter primer pair (Collins *et al.*, 2019; Zhang *et al.*, 2020a; Miya *et al.*, 2020). Interestingly, 31 of 97 fish NIS with established populations in the Mediterranean basin (Zenetos *et al.*, 2022), were missing a 12S rRNA reference sequence (<https://www.ncbi.nlm.nih.gov/nucleotide/>). We have generated four 12S rRNA sequences from species that were not included in existing databases and detected 12 out of 40 fish species listed as NIS in Greece since 2018; 32 of these species are already classified as invasive or established (Zenetos *et al.*, 2018). One species with confirmed records using traditional sampling methods (both visual observations and published records), was not detected in eDNA samples (e.g., Dimitriadis & Sourbès, 2015; Evangelopoulos *et al.*, 2020; Miliou & Loudaros, 2020). This discrepancy was mainly attributed to the incompleteness of reference databases. In the context of the present study, no reference sequences matching the MiFish 12S rRNA barcoding region of *P. forsskali* were found in public databases (NCBI). Despite the increased use of eDNA techniques for rare species detection, the incompleteness of reference databases is the most restricting factor in monitoring and management for NIS in the Mediterranean Sea. Moreover, primer specificity and efficiency for targeting specific groups should be considered. Numerous assays with group- or species-specific primers are designed, promising higher levels of precision and efficiency (Ardura, 2019; Mohammed-Geba *et al.*, 2020).

In this study, we focused on fish NIS detection in the eastern Mediterranean Sea by utilizing primers that primarily amplify fish (Miya *et al.*, 2015); our approach can be used for future research on fish NIS management, particularly as reference databases are constantly updated, improving identification accuracy and rates. Nevertheless, the current approach could be optimized. For example, the combined use of multiple primers could enhance species detection and/or richness as multi-marker approaches often yield more detailed and higher resolution community data by targeting more taxonomic groups (Ferreira *et al.*, 2024; Fontes *et al.*, 2024). Conversely, marker selection could be applied specifically for single species-based applications (Hartle-Mougiou *et al.*, 2024). Protocol optimization could focus on DNA yield by testing filtered seawater volume and/or using inhibitor

removal kits. Finding the optimal volume is extremely beneficial to monitoring strategies, particularly as small volumes can be advantageous due to their minimal physical and logistical requirements, faster sampling, processing times, and equipment simplicity. However, higher water volumes can improve eDNA metabarcoding results and consistency (Bessey *et al.*, 2020; Govindarajan *et al.*, 2022). Additionally, different extraction methods, types of filters, environmental parameters, and protocol costs should be examined (Duarte *et al.*, 2021; Fonseca *et al.*, 2023; Rishan *et al.*, 2023). Finally, a sufficient sample size should provide a stronger basis for statistical analysis, improving our ability to assess the impact of environmental variables on fish NIS detection. Therefore, while eDNA methods can be effective for alien species monitoring, they have not been extensively used in the Mediterranean Sea and protocol standardization is required to maximise their potential for fish NIS detection.

Recommendations for management actions and conservation priorities

Invasive fish species have created profound challenges for local ecosystems and the fishing industry in Greece (Katsanevakis *et al.*, 2020b; Christidis *et al.*, 2022). NIS compete for space and resources with native biodiversity and/or they alter the structure of local habitats (Tsirintanis *et al.*, 2022). Two fish NIS detected in this study (*S. luridus* and *S. rivulatus*) are among the “worst” invasive species in the Mediterranean Sea, whereas eleven have moderate to high impact on biodiversity, ecosystem services, and human health (Katsanevakis *et al.*, 2016; Tsirintanis *et al.*, 2022). Five of the recorded NIS (*L. scleratus*, *P. miles*, *S. luridus*, *S. rivulatus*, and *T. flavimaculosus*) pose serious threats to human health, mainly due to deleterious toxins and direct attacks to humans (Katikou *et al.*, 2009; Tsirintanis *et al.*, 2022). The invasive silver-cheeked toadfish (*L. scleratus*) also damages fishing gears and removes catches, with significant economic losses to coastal small-scale fisheries (Katsanevakis *et al.*, 2018; Christidis *et al.*, 2022). The two siganids (*S. luridus* and *S. rivulatus*), *F. commersonii*, and *P. miles* have also severe destructive impact on local ecosystems; grazing, an intense habitat-altering behaviour of the first two species, and the opportunistic behaviour of the latter, can reshape local marine food-webs and increase predation mortality of already stressed populations, affecting fisheries indirectly (Bariche *et al.*, 2013; Azzurro *et al.*, 2017; Katsanevakis *et al.*, 2018; Batjakas *et al.*, 2023). Despite the impact of invasive fish, only the striped catfish (*Plotosus lineatus*) is included in the European list of invasive alien species of Union Concern, in compliance with the EU Regulation 1143/2014, compromising effective management (Katsanevakis *et al.*, 2023).

Acknowledging the need for effective monitoring and management of NIS distribution is essential (Katsanevakis *et al.*, 2014). Various mitigation measurements have been proposed and/or applied for the restriction or reduction of invasive NIS (e.g., Galil *et al.*, 2019; Kleitou

et al., 2021 and references herein). However, eDNA is not included in official invasive NIS management plans, despite its efficiency (Sepulveda *et al.*, 2020b; Morissette *et al.*, 2021), although its value has been recently recognized by the European Commission (Costello & Trotter, 2023). In countries with increased NIS occurrence, such as Greece, it could produce vast amounts of occurrence data in short time, supporting NIS management. Our results revealed that eDNA can function as an early warning tool for new species introductions as in the case of NIS detected in the North Aegean Sea. Standardised protocols and novel technologies, e.g., automatic samplers positioned in key sites for NIS introductions, could facilitate systematic monitoring and act as early detection tools (Sepulveda *et al.*, 2020a; Aglieri *et al.*, 2023). Coordinated efforts, encompassing research and regulatory measures, are crucial to manage and mitigate risks posed by invasive fish in Greek waters. Public awareness and citizen science, combined with attentive eDNA sampling and SCUBA surveys could provide a holistic approach to monitoring effectively NIS populations and provide solid ground for effective management (Zenetos *et al.*, 2010; Bakker *et al.*, 2017; Parrondo *et al.*, 2018; Bessell *et al.*, 2023). Therefore, collaboration between local communities, researchers, and policymakers is imperative to successfully address this challenge (Packer *et al.*, 2017; Katsanevakis *et al.*, 2020a).

Acknowledgements

This work has been funded by the Operational Programme of Fisheries and Sea (OPFS) 2014-2020, Greece and the European Maritime and Fisheries Fund (EMFF) as part of the project “Monitoring and control of invasive alien species in Greece using innovative techniques under current and future climate conditions – INVASION” (MIS 5049543). OSW is a member of the Marine Biodiversity and Evolution research group, 2021-SGR-01271, funded by the Generalitat de Catalunya. We would like to thank two anonymous reviewers for their constructive suggestions. As the authors are not native English speakers, a Large Language Model (ChatGPT) was used to improve the use of English.

References

Aglieri, G., Quattrocchi, F., Mariani, S., Baillie, C., Spatafora, D. *et al.*, 2023. Fish eDNA detections in ports mirror fishing fleet activities and highlight the spread of non-indigenous species in the Mediterranean Sea. *Marine Pollution Bulletin*, 189, 114792.

Ahmed, D.A., Hudgins, E.J., Cuthbert, R.N., Kourantidou, M., Diagne, C. *et al.*, 2022. Managing biological invasions: the cost of inaction. *Biological Invasions*, 24 (7), 1927-1946.

Akaike, H., 1973. Information theory as an extension of the maximum likelihood principle. p. 267-281. In: *Pro 2nd International Symposium on Information Theory, Tsahkadzor, Armenia, USSR, 2-8 September 1971*. Akademiai Kiado, Budapest.

American Public Health Association (APHA), 1998. *Standard Methods for the Examination of Water and Wastewater* (20th ed.). American Public Health Association, Washington, DC, 1220 pp.

Antich, A., Palacín, C., Zarcero, J., Wangenstein, O.S., Turon, X., 2023. Metabarcoding reveals high-resolution biogeographical and metaphylogeographical patterns through marine barriers. *Journal of Biogeography*, 50 (3), 515-527.

Ardura, A., 2019. Species-specific markers for early detection of marine invertebrate invaders through eDNA methods: Gaps and priorities in GenBank as database example. *Journal for Nature Conservation*, 47, 51-57.

Azzurro, E., Tuset, V. M., Lombarte, A., Maynou, F., Simberloff, D. *et al.*, 2017. External morphology explains the success of biological invasions. *Ecology Letters*, 20 (9), 1169-1179.

Bailey, S., Brown, L., Campbell, M., Canning-Clode, J., Carlton, J. *et al.*, 2020. Trends in the detection of aquatic non-indigenous species across global marine, estuarine and freshwater ecosystems: A 50-year perspective. *Diversity and Distribution*, 26, 1780-1797.

Bakker, J., Wangenstein, O.S., Chapman, D.D., Boussarie, G., Buddo, D. *et al.*, 2017. Environmental DNA reveals tropical shark diversity in contrasting levels of anthropogenic impact. *Scientific Reports*, 7 (1), 1-9.

Bariche, M., Bilecenoglu, M., Kaya, M., Azzurro, E., 2013. Confirmed presence of the invasive alien species *Pterois miles* (Bennett, 1828) in the Mediterranean Sea: a case report. *Mediterranean Marine Science*, 14 (2), 292-294.

Batjakas, I.E., Evangelopoulos, A., Giannou, M., Pappou, S., Papanikola, E. *et al.*, 2023. Lionfish Diet Composition in Three Study Sites in the Aegean Sea (Southern Crete, and Kastellorizo and Nysiros Islands): An Invasive Generalist? *Fishes*, 8, 314.

Bessell, T.J., Appleyard, S.A., Stuart-Smith, R.D., Johnson, O.J., Ling, S.D. *et al.*, 2023. Using eDNA and SCUBA surveys for detection and monitoring of a threatened marine cryptic fish. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 33 (5), 431-442.

Bessey, C., Jarman, S.N., Berry, O., Olsen, Y.S., Bunce, M. *et al.*, 2020. Maximizing fish detection with eDNA metabarcoding. *Environmental DNA*, 2 (4), 493-504.

Blackburn, T.M., Pyšek, P., Bacher, S., Carlton, J.T., Duncan, R.P. *et al.*, 2011. A proposed unified framework for biological invasions. *Trends in ecology & evolution*, 26 (7), 333-339.

Borja, Á., Chust, G., Rodríguez, J.G., Bald, J., Belzunce-Segarra, M.J. *et al.*, 2016. ‘The past is the future of the present’: learning from long-time series of marine monitoring. *Science of the Total Environment*, 566, 698-711.

Borrell, Y.J., Miralles, L., Do Huu, H., Mohammed-Geba, K., Garcia-Vazquez, E., 2017. DNA in a bottle - Rapid metabarcoding survey for early alerts of invasive species in ports. *PLoS One*, 12 (9), e0183347.

Boudouresque, C.F., Blanfuné, A., Fernandez, C., Lejeune, C., Pérez, T. *et al.*, 2017. Marine biodiversity-warming vs. biological invasions and overfishing in the Mediterranean Sea: take care, ‘One Train can hide another’. *MOJ Ecology & Environmental Science*, 2 (4), 1-13.

- Boyer, F., Mercier, C., Bonin, A., Le Bras, Y., Taberlet, P. *et al.*, 2016. obitools: A unix-inspired software package for DNA metabarcoding. *Molecular ecology resources*, 16 (1), 176-182.
- Briski, E., Chan, F.T., Darling, J.A., Lauringson, V., MacIsaac, H.J. *et al.*, 2018. Beyond propagule pressure: importance of selection during the transport stage of biological invasions. *Frontiers in Ecology and the Environment*, 16 (6), 345-353.
- Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L. *et al.*, 2001. Introduction to distance sampling: estimating abundance of biological populations. *Oxford University Press*, New York, NY.
- Burnham, K.P., Anderson, D. R., 2002. *Model Selection and Multimodel Inference*. Springer, New York, 488 pp.
- Chatzimentor, A., Doxa, A., Katsanevakis, S., Mazaris, A.D., 2023. Are Mediterranean marine threatened species at high risk by climate change?. *Global Change Biology*, 29 (7), 1809-1821.
- Christidis, G., Batziakas, E., Peristeraki, P., Kosoglou, I., Tampakakis, K. *et al.*, 2022. The Impact of *Lagocephalus sceleratus* (Gmelin, 1789) on Small-Scale Fisheries in Crete: Preliminary Results. p. 357-361. In: *Proceedings of the Marine and Inland Waters Research Symposium, Porto Heli, Argolida, 16–19 September 2022*. HCMR, Athens.
- Coll, M., Piroddi, C., Steenbeek, J., Kaschner, K., Ben Rais Lasram, F. *et al.*, 2010. The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. *PLoS One*, 5 (8), e11842.
- Collins, R.A., Wangenstein, O.S., O’Gorman, E.J., Mariani, S., Sims, D.W. *et al.*, 2018. Persistence of environmental DNA in marine systems. *Communications Biology*, 1, 185.
- Collins, R.A., Bakker, J., Wangenstein, O.S., Soto, A.Z., Corrigan, L. *et al.*, 2019. Non-specific amplification compromises environmental DNA metabarcoding with COI. *Methods in Ecology and Evolution*, 10 (11), 1985-2001.
- Costello, K.E., Trottet, A., 2023. *Surveillance guidelines for invasive alien species in the marine environment*. Technical note prepared by IUCN for the European Commission, 70 pp.
- Cowart, D.A., Murphy, K.R., Cheng, C.H.C., 2018. Metagenomic sequencing of environmental DNA reveals marine faunal assemblages from the West Antarctic Peninsula. *Marine Genomics*, 37, 148-160.
- Cuthbert, R.N., Diagne, C., Hudgins, E.J., Turbelin, A., Ahmed, D.A. *et al.*, 2022. Biological invasion costs reveal insufficient proactive management worldwide. *Science of the Total Environment*, 819, 153404.
- Deiner, K., Bik, H.M., Mächler, E., Seymour, M., Lacoursière-Roussel, A. *et al.*, 2017. Environmental DNA metabarcoding: Transforming how we survey animal and plant communities. *Molecular Ecology*, 26 (21), 5872-5895.
- Dimitriadis, C., Sourbès, L., 2015. On the occurrence of the Lessepsian immigrants *Alepes djedaba*, *Lagocephalus sceleratus* and *Fistularia commersonii* in Zakynthos Island (SE Ionian Sea, Greece). p. 484-485. In: *New Mediterranean Biodiversity Records (July 2015)*, Tsiamis, K., Aydogan, Ö., Bailly, N., Balistreri, P., Bariche, M. *et al.*, *Mediterranean Marine Science*, 16 (2), 472-488.
- Doble, C.J., Hipperson, H., Salzburger, W., Horsburgh, G.J., Mwita, C. *et al.*, 2020. Testing the performance of environmental DNA metabarcoding for surveying highly diverse tropical fish communities: A case study from Lake Tanganyika. *Environmental DNA*, 2 (1), 24-41.
- Duarte, S., Vieira, P.E., Lavrador, A.S., Costa, F.O., 2021. Status and prospects of marine NIS detection and monitoring through (e) DNA metabarcoding. *Science of the Total Environment*, 751, 141729.
- Evangelopoulos, A., Nikolaou, A., Michailidis, N., Kampouris, T.E., Batjakas, I.E., 2020. Progress of the dispersal of the alien goatfish *Parupeneus forsskali* (Fourmanoir & Guézé, 1976) in the Mediterranean, with preliminary information on its diet composition in Cyprus. *BioInvasions Records*, 9 (2), 209-222.
- Evangelopoulos, A., Karampetsis, D., Christidis, A., Gubili, C., Sapounidis, A. *et al.*, 2024. Non-native fish species in the North Aegean Sea: a review of their distributions integrating unpublished fisheries data. *Frontiers in Marine Science*, 11, 1398037.
- Fediajevaite, J., Priestley, V., Arnold, R., Savolainen, V., 2021. Meta-analysis shows that environmental DNA outperforms traditional surveys, but warrants better reporting standards. *Ecology and Evolution*, 11 (9), 4803-4815.
- Ferreira, A.O., Azevedo, O.M., Barroso, C., Duarte, S., Egas, C. *et al.*, 2024. Multi-marker DNA metabarcoding for precise species identification in ichthyoplankton samples. *Scientific Reports*, 14 (1), 19772.
- Filatov, D.A., 2002. ProSeq: a software for preparation and evolutionary analysis of DNA sequence data sets. *Molecular Ecology Notes*, 2 (4), 621-624.
- Fonseca, V.G., Davison, P.I., Creach, V., Stone, D., Bass, D. *et al.*, 2023. The application of eDNA for monitoring aquatic non-indigenous species: practical and policy considerations. *Diversity*, 15 (5), 631.
- Fontes, J.T., Katoh, K., Pires, R., Soares, P., Costa, F.O., 2024. Benchmarking the discrimination power of commonly used markers and amplicons in marine fish (e) DNA (meta) barcoding. *Metabarcoding and Metagenomics*, 8, e128646.
- Friedlander, A.M., Parrish, J.D., 1998. Habitat characteristics affecting fish assemblages on a Hawaiian coral reef. *Journal of experimental marine biology and ecology*, 224 (1), 1-30.
- Froese, R., Pauly, D., 2023. FishBase. World Wide Web electronic publication, version (02/2014). <http://www.fishbase.org> (Accessed 26 October 2023).
- Frøslev, T.G., Kjølner, R., Bruun, H.H., Ejrnæs, R., Brunbjerg, A.K., *et al.*, 2017. Algorithm for post-clustering curation of DNA amplicon data yields reliable biodiversity estimates. *Nature Communications*, 8, 1188.
- Galanidi, M., Zenetos, A., 2022. Data-Driven Recommendations for Establishing Threshold Values for the NIS Trend Indicator in the Mediterranean Sea. *Diversity*, 14 (1), 57.
- Galil, B.S., Marchini, A., Occhipinti-Ambrogi, A., 2018. East is east and West is west? Management of marine bioinvasions in the Mediterranean Sea. *Estuarine, Coastal and Shelf Science*, 201, 7-16.
- Galil, B.S., Danovaro, R., Rothman, S.B.S., Gevili, R., Goren, M., 2019. Invasive biota in the deep-sea Mediterranean: an emerging issue in marine conservation and management. *Biological Invasions*, 21, 281-288.
- Garrabou, J., Gómez-Gras, D., Medrano, A., Cerrano, C., Pon-

- ti, M. *et al.*, 2022. Marine heatwaves drive recurrent mass mortalities in the Mediterranean Sea. *Global Change Biology*, 28 (19), 5708-5725.
- Giakoumi, S., Katsanevakis, S., Albano, P.G., Azzurro, E., Cardoso, A.C. *et al.*, 2019. Management priorities for marine invasive species. *Science of the total environment*, 688, 976-982.
- Govindarajan, A.F., McCartin, L., Adams, A., Allan, E., Belani, A. *et al.*, 2022. Improved biodiversity detection using a large-volume environmental DNA sampler with in situ filtration and implications for marine eDNA sampling strategies. *Deep Sea Research Part I: Oceanographic Research Papers*, 189, 103871.
- Harrison, J.B., Sunday, J.M., Rogers, S.M., 2019. Predicting the fate of eDNA in the environment and implications for studying biodiversity. *Proceedings of the Royal Society B*, 286 (1915), 20191409.
- Hartle-Mougiou, K., Gubili, C., Xanthopoulou, P., Kasapidis, P., Valiadi, M. *et al.*, 2024. Development of a quantitative colorimetric LAMP assay for fast and targeted molecular detection of the invasive lionfish *Pterois miles* from environmental DNA. *Frontiers in Marine Science*, 11, 1358793.
- Hastie, T.J., Tibshirani, R.J., 1990. Monographs on statistics and applied probability. *Generalized additive models*, 43, 205-208.
- Jerde, C.L., Mahon, A.R., Chadderton, W.L., Lodge, D.M., 2011. "Sight-unseen" detection of rare aquatic species using environmental DNA. *Conservation Letters*, 4 (2), 150-157.
- Jo, T., Murakami, H., Yamamoto, S., Masuda, R., Minamoto, T., 2019. Effect of water temperature and fish biomass on environmental DNA shedding, degradation, and size distribution. *Ecology and evolution*, 9 (3), 1135-1146.
- Karachle, P.K., Oikonomou, A., Pantazi, M., Stergiou, K. I., Zenetos, A., 2022. Can biological traits serve as predictors for fishes' introductions, establishment, and Interactions? The Mediterranean Sea as a case study. *Biology*, 11 (11), 1625.
- Katikou, P., Georgantelis, D., Sinouris, N., Petsi, A., Fotaras, T., 2009. First report on toxicity assessment of the Lessepsian migrant pufferfish *Lagocephalus sceleratus* (Gmelin, 1789) from European waters (Aegean Sea, Greece). *Toxicon*, 54 (1), 50-55.
- Katsanevakis, S., Tempera, F., Teixeira, H., 2016. Mapping the impact of alien species on marine ecosystems: the Mediterranean Sea case study. *Diversity and Distributions*, 22 (6), 694-707.
- Katsanevakis, S., Rilov, G., Edelist, D., 2018. Impacts of marine invasive alien species on European fisheries and aquaculture—plague or boon? *CIESM Monograph*, 50, 125-132.
- Katsanevakis, S., Weber, A., Pipitone, C., Leopold, M., Cronin, M. *et al.*, 2012a. Monitoring marine populations and communities: methods dealing with imperfect detectability. *Aquatic Biology*, 16 (1), 31-52.
- Katsanevakis, S., Bogucarskis, K., Gatto, F., Vandekerckhove, J., Deriu, I. *et al.*, 2012b. Building the European alien species information network (EASIN): a novel approach for the exploration of distributed alien species data. *BioInvasions Records*, 1 (4), 235-245.
- Katsanevakis, S., Coll, M., Piroddi, C., Steenbeek, J., Ben Rais Lasram, F. *et al.*, 2014. Invading the Mediterranean Sea: biodiversity patterns shaped by human activities. *Frontiers in Marine Science*, 1, 32.
- Katsanevakis, S., Coll, M., Frascchetti, S., Giakoumi, S., Goldsborough, D. *et al.*, 2020a. Twelve recommendations for advancing marine conservation in European and contiguous seas. *Frontiers in Marine Science*, 7, 879.
- Katsanevakis, S., Poursanidis, D., Hoffman, R., Rizgalla, J., Rothman, S.B.-S., *et al.*, 2020b. Unpublished Mediterranean records of marine alien and cryptogenic species. *BioInvasions Records*, 9 (2), 165-182.
- Katsanevakis, S., Olenin, S., Puntilla-Dodd, R., Rilov, G., Stæhr, P.A.U. *et al.*, 2023. Marine invasive alien species in Europe: 9 years after the IAS Regulation. *Frontiers in Marine Science*, 10, 1271755.
- Kelly, R.P., Port, J.A., Yamahara, K.M., Crowder, L.B., 2014. Using environmental DNA to census marine fishes in a large mesocosm. *PLoS One*, 9 (1), e86175.
- Kleitou, P., Hall-Spencer, J.M., Savva, I., Kletou, D., Hadjistylli, M. *et al.*, 2021. The case of lionfish (*Pterois miles*) in the Mediterranean Sea demonstrates limitations in EU legislation to address marine biological invasions. *Journal of Marine Science and Engineering*, 9 (3), 325.
- Korakaki, E., Legakis, A., Katsanevakis, S., Koulelis, P. P., Avramidou, E.V. *et al.*, 2021. Invasive alien species of Greece. p. 124-189. In: *Invasive alien species: observations and issues from around the world*. Pullaiah, T., Ielmini, M.R. (Eds). Wiley, Hoboken.
- Lehtiniemi, M., Ojaveer, H., David, M., Galil, B., Gollasch, S. *et al.*, 2015. Dose of truth—monitoring marine non-indigenous species to serve legislative requirements. *Marine Policy*, 54, 26-35.
- Mahé, F., Rognes, T., Quince, C., De Vargas, C., Dunthorn, M., 2015. Swarm v2: highly-scalable and high-resolution amplicon clustering. *PeerJ*, 3, e1420.
- Marchini, A., Galil, B.S., Occhipinti-Ambrogi, A., 2015. Recommendations on standardizing lists of marine alien species: Lessons from the Mediterranean Sea. *Marine Pollution Bulletin*, 101 (1), 267-273.
- Martinez Arbizu, P., 2020. pairwiseAdonis: Pairwise multilevel comparison using adonis. R package version 0.4. <https://github.com/pmartinezarbizu/pairwiseAdonis>
- Mauvisseau, Q., Harper, L.R., Sander, M., Hanner, R.H., Kleyer, H., *et al.*, 2022. The multiple states of environmental DNA and what is known about their persistence in aquatic environments. *Environmental Science & Technology*, 56 (9), 5322-5333.
- Mavruk, S., Bengil, F., Yeldan, H., Manasirli, M., Avsar, D., 2017. The trend of lessepsian fish populations with an emphasis on temperature variations in Iskenderun Bay, the Northeastern Mediterranean. *Fisheries Oceanography*, 26 (5), 542-554.
- McKnight, E.G.W., Shafer, A.B.A., Frost, P.C., 2024. Quantifying the effect of water quality on eDNA degradation using microcosm and bioassay experiments. *Environmental DNA*, 6, e530.
- Miliou, A., Loudaros, E., 2020. First record of *Cheilodipterus novemstriatus* in Rhodes Island (Dodecanese). p. 641-642. In: *New alien Mediterranean biodiversity records (October 2020)*. Ragkousis, M., Abdelali, N., Azzurro, E., Badreddine, A., Bariche, M. *et al.*, 2020, *Mediterranean Marine*

Science, 21 (3), 631-652.

- Miya, M., Gotoh, R.O., Sado, T., 2020. MiFish metabarcoding: a high-throughput approach for simultaneous detection of multiple fish species from environmental DNA and other samples. *Fisheries Science*, 86 (6), 939-970.
- Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J.Y. *et al.*, 2015. MiFish, a set of universal PCR primers for metabarcoding environmental DNA from fishes: detection of more than 230 subtropical marine species. *Royal Society open science*, 2 (7), 150088.
- Mohammed-Geba, K., Sheir, S.K., El-Aziz Hamed, E.A., Galal-Khallaf, A., 2020. Molecular and morphological signatures for extreme environmental adaptability of the invasive mussel *Brachidontes pharaonis* (Fischer, 1870). *Molecular and Cellular Probes*, 53, 101594.
- Morisette, J., Burgiel, S., Brantley, K., Daniel, W.M., Darling, J. *et al.*, 2021. Strategic considerations for invasive species managers in the utilization of environmental DNA (eDNA): steps for incorporating this powerful surveillance tool. *Management of biological invasions*, 12 (3), 747-775.
- Oksanen, J., Simpson, G.L., Blanchet, F.G., Kindt, R., Legendre, P. *et al.*, 2022. *Vegan: community ecology package. R package version 2.6-2*. <http://CRAN.R-project.org/package=vegan>
- Ostberg, C.O., Chase, D.M., 2022. Ontogeny of eDNA shedding during early development in Chinook Salmon (*Oncorhynchus tshawytscha*). *Environmental DNA*, 4 (2), 339-348.
- Packer, J.G., Meyerson, L.A., Richardson, D.M., Brundu, G., Allen, W.J. *et al.*, 2017. Global networks for invasion science: benefits, challenges and guidelines. *Biological Invasions*, 19, 1081-1096.
- Parrondo, M., Clusa, L., Mauvisseau, Q., Borrell, Y.J., 2018. Citizen warnings and post checkout molecular confirmations using eDNA as a combined strategy for updating invasive species distributions. *Journal for Nature Conservation*, 43, 95-103.
- Parsons, T. R., Maita, Y., Lalli, C. M., 1984. *A Manual of Chemical and Biological Methods for Seawater Analysis*. Pergamon Press, Oxford. 173 pp.
- Pastor, F., Valiente, J.A., Khodayar, S., 2020. A warming Mediterranean: 38 years of increasing sea surface temperature. *Remote sensing*, 12 (17), 2687.
- Peterson, B.G., Carl, P., Boudt K., Bennett, R., Ulrich, J. *et al.*, 2020. *PerformanceAnalytics: Econometric Tools for Performance and Risk Analysis. R package version 2.0.4*. <https://CRAN.R-project.org/package=PerformanceAnalytics>
- Ragkousis, M., Sini, M., Koukourouvli, N., Zenetos, A., Katsanevakis, S., 2023. Invading the Greek Seas: spatiotemporal patterns of marine impactful alien and cryptogenic species. *Diversity*, 15 (3), 353.
- Raitsos, D.E., Beaugrand, G., Georgopoulos, D., Zenetos, A., Pancucci-Papadopoulou, A.M. *et al.*, 2010. Global climate change amplifies the entry of tropical species into the eastern Mediterranean Sea. *Limnology and Oceanography*, 55 (4), 1478-1484.
- Rilov, G., Galil, B., 2009. Marine bioinvasions in the Mediterranean Sea—history, distribution and ecology. p. 549-575. In: *Biological invasions in marine ecosystems: ecological, management, and geographic perspectives*. Rilov, G., Crooks, J.A. (Eds.) Springer, Berlin.
- Rishan, S.T., Kline, R.J., Rahman, M.S., 2023. Applications of environmental DNA (eDNA) to detect subterranean and aquatic invasive species: a critical review on the challenges and limitations of eDNA metabarcoding. *Environmental Advances*, 12, 100370.
- Robinson, K.M., Prentice, C., Clemente-Carvalho, R., Hall, K., Monteith, Z.L. *et al.*, 2023. Paired environmental DNA and dive surveys provide distinct but complementary snapshots of marine biodiversity in a temperate fjord. *Environmental DNA*, 5 (3), 597-612.
- Rognes, T., Flouri, T., Nichols, B., Quince, C., Mahé, F., 2016. VSEARCH: a versatile open source tool for metagenomics. *PeerJ*, 4, e2584.
- Roy, H.E., Bacher, S., Essl, F., Adriaens, T., Aldridge, D.C. *et al.*, 2019. Developing a list of invasive alien species likely to threaten biodiversity and ecosystems in the European Union. *Global Change Biology*, 25 (3), 1032-1048.
- Ruppert, K.M., Kline, R.J., Rahman, M.S., 2019. Past, present, and future perspectives of environmental DNA (eDNA) metabarcoding: A systematic review in methods, monitoring, and applications of global eDNA. *Global Ecology and Conservation*, 17, e00547.
- Saito, T., Doi, H., 2021. Degradation modeling of water environmental DNA: Experiments on multiple DNA sources in pond and seawater. *Environmental DNA*, 3, 850-860.
- Sepulveda, A.J., Birch, J.M., Barnhart, E.P., Merkes, C.M., Yamahara, K.M. *et al.*, 2020a. Robotic environmental DNA bio-surveillance of freshwater health. *Scientific Reports*, 10 (1), 14389.
- Sepulveda, A.J., Nelson, N.M., Jerde, C.L., Luikart, G., 2020b. Are environmental DNA methods ready for aquatic invasive species management?. *Trends in ecology & evolution*, 35 (8), 668-678.
- Seymour, M., Durance, I., Cosby, B.J., Ransom-Jones, E., Deiner, K. *et al.*, 2018. Acidity promotes degradation of multi-species environmental DNA in lotic mesocosms. *Communications biology*, 1, 4.
- Stat, M., Huggett, M.J., Bernasconi, R., DiBattista, J.D., Berry, T.E. *et al.*, 2017. Ecosystem biomonitoring with eDNA: metabarcoding across the tree of life in a tropical marine environment. *Scientific Reports*, 7 (1), 12240.
- Strickler, K.M., Fremier, A.K., Goldberg, C.S., 2015. Quantifying effects of UV-B, temperature, and pH on eDNA degradation in aquatic microcosms. *Biological Conservation*, 183, 85-92.
- Taberlet, P., Coissac, E., Hajibabaei, M., Rieseberg, L.H., 2012. Environmental DNA. *Molecular ecology*, 21 (8), 1789-1793.
- Tamburini, M., Keppel, E., Marchini, A., Repetto, M.F., Ruiz, G.M. *et al.*, 2021. Monitoring non-indigenous species in port habitats: first application of a standardized North American protocol in the Mediterranean Sea. *Frontiers in Marine Science*, 8, 700730.
- Thanopoulou, Z., Sini, M., Vatikiotis, K., Katsoupis, C., Dimi-trakopoulos, P.G. *et al.*, 2018. How many fish? Comparison of two underwater visual sampling methods for monitoring fish communities. *PeerJ*, 6, e5066.
- Tsirintanis, K., Azzurro, E., Crocetta, F., Dimiza, M., Froggia, C. *et al.*, 2022. Bioinvasion impacts on biodiversity, ecosystem services, and human health in the Mediterranean

- Sea. *Aquatic Invasions*, 17 (3), 308-352.
- Tsiritanis, K., Sini, M., Ragkousis, M., Zenetos, A., Katsanevakis, S., 2023. Cumulative Negative Impacts of Invasive Alien Species on Marine Ecosystems of the Aegean Sea. *Biology*, 12 (7), 933.
- Tsuji, S., Shibata, N., 2021. Identifying spawning events in fish by observing a spike in environmental DNA concentration after spawning. *Environmental DNA*, 3 (1), 190-199.
- Valdivia-Carrillo, T., Rocha-Olivares, A., Reyes-Bonilla, H., Domínguez-Contreras, J.F., Munguia-Vega, A., 2021. Integrating eDNA metabarcoding and simultaneous underwater visual surveys to describe complex fish communities in a marine biodiversity hotspot. *Molecular Ecology Resources*, 21 (5), 1558-1574.
- Valentini, A., Taberlet, P., Miaud, C., Civade, R., Herder, J. *et al.*, 2016. Next-generation monitoring of aquatic biodiversity using environmental DNA metabarcoding. *Molecular Ecology*, 25 (4), 929-942.
- Venables, W.N., Ripley, B.D., 2002. Random and Mixed Effects. p. 271-300. In: *Modern Applied Statistics with S*. Springer, New York.
- West, K. M., Jones, T. R., Denis-Roy, L., Johnson, O. J., Clausius, E. *et al.*, 2024. Continual Day-Night eDNA Detectability Amidst Diel Reef Species Fluctuations on Diver Transects. *Environmental DNA*, 6 (5), e70018.
- Wood, S. N., Pya, N., Säfken, B., 2016. Smoothing parameter and model selection for general smooth models. *Journal of the American Statistical Association*, 111 (516), 1548-1563.
- Zaiko, A., Pochon, X., Garcia-Vazquez, E., Olenin, S., Wood, S. A., 2018. Advantages and limitations of environmental DNA/RNA tools for marine biosecurity: management and surveillance of non-indigenous species. *Frontiers in Marine Science*, 5, 322.
- Zangaro, F., Saccomanno, B., Tzafesta, E., Bozzeda, F., Specchia, V. *et al.*, 2021. Current limitations and future prospects of detection and biomonitoring of NIS in the Mediterranean Sea through environmental DNA. *NeoBiota*, 70, 151-165.
- Zenetos, A., Gofas, S., Verlaque, M., Cinar, M. E., Garcia Raso, J. E. *et al.*, 2010. Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Mediterranean Marine Science*, 11 (2), 381-493.
- Zenetos, A., Koutsogiannopoulos, D., Ovalis, P., Poursanidis, D. 2013. The role played by citizen scientists in monitoring marine alien species in Greece. *Cahiers de Biologie Marine*, 54 (3), 419-426.
- Zenetos, A., Arianoutsou, M., Bazos, I., Balopoulou, S., Corsini-Foka, M., *et al.*, 2015. ELNAIS : A collaborative network on Aquatic Alien Species in Hellas (Greece). *Management of Biological Invasions*, 6 (2), 185-196.
- Zenetos, A., Corsini-Foka, M., Crocetta, F., Gerovasileiou, V., Karachle, P. K. *et al.*, 2018. Deep cleaning of alien and cryptogenic species records in the Greek Seas (2018 update). *Management of Biological Invasions*, 9 (3), 209-226.
- Zenetos, A., Albano, P. G., Garcia, E. L., Stern, N., Tsiamis, K. *et al.*, 2022. Established non-indigenous species increased by 40% in 11 years in the Mediterranean Sea. *Mediterranean Marine Science*, 23 (1), 196-212.
- Zhang, S., Zhao, J., Yao, M., 2020a. A comprehensive and comparative evaluation of primers for metabarcoding eDNA from fish. *Methods in Ecology and Evolution*, 11 (12), 1609-1625.
- Zhang, Y., Pavlovska, M., Stoica, E., Prekrasna, I., Yang, J. *et al.*, 2020b. Holistic pelagic biodiversity monitoring of the Black Sea via eDNA metabarcoding approach: From bacteria to marine mammals. *Environment International*, 135, 105307.
- Ziegler, P. E., 2013. Influence of data quality and quantity from a multiyear tagging program on an integrated fish stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences*, 70 (7), 1031-1045.

Supplementary Data

The following supplementary information is available online for the article:

Table S1. Reference sequences for non-indigenous fish species (NIS) not included in the DUFA 12S rRNA gene database; 75 from retrieved from the National Center for Biotechnology Information (NCBI) nucleotide database (excluding unverified entries) and four barcoded for this study (Accession Number PQ638932-5).

Table S2. Comparison of Generalised Additive Models for fish non-indigenous species (NIS) detection through eDNA across sites of known occurrence in Greece, D: detected/not detected; TSS: total suspended solids; TDN: total dissolved nitrogen, $R^2(\text{adj})$: adjusted R^2 ; DE: deviance explained, df: degrees of freedom; AIC: Akaike information criterion; in bold the model with the lowest AIC value; s(): smooth functions of the predictor variables using thin plate regression splines.

Table S3. Environmental variables recorded at each sampling site in Greece, DO: dissolved oxygen; chl-a: chlorophyll-a; TSS: total suspended solids; warm period: May-October; cold period: November-April.

Table S4. PERMANOVA results of the geographic region of sampling (region), period, and type of substrate effect on the occurrence of non-indigenous fish species in Greek territorial waters using environmental DNA analysis; * statistically significant; df: degrees of freedom; SS: sum of squares.

Table S5. Best Generalised Additive Model results for the detection of marine non-indigenous species (NIS) fish across sites of known occurrence in Greece using 12S rRNA on seawater environmental DNA samples; * statistically significant.

Table S6. Reference list of non-indigenous species records in Greece.

Fig. S1: Number of reads during bioinformatic analysis stages; MOTU: molecular operational taxonomic units; NIS: non-indigenous species; *: at least 0.97 best-identity score; percentages refer to the reads of each category compared to the total paired-end reads.

Fig. S2: Non-indigenous fish species (NIS) communities detected in Greece using environmental DNA metabarcoding upon oc-

currence data. Blank circles represent sites where NIS fishes were not detected during sampling.

Fig. S3: Environmental parameters and depth recorded from February 2021 to March 2022; TSS: total suspended solids.

Fig. S4: Pearson correlations of environmental parameters recorded *in-situ* during seawater eDNA sampling expeditions from February 2021 to March 2022 in Greece, DO: dissolved oxygen; chl-a: chlorophyll-a; TSS: total suspended solids; red symbols indicate p-value levels: <0.001 ~ "****", 0.001-0.010 ~ "***", 0.010- 0.050 ~ "**", 0.050- 0.100 ~ ".", >0.100 ~ "".

Fig. S5: NMDS ordination of the marine-fish non-indigenous communities occurrence data in water samples using eDNA metabarcoding, for the effect of a) sampling region (North Aegean/Central Aegean/Ionian and Levantine Sea) and period (cold: November-April/warm: May-October) and b) substrate type (hard/soft). The stress of the final configurations was equal to 0.097.

Fig. S6: Occurrence map of non-indigenous fish species recorded through underwater visual surveys (SCUBA; yellow bullets); black pyramids indicate all sampling sites.

Fig. S7: Generalised Additive Model plots demonstrating the effect of temperature and total dissolved nitrogen in the detectability of marine non-indigenous fish species in Greece.