

## Mediterranean Marine Science

Vol 25, No 3 (2024)

Mediterranean Marine Science



**Comparative feeding habits of the invasive non-indigenous devil firefish *Pterois miles* and the indigenous scorpionfishes *Scorpaena porcus*, *Scorpaena scrofa*, and *Scorpaena notata* on the southwest coast of Türkiye, eastern Mediterranean**

*FIKRET ÖNDES, UĞUR ÖZDEN, VAHIT ALAN, ERHAN IRMAK, HARUN GÜÇLÜSOY*

doi: [10.12681/mms.38039](https://doi.org/10.12681/mms.38039)

### To cite this article:

ÖNDES, F., ÖZDEN, U., ALAN, V., IRMAK, E., & GÜÇLÜSOY, H. (2024). Comparative feeding habits of the invasive non-indigenous devil firefish *Pterois miles* and the indigenous scorpionfishes *Scorpaena porcus*, *Scorpaena scrofa*, and *Scorpaena notata* on the southwest coast of Türkiye, eastern Mediterranean. *Mediterranean Marine Science*, 25(3), 753–767. <https://doi.org/10.12681/mms.38039>

## Comparative feeding habits of the invasive non-indigenous devil firefish *Pterois miles* and the indigenous scorpionfishes *Scorpaena porcus*, *Scorpaena scrofa*, and *Scorpaena notata* on the southwest coast of Türkiye, eastern Mediterranean

Fikret ÖNDES<sup>1</sup>, Uğur ÖZDEN<sup>2</sup>, Vahit ALAN<sup>3,4</sup>, Erhan IRMAK<sup>2</sup> and Harun GÜÇLÜSOY<sup>4,5</sup>

<sup>1</sup>Department of Fisheries and Seafood Processing Technology, Faculty of Fisheries, İzmir Kâtip Çelebi University, İzmir, Türkiye

<sup>2</sup>Department of Basic Sciences, Faculty of Fisheries, İzmir Kâtip Çelebi University, İzmir, Türkiye

<sup>3</sup>Department of Fisheries, Graduate School of Natural and Applied Sciences, İzmir Kâtip Çelebi University, İzmir, Türkiye

<sup>4</sup>Mediterranean Conservation Society, İzmir, Türkiye

<sup>5</sup>Living Marine Resources Programme, Institute of Marine Sciences and Technology, Dokuz Eylül University, İzmir, Türkiye

Corresponding author: Fikret ÖNDES; [fikret.ondes@ikcu.edu.tr](mailto:fikret.ondes@ikcu.edu.tr)

Contributing Editor: Paraskevi K. KARACHLE

Received: 05 June 2024; Accepted: 06 November 2024; Published online: 03 December 2024

### Abstract

The feeding habits of the invasive alien devil firefish (*Pterois miles*) and the indigenous scorpaenids *Scorpaena porcus*, *Scorpaena scrofa*, and *Scorpaena notata* in the eastern Mediterranean Sea were investigated based on stomach content analysis. Furthermore, we examined whether these unfamiliar species feed upon similar prey. Specimens were captured by professional fishers using trammel nets on the southwestern coast of Türkiye in the winter and summer of 2021. A total of 608 stomachs (286 *P. miles*, 112 *S. porcus*, 136 *S. scrofa*, and 74 *S. notata*) were investigated. The results demonstrate that the diet of *P. miles* included a total of 45 taxa, with 33 identified at the species level (27 indigenous, 6 non-indigenous), while those of the three congeneric scorpaenids included a total of 46 taxa and 37 identified species (30 indigenous, 7 non-indigenous). For *P. miles*, the highest index of relative importance (%IRI) values were determined for fish in both seasons (83.44% in winter, 93.82% in summer), with the damselfish *Chromis chromis* being the most preferred species. *S. scrofa* fed mainly on fish in winter and cephalopods in summer, whereas the diets of *S. porcus* and *S. notata* included mainly crustaceans in both seasons. In winter, *P. miles* exhibited a higher percentage of empty stomachs, demonstrating a significant difference in the vacuity index in different sampling seasons, while other scorpaenids did not exhibit statistically significant differences. This study also provides the northernmost record of the non-indigenous crab species *Gonioinfradens giardi* and the second record of the squat lobster *Dactylonida curvimana* in Türkiye. It also provides new reports of rarely observed fish species for the eastern Mediterranean, including *Bellottia apoda* and *Odondebuena balearica*.

**Keywords:** stomach content; diet; marine invasion; scorpaenids; lionfish; *Gonioinfradens giardi*.

### Introduction

Studies on the feeding ecology of fish species are important for stock assessment, trophic models, and the provision of fundamental information to better understand the complex nature of aquatic ecosystems (Başçınar & Sağlam, 2009). Furthermore, studies related to the feeding habits of non-indigenous species (NIS), particularly invasive alien species (IAS), contribute to a better understanding of their impacts on food webs in invaded environments (David *et al.*, 2017). Notably, IAS not only cause ecological problems such as radical changes in food webs and the loss of biodiversity but they have also been reported to cause economic and even cultural losses (Bonanno, 2016; Galanidi *et al.*, 2023). The problem of biological invasions in aquatic ecosystems has been eval-

uated at the global scale and has been noted as a major threat in recent years –especially in the Mediterranean basin (e.g., Çinar *et al.*, 2021; Zenetos *et al.*, 2022).

The Convention on Biological Diversity (CBD), the European Union (EU), and various conventions such as the Barcelona Convention and state regulations in the Mediterranean region have long recognized the need to address the issue of IASs through international cooperation, legislation, and management strategies (Shine, 2007). For example, the EU Regulation on the Prevention and Management of the Introduction and Spread of Invasive Alien Species (No. 1143/2014) came into force in 2015 (Anonymous, 2014). This regulation provides a framework for preventing, minimizing, and mitigating the adverse impacts of IASs on biodiversity and ecosystems within the EU. From 1990 to 2017, the observed

cost of biological invasions was calculated at approximately US\$3.6 billion, while the projected cost was approximately \$27.3 billion for Mediterranean countries (Kourantidou *et al.*, 2021).

Türkiye's Mediterranean coastal areas are important, serving as a critical pathway for the introduction and spread of NIS and IAS across the Mediterranean region. This region is known for its incidence of invasions caused by the ongoing influx of species via the Suez Canal (known as Lessepsian migration), which has led to significant ecological and economic consequences for the area (Çinar *et al.*, 2021; Vagenas *et al.*, 2024).

Species of the genus *Pterois* can have a detrimental impact on the ecosystems they invade (Del Río *et al.*, 2023). For instance, the red lionfish *Pterois volitans* was imported to the United States via the ornamental fish trade, accidentally reached the sea, and began to invade the Caribbean and the Gulf of Mexico in 1985 (Trégarot *et al.*, 2015; South *et al.*, 2017). *P. volitans* rapidly spread to large areas, and its invasion resulted in a high consumption rate of the juveniles of commercially important and reef-associated species, devastating indigenous reef communities due to predation (Albins, 2013; Albins & Hixon, 2013). In the Mediterranean, the devil firefish, *Pterois miles* has been recognized as an IAS (Kletou *et al.*, 2016). The species originates from the Indian Ocean and the Red Sea (Schultz, 1986). Since the early 2020s, *P. miles* has spread to large areas in the eastern Mediterranean (Savva *et al.*, 2020; Kletou *et al.*, 2021; Ulman *et al.*, 2022). While the first record of the devil firefish in the Mediterranean (Haifa Bay) dates back to 1991 (Golani & Sonin, 1992), the first individual observed in Turkish waters was reported nearly a decade later (Turan *et al.*, 2014).

In recent years, *P. miles* has emerged as a target for small-scale fisheries in Türkiye. It is regularly sold at local fish markets and has become increasingly popular as a consumed seafood product (Öndes & Ünal, 2023). In October 2024, market prices fluctuated within a range of 10 to 11 euro/kg (AKD, unpublished data).

As with other IAS, *Pterois* spp. (*P. miles* and *P. volitans*) are represented by high fecundity, with these species spawning throughout the year in the Western North Atlantic, while the spawning of *P. miles* in the Mediterranean was observed in summer and autumn only (Morris, 2009; Morris & Whitfield, 2009; Savva *et al.*, 2020; Mouchlianitis *et al.*, 2022; Kondylatos *et al.*, 2024). The estimated annual fecundity of *Pterois* spp. per female has been reported as more than 2 million eggs in the Western North Atlantic (Morris, 2009; Morris & Whitfield, 2009), while the batch fecundity of *P. miles* has been calculated at between 3225 and 63149 oocytes in the Mediterranean (Mouchlianitis *et al.*, 2022). The invasion success of these species is driven by several key factors. Their high reproductive rate, early maturation, and effectiveness as predators –aided by successful feeding strategies– allow them to efficiently compete for space. Additionally, their ability to avoid predators through morphological traits (e.g., venomous spines), as well as their rapid growth rate, further enhance their capacity to establish and spread

(Morris *et al.*, 2009; Dimitriadis *et al.*, 2020; Kondylatos *et al.*, 2024). The presence of very few species that can prey on this species in the Mediterranean increases the concern of researchers and conservationists (Kletou *et al.*, 2019; Ulman *et al.*, 2021; Loya-Cancino *et al.*, 2023; Samourhani *et al.*, 2024). Hence, they pose a serious risk in a fragile ecosystem such as the Mediterranean ecosystem, where apex predators are decreasing while epidemic diseases, mass deaths, and anthropogenic effects are increasing (Ferretti *et al.*, 2008; Irmak & Engin, 2015; Garrabou *et al.*, 2019; Moullec *et al.*, 2019).

It is well known that scorpionfish are opportunistic carnivorous species that primarily feed on benthic invertebrates and fishes (Harmelin-Vivien *et al.*, 1989; Başçınar & Sağlam, 2009). They are also known for their characteristic ambush predation, where they move slowly, engage in “blowing” behavior to surprise their prey and use their large pectoral fins to corner their prey (Morris & Whitfield, 2012; Albins, 2013; Díaz-Ferguson & Hunter, 2019). Some weedy scorpaenids display camouflage and use this to both hunt their prey and protect against predators (Marshall *et al.*, 2019). Thus, it is obvious that their foraging behaviors can show variations depending on species availability.

Although numerous studies have focused on the diets of *P. miles* and *P. volitans* in the Western Atlantic and Caribbean (e.g., Morris & Akins, 2009; Dahl & Patterson, 2014; Eddy *et al.*, 2016; Harms-Tuohy *et al.*, 2016; Peake *et al.*, 2018; Sancho *et al.*, 2018; Acero *et al.*, 2019), the number of studies conducted in the Mediterranean remains limited (Zannaki *et al.*, 2019; Agostino *et al.*, 2020; Savva *et al.*, 2020; Batjakas *et al.*, 2023). As such, the impacts of this invasive species on the fragile Mediterranean ecosystem are not yet well understood (Kletou *et al.*, 2016; Poursanidis *et al.*, 2020; Phillips *et al.*, 2024).

Regarding the feeding habits of other scorpaenids, such as the black scorpionfish *Scorpaena porcus*, red scorpionfish *Scorpaena scrofa*, and small red scorpionfish *Scorpaena notata*, observations have been mainly documented from the western and central Mediterranean (Harmelin-Vivien *et al.*, 1989; Morte *et al.*, 2001; Rafrafi-Nouira *et al.*, 2016; Compaire *et al.*, 2018; Omri *et al.*, 2019), with no specific studies on the diets of these species in the eastern Mediterranean. Moreover, no studies have shown whether these species, distributed in the same or similar habitat and depth, feed on the same or different prey, or whether competition exists among them. To alleviate this deficiency to some extent, we aimed to determine and compare the diets of four scorpaenids in the eastern Mediterranean through the present study.

## Materials and Methods

Specimens were obtained during the ordinary activities of commercial fishers operating in the winter (January and February) and summer (July and August) of 2021 between Fethiye-Göcek and Gökova in the eastern Mediterranean (Fig. 1). These operations were performed by



**Fig. 1:** Map of the study area.

several small-scale fishing boats (less than 12 m in length) using trammel nets in rocky habitats in areas shallower than 100 m. The mesh sizes of the nets varied between 40 and 60 mm, and their lengths ranged between 500 and 2000 m. The collected specimens were immediately transferred to the İzmir Katip Çelebi University Fisheries Biology Laboratory using an isothermal box with ice to avoid digestion. In the laboratory, total weight (TW) (including stomach and intestines), total length (TL), sex, and stomach fullness (empty or full) were recorded for each specimen. For TW and TL determinations, an electronic balance (accuracy 0.001g) and caliper (accuracy 0.01 mm) were used, respectively. Sex was determined by gonadal examination based on macroscopic investigation (Avşar, 2016). Stomachs were taken and weighed, with food items being separated and counted in Petri dishes containing 70% ethanol (Rajasilta *et al.*, 2014). Prey items were evaluated under an Olympus SZ2-LGB stereo microscope, whereas larger ones were identified without a microscope. The weight of each prey item was measured and recorded. All prey items were identified to the lowest possible taxonomic level (Hureau & Monod, 1973; Whitehead *et al.*, 1986; Galil *et al.*, 2002; Bariche, 2012; Golani *et al.*, 2021).

Feeding intensity was evaluated with the use of the vacuity index (VI) of the sampled stomach, which corresponds to the percentage of empty stomachs (ES) with respect to all analyzed stomachs (TS) (Ouakka *et al.*, 2017).

$$VI = ES / TS \times 100$$

To understand the contribution of each prey type to the diets of scorpaenids, three simple methods –numerical, gravimetric, and frequency of occurrence– were used (Hyslop, 1980). The percentages of numerical abundance (%N) were calculated using the following equation:

$$\%N = (i/TNP) \times 100$$

$i$  = number of prey items, TNP = total number of prey items

Similarly, a gravimetric method was used, as follows:

$$\%W = (i/TWAP) \times 100$$

$i$  = weight of prey item, TWAP = total weight of all prey items

Percentage frequency of occurrence (%F) was calculated as:

$$\%F = (i/TNFS) \times 100$$

$i$  = number of stomachs containing a specific prey item, TNFS = total number of full stomachs

(Hyslop, 1980; Rafrafi-Nouira *et al.*, 2016).

The index of relative importance (IRI) identified by Pinkas *et al.* (1971), and modified by Hacunda (1981), was calculated using the following formula:

$$IRI = \%F \times (\%N + \%W)$$

This index is expressed as:

$$\%IRI = (IRI / \sum IRI) \times 100$$

This index was selected since it helps to avoid bias and presents a clearer picture of the relative importance of prey items in the diet.

Based on the mean TL, two size classes were determined for each species: small (TL lower than the mean value) and large (TL higher than the mean value). Sexual, seasonal, and size-related differences in the vacuity index (VI) and percentage frequency of occurrence (%F) were assessed for each species using the chi-square test ( $\chi^2$ ) (Sokal & Rohlf, 1981) and the R software (R Core Team, 2022).

## Results

In this study, a total of 608 individuals (286 *P. miles*, 112 *S. porcus*, 136 *S. scrofa*, and 74 *S. notata*) were collected. The TL of *P. miles* individuals varied from 18 to



37 cm (mean  $\pm$  SD = 26  $\pm$  4 cm). Similarly, the TL of *S. scrofa* ranged from 15 to 36 cm (mean  $\pm$  SD = 23  $\pm$  4 cm). On the other hand, *S. notata* specimens ranged from 6 to 21 cm TL (mean  $\pm$  SD = 16  $\pm$  3 cm), while *S. porcus* ranged from 11 to 24 cm TL (mean  $\pm$  SD = 17  $\pm$  3 cm).

For *P. miles*, *S. porcus*, *S. scrofa*, and *S. notata*, the VI (for the total number of each species, regardless of sex, season, and size class) was determined as 39, 58, 78, and 48%, respectively. The results indicate that VI of *P. miles* was significantly higher in winter than in summer (Table 1). Notably, there was no significant difference in the VI of other scorpaenids relating to the sampling season. On the other hand, except for *S. porcus*, the VI of scorpaenids did not show significant variations between sexes. In *S. porcus*, the VI of males was significantly higher than that of the females. Moreover, there was no significant difference in VI with regard to the size groups for any of the species (Table 1).

The overall results of the qualitative and quantitative analyses of prey ingested for each species are presented in Tables 2, 3, 4, and 5. The diet of *P. miles* was diverse when compared to those of other scorpaenids in the present study. Of the 45 total taxa identified from the stomachs, 33 were identified at the species level. While stomachs of *S. porcus* specimens included 21 taxa (14 identified at the species level), *S. notata* included 25 taxa (20 identified at the species level). The lowest diet variety was observed in *S. scrofa*, with 19 taxa (11 identified at the species level). Except for *S. scrofa*, the number of taxa in other scorpaenids was found to be higher in summer than in winter (Tables 2, 3, 4, and 5). Additionally, a


total of 12 species were found to be consumed by *P. miles* and at least one of the other three scorpaenids. These species were *Gonioinfradens giardi*, *Charybdis longicollis*, *Pilumnus* sp., *Galathea* sp., *Munida* sp., *Octopus vulgaris*, *Spicara* sp., *Serranus hepatus*, *Coris julis*, *Siganus luridus*, *Odondebuena balearica*, and *Bregmaceros nectabanus*.

Concerning the %F in *P. miles*, 84.29% of the full stomachs included Teleostei, followed by Crustacea with 36.64%, while Mollusca represented 2.09% and Annelida only 0.52%. Additionally, for *P. miles*, the %F of fish was higher in summer, while the %F of Crustacea was higher in winter (Fig. 2). The %F of Crustacea and Teleostei in *P. miles* showed significant differences depending on the sampling season ( $X^2$  test,  $P = 0.04$  for Crustacea,  $P = 0.01$  for Teleostei). Moreover, the stomachs of *S. porcus* and *S. scrofa* mainly included Crustacea (%F = 82.61 and 53.33, respectively), and for both species, the %F of Crustacea was higher in winter (Fig. 2). On the contrary, the %F of both Crustacea and Teleostei was higher in summer in *S. notata*, with this species not being observed to feed on Annelida and Mollusca throughout the study (Fig. 2). On the other hand, %F was very similar in both size groups of fish species—except for *S. notata*. The %F of main prey taxa (Annelida, Crustacea, Mollusca, and Teleostei) in the other three indigenous scorpaenids did not show significant differences by sampling season ( $X^2$  test,  $P > 0.05$ ). Moreover, the %F of main prey taxa in four species of the Scorpaenidae family did not show significant differences between sexes ( $X^2$  test,  $P > 0.05$ ). Similarly, the %F of main prey taxa did not show significant differences

**Table 1.** Summary of  $X^2$  test related to the sexual, seasonal and size-related differences in the vacuity index (VI) of *Pterois miles*, *Scorpaena porcus*, *Scorpaena scrofa*, and *Scorpaena notata* in winter and summer 2021 in the Mediterranean coasts of Türkiye.

Species	Factor	$X^2$	df *	P-value
Season				
<i>Pterois miles</i>		23.85	1	< 0.001
<i>Scorpaena porcus</i>		1.044	1	0.307
<i>Scorpaena scrofa</i>		0.773	1	0.379
<i>Scorpaena notata</i>		0.053	1	0.818
Sex				
<i>Pterois miles</i>		0.12	1	0.729
<i>Scorpaena porcus</i>		7.063	1	0.008
<i>Scorpaena scrofa</i>		0.001	1	0.978
<i>Scorpaena notata</i>		1.145	1	0.285
Size class				
<i>Pterois miles</i>		0.03	1	0.867
<i>Scorpaena porcus</i>		0.79	1	0.375
<i>Scorpaena scrofa</i>		1.43	1	0.232
<i>Scorpaena notata</i>		0.07	1	0.796

\*df = degrees of freedom.



**Fig. 2:** Percentage frequency of occurrence (%F) of A) *Pterois miles*, B) *Scorpaena porcus*, C) *Scorpaena scrofa*, and D) *Scorpaena notata* on the southwest coast of Türkiye in the winter and summer of 2021. Meanings of abbreviations: ANN = Annelida, CRU = Crustacea, MOL = Mollusca, PIS = Pisces.


in a size classes  $X^2$  test,  $P > 0.05$ )—except for the %F of Teleostei in *S. notata*. There was a significant difference in the %F of Teleostei among the size classes of *S. notata* ( $X^2$  test,  $P = 0.01$ ). Small specimens of the latter species preferred mainly Crustacea as prey (%F= 90), while the highest %F in large specimens was in Teleostei with 68%.

The %IRI results also demonstrated that *P. miles* fed primarily on Teleostei in both seasons, while *S. porcus* and *S. notata* preferred to consume mainly Crustacea in the winter and summer. On the other hand, the main diet of *S. scrofa* consisted of fish in the winter, shifting to cephalopods in the summer (Fig. 3). In particular, *P. miles* demonstrated high predation (%IRI=80.15) on *Chromis chromis* in the summer (Table 2). TL of *C. chromis* varied between 1.5 and 8.2 cm (mean  $\pm$  SD = 4.8  $\pm$  2.4 cm). The highest %IRI of *S. porcus* was observed for the NIS of crab *Goniinfradens giardi* (%IRI=42.09) in the winter and unidentified crabs (%IRI=53.26) in the summer (Table 3). The highest %IRI (45.51) of *S. scrofa* was for unidentified fish species in the winter, while the highest %IRI (61.88) in the summer was determined for the common octopus *Octopus vulgaris* (Table 4). The main prey of *S. notata* consisted of Penaeidae and Brachyuran crustaceans (Table 5). Additionally, the %N and %W values for each prey item are shown in Tables 2, 3, 4, and 5.

Regarding the size classes, %IRI results indicate that

marked differences were not detected between small and large individuals of *P. miles* and *S. porcus*. Unlike these species, small individuals of *S. notata* mainly fed on Crustacea, while the highest %IRI for large individuals was Teleostei. Moreover, individuals from the small size classes of *S. scrofa* and *S. porcus* did not feed on Polychaeta (Table 6).

This study also provided the northernmost record of the non-indigenous crab species *Goniinfradens giardi* on Türkiye's coasts. A total of 13 specimens of this species were obtained from the stomachs of *P. miles* and *S. porcus*. Additionally, a specimen of the spider lobster *Dactylonida curvimana* was found in the stomach of *S. scrofa*, with this observation constituting the second record of this species in Turkish waters. Furthermore, fish species rarely reported along the Mediterranean coasts of Türkiye were found in the stomachs of *S. scrofa* (*Bellottia apoda*), and in the diets of *P. miles* and *S. notata* (*Odondebuena balearica*) (Tables 2, 4, and 5). The results indicate that a total of 59 identified prey species were found in this study, among which 10 are NIS (*Macrophthalmus graeffei*, *Goniinfradens giardi*, *Charybdis helleri*, *Charybdis longicollis*, *Thalamita poissonii*, *Metapenaeopsis aegyptia*, *Siganus luridus*, *Cheilodipterus novemstriatus*, *Parupeneus forsskali*, and *Bregmaceros nectabanus*).



**Fig. 3:** The index of relative importance (%IRI) of A) *Pterois miles*, B) *Scorpaena porcus*, C) *Scorpaena scrofa*, and D) *Scorpaena notata* on the southwest coast of Türkiye in the winter and summer of 2021. Meanings of abbreviations: ANN = Annelida, CRU = Crustacea, MOL = Mollusca, PIS = Pisces.

**Table 2.** Overall results of the qualitative and quantitative analyses of prey ingested for *Pterois miles* in winter and summer 2021 in the southwest coast of Türkiye.

Preys of <i>P. miles</i>	Winter				Summer			
	%F	%N	%W	%IRI	%F	%N	%W	%IRI
POLYCHAETA	-	-	-	-	0.97	2.60	0.11	0.04
Unidentified Polychaeta	-	-	-	-	0.97	2.60	0.11	0.04
CRUSTACEA	44.32	41.4	16.07	16.27	30.10	55.41	16.07	6.01
<b>Isopoda</b>								
Unidentified Isopod	-	-	-	-	0.97	0.43	0.06	0.01
<b>Mysida</b>								
Unidentified Mysid	7.41	5.86	0.18	1.34	1.94	0.87	0.11	0.03
<b>Stomatopoda</b>								
Unidentified Stomatopod	1.23	0.39	0.44	0.03	0.97	1.30	0.91	0.03
<b>Decapoda</b>								
* <sup>a</sup> <i>Gonioinfradens giardi</i>	3.70	2.34	3.13	0.61	-	-	-	-
* <i>Charybdis helleri</i>	-	-	-	-	0.97	0.87	1.43	0.03
* <i>Charybdis longicollis</i>	1.23	0.78	0.32	0.04	-	-	-	-
<i>Pilumnus</i> sp.	1.23	0.39	0.19	0.02	-	-	-	-
<i>Paguristes</i> sp.	1.23	0.39	0.05	0.02	-	-	-	-

Continued

Table 2 continued

Preys of <i>P. miles</i>	Winter				Summer			
	%F	%N	%W	%IRI	%F	%N	%W	%IRI
<i>Galathea nexa</i>	1.23	0.39	0.16	0.02	0.97	1.30	0.12	0.02
<i>Galathea</i> sp.	7.41	2.34	1.08	0.76	7.77	9.52	1.87	1.22
<i>Munida rugosa</i>	1.23	0.39	0.21	0.02	-	-	-	-
<i>Munida</i> sp.	2.47	2.34	0.52	0.21	2.91	6.49	1.98	0.34
<i>Scyllarus arctus</i>	1.23	3.91	0.70	0.17	6.80	21.65	4.16	2.41
Unidentified Brachyura	4.94	1.56	0.95	0.37	3.88	2.60	0.93	0.19
Penaeidae	16.05	16.02	6.73	10.96	5.83	5.19	2.97	0.65
* <i>Metapenaeopsis aegyptia</i>	1.23	0.39	0.31	0.03	-	-	-	-
Unidentified Crustacea	11.11	3.91	1.10	1.67	11.65	5.19	1.53	1.08
MOLLUSCA	4.55	3.51	1.23	0.3	-	-	-	-
<b>Gastropoda</b>								
<i>Conus ventricosus</i>	1.23	0.39	0.01	0.01	-	-	-	-
<i>Conomurex persicus</i>	1.23	0.39	0.02	0.02	-	-	-	-
Unidentified Gastropod	3.70	1.56	0.03	0.18	-	-	-	-
<b>Cephalopoda</b>								
<i>Octopus vulgaris</i>	1.23	1.17	1.17	0.09	1.94	0.87	1.61	0.07
TELEOSTEI	76.14	55.06	82.68	83.44	91.26	93.07	104.08	93.82
Clupeidae	-	-	-	-	0.97	0.87	0.66	0.02
<i>Chromis chromis</i>	1.23	0.39	11.23	0.43	58.25	41.99	58.16	80.15
<i>Scorpaena maderensis</i>	11.11	19.14	5.07	8.07	1.94	0.87	10.52	0.30
<i>Scorpeana</i> sp.	4.94	2.73	0.43	0.47	4.85	13.42	4.58	1.20
<i>Spicara maena</i>	1.23	0.39	12.63	0.48	-	-	-	-
<i>Spicara smaris</i>	2.47	0.78	11.03	0.88	0.97	0.43	3.19	0.05
<i>Spicara</i> sp.	2.47	0.78	5.36	0.46	-	-	-	-
<i>Apogon imberbis</i>	-	-	-	-	2.91	1.73	2.57	0.17
<i>Serranus cabrilla</i>	1.23	0.39	5.90	0.23	-	-	-	-
<i>Serranus hepatus</i>	-	-	-	-	0.97	0.43	2.49	0.04
<i>Serranus</i> sp.	-	-	-	-	0.97	0.87	0.15	0.01
<i>Thalassoma pavo</i>	-	-	-	-	0.97	0.43	0.07	0.01
<i>Coris julis</i>	-	-	-	-	0.97	0.43	1.53	0.03
* <i>Siganus luridus</i>	-	-	-	-	0.97	0.43	2.39	0.04
Blennidae	1.23	0.39	0.65	0.04	0.97	1.73	1.71	0.05
Gobiidae	9.88	8.59	3.37	3.55	6.80	9.09	1.80	1.02
<i>Deltentosteus quadrimaculatus</i>	2.47	0.78	0.78	0.12	-	-	-	-
$\gamma$ <i>Odondebuena balearica</i>	-	-	-	-	0.97	0.43	0.54	0.01
<i>Gobius paganellus</i>	-	-	-	-	1.94	0.87	1.98	0.08
<i>Sparisoma cretense</i>	1.23	0.39	0.20	0.02	-	-	-	-
* <i>Bregmaceros nectabanus</i>	-	-	-	-	1.94	0.87	0.55	0.04
* <i>Parupeneus forsskali</i>	-	-	-	-	0.97	0.43	3.42	0.05
Unidentified Fish	49.38	20.31	26.03	68.69	30.10	17.75	7.77	10.55

\* non-indigenous species,  $\alpha$  northernmost record of species,  $\gamma$  rare species.



**Table 3.** Overall results of the qualitative and quantitative analyses of prey ingested for *Scorpaena porcus* in winter and summer 2021 in the southwest coast of Türkiye.

Preys of <i>S. porcus</i>	Winter				Summer			
	%F	%N	%W	%IRI	%F	%N	%W	%IRI
POLYCHAETA	4.55	3.57	0.98	0.69	-	-	-	-
Unidentified Polychaeta	4.55	3.57	0.98	0.69	-	-	-	-
CRUSTACEA	85.71	78.57	78.18	84.88	80	82	60.34	90
<b>Stomatopoda</b>								
<i>Rissoides desmaresti</i>	-	-	-	-	4.0	4	2.30	0.68
<i>Squilla mantis</i>	-	-	-	-	4.0	2	19.42	2.30
<i>Squilla</i> sp.	4.55	3.57	10.76	2.18	-	-	-	-
<b>Decapoda</b>								
* <i>Gonioinfradens giardi</i>	27.27	25.00	21.12	42.09	-	-	-	-
* <i>Thalamita poissonii</i>	4.55	7.14	14.78	3.33	20.0	20	13.54	18.00
<i>Xantho</i> sp.	4.55	3.57	0.49	0.62	-	-	-	-
Portunidae	-	-	-	-	4.0	2	1.20	0.34
<i>Galathea intermedia</i>	-	-	-	-	4.0	2	1.01	0.32
<i>Galathea squamifera</i>	4.55	3.57	4.77	1.27	-	-	-	-
<i>Galathea</i> sp.	4.55	3.57	1.50	0.77	-	-	-	-
Unidentified Brachyura	18.18	17.86	21.17	23.75	40.0	30	19.62	53.26
Penaeidae	-	-	-	-	24.0	20	3.09	14.87
Unidentified Crustacea	18.18	14.29	3.59	10.87	4.0	2	0.16	0.23
MOLLUSCA	-	-	-	-	8.0	4	0.09	0.88
<b>Bivalvia</b>								
Unidentified Bivalvia	-	-	-	-	8.0	4	0.09	0.88
TELEOSTEI/FISHES	23.81	17.86	20.83	14.42	24	14	39.58	9.13
* <i>Siganus luridus</i>	4.55	3.57	16.40	3.04	-	-	-	-
<i>Sardina pilchardus</i>	-	-	-	-	8.0	4	23.19	5.84
<i>Lesueurigobius friesii</i>	-	-	-	-	4.0	2	3.40	0.58
<i>Spicara</i> sp.	-	-	-	-	4.0	2	11.37	1.44
<i>Symphodus</i> sp.	-	-	-	-	4.0	2	1.40	0.37
Unidentified Fish	18.18	14.29	4.43	11.38	8.0	4	0.22	0.90

\* non-indigenous species,  $\alpha$  northernmost record of species.

## Discussion

The present study presents the first comparison of the feeding habits of four scorpionfish species from the eastern Mediterranean. *Pterois miles* feeds on more species than the other three scorpaenids, including members of the family Scorpaenidae, and it presents high predation on species such as *C. chromis* during certain periods. *P. miles* is thriving in the studied areas, thereby posing a great threat to the Mediterranean ecosystem (Savva *et al.*, 2020; Kleitou *et al.*, 2021).

Many studies have emphasized how destructive the effects of its congener (*Pterois volitans*) are in the Atlantic and Caribbean. Based on an experimental study, a significant decline in the recruitment of indigenous fish species (average of 79% over a 5-week duration) was observed due to predation by *P. volitans* (Albins & Hixon, 2008). This species poses a major threat to coral reefs

and herbivore species, feeding on more than 70 different species and causing devastating problems in food webs (Valdez-Moreno *et al.*, 2012; Albins & Hixon, 2013). Its high predation on juvenile snappers and groupers has caused marked reductions in the stocks of these economically important species in the Gulf of Mexico (Huth *et al.*, 2018). The results of our study suggest that *P. miles* poses a threat to the Mediterranean ecosystem and that if its population increases excessively, it has the potential to have an impact on the Mediterranean ecosystem similar to that of the *P. volitans* in the Atlantic. The combination of increasing impacts of climate change and the mismanagement of fisheries has led to serious changes in the Mediterranean ecosystem, resulting in declines in stocks of indigenous species in the eastern Mediterranean; for example, more than 90% of the catch composition of small-scale fishers' nets now consists of NISs (Öndes & Ünal, 2023). Similarly, a recent study conducted in the

**Table 4.** Overall results of the qualitative and quantitative analyses of prey ingested for *Scorpaena scrofa* in winter and summer 2021 in the southwest coast of Türkiye.

Preys of <i>S. scrofa</i>	Winter				Summer			
	%F	%N	%W	%IRI	%F	%N	%W	%IRI
POLYCHAETA	4.35	3.23	0.22	0.68	-	-	-	-
Unidentified Polychaeta	4.35	3.23	0.22	0.68	-	-	-	-
CRUSTACEA	60.87	51.64	32.73	38.34	28.57	37.5	6.7	13.44
<b>Mysida</b>								
Unidentified Mysid	4.35	3.23	0.16	0.66	-	-	-	-
<b>Decapoda</b>								
* <i>Charybdis longicollis</i>	4.35	3.23	8.47	2.29	-	-	-	-
<sup>β</sup> <i>Dactylonida curvimana</i>	4.35	3.23	9.16	2.43	-	-	-	-
<i>Munida</i> sp.	4.35	3.23	1.90	1.00	-	-	-	-
<i>Scyllarides latus</i>	4.35	3.23	0.60	0.75	-	-	-	-
Unidentified Brachyura	13.04	9.68	1.97	6.84	14.29	12.5	4.11	5.05
Penaeidae	8.70	6.45	7.95	5.64	14.29	25	2.59	8.39
Unidentified Crustacea	21.74	16.13	2.30	18.05	-	-	-	-
MOLLUSCA	8.70	6.46	27.63	6.67	28.57	25	76.72	61.88
<b>Gastropoda</b>								
Unidentified Gastropod	4.35	3.23	9.76	2.54	-	-	-	-
<b>Cephalopoda</b>								
<i>Octopus vulgaris</i>	4.35	3.23	17.87	4.13	28.57	25	76.72	61.88
TELEOSTEI/FISHES	47.83	41.96	39.65	54.98	42.86	37.5	16.58	24.68
<i>Engraulis encrasicolus</i>	-	-	-	-	14.29	12.5	14.55	8.23
* <i>Cheilodipterus novemstriatus</i>	4.35	3.23	6.70	1.94	-	-	-	-
<i>Coris julis</i>	4.35	3.23	17.87	4.13	-	-	-	-
<sup>γ</sup> <i>Bellottia apoda</i>	4.35	3.23	2.46	1.11	-	-	-	-
<i>Chlopsis bicolor</i>	4.35	3.23	0.71	0.77	-	-	-	-
<i>Conger conger</i>	4.35	3.23	0.96	0.82	-	-	-	-
Congeridae	4.35	3.23	0.34	0.70	-	-	-	-
Unidentified Fish	30.43	22.58	10.61	45.51	28.57	25	2.03	16.45

\* non-indigenous species, <sup>β</sup> second record of species, <sup>γ</sup> rare species.

eastern Mediterranean (Rhodes Island) highlighted the increased abundance, biomass, and occurrence frequency of IAS in gillnet and trammel net fisheries (Kondylatos *et al.*, 2023). Öndes & Ünal (2023) also declared that *P. miles* was the most commonly caught species in 36 mm trammel nets on the southern coasts of Türkiye, which provides clear evidence of the increased abundance of this species in the region. Notably, only a few species –including dusky grouper (*Epinephelus marginatus*), white grouper (*Epinephelus aeneus*), common octopus (*O. vulgaris*), and silver-cheeked toadfish *Lagocephalus sceleratus*– were reported as predators of *P. miles* (Ulman *et al.*, 2021) in this region. In recent years, this species has begun to be consumed by locals and has become the target species of some small-scale fishing communities. Supporting more selective fishing pressure on the species and expanding its market is of great importance in combating this IAS and minimizing the negative effects it may have on the populations of remaining native species

(Ünal *et al.*, 2022; Olguner *et al.*, 2024).

The present study demonstrated that the main prey groups of *P. miles* were Teleostei, followed by Crustacea. Mollusca and Annelida were rarely observed in the investigated stomachs. These findings are very similar to those of other studies conducted in the eastern Mediterranean (Zannaki *et al.*, 2019; Agostino *et al.*, 2020; Savva *et al.*, 2020; Batjakas *et al.*, 2023; Koilakos *et al.*, 2024). For example, a recently published study evaluated the diet of devil firefish in three sites of the Aegean Sea, Greece, and indicated that this species mainly preyed upon teleosts (4 to 83% numerical abundance) and decapods (12 to 95% numerical abundance) (Batjakas *et al.*, 2023). Earlier, Zannaki *et al.* (2019) reported that the diet of *P. miles* in Rhodes, north-eastern Aegean Sea, mainly consisted of Teleostei (78.5% in number, approximately 94.7% in biomass). Similarly, the percentage of prey index of *P. miles* included 87.39% fishes and 12.61% crustaceans in Cyprus, SE Mediterranean (Savva *et al.*, 2020). The

**Table 5.** Overall results of the qualitative and quantitative analyses of prey ingested for *Scorpaena notata* in winter and summer 2021 in the southwest coast of Türkiye.

Preys of <i>S. notata</i>	Winter				Summer			
	%F	%N	%W	%IRI	%F	%N	%W	%IRI
CRUSTACEA	66.67	80.02	23.51	61.32	83.33	76.82	25.69	76.21
<b>Mysida</b>								
Unidentified Mysid	6.67	34.29	5.30	12.27	-	-	-	-
<b>Stomatopoda</b>								
<i>Rissoides desmaresti</i>	6.67	2.86	6.70	2.96	-	-	-	-
<i>Squilla</i> sp.	6.67	2.86	1.13	1.23	-	-	-	-
<b>Decapoda</b>								
<i>Pilumnus hirtellus</i>	-	-	-	-	4.17	1.79	1.30	0.29
<i>Pilumnus</i> sp.	-	-	-	-	4.17	1.79	1.36	0.29
<i>Ilia nucleus</i>	-	-	-	-	4.17	1.79	2.89	0.44
* <i>Thalamita poissonii</i>	6.67	2.86	1.26	1.28	-	-	-	-
<i>Xantho poressa</i>	-	-	-	-	4.17	1.79	1.59	0.32
<i>Xantho</i> sp.	6.67	2.86	0.24	0.96	-	-	-	-
<i>Pisa tetraodon</i>	-	-	-	-	4.17	1.79	0.32	0.20
<i>Ethusa mascarone</i>	-	-	-	-	4.17	1.79	0.27	0.19
<i>Goneplax rhomboides</i>	-	-	-	-	4.17	1.79	2.07	0.36
* <i>Macrophthalmus graeffei</i>	-	-	-	-	4.17	1.79	2.07	0.36
<i>Galathea</i> sp.	6.67	2.86	0.65	1.09	4.17	1.79	1.39	0.30
Penaeidae	20.00	11.43	4.34	14.66	50.00	48.21	9.33	64.69
Unidentified Brachyura	26.67	17.14	3.79	25.95	25.00	12.50	3.10	8.77
Unidentified Crustacea	6.67	2.86	0.10	0.92	-	-	-	-
TELEOSTEI/FISHES	40	20.01	76.49	38.69	45.83	23.24	74.32	23.78
<i>Serranus hepatus</i>	6.67	2.86	30.50	10.34	4.17	1.79	3.57	0.50
<i>Sardina pilchardus</i>	6.67	2.86	24.34	8.43	4.17	1.79	14.17	1.49
<i>Engraulis encrasicolus</i>	6.67	2.86	12.89	4.88	16.67	8.93	39.12	18.00
<i>Lesueurigobius friesii</i>	-	-	-	-	8.33	5.36	7.00	2.31
<i>Lesueurigobius suerii</i>	-	-	-	-	4.17	1.79	2.07	0.36
* <i>Bregmaceros nectabanus</i>	-	-	-	-	4.17	1.79	8.36	0.95
<sup>γ</sup> <i>Odondebuena balearica</i>	6.67	2.86	3.15	1.86	-	-	-	-
Unidentified Fish	20.00	8.57	5.61	13.18	4.17	1.79	0.03	0.17

\* non-indigenous species,  $\gamma$  rare species.

**Table 6.** The size related differences in the percentage index of relative importance (%IRI) of *Pterois miles*, *Scorpaena porcus*, *Scorpaena scrofa*, and *Scorpaena notata* in winter and summer 2021 in the Mediterranean coasts of Türkiye.

Species	Size Class	Main Prey Taxa			
		Polychaeta	Crustacea	Mollusca	Teleostei
<i>P. miles</i>	Class 1 ( $\leq 26$ cm)	0.046	10.06	0.104	89.78
	Class 2 ( $> 26$ cm)	0	6.38	0.1	93.51
<i>S. porcus</i>	Class 1 ( $\leq 17$ cm)	0	89.03	1.17	9.79
	Class 2 ( $> 17$ cm)	2.42	86.62	0	10.96
<i>S. scrofa</i>	Class 1 ( $\leq 23$ cm)	0	23.03	27.02	49.95
	Class 2 ( $> 23$ cm)	1.58	36.54	15.74	46.14
<i>S. notata</i>	Class 1 ( $\leq 16$ cm)	0	88.39	0	11.60
	Class 2 ( $> 16$ cm)	0	44.81	0	55.19

latter authors also noted that the frequency of occurrence of accidental prey, such as fishing lines, plastics, and dry leaves of *Posidonia oceanica* was 11.11%, while that of parasites (Nematoda and Trematoda) was 29.63%.

*P. miles* prey heavily on certain key species within the Mediterranean ecosystem. For instance, our results indicate that *P. miles* fed mainly on damselfish *Chromis chromis* in summer. Likewise, Savva *et al.* (2020) highlighted that *C. chromis* was one of the most frequent prey items (F = 27.16%) of *P. miles* in Cyprus. Similar results were reported in another study conducted in Cyprus (Agostino *et al.*, 2020). In Rhodes, *C. chromis* was identified as one of the primary fish species consumed by *P. miles* (Zannaki *et al.*, 2019). Since this species is the most abundant fish in rocky areas of the Mediterranean, it has a key role in transferring carbon, nitrogen, and phosphorus directly from the pelagic system to the littoral, while also serving as an important predator of zooplankton and prey item of larger fishes and seabirds (Pinnegar, 2018). In line with this assumption –and based on the findings related to the predation of *P. miles*– the decline in this species' population may also be attributed to the devil firefish invasion. Nevertheless, feeding pressure from *P. miles* on this key species can show seasonal variations. For instance, our results highlighted that predation on the species reduced drastically in the winter season.

Moreover, in the present study, 12 of the species detected in the stomach of *P. miles* were also found in the diet of at least one of the other scorpaenids. For example, in the winter, *P. miles* placed relatively high pressure on Penaeidae, while in the summer, *S. notata* exhibited the highest level of predation on the latter. Nevertheless, *P. miles* not only compete for food with other scorpaenids but also feed on them (Table 2).

Seasonal variations in the diet of *S. scrofa* were observed in the present study. This species mainly preferred Mollusca in the summer, while Crustacea and Teleostei were common prey groups in the winter. Notably, this species demonstrated high predation on *O. vulgaris* in the summer. However, the level of predation was significantly lower in the winter. This may be related to the octopus' migration to shallow areas during the spring and summer months due to its reproductive cycle; thus, these areas also included small-sized octopus specimens during that period (Hernández-García *et al.*, 2002; Katsanevakis & Verriopoulos, 2004). The extant literature provides little information on the feeding biology of *S. scrofa*, and there are not many studies with which we can compare our findings. One available study performed in western Libya reported that the most important prey groups of *S. scrofa* were the bony fish and crustaceans, contributing by 48.02 and 44.19% respectively (Shahrani & Shakman, 2015). Similarly, Bradai & Bouain (1990) investigated the feeding habits of *S. scrofa* from the Gulf of Gabes, Tunisia, and reported that bony fish were the most abundant prey.

The present study noted that *S. notata* fed mainly on crustaceans and less commonly on fish. In particular, Penaeidae was represented with high %IRI values in the summer, while brachyuran crustaceans were preferred by a significant number of specimens in both seasons. Sim-

ilarly, a previous study performed in the seagrass beds near Marseilles, France noted that this species mostly fed on Caridae (Harmelin-Vivien *et al.*, 1989). Another study notified that the main prey items observed in the stomachs of *S. notata* in the western Mediterranean were decapod crustaceans (e.g., *Pisidia longimana*, *Pilumnus hirtellus*, *Processa mediterranea*, and *Alpheus glaber*) (Morte *et al.*, 2001). Additionally, Mokrane *et al.* (2015) reported very similar results for the Algerian coast, where the diet of *S. notata* mainly consisted of Brachyura and Natantia. Furthermore, the main prey groups of *S. notata* in the eastern Adriatic Sea were reported as Crustacea, Mollusca, Polychaeta, and Teleostei; among them, decapod crustaceans (e.g., of the genera *Alpheus*, *Galathea*, *Liocarcinus*, and *Pilumnus*) were dominant (Šantić *et al.*, 2021).

Our study showed that like *S. notata*, the highest %F and %IRI values for *S. porcus* were found for crustaceans, followed by fish species. The same situation was reported in the SW Mediterranean (Algeria and Tunisia) (Rafrafi-Nouira *et al.*, 2016; Omri *et al.*, 2019). It was reported that the diet composition of *S. porcus* showed variations depending on the season, size, and sex, as well as Palaemonidae, which was the most common prey group in the NE Atlantic (Compaire *et al.*, 2018). Unlike findings from the Mediterranean Sea, this species also had relatively high predation with %IRI = 35 for one of the economically important species in the Black Sea: red mullet (*Mullus barbatus*) (Başçınar & Sağlam, 2009).

In the present study, the performed transportation and handling procedures, as well as storage conditions and time between collection and dissection, aimed to minimize any further digestion in the collected specimens. Thus, the relatively high vacuity index values found in other species when compared to *P. miles* in both seasons depict the true status of our specimens in their environment at the moment of capture. Studies conducted on the feeding of *S. porcus* in various regions of the Mediterranean have reported that the highest VI varied between 42 and 57% (Morte *et al.*, 2001; Castriota *et al.*, 2012; Rafrafi-Nouira *et al.*, 2016; Omri *et al.*, 2019), while that for *S. notata* varied between 41 and 69% (Morte *et al.*, 2001; Castriota *et al.*, 2012). Notably, these values are similar to our findings. It should also be noted that the sampling method can influence VI because fish may also regurgitate their stomach contents due to the stress experienced during fishing operations (e.g., bottom trawling) (Compaire *et al.*, 2018). Moreover, our study indicated that the empty stomach ratio in *P. miles* was significantly higher in winter, and other scorpionfish species did not show significant seasonal differences. Similarly, Compaire *et al.* (2018) did not find significant seasonal variations in the VI of *S. porcus*, whereas Morte *et al.* (2001) noted significant differences in *S. porcus* and *S. notata*, with the maximum values for both species occurring in summer.

The current study also contributed to the enrichment of our knowledge on the distribution of some NISs in Türkiye through the recording of their northernmost and second records. For example, this study reports the northernmost record of the brachyuran crab *Gonioinfradens*



*giardi*, which is found to constitute a significant part of the winter diet of *P. miles* in Türkiye. Based on the use of both morphological and molecular analyses, Galil *et al.* (2018) reported that the species previously reported as *Gonioinfradens paucidentata* in the Mediterranean was incorrectly identified, noting that *G. giardi* is the correct name of the species. *G. paucidentata* was first reported from Kaş, Türkiye in 2009 (Karhan & Yokeş, 2012). Additionally, the spider lobster *D. curvimana* was first detected in Turkish waters in the Gulf of Antalya, Levantine Sea in 2007 (Irmak & Öndes, 2019). In the current study, the single *D. curvimana* individual obtained from the stomach of the *S. scrofa* in the winter of 2021 constitutes the second record for Turkish waters. This species has also been reported in Rhodes and Cyprus in the eastern Mediterranean (Corsini-Foka & Pancucci-Papadopoulou, 2012; Chartosia *et al.*, 2018). Furthermore, among the fish included in scorpionfish diets, there are some species rarely reported from Türkiye. For example, previous studies have reported the presence of the species *Bellottia apoda* (Gramitto *et al.*, 2011) and *Odondebuena balearica* (Stern *et al.*, 2019) in the SE Mediterranean. Notably, 17% of the species identified in the present study are alien to the Mediterranean.

In conclusion, this study provides important information on the diets of four scorpaenids in the eastern Mediterranean. *P. miles*, an IAS, exhibited higher dietary diversity when compared to other species. Notably, it even exhibited predation on other scorpaenids. Concerning diet overlap, a total of 12 species –including some economically important species such as *Octopus vulgaris* and *Siganus luridus*– were observed to be consumed by *P. miles* and at least one of the other three scorpaenids studied here. Furthermore, it is thought that competition may exist between the three native species (especially *S. notata*) and *P. miles* in terms of predation on the Penaeidae family. In future studies in the region, more detailed evaluations can be conducted by examining the diets of other predators that inhabit the same habitat as these four species. Additionally, applying trophic models should enhance the analysis.

## Acknowledgements

This study was partially funded by the Mediterranean Conservation Society. We are grateful to the commercial fishers who provided the specimens of scorpaenids. Special thanks to Dr. Tunca Olguner for the preparation of the base map. We would also like to thank the Ministry of Agriculture and Forestry of the Republic of Türkiye for research permission. We thank our anonymous reviewers for their comments and suggestions. This study is dedicated to the memory of the late fisher Mr. Mehmet Aytuğ.

## References

Acero, P.A., Bustos-Montes, D., Pabón Quintero, P., Polo-Silva, C.J., Muñoz, A.S., 2019. Feeding habits of *Pterois volitans*:

A real threat to Caribbean Coral Reef Biodiversity. Makowski, C., Finkl, C. (Eds). In: Impacts of invasive species on coastal environments. Coastal Research Library, Springer, Switzerland.

- Agostino, D.D., Jimenez, C., Reader, T., Hadjioannou, L., Heyworth, S. *et al.*, 2020. Behavioural traits and feeding ecology of Mediterranean lionfish and naiveté of native species to lionfish predation. *Marine Ecology Progress Series*, 638, 123-135.
- Albins, M.A., Hixon, M.A., 2008. Invasive Indo-Pacific lionfish *Pterois volitans* reduce recruitment of Atlantic coral-reef fishes. *Marine Ecology Progress Series*, 367, 233-238.
- Albins, M.A., 2013. Effects of invasive Pacific red lionfish *Pterois volitans* versus a native predator on Bahamian coral-reef fish communities. *Biological Invasions*, 15, 29-43.
- Albins, M.A., Hixon, M.A., 2013. Worst case scenario: potential long-term effects of invasive predatory lionfish (*Pterois volitans*) on Atlantic and Caribbean coral-reef communities. *Environmental Biology of Fishes*, 96, 1151-1157.
- Anonymous, 2014. Regulation (EU) No. 1143/2014 of the European Parliament and of the Council on the prevention and management of the introduction and spread of invasive alien species. Official Journal of the European Union L 317, 4 November 2014, 35-55.
- Avşar, D., 2016. Balıkçılık biyolojisi ve popülasyon dinamiği. Akademisyen Kitabevi, Ankara, 289 pp (In Turkish).
- Bariche, M. 2012. Field identification guide to the living marine resources of the Eastern and Southern Mediterranean. FAO Species Identification Guide for Fishery Purposes. Rome, 610 pp.
- Başçınar, N.S., Sağlam, H., 2009. Feeding habits of black scorpionfish *Scorpaena porcus*, in the South-Eastern Black Sea. *Turkish Journal of Fisheries and Aquatic Sciences*, 9 (1), 99-103.
- Batjakas, I.E., Evangelopoulos, A., Giannou, M., Pappou, S., Papanikola, E. *et al.*, 2023. Lionfish diet composition at three study sites in the Aegean Sea: an invasive generalist? *Fishes*, 8 (6), 314.
- Bonanno, G., 2016. Alien species: to remove or not to remove? That is the question. *Environmental Science & Policy*, 59, 67-73.
- Bradai, M.N., Bouain, A., 1990. Régime alimentaire de *Scorpaena porcus* et de *S. scrofa* (Teleostei, Scorpaenidae) du Golfe de Gabès, Tunisie. *Cybium*, 14 (3), 207-216.
- Castriota, L., Falautano, M., Finioia, M.G., Consoli, P., Peda, C. *et al.*, 2012. Trophic relationships among scorpaeniform fishes associated with gas platforms. *Helgoland Marine Research*, 66, 401-411.
- Chartosia, N., Anastasiadis, D., Bazairi, H., Crocetta, F., Deidun, A. *et al.*, 2018. New Mediterranean biodiversity records. *Mediterranean Marine Science*, 19 (2), 398-415.
- Compaire, J.C., Casademont, P., Cabrera, R., Gómez-Cama, C., Soriguer, M.C., 2018. Feeding of *Scorpaena porcus* (Scorpaenidae) in intertidal rock pools in the Gulf of Cadiz (NE Atlantic). *Journal of the Marine Biological Association of the United Kingdom*, 98 (4), 845-853.
- Corsini-Foka, M., Pancucci-Papadopoulou, M.A., 2012. Inventory of Crustacea Decapoda and Stomatopoda from Rhodes Island (Eastern Mediterranean Sea), with emphasis on rare and newly recorded species. *Journal of Biological Re-*



- search, 18, 359.
- Çinar, M.E., Bilecenoglu, M., Yokeş, M.B., Öztürk, B., Taşkin, E. *et al.*, 2021. Current status (as of end of 2020) of marine alien species in Turkey. *PLoS One*, 16 (5), e0251086.
- Dahl, K.A., Patterson III, W.F., 2014. Habitat-specific density and diet of rapidly expanding invasive red lionfish, *Pterois volitans*, populations in the northern Gulf of Mexico. *PLoS one*, 9 (8), e105852.
- David, P., Thebault, E., Anneville, O., Duyck, P. F., Chapuis, E. *et al.*, 2017. Impacts of invasive species on food webs: a review of empirical data. *Advances in Ecological Research*, 56, 1-60.
- Del Río, L., Navarro-Martínez, Z.M., Cobián-Rojas, D., Chevalier-Montegudo, P.P., Angulo-Valdes, J.A. *et al.*, 2023. Biology and ecology of the lionfish *Pterois volitans*/*Pterois miles* as invasive alien species: a review. *Peer J*, 11, e15728.
- Díaz-Ferguson, E.E., Hunter, M.E., 2019. Life history, genetics, range expansion and new frontiers of the lionfish (*Pterois volitans*, Perciformes: Pteroidae) in Latin America. *Regional Studies in Marine Science*, 31, 100793.
- Dimitriadis, C., Galanidi, M., Zenetos, A., Corsini-Foka, M., Giovos, I. *et al.*, 2020. Updating the occurrences of *Pterois miles* in the Mediterranean Sea, with considerations on thermal boundaries and future range expansion. *Mediterranean Marine Science*, 21 (1), 62-69.
- Eddy, C., Pitt, J., Morris Jr, J.A., Smith, S., Goodbody-Gringley, G. *et al.*, 2016. Diet of invasive lionfish (*Pterois volitans* and *P. miles*) in Bermuda. *Marine Ecology Progress Series*, 558, 193-206.
- Ferretti, F., Myers, R.A., Serena, F., Lotze, H.K., 2008. Loss of large predatory sharks from the Mediterranean Sea. *Conservation Biology*, 22 (4), 952-964.
- Galanidi, M., Aissi, M., Ali, M., Bakalem, A., Bariche, M. *et al.*, 2023. Validated inventories of non-indigenous species (NIS) for the Mediterranean Sea as tools for regional policy and patterns of NIS spread. *Diversity*, 15 (9), 962.
- Galil, B.S., Frogliani, C., Noel, P.Y., 2002. CIESM Atlas of Exotic Species in the Mediterranean. Volume 2: Crustaceans: decapods and stomatopods. F. Briand (Ed.), Monaco, 192 pp.
- Galil, B.S., Douek, J., Gevili, R., Goren, M., Yudkovsky, Y. *et al.*, 2018. The resurrection of *Charybdis* (*Gonioinfradens*) *giardi* (Nobili, 1905), newly recorded from the SE Mediterranean Sea. *Zootaxa*, 4370 (5), 580-590.
- Garrabou, J., Gómez-Gras, D., Ledoux, J.B., Linares, C., Bensoussan, N. *et al.*, 2019. Collaborative database to track mass mortality events in the Mediterranean Sea. *Frontiers in Marine Science*, 6, 707.
- Golani, D., Sonin, O., 1992. New records of the Red Sea fishes, *Pterois miles* (Scorpaenidae) and *Pteragogus pelycus* (Labridae) from the eastern Mediterranean Sea. *Japanese Journal of Ichthyology*, 39 (2), 167-169.
- Golani, D., Azzurro, E., Dulčić, J., Massutí, E., Orsi-Relini, L., 2021. Atlas of exotic fishes in the Mediterranean Sea. CIESM Publishers, Monaco, 365 pp.
- Gramitto, M.E., Deval, M.C., Saygu, I., 2011. First record of two deep-water fish, *Bellottia apoda* and *Symphurus ligulatus* in the Turkish Mediterranean Sea. *Cybio*, 35 (1), 75-76.
- Hacunda, J.S., 1981. Trophic relationships among demersal fishes in a coastal area of the gulf of Maine. *Fishery Bulletin*, 79 (4), 775.
- Harmelin-Vivien, M.L., Kaim-Malka, R.A., Ledoyer, M., Jacob-Abraham, S.S., 1989. Food partitioning among scorpaenid fishes in Mediterranean seagrass beds. *Journal of Fish Biology*, 34 (5), 715-734.
- Harms-Tuohy, C.A., Schizas, N.V., Appeldoorn, R.S., 2016. Use of DNA metabarcoding for stomach content analysis in the invasive lionfish *Pterois volitans* in Puerto Rico. *Marine Ecology Progress Series*, 558, 181-191.
- Hernández-García, V., Hernández-López, J.L., Castro-Hdez, J.J., 2002. On the reproduction of *Octopus vulgaris* off the coast of the Canary Islands. *Fisheries Research*, 57 (2), 197-203.
- Hureau, J.C., Monod, T. (Eds.), 1973. Check-list of the fishes of the north-eastern Atlantic and of the Mediterranean. UNESCO, Paris.
- Huth, W.L., McEvoy, D.M., Morgan, O.A., 2018. Controlling an invasive species through consumption: the case of lionfish as an impure public good. *Ecological Economics*, 149, 74-79.
- Hyslop, E.J., 1980. Stomach contents analysis—a review of methods and their application. *Journal of Fish Biology*, 17 (4), 411-429.
- Irmak, E., Engin, S., 2015. A newly established population of the Indian Ocean Twospot Cardinalfish, *Cheilodipterus novemstriatus* (Rüppell, 1838), in the Northern Levantine Sea (Osteichthyes: Apogonidae). *Zoology in the Middle East*, 61 (2), 186-188.
- Irmak, E., Öndes, F., 2019. Occurrence of the squat lobster *Munida curvimana* A. Milne Edwards & Bouvier, 1894 and Columbus crab *Planes minutus* (Linnaeus, 1758) in Turkey. *Journal of the Black Sea/Mediterranean Environment*, 25 (2), 150-159.
- Karhan, S.Ü., Yokeş, M.B., 2012. An earlier record of the Indo-Pacific swimming crab, *Gonioinfradens paucidentatus* (A. Milne-Edwards, 1861) (Decapoda, Brachyura, Portunidae) off the Mediterranean coast of Turkey. *Crustaceana*, 85 (1), 117-121.
- Katsanevakis, S., Verriopoulos, G., 2004. Den ecology of *Octopus vulgaris* Cuvier, 1797, on soft sediment: availability and types of shelter. *Scientia Marina*, 68 (1), 147-157.
- Kleitou, P., Savva, I., Kletou, D., Hall-Spencer, J.M., Antoniou, C. *et al.*, 2019. Invasive lionfish in the Mediterranean: Low public awareness yet high stakeholder concerns. *Marine Policy*, 104, 66-74.
- Kleitou, P., Hall-Spencer, J.M., Savva, I., Kletou, D., Hadjistylli, M. *et al.*, 2021. The case of lionfish (*Pterois miles*) in the Mediterranean Sea demonstrates limitations in EU legislation to address marine biological invasions. *Journal of Marine Science and Engineering*, 9 (3), 325.
- Kletou, D., Hall-Spencer, J.M., Kleitou, P., 2016. A lionfish (*Pterois miles*) invasion has begun in the Mediterranean Sea. *Marine Biodiversity Records*, 9, 1-7.
- Koilakos, S.M., Georgatis, I., Leonardos, I., 2024. Feeding Strategies and Biological Traits of the Lessepsian Migrant *Pterois miles* (Bennett, 1828) in the Messenian Gulf, SW Greece. *Fishes*, 9 (10), 380.
- Kondylatos, G., Vagenas, G., Kalaentzis, K., Mavrouleas, D., Conides, A. *et al.*, 2023. Exploring the Structure of Static Net Fisheries in a Highly Invaded Region: The Case of Rhodes Island (Eastern Mediterranean). *Sustainability*,

- 15, 14976.
- Kondylatos, G., Theocharis, A., Mandalakis, M., Avgoustinaki, M., Karagyaurova, T. *et al.*, 2024. The Devil Firefish *Pterois miles* (Bennett, 1828): Life History Traits of a Potential Fishing Resource in Rhodes (Eastern Mediterranean). *Hydrobiology*, 3 (1), 31-50.
- Kourantidou, M., Cuthbert, R.N., Haubrock, P.J., Novoa, A., Taylor N.G. *et al.*, 2021. Economic costs of invasive alien species in the Mediterranean basin. *NeoBiota*, 67, 427-458.
- Loya-Cancino, K.F., Ángeles-González, L.E., Yañez-Arenas, C., Ibarra-Cerdeña, C.N., Velázquez-Abunader, I. *et al.*, 2023. Predictions of current and potential global invasion risk in populations of lionfish (*Pterois volitans* and *Pterois miles*) under climate change scenarios. *Marine Biology*, 170 (3), 27.
- Marshall, N.J., Cortesi, F., de Busserolles, F., Siebeck, U.E., Cheney, K.L., 2019. Colours and colour vision in reef fishes: Past, present and future research directions. *Journal of Fish Biology*, 95 (1), 5-38.
- Mokrane, Z., Laribi, H., Touahria, N., Boufersaoui, S., Zerouali, F., 2015. Diet in relation to the tract structure and the chitinolytic activity in *Scorpaena notata* (Rafinesque, 1810) at the Algerian coast. *Cahiers de Biologie Marine*, 56, 245-252.
- Morris, J.A., 2009. The biology and ecology of invasive Indo-Pacific lionfish. Ph.D. Thesis. North Carolina State University, Raleigh, NC. 168pp.
- Morris, J.A., Akins, J.L., 2009. Feeding ecology of invasive lionfish (*Pterois volitans*) in the Bahamian archipelago. *Environmental Biology of Fishes*, 86, 389-398.
- Morris, J.A., Akins, J.L., Barse, A., Cerino, D., Freshwater, D.W. *et al.*, 2009. Biology and ecology of the invasive lionfishes, *Pterois miles* and *Pterois volitans*. *Proceedings of the 61st Gulf Caribbean Fisheries Institute*, 29, 409-414.
- Morris, J.A., Whitfield, P., 2012. Biology, ecology, control and management of the invasive Indo-Pacific lionfish: An updated integrated assessment. Monograph or Serial issue. Beaufort, NC: NOAA/National Ocean Service. Center for Coastal Fisheries and Habitat Research.
- Morte, S., Redon, M.J., Sanz-Brau, A., 2001. Diet of *Scorpaena porcus* and *Scorpaena notata* (Pisces: Scorpaenidae) in the western Mediterranean. *Cahiers de Biologie Marine*, 42 (4), 333-344.
- Mouchlianitis, F.A., Kalaitzi, G., Kleitou, P., Savva, I., Kletou, D. *et al.*, 2022. Reproductive dynamics of the invasive lionfish (*Pterois miles*) in the Eastern Mediterranean Sea. *Journal of Fish Biology*, 100 (2), 574-581.
- Moullec, F., Barrier, N., Drira, S., Guilhaumon, F., Marsaleix, P. *et al.*, 2019. An end-to-end model reveals losers and winners in a warming Mediterranean Sea. *Frontiers in Marine Science*, 6, 345.
- Olguner, M.T., Tosunoğlu, Z., Ünal, V., Kök, F. 2024. Impact of invasive edible species on Mediterranean small-scale fisheries. *Fish Forum* 2024, 19-13 February 2024, Antalya.
- Omri, N., Derbal, F., Kara, M.H., 2019. Diet of the black scorpionfish *Scorpaena porcus* (Scorpaenidae) of the gulf of Annaba, Algeria. *Cybium*, 43 (2), 179-186.
- Ouakka, K., Yahyaoui, A., Mesfioui, A., El Ayoubi, S., 2017. Stomach fullness index and condition factor of European sardine (*Sardina pilchardus*) in the south Moroccan Atlantic coast. *Aquaculture, Aquarium, Conservation & Legislation*, 10 (1), 56-63.
- Öndes, F., Ünal, V., 2023. The dominance of non-indigenous species in the catch composition of small-scale fisheries: A case study from the Kaş-Kekova Special Environmental Protection Area, Türkiye, Eastern Mediterranean. *Acta Ichthyologica et Piscatoria*, 53, 27-35.
- Peake, J., Bogdanoff, A.K., Layman, C.A., Castillo, B., Reale-Munroe, K. *et al.*, 2018. Feeding ecology of invasive lionfish (*Pterois volitans* and *Pterois miles*) in the temperate and tropical western Atlantic. *Biological Invasions*, 20, 2567-2597.
- Phillips, E.W., Bottacini, D., Schoonhoven, A.N., Kamstra, Y.J., De Waele, H. *et al.*, 2024. Limited effects of culling on the behavior of invasive lionfish (*Pterois miles*) in the Mediterranean. *Journal of Fish Biology*, 1-10.
- Pinkas, L., 1971. Food habits of albacore, bluefin tuna, and bonito in California waters. *Fishery Bulletin*, 152, 1-139.
- Pinnegar, J.K., 2018. Why the damselfish *Chromis chromis* is a key species in the Mediterranean rocky littoral—a quantitative perspective. *Journal of Fish Biology*, 92 (3), 851-872.
- Poursanidis, D., Kalogirou, S., Azzurro, E., Parravicini, V., Bariche, M. *et al.*, 2020. Habitat suitability, niche unfilling and the potential spread of *Pterois miles* in the Mediterranean Sea. *Marine Pollution Bulletin*, 154, 111054.
- Rafrafi-Nouira, S., El Kamel-Moutalibi, O., Boumaïza, M., Reynaud, C., Capapé, C., 2016. Food and feeding habits of black scorpionfish, *Scorpaena porcus* (Osteichthyes: Scorpaenidae) from the northern coast of Tunisia (Central Mediterranean). *Journal of Ichthyology*, 56, 107-123.
- Rajasilta, M., Hänninen, J., Vuorinen, I., 2014. Decreasing salinity improves the feeding conditions of the Baltic herring (*Clupea harengus membras*) during spring in the Bothnian Sea, northern Baltic. *ICES Journal of Marine Science*, 71 (5), 1148-1152.
- R Core Team, 2022. R: A language and environment for statistical computing. R foundation for statistical computing, Vienna, Austria.
- Samourdani, A., Ketsilis-Rinis, V., Koutsidi, M., Lazaris, A., Peristeraki, P. *et al.*, 2024. Ecology and behaviour of the invasive lionfish *Pterois miles* colonizing coastal areas of the central Mediterranean. *Estuarine, Coastal and Shelf Science*, 108796.
- Sancho, G., Kingsley-Smith, P.R., Morris, J.A., Toline, C.A., McDonough, V. *et al.*, 2018. Invasive lionfish (*Pterois volitans/miles*) feeding ecology in Biscayne National Park, Florida, USA. *Biological Invasions*, 20, 2343-2361.
- Šantić, M., Pallaoro, A., Rađa, B., 2021. Feeding habits of *Scorpaena notata* (Scorpaenidae) from eastern Adriatic Sea. *Cybium*, 45 (3), 217-224.
- Savva, I., Chartosia, N., Antoniou, C., Kleitou, P., Georgiou, A. *et al.*, 2020. They are here to stay: the biology and ecology of lionfish (*Pterois miles*) in the Mediterranean Sea. *Journal of Fish Biology*, 97 (1), 148-162.
- Schultz, E.T., 1986. *Pterois volitans* and *Pterois miles*: two valid species. *Copeia*, 686-690.
- Shahrani, S.E., Shakman, E., 2015. Food and feeding habits of *Scorpaena scrofa* (Linnaeus 1758) in the western part of Libya. *Libyan Journal of Marine Science*, 14, 5-15.
- Shine, C., 2007. Invasive species in an international context:

- IPPC, CBD, European Strategy on Invasive Alien Species and other legal instruments. *EPPO bulletin*, 37 (1), 103-113.
- Sokal, R.R., Rohlf F.J., 1981. Biometry. The principles and practices of statistics in biological research, 2nd ed. San Francisco: W.H. Freeman.
- South, J., Dick, J.T., McCard, M., Barrios-O'Neill, D., Anton, A., 2017. Predicting predatory impact of juvenile invasive lionfish (*Pterois volitans*) on a crustacean prey using functional response analysis: effects of temperature, habitat complexity and light regimes. *Environmental Biology of Fishes*, 100, 1155-1165.
- Stern, N., Badreddine, A., Bitar, G., Crocetta, F., Deidun, A. *et al.*, 2019. New Mediterranean Biodiversity Records (July 2019). *Mediterranean Marine Science*, 20 (2), 409-426.
- Trégarot, E., Fumaroli, M., Arqué, A., Hellio, C., Maréchal, J.P., 2015. First records of the red lionfish (*Pterois volitans*) in Martinique, French West Indies: monitoring invasion status through visual surveys. *Marine Biodiversity Records*, 8, e1.
- Turan, C., Ergüden, D., Gürlek, M., Yağlıoğlu, D., Uyan, A. *et al.*, 2014. First record of the Indo-Pacific lionfish *Pterois miles* (Bennett, 1828) (Osteichthyes: Scorpaenidae) for the Turkish marine waters. *Journal of Black Sea/Mediterranean Environment*, 20 (2), 158-163.
- Ulman, A., Harris, H.E., Doumpas, N., Deniz Akbora, H., Mabruk, A. *et al.*, 2021. Low pufferfish and lionfish predation in their native and invaded ranges suggests human control mechanisms may be necessary to control their Mediterranean abundances. *Frontiers in Marine Science*, 8, 670413.
- Ulman, A., Ali, F. Z., Harris, H. E., Adel, M., Mabruk, S.A.A. *et al.*, 2022. Lessons from the Western Atlantic lionfish invasion to inform management in the Mediterranean. *Frontiers in Marine Science*, 9, 865162.
- Ünal, V., Tıraşın, E.M., Tosunoğlu, Z., 2022. An ecosystem approach to fisheries management for small-scale fisheries in Gökova marine protected area, Turkey: challenges encountered during the transition process. Vasconcellos, M., Ünal, V. (Eds). Transition towards an ecosystem approach to fisheries in the Mediterranean Sea - Lessons learned through selected case studies. FAO Fisheries and Aquaculture Technical Paper No. 681. Rome, 117-136 pp.
- Vagenas, G., Karachle, P.K., Oikonomou, A., Stoumboudi, M.T., Zenetos, A., 2024. Decoding the spread of non-indigenous fishes in the Mediterranean Sea. *Scientific Reports*, 14, 6669.
- Valdez-Moreno, M., Quintal-Lizama, C., Gómez-Lozano, R., García-Rivas, M.D.C., 2012. Monitoring an alien invasion: DNA barcoding and the identification of lionfish and their prey on coral reefs of the Mexican Caribbean. *PloS one*, 7 (6), e36636.
- Whitehead, P.J.P., Bauchot, M.L., Hureau, J.C., Nielsen, J., Tortonese, E. (Eds), 1986. Fishes of the North-eastern Atlantic and the Mediterranean. UNESCO, Paris. Vols. I-III: 1473 pp.
- Zannaki, K., Corsini-Foka, M., Kampouris, T.E., Batjakas, I.E., 2019. First results on the diet of the invasive *Pterois miles* (Actinopterygii: Scorpaeniformes: Scorpaenidae) in the Hellenic waters. *Acta Ichthyologica et Piscatoria*, 49 (3), 311-317.
- Zenetos, A., Albano, P.G., Garcia, E.L., Stern, N., Tsiamis, K. *et al.*, 2022. Established non-indigenous species increased by 40% in 11 years in the Mediterranean Sea. *Mediterranean Marine Science*, 23, 196-212.