



Mediterranean Marine Science

Vol 26, No 2 (2025)

Special issue, 2025 Marine Animal Forest of the World (MAF WORLD)



Polyp number estimation through photogrammetry: a proof of concept using the example of Corallium rubrum

LUCREZIA DI FABIO, BRUNA GIORDANO, LAURINE MOIRAND, LORENZO BRAMANTI

doi: <u>10.12681/mms.38449</u>

To cite this article:

DI FABIO, L., GIORDANO, B., MOIRAND, L., & BRAMANTI, L. (2025). Polyp number estimation through photogrammetry: a proof of concept using the example of Corallium rubrum. *Mediterranean Marine Science*, *26*(2), 306–311. https://doi.org/10.12681/mms.38449

Contribution to the Special Issue: Marine Animal Forest of the World (MAF WORLD)

Polyp number estimation through photogrammetry: a proof of concept using the example of *Corallium rubrum*

Lucrezia Di FABIO^{1,3}, Bruna GIORDANO^{2,3}, Laurine MOIRAND³, and Lorenzo BRAMANTI³

¹Department of Life and Environmental Sciences (DiSVA), Polytechnic University of Marche, Ancona, Italy

²University of Cagliari, Department of Life and Environmental Sciences, Cagliari, Italy

³CNRS-Sorbonne Université, Observatoire Océanologique de Banyuls-sur-Mer, Laboratoire d'Ecogéochimie des Environnements Ben-

thiques (LECOB), Banyuls-sur-Mer, France

Corresponding author: Bruna GIORDANO; bruna.giordano@unica.it

Contributing Editor: Federica COSTANTINI

Received: 16 June 2024; Accepted: 04 October 2024; Published online: 21 March 2025

Abstract

The Mediterranean red coral (*Corallium rubrum*) is a gonochoric octocoral facing population decline due to overexploitation. Effective management and restocking of this species require accurate estimates of reproductive potential, such as fecundity, which depends on the number of polyps. In the present study, we propose a novel method for estimating the number of polyps in *C. rubrum* colonies based on polyp density and colony surface. The latter can be determined using existing structure from motion (SfM) techniques. Here, we describe a protocol developed to produce reliable estimates of polyp density, based on counting the number of polyps on branch sections with known surface areas. The mean polyp density, determined on colonies collected in the NW Mediterranean at 25-30 m depth, was 16.9 ± 4.5 polyps cm⁻². Hence, the total number of polyps in a colony can be calculated by multiplying this density with the colony's total surface area. Method accuracy was assessed by manually counting the exact number of polyps on small-sized colonies and comparing the result to estimates obtained from both our new method and an existing approach based on colony basal diameter. While the latter incurred a mean estimation error of 29%, our new approach yielded a considerably improved accuracy with a mean estimation error of 9%. Further research is needed to validate this new approach accors different red coral populations and potentially other coral species.

Keywords: Ocotcoral; Mediterranean Red Coral; Fecundity; Polyps; Structure from Motion.

Introduction

Mediterranean red coral (*Corallium rubrum*) is an octocoral belonging to the family of Corallidae, the socalled precious corals (Tsounis *et al.*, 2010). It is an iconic species endemic to the Mediterranean Sea and adjacent Atlantic rocky shores, whose bathymetric distribution spans depths ranging from 10 to 1000 m (Zibrowius *et al.*, 1984; Knittweis *et al.*, 2016). Red coral is a key component of Mediterranean marine ecosystems where it forms Marine Animal Forest (MAF, *sensu* Rossi *et al.*, 2017). Furthermore, it adds a three-dimensional complexity to the substrate and serves as both a refuge and nursery ground for several species (Maggioni & Bramanti, 2022).

Due to the use of its red calcareous skeleton in the manufacture of jewelry, this species has long been subjected to intensive exploitation (Tsounis *et al.*, 2013;

Santangelo & Bramanti, 2010). In recent years, the combined effects of marine heatwaves and overfishing have led to the endangerment of shallow populations (10-50 m depth; Santangelo *et al.*, 2007; Viladrich *et al.*, 2022; Garrabou *et al.*, 2022). Despite the implementation of various management and conservation measures, the species' slow growth rate and reproductive strategy require more active interventions such as local restocking, particularly for populations that have experienced significant local depletion.

Effective restoration and restocking plans depend on a comprehensive understanding of the life-history characteristics of the target species as well as knowledge of the reproductive potential. For modular organisms, such as corals, the reproductive potential is intrinsically linked to the number of reproductive modules within colonies, i.e., the polyps (Sakai, 1998). Therefore, methods capable

of delivering accurate polyp counts are needed to obtain reliable estimates of reproductive potential (Beiring & Lasker, 2000). In C. rubrum, polyps are not evenly distributed along a colony but exhibit higher densities in primary branches and lower densities in the basal section (Benedetti et al., 2020). For some time, a common approach to estimate the total number of polyps in C. rubrum was based on the relationship between colony basal diameter and the number of polyps (Santangelo et al., 2007; Bramanti et al., 2009; Bramanti et al., 2015). This method may not accurately estimate the total number of polyps as it relies on an indirect relationship linking basal diameter to colony size. A better approach would be to use colony surface area as a proxy for the number of polyps, as both are directly correlated. Moreover, the previous methods do not account for the sex of the colonies, neglecting any differences in polyp distributions between males and females.

Structure from motion (SfM) is a photogrammetric technique that uses 2D images to calculate accurate dimensional data allowing 3D reconstructions of objects (Aston et al., 2022). It has emerged as a promising tool that allows the precise determination of coral surface areas, both in field and aquarium settings, without having to destroy the coral (Million et al., 2021; Lange & Perry, 2020; Conley & Hollander, 2021). Other methodologies such as paraffin dipping with double wax (Stimson & Kenzie, 1991) or single wax (Veal et al., 2010) typically involve the removal of live tissue, which is a destructive process that can produce irreversible damage or alterations to the coral skeleton or even lead to the death of the entire organism. In comparison, SfM offers several advantages, particularly due to its non-invasive nature. By capturing multiple overlapping images from different angles, the photogrammetric technique enables the creation of detailed 3D reconstructions of coral colonies. Moreover, it provides a fast, easy, and inexpensive method (Lange & Perry, 2020) for obtaining precise measurements of coral surface area across a wide range of morphologies (Ferrari et al., 2017).

In the present paper, we test a non-destructive method for estimating the number of polyps in *C. rubrum* colonies. The approach uses photogrammetric measurements of colony surface area and manual counts of the number of polys in a representative area. By multiplying the resulting polyp density with the surface area, the total number of polyps in a colony can be estimated. In combination with knowledge of their fecundity (number of larvae released per polyp) this yields precise estimates of their reproductive potential. Due to its non-destructive nature, this technique could be applied at the population level, providing a fundamental parameter for population and meta-population dynamics studies (Bramanti *et al.*, 2009; Guizien & Bramanti, 2014), which can in turn inform conservation and restoration programs.

Material and Methods

Experimental design

C. rubrum colonies were collected by scuba diving in two locations (Banyuls-sur-Mer, France; Cap de Creus, Spain) from depths between 25 and 35 m during the summers of 2021 and 2022. For all individuals, sex was determined both morphologically (following Santangelo *et al.*, 2003) and genetically (following Pratlong *et al.*, 2017). We selected 28 colonies of both sexes with morphologies that included several branching orders to test for differences in polyp density between 1) sexes and 2) branching orders.

3-D coral reconstruction and surface measurements

After sample collection, each live colony was placed at the center of a black circular tank filled with seawater and screwed onto a cubic support with a visible size scale. Photographs of each coral colony were taken under uniform lightening from two angles (90° and 45°) using an Olympus Tough TG-6, maintaining roughly 80% overlap between 2 consecutive images. Approximately 110-150 photos were necessary to construct an accurate 3D model of a colony using Agisoft Metashape software (Fig. 1A). This 3D model generation involves the following steps: 1) Alignment: align pictures to determine camera positions and orientations; 2) Point cloud model: generate a sparse point cloud model to capture the basic structure of the scene; 3) Dense cloud model: produce a dense point cloud model with more detailed geometry; 4) Mesh model: reconstruct a polygonal mesh representing the object; 5) Textured model: map textures onto the mesh to enhance visual quality. Then, we used Meshlab v2016.12, an open-source software for processing and editing 3D meshes, to calculate the surface areas of the 3D models. Models were cleaned using the "Select Vertices" tool in Meshlab to remove the background, non-coral elements (e.g., PVC support), and any portions of coral that appeared dead or damaged. Gaps that resulted from the removal of dead portions or overgrowth were left uncorrected to avoid adding to the surface area. Finally, the surface area was determined using Filter > Quality Measures and Computations > Computer Geometric Measures (Fig. 1B).

Polyp Density Estimation

Polyp density was determined by randomly selecting between 2 and 4 fragments from each of the 28 coral colonies (106 fragments in total) and measuring their surface area with MeshLab, as previously described. Then, we counted the number of polyps on the selected branches



Fig. 1: The figure shows the steps to estimate the polyps' density. A) Screenshot of the 3D model reconstruction through Agisoft Metashape. It shows how the software aligns pictures to determine camera positions and orientations. B) Screenshot of a 3D model imported in MeshLab, showing a refined model of the actual living surface of the coral. C) View of a coral branch with closed polyps at the stereomicroscope (scale bar 1 mm). D) Macro of a coral branch with open polyps (scale bar 1 mm).

using a stereomicroscope (Fig. 1C, D). Polyp density then simply is the number of counted polyps divided by the measured surface area. Polyp counts for each branch were performed in triplicate by two different observers.

According to Benedetti *et al.* (2020), polyp density decreases with branch order, with densities being higher on apical branches. For this reason, we considered three orders of branching: 1st to 3rd, counting from the apex toward the base. To test for statistically significant differences in polyp density between different branch orders, we used a one-way ANOVA (R "stats" package), using the branching level (1st to 3rd) as fixed factor. When a significant effect of branching was found (p < 0.05), pairwise post-hoc tests were run (Tukey test with Bonferroni correction). Moreover, we performed independent t-tests to test for differences in polyp density between male and female colonies and to check for the existence of an observer effect in the polyp counts.

The weighted average polyp density across the three branching orders was computed using the proportions of surfaces areas occupied by the three branch types with respect to the total colony surface area (all determined using MeshLab, see above) for 17 colonies out of a total of 28. These proportions were used as weights to calculate the mean density:

$$N_{polyps} = d_{tot} x S_{tot} =$$

$$S_{1} x d_{1} + S_{2} x d_{2} + S_{3} x d_{3} =$$

$$S_{tot} (w_{1} x d_{1} + w_{2} x d_{2} + w_{3} x d_{3})$$

$$d_{tot} = (w_{1} x d_{1} + w_{2} x d_{2} + w_{3} x d_{3})$$

where d_1 , d_2 , and d_3 are the respective densities of the first, second, and third branching order while d_{iot} is the overall density. w_1 , w_2 , w_3 , are the weights calculated based on the proportion of the are surface area of branches in each order with respect to the total surface area S_{iot} . $S_{S1/2/3}$ are the respective surface areas of the 1st/2nd/3rd branch order.

Finally, the above approach was validated by manually counting the total number of polyps (Npolyps in the above equations) in a total of 6 colonies and comparing the result to the estimates obtained from the weighted model. For comparison, we also estimated Npolyps using the Santangelo *et al.* (2003) method which is based on the colony basal diameter, and calculated the mean error between the estimated (*est*) and measured (*obs*) values of Npolyps for both methods as $(N_{obs}-N_{est})/N_{obs}$. This validation was used to test the robustness of our methodology.

Results

Mean polyp density was 20.0 ± 4.30 , 14.4 ± 5.07 , and 11 ± 3.97 polyps cm⁻² (mean \pm SD) for the 1st, 2nd, and 3rd branching order, respectively. ANOVA results showed that these differences are statistically significant (Fig. 2; Table 1, F = 41.45; p = 1.94e⁻¹⁴). The post-hoc Tukey test highlighted significant differences between each pair of branches (Table 1).

Results of the t-tests showed that polyp density did not differ significantly between male and female colonies (Fig. 2), with 17.1 ± 6.0 and 16.4 ± 4.6 polyps cm⁻², respectively (t = 0.703, df = 53.4, p = 0.48). Both human observers obtained similar polyp density estimates (21.0 \pm 3.5 and 21.5 \pm 4.6 polyps cm⁻², respectively), with no significant differences detected by the t-test (t = 1.61, df = 61.2, p = 0.112).

The 1^{st} , 2^{nd} , and 3^{rd} order branches represent on average 54%, 30%, and 16% of the total surface area of a red coral colony, respectively. This resulted in a mean overall density of:

$$(w_1 x d_1 + w_2 x d_2 + w_3 x d_2) = 16.9 \pm 4.5$$
 polyps cm⁻²

The mean error was 9% for our new method, while the Santangelo *et al.* (2003) approach produced estimates with a mean error of 29%. This indicates a systematic underestimation by both methods, although the error is considerably improved with the new approach.

Discussion

In the present study, we tested a novel method for estimating the number of polyps in *C. rubrum* colonies based on generating 3D models of the colonies from hundreds of photographs to accurately determine the colony's surface area.

To be able to estimate the total number of polyps, the overall polyp density must be known, which we deter-



Fig. 2: Boxplot showing the mean polyps' density per branching order and sex. M= males, F=females.

mined through manual counts, yielding a mean overall polyp density across all samples of 16.9 ± 4.5 polyps cm⁻². Since Benedetti et al. (2020) found that polyp density varied with branch order, we determined the polyp density for the top three branches (1st to 3rd counted from the apex toward the base), which yielded statistically significant differences with densities decreasing with increasing branch order (i.e., 1^{st} order > 2^{nd} order > 3^{rd} order). While this trend is in agreement with observations by Bendetti et al. (2020), the absolute values differ. For the first and third order branches, our approach yielded polyp densities of 20.0 \pm 4.30 and 11 \pm 3.97 polyps cm^-2, respectively, while the corresponding values of Benedetti et al. (2020) are 16.1 \pm 1.7 and 8.0 \pm 1.2 polyps cm⁻², which are likely underestimations due to the approximation of the branch shape to a truncated cone. These polyp densities were used to calculate the weighted average stated above. In contrast, no significant sex-specific differences in polyp densities were found, suggesting that polyp density estimates can be performed on randomly selected C. rubrum colonies without accounting for their sex.

Table 1. (A) Results of the one-way ANOVA test comparing the polyps' density between branching order. df = degrees of freedom; SS = Sums of squares value; MS = mean square value; F = F ratio; * indicate p < .05; ** indicate p < .01; *** p < 0.001. (B) Tukey post hoc pairwise comparisons. Diff = Difference in density; lwr and upr = lower and upper bounds of 95% confidence interval; p adj = p adjusted for multiple comparisons.

A-One-way ANOVA	df	SS	MS	F value	p (> F)
branching	2	1646	823.1	41.45	1.94e-14 ***
Residuals	121	2403	19.9		
B-Tukey post-hoc		diff	lwr	lwr	p adj
2nd-1st		5.616875	-7.873213	-3.3605371	0.0000001
3rd-1st		-9.020000	-11.604411	-6.4355888	0.0000000
3rd-2nd		-3.403125	-6.331580	-0.4746696	0.0183324

When comparing our non-destructive method with an approach based on live tissue samples that calculates polyp densities from colony basal diameter (Santangelo et al., 2007; Cánovas-Molina et al., 2009; Bramanti et al., 2015), results showed that the new method performs better with considerably reduced errors of 9% vs 29%. The higher accuracy of the new approach is likely due to the fact that we directly measure the surface area (using photogrammetry and 3D reconstructions) instead of having to infer it from the basal diameter. Nevertheless, the colony basal diameter can be determined fairly quickly which renders this approach a viable option for the analysis of extensive datasets, such as those generated by ROV surveys (Cau et al., 2016; Carugati et al., 2022), for which our more time-consuming photogrammetric method may be impractical. Our method seems more suited to punctual research when more accurate estimations of polyp numbers are needed.

In C. rubrum, as in all marine anthozoans, the production of gametes is a function of both module (polyp) fertility and the number of fertile polyps per colony (Sakai, 1998). An accurate determination of the number of polyps is therefore crucial to obtain reliable estimates of key reproductive traits such as fecundity, which is an important parameter in ecosystem models that aim to generate forecast of the population dynamics of modular organisms (Cant et al., 2024; Bramanti et al., 2009; Santangelo et al., 2007). Furthermore, this new approach could be applied to standardize results across several manipulative experiments, such as in ecotoxicological and physiological studies, which depend on normalized polyp numbers rather than nubbins or surface area (e.g., Lange et al., 2023). For instance, this method could be used in studies that aim to accurately determine the prey capture rate per polyp (e.g., Chapron et al., 2021), which is important to gauge the energy expenditure in feeding. Moreover, studies that assess prey capture by dissecting polyps (e.g., Tsounis et al., 2005) could take advantage of our methodology to obtain estimations at the colony level. In general, the present method which was developed for C. rubrum can be adapted to other cnidarians, providing a valuable, non-destructive approach for studying coral ecology, conservation, and restoration.

Conclusion

Our study introduced a new method for the accurate estimation of polyp numbers in *C. rubrum* based on 3D photogrammetric reconstructions of live colonies. We found an overall mean polyp density of 16.9 ± 4.5 polyps cm⁻², with significant differences between branching orders (i.e., higher densities in 1st-order branches). This approach constitutes an improvement with regard to existing methods as it allows a more precise estimations of polyps number, crucial for understanding reproductive dynamics and informing conservation plans. While our method is more time-consuming than the approach based on basal diameter measurements, its higher accuracy makes it ideal for more focused studies on a limited number of colonies. This method will lead to an improved understanding of *C. rubrum* ecology and offers a valuable tool for broader marine anthozoan research and conservation applications.

Acknowledgements

This paper is part of the BG PhD thesis funded by the Doctorate School of the University of Cagliari and COST Action CA20102 Marine Animal Forests of the World (MAF-WORLD, Short-term scientific mission). The present study is part of the project ROMERO which benefits from the support of Fondation Prince Albert II de Monaco. LB is funded by the "BNP PARIBAS climate change and biodiversity initiative" through the DEEPLIFE project. LDF was funded by the "ERAS-MUS+ student mobility for traineeships" program.

References

- Aston, E.A., Duce, S., Hoey, A.S., Ferrari, R., 2022. A Protocol for Extracting Structural Metrics From 3D Reconstructions of Corals. *Frontiers in Marine Science*, 9, 854395.
- Beiring, E., Lasker, H., 2000. Egg production by colonies of a gorgonian coral. *Marine Ecology Progress Series*, 196, 169-177.
- Benedetti, M.C., Bramanti, L., Priori, C., Erra, F., Iannelli, M. et al., 2020. Polyp longevity in a precious gorgonian coral: hints toward a demographic approach to polyp dynamics. *Coral Reefs*, 39, 1125-1136.
- Bramanti, L., Iannelli, M., Santangelo, G., 2009. Mathematical modelling for conservation and management of gorgonians corals: youngs and olds, could they coexist? *Ecological Modelling*, 220, 2851-2856.
- Bramanti, L., Iannelli, M., Fan, T. Y., Edmunds, P. J., 2015. Using demographic models to project the effects of climate change on scleractinian corals: Pocillopora damicornis as a case study. *Coral Reefs*, 34, 505-515.
- Cánovas-Molina, A., Montefalcone, M., Bavestrello, G., Cau, A., Bianchi, C.N. *et al.*, 2016. A new ecological index for the status of mesophotic megabenthic assemblages in the mediterranean based on ROV photography and video footage. *Continental Shelf Research*, 121, 13-20.
- Carugati, L., Moccia, D., Bramanti, L., Cannas, R., Follesa, M.C. *et al.*, 2022. Deep-Dwelling Populations of Mediterranean Corallium rubrum and Eunicella cavolini: Distribution, Demography, and Co-Occurrence. *Biology*, 11, 333.
- Cau, A., Bramanti, L., Cannas, R., Follesa, M.C., Angiolillo, et al., 2016. Habitat constraints and self-thinning shape Mediterranean red coral deep population structure: implications for conservation practice. Scientific Reports, 6 (1), 23322.
- Chapron, L., Galand, P.E., Pruski, A.M., Peru, E., Vétion, G. et al., 2021. Resilience of cold-water coral holobionts to thermal stress. *Proceedings of the Royal Society B*, 288 (1965), 20212117.
- Conley, D.D., Hollander, E.N.R., 2021. A Non-destructive Method to Create a Time Series of Surface Area for Coral Using 3D Photogrammetry. *Frontiers in Marine Science*,

8,660846.

- Ferrari, R., Figueira, W.F., Pratchett, M.S., Boube, T., Adam, A. et al., 2017. 3D photogrammetry quantifies growth and external erosion of individual coral colonies and skeletons. *Scientific Reports*, 7, 16737.
- Garrabou, J., Gómez-Gras, D., Medrano, A., Cerrano, C., Ponti, M. *et al.*, 2022. Marine heatwaves drive recurrent mass mortalities in the Mediterranean Sea. *Global Change Biol*ogy, 28, 5708-5725.
- Guizien, K., Bramanti, L., 2014. Modelling ecological complexity for marine species conservation: the effect of variable connectivity on species spatial distribution and age-structure. *Theoretical Biology Forum*, 107, 47-56.
- Knittweis, L., Aguilar, R., Alvarez, H., Borg, J.A., Evans, J. et al, 2016. New depth record of the precious red coral Corallium rubrum for the Mediterranean. Rapport de La Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée, Vol 41, 467 pp.
- Lange, I.D., Perry, C.T., 2020. A quick, easy and non-invasive method to quantify coral growth rates using photogrammetry and 3D model comparisons. *Methods in Ecology and Evolution*, 11, 714-726.
- Lange, K., Maguer, J.F., Reynaud, S., Ferrier-Pagès, C., 2023. Nutritional ecology of temperate octocorals in a warming ocean. *Frontiers in Marine Science*, 10, 1236164.
- Maggioni, F., Bramanti L., 2022. Mediterranean red coral as a spawning ground for cuttlefish. *Global Journal of Ecology*, 7 (2), 120-121.
- Million, W.C., O'Donnell, S., Bartels, E., Kenkel, C.D., 2021. Colony-Level 3D Photogrammetry Reveals That Total Linear Extension and Initial Growth do not scale with complex morphological growth in the branching coral, Acropora cervicornis. *Frontiers in Marine Science*, 8, 646475.
- Pratlong, M., Haguenauer, A., Chenesseau, S., Brener, K., Mitta, G. *et al.*, 2017. Evidence for a genetic sex determination in Cnidaria, the Mediterranean red coral (Corallium rubrum). *Royal Society Open Science*, 4 (3), 160880.
- Rossi, S., Bramanti, L., Gori, A., Orejas, C., 2017. An overview of the animal forests of the world. p 1-26. In: *Marine* animal forests. Rossi, S., Bramanti, L., Gori, A., Orejas, C.

(Eds). Springer International Publishing.

- Sakai, K., 1998. Effect of colony size, polyp size, and budding mode on egg production in a colonial coral. *The Biological Bulletin*, 195, 319-325.
- Santangelo, G., Bramanti, L., 2010. Quantifying the decline in Corallium rubrum populations. Marine *Ecology Progress Series*, 418, 295-297.
- Santangelo, G., Bramanti, L., Iannelli, M., 2007. Population dynamics and conservation biology of the over-exploited Mediterranean red coral. *Journal of Theoretical Biology*, 244, 416-423.
- Santangelo, G., Carletti, E., Maggi, E., Bramanti, L., 2003. Reproduction and population sexual structure of the overexploited Mediterranean red coral Corallium rubrum. *Marine Ecology Progress Series*, 248, 99-108.
- Stimson, J., Kinzie R.A., 1991. The temporal pattern and rate of release of zooxanthellae from the reef coral Pocillopora damicornis (Linnaeus) under nitrogen-enrichment and control conditions. *Journal of experimental marine Biology* and Ecology, 153 (1), 63-74.
- Tsounis, G., Rossi S., Laudien J., Bramanti L., Fernandez N. et al., 2005. Diet and seasonal prey capture rates in Mediterranean red coral (*Corallium rubrum* L. 1758). *Marine Biology*, 149 (2), 313-325.
- Tsounis, G., Rossi, S., Grigg, R., Santangelo, G., Bramanti, L. et al., 2010. The exploitation and conservation of precious corals. Oceanography and marine biology: an annual review, 48, 161-212.
- Tsounis, G., Rossi, S., Bramanti, L., Santangelo, G., 2013. Management hurdles for sustainable harvesting of Corallium rubrum. *Marine Policy*, 39, 361-364.
- Veal, C.J., Carmi, M., Fine, M., Hoegh-Guldberg, O., 2010. Increasing the accuracy of surface area estimation using single wax dipping of coral fragments. *Coral Reefs*, 29, 893-897.
- Viladrich, N., Linares, C., Padilla Gamiño, J.L., 2022. Lethal and sublethal effects of thermal stress on octocorals early life-history stages. *Global Change Biology*, 28, 7049-7062.
- Zibrowius, H., Monteiro Marques, V., Grasshoff, M., 1984. La répartition du Corallium rubrum dans l'Atlantique (Cnidaria: Anthozoa: Gorgonaria). *Tethys*, 11 (2), 163-170.