



# **Mediterranean Marine Science**

# early view



Application of a new multi-metric index with a comparative analysis for assessing the environmental status of Posidonia oceanica meadows (Algeria, Southern Mediterranean)

AMINA SENGOUGA, SALIMA BOUMAZA, ALI MISRAOUI, MOHAMED ZERROUKI, RIHEM BOUDJADJA, FERIAL LOUANCHI, RACHID SEMROUD

doi: <u>10.12681/mms.39215</u>

# To cite this article:

SENGOUGA, A., BOUMAZA, S., MISRAOUI, A., ZERROUKI, M., BOUDJADJA, R., LOUANCHI, F., & SEMROUD, R. (2025). Application of a new multi-metric index with a comparative analysis for assessing the environmental status of Posidonia oceanica meadows (Algeria, Southern Mediterranean). *Mediterranean Marine Science*, early view. https://doi.org/10.12681/mms.39215

# Application of a new multi-metric index with a comparative analysis for assessing the environmental status of *Posidonia oceanica* meadows (Algeria, Southern Mediterranean)

# Amina SENGOUGA<sup>1</sup>, Salima BOUMAZA<sup>1</sup>, Ali MISRAOUI<sup>1</sup>, Mohamed ZERROUKI <sup>2</sup>, Rihem BOUDJADJA<sup>1</sup>, Ferial LOUANCHI <sup>2</sup>, and Rachid SEMROUD<sup>1</sup>

 <sup>1</sup>LCVRM: Laboratoire de Conservation et de Valorisation des Ressources Marines, Ecole Nationale Supérieure des Sciences de la Mer et de l'Aménagement du Littoral, Campus Universitaire de Dely Ibrahim, Bois des Cars, BP 19, 16320, Algiers, Algeria
 <sup>2</sup>ECOSYSMarL: Laboratoire des Écosystèmes Marins et Littoraux, Ecole Nationale Supérieure des Sciences de la Mer et de l'Aménagement du Littoral, Campus Universitaire de Dely Ibrahim, Bois des Cars, BP 19, 16320, Algiers, Algeria

Corresponding author: Amina SENGOUGA; sengouga.am@gmail.com

Contributing Editor: Panayota KOULOURI

Received: 28 October 2024; Accepted: 18 June 2025; Published online: 15 July 2025

# Abstract

Several biotic indices have been developed over the last two decades to assess water and habitat quality in the Mediterranean Sea, with *Posidonia oceanica* serving as a Biological Quality Element (BQE). To contribute to this effort, we created a new Multi-metric Index for the Algerian coast based on *P. oceanica* (MIAPo), adapted from existing indices and calibrated at eight sites with varying human pressures. This study also intends to compare the performance of MIAPo and PREI (*P. oceanica* Rapid Easy Index) in assessing the ecological status of coastal Algerian waters. MIAPo was built using multivariate analyses that combined five metrics: lower limit depth, leaf area index, epiphytic load, leaf tannin cell density, and leaf nitrogen content. To accurately interpret MIAPo and PREI, a Pressure Index (PI) was calculated, taking into account urbanization, agriculture, coastal population, uncontrolled dumping, fishing activity, rivers, industrial activity, and urban effluents. According to the Water Framework Directive (WFD) classification, the indices MIAPo and PREI demonstrated high comparability in the ecological status assigned to the studied sites (62.5% agreement), which was validated using Spearman's correlation coefficient (R = 0.88; p = 0.004). In addition, MIAPo and PREI show a strong correlation with the PI (R = -0.96; p = 0.0002 and R = -0.80; p = 0.02, respectively). The proposed index, MIAPo, provides a more comprehensive measure than PREI for the assessment and monitoring of Algerian coastal waters. Nonetheless, more field data collection is needed to expand the use of this valuable tool on a broader geographical scale.

Keywords: Posidonia oceanica; Biological indicator; Pressure-impact relationship; Water quality; Ecological status.

#### Introduction

The biological indicator Posidonia oceanica is widely used as a Biological Quality Element (BQE) under the Water Framework Directive (WFD, 2000/60/EEC). Several multi-metric indices have been proposed to assess the ecological quality of coastal waters based on plant characteristics and/or meadow structure: the "PosSte" (Buia et al., 2004), the "POMI" (Romero et al., 2007), the "Valencian CS" (Fernández Torquemada et al., 2008), the "PREI" (Gobert et al., 2009), the "BiPo" (Lopez y Royo et al., 2010a) and "the Vitality Index" (Pergent et al., 2015) (Table 1). To comply with the WFD, these indices must meet a set of required criteria, including (i) a significant relationship between the Ecological Quality Ratio (EQR) and anthropogenic pressures and (ii) the use of a common scale of five ecological status classes: High, Good, Moderate, Poor and Bad.

Several studies analyzed and compared these multi-metric indices (including the number and type of metrics, survey depth, metric aggregation method, and reference condition definition) as well as their performance across pressure gradient (Table 1). BiPo and PREI appear to be the most widely adopted tools for quality water assessment due to their rapid field-based protocols, ease of use, and low cost (Table 1). On the other hand, the performance of multi-metric indices based on *P. oceanica* metrics to assess water quality appears to be related to the number of metric types used, as demonstrated by Bennett *et al.* (2011) and Mascaró *et al.* (2012).

Moreover, the intricate nature of seagrass habitats has led numerous studies to underscore the importance of employing multiple indices derived from various metrics to evaluate the ecological status of *P. oceanica* meadows (Montefalcone *et al.*, 2006; Martínez-Crego *et al.*, 2010; Lopez y Royo *et al.*, 2011; Güreşen *et al.*, 2020a; Manci-

	Regions of development and other application regions	Methods	Metrics and depths	Reference con- ditions	Pressures tested
Poste III system for C. coastal water y classification using <i>P. oce-</i> <i>anica</i>	<b>aly</b> (Buia <i>et al.</i> , 2004) atalonia (Spain) and Corsica (France) Ischia (Italy) (Lopez Royo <i>et al.</i> , 2011)	Combination of 5 metrics in a data warehouse	<u>At intermediate depth (15 m)</u> Shoot density, intermediate leaf width, leaf production, rhizome production, rhizome elongation.	Existing pris- tine/undis- turbed sites	~
POMI: C Posidonia C: oceanica al Multivariate Sé Index Pe	atalonia (Spain) (Romero <i>et al.</i> , 2007) atalonia (Spain) and Corsica (France) (Lopez y Royo <i>et</i> ., 2011); uronikos (Greece) (Gerakaris <i>et al.</i> , 2017) OMI 9: Catalonia (Spain) (Bennett <i>et al.</i> , 2011) OMI 5: Catalonia, Balearic Islands and Croatia (Mas- ró <i>et al.</i> , 2012)	Combination of 14 metrics using PCA	At intermediate depth (15 m) Shoot density <sup>9</sup> , meadow cover <sup>9,5</sup> , percentage of plagiotropic rhizomes, shoot leaf area <sup>9,5</sup> , percentage of foliar necrosis <sup>9</sup> , N and P rhi- zomes content, sucrose rhizomes content <sup>9,5</sup> , $\delta^{15}N^{9,5}$ and $\delta^{34}S^9$ isotopic ratio in rhizomes, N epiphyte content <sup>9,5</sup> , Cu, Pb <sup>9</sup> and Zn in rhizomes.	A composite site from exist- ing values and expert judg- ment	Coastline con- structions, beach regeneration, ur- ban sewage, agri- cultural and urban soil use, tourism, fishing, recre- ational, fishing , commercial and industrial harbors
CS Valencian V: Se Th	<b>ilencia (Spain)</b> (Fernández Torquemada <i>et al.</i> , 2008) tronikos Gulf (Greece) (Gerakaris <i>et al.</i> , 2017) emiti Islands (Italy) (Tursi <i>et al.</i> , 2022a)	Combination of 9 metrics using PCA	At intermediate depth (14-17 m) Shoot density, meadow cover, dead matte cover, percentage of plagiotropic rhizomes, rhizome baring/burial, shoot leaf area, per- centage of foliar necrosis, herbivore pres- sure, leaf epiphyte biomass.	Virtual refer- ence site from existing values and expert judgment	Coastal construc- tions, beach re- generation, urban and industrial sewage, rivers and channels, agricultural soil use

	-
2	2
	2
	ŝ
	Ξ
- 1	3
	2
	S
5	ũ
~	
	ð
- 7	-
- 1	0
Ľ	3
	-

nunuco i enunu	lett				
Indices	Regions of development and other application regions	Methods	Metrics and depths	Reference con- ditions	Pressures tested
BiPo:	Corsica (France) (Lopez y Royo et al., 2010a)	Normalization	At the lower limit	Virtual refer-	Land use, indus-
Biotic Index using the sea-	Catalonia (Spain), Corsica (France), Ischia (Italy) (Lopez y Royo <i>et al.</i> , 2011)	and combination by averaging 4 metrics	Lower limit depth and type	ence sites from existing values	trial activity, river discharges, port
grass Posido- nia oceanica	Corsica (France) (Pergent et al., 2015)		Shoot density shoot length		activities and coastal planning.
	Tipaza, Algiers (Algeria) (Boumaza et al., 2015)				
	Mostaganem and Oran (Algeria) (Bentaallah, 2017)				
	Saronikos Gulf (Greece) (Gerakaris et al., 2017)				
	Vasiliko (Cyprus) (Kletou <i>et al.</i> , 2020)				
	Gökçeada (Turkey) (Güreşen <i>et al.</i> , 2020a)				
	Gökçeada (Turkey) and Corsica (France) (Güreşen <i>et al.</i> , 2020b)				
	Tremiti Islands (Italy) (Tursi <i>et al.</i> , 2022a)				

Corsica and PACA (Gobert et al., 2009)NormalizationAt the lower limitItalia (Rende et al., 2011; Rigo et al., 2019; Oprandi et al., 2021)NormalizationAt the lower limitA con site fr by averaging 5A con cover limit depth and typeA con site fr ing va2021)Tipaza, Algiers (Algeria) (Boumaza et al., 2015)MetricsA tintermediate depth (15 m): Shoot density, shoot leaf area, ratio between epiphytic and leaf biomassA con site fr ing va2021)Liguria (Italy) (Mancini et al., 2020)Shoot density, shoot leaf area, ratio between epiphytic and leaf biomassA con site fr ing vaSicily(Bellissimo et al., 2020)Sicily(Bellissimo et al., 2020)A con sicily(Bellissimo et al., 2020)A con sicily(Bellissimo et al., 2020)Tremiti Islands (Italy) (Tursi et al., 2023)A con sicily (Bellissimo et al., 2020)A con sicily (Bellissimo et al., 2020)	<pre>nposite Fish farming, in- om exist- dustrial develop- lues and ment, agriculture,</pre>	ing tourism, fishing, commercial ports	and urbanization					
Corsica and PACA (Gobert et al., 2009)Normalization $\underline{At the lower limit}$ Italia (Rende et al., 2011; Rigo et al., 2019; Oprandi et al.)Domination $\underline{At the lower limit}$ 2021)Tipaza, Algiers (Algeria) (Boumaza et al., 2015)py averaging 5 $\underline{At intermediate depth (15 m)}$ Tipaza, Algiers (Algeria) (Boumaza et al., 2017)metrics $\underline{At intermediate depth (15 m)}$ Saronikos Gulf (Greece) (Gerakaris et al., 2017)Ehoot density, shoot leaf area, ratio betweLiguria (Italy) (Mancini et al., 2020)Yasiliko (Cyprus) (Kletou et al., 2020)Vasiliko (Cyprus) (Kletou et al., 2020)Sicily(Bellissimo et al., 2020)Sicily(Bellissimo et al., 2020)Finati Islands (Italy) (Tursi et al., 2023)	A com site fr ing va	model						
Corsica and PACA (Gobert et al., 2009)NormalizationItalia (Rende et al., 2011; Rigo et al., 2019; Oprandi et al., 2021)by averaging 52021)metricsTipaza, Algiers (Algeria) (Boumaza et al., 2015)py averaging 5Saronikos Gulf (Greece) (Gerakaris et al., 2017)metricsLiguria (Italy) (Mancini et al., 2020)Vasiliko (Cyprus) (Kletou et al., 2020)Vasiliko (Cyprus) (Kletou et al., 2020)Sicily(Bellissimo et al., 2020)Sicily(Bellissimo et al., 2020)TipazaVasiliko (Cyprus) (Kletou et al., 2020)TipazaSicily(Bellissimo et al., 2020)TipazaTremiti Islands (Italy) (Tursi et al., 2022a)	At the lower limit Lower limit depth and type	At intermediate depth (15 m): Shoot density, shoot leaf area, ratio betwe	epiphytic and leaf biomass					
Corsica and PACA (Gobert <i>et al.</i> , 2009) Italia (Rende <i>et al.</i> , 2011; Rigo <i>et al.</i> , 2019; Oprandi <i>et al.</i> , 2021) Tipaza, Algiers (Algeria) (Boumaza <i>et al.</i> , 2015) Saronikos Gulf (Greece) (Gerakaris <i>et al.</i> , 2017) Liguria (Italy) (Mancini <i>et al.</i> , 2020) Vasiliko (Cyprus) (Kletou <i>et al.</i> , 2020) Vasiliko (Cyprus) (Kletou <i>et al.</i> , 2020) Sicily(Bellissimo <i>et al.</i> , 2020)2000/60/EC Tremiti Islands (Italy) (Tursi <i>et al.</i> , 2023)	Normalization and combination by averaging 5	metrics						
	<b>Corsica and PACA</b> (Gobert <i>et al.</i> , 2009) <i>Italia</i> (Rende <i>et al.</i> , 2011; Rigo <i>et al.</i> , 2019; Oprandi <i>et al.</i> ,	<sup>2021)</sup> Tipaza, Algiers (Algeria) (Boumaza <i>et al.</i> , 2015)	Saronikos Gulf (Greece) (Gerakaris et al., 2017)	Liguria (Italy) (Mancini <i>et al.</i> , 2020)	Vasiliko (Cyprus) (Kletou <i>et al.</i> , 2020)	Sicily(Bellissimo et al., 2020)2000/60/EC	Tremiti Islands (Italy) (Tursi et al., 2022a)	

Pressures tested	Distance to cit- ies and harbors, trawling, urban and industrial discharges
Reference con- ditions	Exposed and non-exposed (or less ex- posed) sites from the study area
Metrics and depths	At the lower limit Lower limit depth and type, meadow cover, percentage of plagiotropic rhizomes, leaf production, rhizome elongation At intermediate depth (15 m) Shoot density, shoot leaf area
Methods	Averaging of 8 metrics
Regions of development and other application regions	Corsica (France) (Pergent <i>et al.</i> , 2015) Gökçeada (Turkey) and Corsica (France) (Güreşen <i>et al.</i> , 2020b) Tremiti Islands (Italy) (Tursi <i>et al.</i> , 2022a);
Indices	Vitality Index

ni *et al.*, 2020; Tursi *et al.*, 2022a). The study conducted by Boumaza *et al.* (2015) on *P. oceanica* meadows along the central Algerian coast indicates that the PREI most accurately represents the anticipated ecological status. In contrast, the BiPo exhibits a diminishing correlation with human pressures. Kletou *et al.* (2020) established a strong correlation between human pressures and both PREI and BiPo in Vasiliko Bay, Cyprus. Significant correlations between BiPo and anthropogenic pressures have been identified along the Western Algerian coast (Bentaallah, 2017) and the Turkish coasts (Güreşen *et al.*, 2020b).

Using multi-metric indices outside their intended regional range requires adaptation, calibration, and validation to ensure effectiveness (Martínez-Crego et al., 2010; Pergent et al., 2015; Dallas, 2021). The selection of appropriate multi-metric indices for assessing ecological status is thus critical for coastal water quality management and monitoring. The challenge of implementing a multi-metric index based on P. oceanica, which is shared throughout the Mediterranean basin, appears to be essential and requires the collaboration of all scientists, as highlighted by Di Camillo et al. (2023) in the case of coralligenous reefs. As part of ongoing efforts to harmonize monitoring tools across the Mediterranean region, the purpose of this work is to present a new multi-metric index called the Multi-metric Index for the Algerian coast based on P. oceanica (MIAPo), developed for the first time along the southern Mediterranean coasts. It incorporates tannin cells density as an additional descriptor alongside those commonly used. The sensitivity of this cellular indicator to anthropogenic pressures has been tested and demonstrated by Boumaza et al. (2022). Its inclusion aligns with current recommendations advocating for the integration of complementary metrics from the community and population levels to biochemical and cellular levels to achieve a more comprehensive ecosystem assessment (Martínez-Crego et al., 2010; Martinez-Haro et al., 2015) and rapid time responses (Ferrat et al., 2003; Romero et al., 2016). Notably, this parameter stands out from other metrics at a similar level due to its ease and rapid measurement, without the need for advanced equipment. The proposed index is based on previous multi-metric indices listed in Table 1 and attempts to assess the ecological status of Algerian coastal waters. It was compared to the previously existing multi-metric index PREI (Gobert et al., 2009). We estimated the robustness of MIAPo and PREI by investigating their relationships with cumulative multiple human pressures represented by a Pressure Index (PI).

# **Materials and Methods**

# Metrics selection and P. oceanica indices application

The most important aspect of the multi-metric index development procedure is metric selection, which prevents an incorrect interpretation of the biological response to environmental degradation (Stoddard *et al.*, 2008; Hawkins *et al.*, 2010). MIAPo metrics are cho-

Table 1 continued

sen based on our previous studies within the P. oceanica meadows in the studied geographic area (Boumaza et al., 2012, 2014, 2015, 2022; Sengouga, 2017; Sengouga et al., 2018, 2019). These studies helped us identify the most relevant metrics, particularly those that appeared to be sensitive to anthropogenic pressures. Five metrics were chosen (Table 2) based on recommendations from Martínez-Crego et al. (2010) and Romero et al. (2016). The researchers emphasized the importance of using metrics at the population and community levels to maintain ecological integrity and biomarkers to detect disturbances early. Each type of metric was represented by one metric on one side. This method is simple and avoids the problems that come with weighing one type more than another. On the other hand, the performance characteristics of the chosen metrics were related to their ability to distinguish between the least-disturbed and most-disturbed sites (Stoddard et al., 2008). Furthermore, based on the 0.71 threshold established by Stoddard et al. (2008), a redundancy test was carried out using Spearman's correlation analysis and showed that the selected metrics were not redundant (Table 3). The PREI was elaborated according to Gobert et al. (2009), and it differed from MIAPo in the metrics and their types (Table 2).

# Reference levels and reference conditions

According to Romero et al. (2007), reference levels, which referred to both the upper and lower limits of a metric's range, were used in the normalization process required to generate MIAPo. The "hypothetical reference site" was determined by averaging the two higher and two lower values for metrics that decreased and increased in response to anthropogenic disturbances, respectively (Table 2). The "hypothetical worst site" was established using the reverse procedure. In the case of the lower limit depth, the maximum and minimum values observed in the prospected geographical zone were used to elaborate on the previous "hypothetical sites". The reference condition for generating the PREI index was defined by Gobert et al. (2009). We adapted this approach to our studied area by averaging the three highest values (after discarding the maximum) for shoot density and leaf surface area, 0 dry weight for epiphytic leaf biomass E/L (considering that healthy seagrass has non-epiphyted leaves), and the maximum value for the lowest limit depth recorded in the studied sites.

` •	·			
Types of metrics	Metrics	MIAPo	PREI	Responses
Community	Epiphytic leaf biomass (E/L) (mg.mg <sup>-1</sup> ) Epiphytic load (mg.cm <sup>-2</sup> )	+	+	Increase Increase
Population	Shoot density (shoot.m <sup>-2</sup> ) Leaf area index (m <sup>2</sup> .m <sup>-2</sup> )	+	+	Decrease Decrease
Lower limit	Type of lower limit Depth of lower limit (m)	+	+ +	Typology modification Decrease
Individual	Shoot leaf surface (cm <sup>2</sup> /shoot)		+	Decrease
Cellular	Leaf tannin cell density (cells.mm <sup>-2</sup> )	+		Increase
Biochemical	Leaf nitrogen content (% DW)	+		Increase

 Table 2. Metrics used for MIAPo and PREI indices, categorized by metric type and expected response to anthropogenic disturbance (based on previous studies).

**Table 3.** Redundancy test using Spearman's correlation analysis for the MIAPo selected metrics. Spearman correlation coefficients (R) are in bold; p-values are in regular font.

			Spearman correl	ation R		
		Epiphytic load	Leaf area index	Depth of lower limit	Leaf nitrogen content	Leaf tannin cell density
0	Epiphytic load		-0.52	-0.19	0.50	0.45
value	Leaf area index	0.1827		0.49	-0.12	-0.02
Ъ	Depth of lower limit	0.6494	0.2166		-0.37	-0.44
	Leaf nitrogen content	0.207	0.7789	0.3652		0.43
	Leaf tannin cell density	0.2604	0.9554	0.2715	0.2894	

#### **Ecological Quality Ratio**

To ensure comparable values of MIAPo and PREI and highlight concordant or discordant responses, MIAPo's EQR was estimated by normalizing data between 0 and 1 (PREI already represents the EQR value). An arbitrary value of 0.1 was assigned to the boundary of the "bad" status and the remaining interval from 0.1 to 1 was divided into four equal intervals as required by the WFD (Romero et al., 2007; Gobert et al., 2009). This EQR was calculated as described by Romero et al. (2007), Oliva et al. (2012) and García-Marín et al. (2013) for their indices. Thus, we combined the MIAPo metrics into a single scale through a principal component analysis (PCA) based on the correlation matrix to account for differences in measurement scales (Legendre & Legendre, 1998). The two hypothetical sites corresponding to the extreme conditions (reference and worst) were included as supplementary objects. The EQR was calculated as follows:

$$EQR'_{x} = (Cl_{x} - Cl_{worst}) / (Cl_{Reference} - Cl_{worst})$$
(1)

$$EQR_{r} = (EQR'_{r} + 0.11)/(1 + 0.10)$$
(2)

where, CI is the PCA's first component scores; EQR<sub>x</sub> is the ecological quality ratio of the site  $\varkappa$ ;  $CI_{\varkappa}$  is the score of the site  $\varkappa$ ;  $CI_{reference}$  is the score of the hypothetical "reference" site;  $CI_{worst}$  is the score of the hypothetical "worst" site.

# Study area, sampling collection and metrics measurements

This study was carried out along the central Algerian coast (Fig. 1). Eight P. oceanica meadows from two provinces were analyzed, each with different levels and types of anthropogenic pressure. Four sites in the western part of the study area are in Tipaza province (Chenoua, Kouali, Berrard, and Bou Ismail), while four sites in the eastern part are in Algiers province (El Djamila, Ain Benian, Rais Hamidou, and Aguelli). The available state data on the various human activities in these two provinces, obtained from publicly available and several studies on the region (PNUE/PAM/CAR PAP, 2006; Bakalem et al., 2009; Mangos & Claudot, 2013; Haouchine et al., 2015), indicate that Tipaza is renowned for its significant agricultural and forestry potential as well as tourism and fishing vocations. Algiers, the country's capital, is the most densely populated area, with the highest concentration of services, facilities, infrastructure, industry, and major urban developments.

For the assessment of MIAPo and PREI metrics, the following data were collected in the field between July and August 2018 at approximately 12 m depth (excluding the meadow lower limit depth and typology) to avoid the masking effects of depth and seasonal variability. This intermediate depth was selected considering the lower limits of *P. oceanica* meadows in the geographical area, which range from 14 to 20 m and mostly coincide with the lower limit of the rocky substrate. The WFD recommends using the protocols at shallower depths when meadows do not reach the standard depth of 15 m (Blouet *et al.*,



Fig. 1: Map of the study area with studied municipalities and location of the 8 P. oceanica meadows investigated.

2011), and where the conditions are optimal for meadow development (PNUE/PAM CAR/ASP, 2019):

Metrics 1-2: The lower limit depth was recorded *in situ* and classified according to Meinesz & Laurent's (1978) typology.

Metric 3: Shoot density (shoots. m<sup>-2</sup>) was estimated using 10 randomly located 0.16 m<sup>2</sup> quadrats. This sampling effort provides the minimum standard error required for an accurate estimation of mean seagrass density (Panayotidis *et al.*, 1981; Pergent *et al.*, 1995; Pergent-Martini *et al.*, 2005; Lopez Y Royo *et al.*, 2010b).

Metric 4-5: Leaf area per shoot (cm<sup>2</sup>) was estimated from leaf morphometrics of 15 shoots collected and leaf area index (LAI) from shoot leaf area and shoot density (m<sup>2</sup>.m<sup>-2</sup>) (Buia *et al.*, 2004; Pergent *et al.*, 1995). To minimize the impact of destructive sampling methods (Buia *et al.*, 2004; Rotini *et al.*, 2013; PNUE/PAM CAR/ASP, 2019), we adopted a relatively limited sampling effort of 15 shoots, which is considered sufficient for this purpose (Pergent *et al.*, 1995; Pergent-Martini *et al.*, 2005). Furthermore, the PREI assessment in this work is based on a previous multi-year monitoring (Boumaza *et al.*, 2015; Sengouga, 2017; Sengouga *et al.*, 2019) that showed consistent PREI-anthropogenic pressure trends when using 20 shoots in 2011 and 15 shoots in both 2015 and 2018.

Metrics 6-7: Epiphyte biomass separated from each shoot (15 shoots) was measured after drying (70°C for 72 h) and standardized to leaf dry mass (mg.mg<sup>-1</sup>DW) (E/L) (Terrados & Pons, 2008) and to leaf area (mg DW.cm<sup>-2</sup>) (Boumaza *et al.*, 2014) to obtain epiphytic load.

Metric 8: Leaf tannin cell density (cells. mm<sup>-2</sup>) was carried out under a microscope, over cross-sections at the basal, the middle and the apical region of all adult and intermediate leaves (*sensu* Giraud, 1979) of 5 collected shoots, as described by Boumaza *et al.* (2022).

Metric 9: Leaf nitrogen content was analyzed in all adult and intermediate leaves from 9 collected shoots, which were deemed sufficient based on sample sizes used in previous studies (Delgado *et al.*, 1999; Fourqurean *et al.*, 2007; Leoni *et al.*, 2007; Scartazza *et al.*, 2017). The epiphyte-free leaves were lyophilized and ground into powder. The nitrogen content was determined using the Kjeldahl method with three or two subsamples (0.2 g) from each aliquot of ground material and expressed as %DW. This method has already been used with seagrasses (Duarte, 1990; Dawson & Dennison, 1996; Udy & Dennison, 1997; Walker *et al.*, 2004; Zhang *et al.*, 2011). Ammonium chloride was used as a reference material and was analyzed in each series of six samples. The tests produced a Root Mean Square Error of 0.02% DW.

# Pressure assessment

To relate the status of *P. oceanica* meadows to human activities, we quantified the key pressures in the studied geographical area using publicly available and satellite data, focusing on those that affect water quality. Eight quantitative data-driven pressures were considered: urbanization (% of urbanized area to municipality area), coastal population (% of municipality demography to province demography), agriculture (% of agricultural area to municipality area), fishing (number of fishing vessels per harbor), urban effluents (number of discharge points per municipality), uncontrolled dumping (% of wastewater volume not connected to the sewer system per municipality), rivers and industry (distance from the nearest river or industrial zone). In the case of industrial activities, river and fishing harbors, a distance of 10 km was considered significant (Comeleo et al., 1996; Rodriguez et al., 2007; Lopez y Royo et al., 2009a; Parravicini et al., 2012), implying that river mouths, industrial zones, and harbors located more than 10 km away from a site are not considered significant pressures.

PI was calculated using a rapid approach inspired by prior research on the vulnerability of coastal areas and rivers (Leopold, 1969; Bodéré *et al.*, 1991; Hallegouët *et al.*, 1997; Larid, 2002). It represents a cumulative of selected key pressures. It entails assigning scores ranging from 1 (no effect) to 5 (high effect) to various anthropogenic pressure values (Table 4) to categorize the pressures within this scale in two steps:

(i) definition of "high effect" and "no effect" conditions. In fact, for percentage-based pressures (urbanization, coastal population, agriculture, and uncontrolled dumping), "high effect" conditions correspond to 100%, while "no effect" conditions correspond to an arbitrary value of less than 5%, as assigned by (Larid, 2002). For pressures represented as distances (industrial activity and rivers), "high effect" conditions correspond to 0 km and "no effect" conditions correspond to distances greater than 10 km, as predicted by Comeleo *et al.* (1996), Rodriguez *et al.*(2007), Lopez y Royo *et al.*(2009a) and Parravicini *et al.* (2012). For pressures represented as activity counts (fishing activity and urban effluents), the lowest recorded values

Table 4. Definition of "high effect" and "no effect" conditions of the selected key pressures.

Units		Percenta	uges (%)		Distanc	es (km)	Nun	ıbers
Pressures	Urbaniza- tion	Coastal population	Agricul- ture	Uncon- trolled dumping	Industrial activity	Rivers	Fishing activity	Urban ef- fluents
No effect		< 5	%		> 1(	) km	0	0
High effect		100	) %		01	κm	339	11

represent "no effect" conditions, while the highest recorded values represent "high effect" conditions.

(ii) pressure categorization (Table 5) is achieved by dividing the interval between "no effect" condition values and "high effect" condition values into 5 intervals of equal amplitude.

PI, which ranges from 0 (no human impact) to 1 (high human impact), is calculated as follows:

$$PI = \frac{\sum_{1}^{N} S_n}{5 \sum_{1}^{N} P_n} \tag{3}$$

where,  $S_n$  is the rate for each pressure, varying from 1 to 5, and  $P_n$  is the number of the considered pressures.

#### Statistical analysis

Statistical analyses were carried out in the R4.2.0 environment, with all tests performed at the 95% significance level. When the parametric assumptions were verified, one-way ANOVA was used to test differences between sampling sites. Otherwise, a non-parametric Kruskal-Wallis test was used. The Shapiro-Wilk and Bartlett tests were used to prove distribution normality and variance homogeneity, respectively. Metrics used to construct MIAPo were combined using PCA. Then, hierarchical clustering was used to determine whether the ordination of the studied sites along the first PCA axis was consistent with meadow quality. The number of clusters was decided using the number of ecological statuses obtained from the MIAPo. The relationship between MIAPo and PREI was investigated using Spearman's rank correlation, and an absolute average class difference (AACD) analysis was carried out to assess the agreement between these two indices, as described by Lopez y Royo *et al.* (2011). The Spearman's rank correlation was also used to determine how well MIAPo, PREI, and their metrics reflect human pressures. As the pressure data were evaluated at the municipality level, we averaged the EQRs and metrics of sites within the same municipality to allow for correlation estimation.

# Results

#### Ecological status class evaluation using MIAPo

Table 6 summarizes the mean ( $\pm$  SE) values per site for each metric used in the MIAPo calculation. Statistically significant differences among sites for all the metrics can be noted, allowing their use for the MIAPo concept, as indicated by Stoddard *et al.* (2008).

Table 5. Anthropogenic pressure (Pn), data sources, and scores (Sn) by value interval.

<b>D</b> ( <b>D</b> )	D-4-			Scores (S	n)	
Pressures (Pn)	Data sources	1	2	3	4	5
Urbanization (%)	• Cartography using Google Earth satellite imagery					
Coastal population	• Website of the Province of Tipaza					
(%)	• Statistical yearbook of the prov- ince of Algiers	- 5	5 00 7	20 0 52 5	50 6 76 0	76.2 100
Agriculture (%)	• Cartography using Google Earth satellite imagery	< 3	3-28.7	28.8-32.3	32.0-70.2	/0.3-100
Uncontrolled dump- ing (%)	<ul> <li>Website of the Province of Tipaza</li> <li>Statistical yearbook of the prov- ince of Algiers</li> </ul>					
Fishing activity (number)	<ul> <li>Website of the Province of Tipaza</li> <li>The Directorate of Fisheries and Fisheries Production of the prov- ince of Algiers</li> </ul>	< 84	84-168	169-253	254-338	≥ 339
Urban effluents (num- ber)	<ul> <li>National Coastal Commission of Tipaza</li> <li>Public establishment for urban hygiene and environmental pro- tection of the province of Algiers</li> </ul>	< 3	3-5	6-8	9-10	≥11
Industrial activity (Km) Rivers (Km)	<ul> <li>Ministry of Industry and pharma- ceutical production, and cartog- raphy</li> <li>Cartography using Google Earth satellite imagery</li> </ul>	> 10	10-7.5	7.6-5	5.1-2.4	≤ 2.5

Two axes, PCI and PCII, of 49.5% and 21%, respectively, were extracted using PCA, explaining 70.5% of the variance in *P. oceanica* metrics (Fig. 2). PCI was significantly correlated with most of the metrics, indicating that this axis is effective at indicating the state of meadow health. Lower limit depth and leaf area index were the most correlated with PCI, with the highest negative loading scores (-0.83 and -0.81, respectively), indicating a good status of meadows. In contrast, epiphytic load, leaf tannin cell density, and leaf nitrogen content were positively correlated with PCI, indicating a poor status of meadow health (Fig. 2). The PCA applied to the studied sites illustrates their logical ordination along the first axis, following a quality gradient from the 'optimal' site to the 'worst' site (Fig. 3). The sampling sites were grouped into three statistically significant clusters (Fig. 3).

(i) Cluster 1 (high status) consists of the Kouali and Berrard sites in Tipaza Province. These sites are distinguished by the lowest leaf nitrogen content, the highest lower limit depth, and the leaf area index (the latter only in Kouali) (Table 6).

(ii) Cluster 3, which corresponds to the moderate status at the El Djamila site (Algiers Province), has the

**Table 6.** Mean values of the selected metrics for MIAPo calculation (Mean values  $\pm$ SE) and the P-values from One-Way ANOVA and Kruskal-Wallis (K-W) tests applied to *P. oceanica* metrics for the 8 studied sites.

Sites	Tannin cell density (cells.mm <sup>-2</sup> )	Leaf nitrogen content (%DW)	Epiphytic load (mg DW .cm <sup>-2</sup> )	Leaf area index (m <sup>2</sup> .m <sup>-2</sup> )	Lower limit depth (m)
Chenoua	$44.54\pm8.10$	$2.29\pm0.04$	$1.32\pm0.10$	$4.60\pm0.43$	17
Kouali	$17.12\pm2.91$	$1.83\pm0.04$	$1.23\pm0.11$	$7.39\pm 0.57$	20
Berrard	$14.94\pm3.66$	$1.71\pm0.03$	$0.71\pm0.10$	$4.97\pm0.33$	19
Bou Ismail	$22.03 \pm 1.97$	$2.55\pm0.02$	$1.83\pm0.18$	$4.64\pm0.47$	18
El Djamila	$33.17\pm6.47$	$1.91\pm0.09$	$1.69\pm0.20$	$2.52\pm0.26$	14
Ain Benian	$11.51\pm1.84$	$2.18\pm0.02$	$1.53\pm0.12$	$3.53\pm0.34$	18
Rais Hamidou	$19.58 \pm 1.94$	$2.28\pm0.07$	$0.81\pm0.10$	$5.91\pm0.43$	17
Aguelli	$11.46\pm2.65$	$2.17\pm0.02$	$1.15\pm0.11$	$4.45\pm0.29$	17
Optimal	5.99	1.69	0.07	6.65	20
worst	68.68	2.57	3.20	2.25	14
<i>p</i> -value	2.0 E-04	2.3 E-08	3.9 E-07	2.3 E-12	/
Test	K-W	ANOVA	K-W	ANOVA	/



Fig. 2: Principal Component Analysis plot of P. oceanica metrics and their loadings on the first axis (in brackets).



*Fig. 3:* (A) Principal Component Analysis plot with site scores. (B) Hierarchical clustering dendrogram, highlighting the 3 main clusters corresponding to the 3 types of ecological statuses obtained from MIAPo.

lowest values for the leaf area index and lower limit depth.

(iii) Cluster 2 represents a good status and includes the remaining sites.

The EQR calculation classified the studied sites into three levels of status: high, good, and moderate, with values ranging from 0.44 to 0.86 (Table 7). The Algiers province meadows had a lower mean EQR than the Tipaza province meadows ( $0.62 \pm 0.13$  and  $0.71 \pm 0.16$ , respectively). Only one meadow was designated as moderate (El Djamila), five as good (Chenoua, Bou Ismail, Ain Benian, Rais Hamidou, and Aguelli), and two as high (Kouali and Berrard).

# Ecological status class evaluation using PREI and comparability with MIAPo

Table 8 summarizes the mean values per site ( $\pm$  SE) for each metric used in the PREI calculation. The data analysis revealed statistically significant differences across sites for all metrics.

The AACD shows that MIAPo and PREI are highly comparable (62.5% agreement). Only three sites, Berrard, Bou Ismail, and El Djamila, changed their MIAPo

Mediterr. Mar. Sci., 26/3, 2025, Early view

status from "high, good, and moderate" to "good, moderate, and poor" by the PREI (Fig. 4). Furthermore, a highly significant correlation was found between MIAPo and PREI (R = 0.88, p = 0.004; Fig. 5), while MIAPo consistently produced higher values than PREI across all sites (EQR mean values: 0.665 and 0.605 respectively), with the exception of the Chenoua site (Fig. 4).

#### Evaluation of anthropogenic pressures

Table 9 lists the results of each site's human pressure evaluation. The PI values revealed that sites in Algiers province experience higher pressures than sites in Tipaza province.

The Spearman correlations between the PREI EQR and MIAPo EQR values and the PI are statistically significant (R = -0.80; p = 0.02 and R = -0.96; p = 0.0002, respectively; Fig. 6).

#### Responses of metrics to human pressures

Correlations between *P. oceanica* metrics and the PI are statistically significant for population and community

**Table 7.** The scores on the first Principal Component Analysis axis and the resulting EQR values from each sampling site for theMIAPo.

Sites		MIAPo	
Siles	CI	EQR	Status
Chenoua	1.042	0.555	Good
Kouali	-2.143	0.863	High
Berrard	-1.94	0.832	High
Bou Ismail	0.944	0.575	Good
El Djamila	2.961	0.443	Moderate
Ain Benian	0.087	0.646	Good
Rais Hamidou	-0.679	0.726	Good
Aguelli	-0.272	0.684	Good
Optimal	-4.069	1.000	
Worst	6.692	0.099	

Table 8. Mean	values of the	he metrics	used for PR	EI calculation	(Mean	values $\pm$	= SE), P-	values	from (	One-Way	ANOVA	A and
Kruskal-Wallis	(K-W) tests	applied to A	P. oceanica	netrics and PR	EI EQR	in the 8	studied s	ites.				

Sites	Shoot density (shoots. m <sup>-2</sup> )	Leaf surface area (cm². Shoot <sup>-1</sup> )	E/L (mg.mg <sup>-1</sup> DW)	Low (1	ver limit depth m) and type	PREI EQR	Status
Chenoua	$169.38\pm8.2$	$271.68\pm25.5$	$0.31\pm0.03$	17	Stable	0.616	Good
Kouali	$211.88\pm20.2$	$348.73\pm26.8$	$0.32\pm0.02$	20	Stable	0.826	High
Berrard	$195.00\pm13.2$	$254.99 \pm 16.8$	$0.17\pm0.02$	19	Stable	0.734	Good
Bou Ismail	$217.50\pm21.9$	$213.42\pm21.6$	$0.54\pm0.04$	18	Regressive	0.509	Moderate
El Djamila	$119.38\pm7.2$	$211.26\pm21.8$	$0.39\pm0.05$	14	Regressive	0.270	Poor
Ain Benian	$151.88\pm8.5$	$232.58\pm22.5$	$0.38\pm0.03$	18	Stable	0.622	Good
Rais Hamidou	$187.50\pm21.6$	$315.27\pm22.9$	$0.23\pm0.03$	17	Stable	0.662	Good
Aguelli	$155.00\pm13.1$	$287.37 \pm 18.4$	$0.29\pm0.03$	17	Stable	0.616	Good
Optimal	300	467.76	0	20			
Worst	0	0	1	14			
<i>p</i> -value	2.4 E-04	8.7 E-05	3.8 E-08	/			
Test	K-W	ANOVA	K-W	/			



Fig. 4: EQR values and ecological status classification of sites according to MIAPo and PREI indices.



Fig. 5: Significant positive correlation between the PREI EQR and the MIAPo EQR (R = 0.88, p = 0.004).

Table 9. The rates assigned for each	pressure and the results of the	pressure index (PI) ca	alculated by municipality.
--------------------------------------	---------------------------------	------------------------	----------------------------

Provinces		Tipaza			Algiers	
Municipalities	Tipaza	Ain Tagourait	Bou Ismail	Ain Benian	Rais Hamidou	Reghaia
Sites	Chenoua/ Kouali	Berrard	Bou Ismail	El Djamila / Ain Benian	Rais Hamidou	Aguelli
			Rates (R	n)		
Urbanization	2	2	5	5	5	4
Agriculture	2	3	1	4	1	4
Coastal population	1	1	2	1	1	1
Uncontrolled dumping	2	2	2	1	1	1
Fishing activity	2	2	2	5	4	2
Rivers*	3	1	1	4	1	4
Industrial activity*	1	2	4	4	5	3
Urban effluents	5	3	4	2	2	1
PI	0.45	0.38	0.50	0.60	0.48	0.48

\* Despite the fact that the distances between sites within the same municipality vary, the outcome scores remain consistent.



Fig. 6: Significative negative Spearman correlation between the pressure index (PI) and both PREI EQR and MIAPo EQR.

types, with the exception of shoot density (Table 10).

#### Discussion

The simultaneous use of two multi-metric indices, MIAPo and PREI, provides comparable results in the classification of the water quality of Algerian central coastal waters despite the differences they present. Metrics were chosen and combined using an average for the PREI and a PCA for the MIAPo. In fact, different studies discovered similarities when comparing such disparate indices (Gerakaris et al., 2017; Kletou et al., 2020; Tursi et al., 2022a). Another notable difference between MIAPo and PREI lies in how they define reference conditions. In fact, while the worst condition of the PREI corresponds to extreme values for a recorded recent meadow die-off (e.g., 0 shoot.m<sup>-2</sup> for shoot density and 1 mg.mg<sup>-1</sup> DW for epiphytic load), the MIAPo uses only values of field data from local scale to define both optimal and worst conditions (e.g., 2.25 m<sup>2</sup>.m<sup>-2</sup> for leaf area index). Defining reference conditions can impact the precision

and robustness of biotic indices (Martínez-Crego *et al.*, 2010; Romero *et al.*, 2016) and lead to incorrect interpretation of results (Borja *et al.*, 2012).

The proposed MIAPo has the potential to be a useful tool for assessing the ecological status of Algerian coastal waters for the following reasons: (i) The MIAPo is sensitive to human pressures (R = -0.96) and can accurately reflect changes in ecological stress. (ii) Using hierarchical classification, the studied sites were arranged into clusters that corresponded exactly to their ecological status. This implies that the spatial variability of the selected metrics within the study area is clearly due to differences in meadow quality. (iii) It integrates a small number of metrics combining insights provided by the maximum variety of metric types. (iv) Notably, it integrates leaf tannin cell density as an easily applicable cellular indicator, adding it to the commonly used descriptors, thereby improving both cost-effectiveness and early detection. (v) To date, it remains the only index developed within the southern Mediterranean coast. (vi) The MIAPo has high quantitative (R = 0.88) and qualitative (62.5%) agreement with the PREI, which has already been used and verified in

**Table 10.** Spearman's rank correlation between the pressure index (PI) and the studied metrics (MIAPo and PREI metrics are superscripted by 'M' and 'P', respectively). Significative correlations are in bold.

Types of motion	Metrics	Spearman's rank	Spearman's rank correlation with PI		
Types of metrics		R	<i>p</i> -value		
	Epiphytic leaf biomass <sup>P</sup>	0.7239	0.04		
Community	Epiphytic load <sup>M</sup>	0.7239	0.04		
D 1.1	Shoot density <sup>P</sup>	-0.6258	0.1		
Population	Leaf area index <sup>M</sup>	-0.7976	0.02		
Lower limit	Depth of lower limit <sup>P,M</sup>	-0.9203	0.001		
Individual	Shoot leaf surface <sup>P</sup>	-0.5767	0.13		
Cellular	Leaf tannin cell density M	0.0491	0.91		
Biochemical	Leaf nitrogen content <sup>M</sup>	0.135	0.75		

the same region (Boumaza *et al.*, 2015; Sengouga, 2017; Sengouga *et al.*, 2019).

In terms of the sensitivity of the two applied indices to human pressure, the results show a stronger correlation between the MIAPo and the PI than the PREI (R = -0.96and -0.80, respectively). This relative difference is most likely due to differences in metrics and their types (Lopez y Royo et al., 2011; Gerakaris et al., 2017; Mancini et al., 2020). In fact, we have included additional metric types to supplement those already included in the PREI: cellular and biochemical metrics (leaf nitrogen content and leaf tannin cell density). Martínez-Crego et al. (2010) and Roca et al. (2015) suggested that incorporating these types of metrics can improve stress index sensitivity and early detection efficacy. Furthermore, several studies have supported the use of these indicators in monitoring programs (Dumay et al., 2004; Pérez et al., 2008; Boumaza et al., 2012, 2014, 2022; Jones et al., 2018; Helber et al., 2021; Kerninon et al., 2021; Jiménez-Casero et al., 2023).

In terms of metric adequacy, the population and community metrics (lower limit depth, leaf area index, and epiphytic load) had strong loadings (higher than 0.72; Fig. 2) on the PCA first axis, which was supported by their high relationship with the PI. Gerakaris et al. (2017) highlighted the strongest relationships between population and community metrics and human pressures in their research. These types of metrics are more useful for assessing ecosystem conditions because they have longer response times and are more integrative (Roca et al., 2015; Romero et al., 2016). As for monitoring P. oceanica meadows, leaf area index (Pergent et al., 1995; Pergent-Martini et al., 2005; Koçak et al., 2011; Dural et al., 2012; Tursi et al., 2022b), lower limit depth (Meinesz & Laurent, 1978; Pergent et al., 1995; Pergent-Martini et al., 2005; Boudouresque et al., 2009; Descamp et al., 2011; Tursi et al., 2022b) and the epiphytic load (Pergent et al., 1995; Delgado et al., 1999; Dimech et al., 2002; Cancemi et al., 2003; Pergent-Martini et al., 2005) are commonly used to reflect the light condition, water transparency and nutrient loading. In fact, the site with the lowest EQR, El Djamila, is characterized by a set of human activities that are the primary cause of water turbidity and enrichment: a highly urbanized coastal line, proximity to an important river, and proximity to a fishing and pleasure port with the largest number of vessels in the area.

In contrast, biochemical and cellular metrics are less effective than population and community metrics in reflecting human pressure (p > 0.05), as also reported by Gerakaris *et al.* (2017). However, our PCAfindings show that these indicators can provide insight into the variability of meadow health (Fig. 2). The effectiveness of leaf tannin cell density and leaf nitrogen content as early indicators was perceived at Chenoua, the only site where the EQR assessed by the PREI was higher than that assessed by the MIAPo. While the three commonly used metrics had moderate values at Chenoua site, these two additional indicators showed signs of disturbance. This finding supports previous recommendations to include more sensitive and complex indicators, particularly those of biochemical and cellular types, for earlier detection of environmental stress (Ferrat et al., 2003). However, this pattern should be confirmed across a large number of sites and supplemented with in situ measurements of abiotic parameters. In fact, Martínez-Crego et al. (2010) highlighted the limitation of indices based solely on these types of metrics to assess the ecological integrity of the habitat and recommended a combination of metrics from the biochemical to the community level to offer a good diagnostic tool in the context of bioindication (Martínez-Crego et al., 2010; Personnic et al., 2014; Martinez-Haro et al., 2015; Romero et al., 2016). As a result, one of the benefits of the MIAPo is that it ensures the complementarity of metrics from community, population, biochemical, and cellular types to provide a comprehensive assessment of the ecosystem.

In terms of study area classification, the results were expected, and the sites of the capital, Algiers, were classified as the most disturbed zone, with the highest values of PI and the lowest mean EQRs with both PREI (Algiers: 0.54; Tipaza: 0.67) and MIAPo (Algiers: 0.62; Tipaza: 0.71), which is obviously due to the concentration of human activities in this area. To ensure the survival of *P. oceanica* meadows, it is strongly advised that measures be taken to reduce the impact of human activity in this area. Similarly, several studies using BiPo (Lopez y Royo *et al.*, 2010a; Boumaza *et al.*, 2015; Pergent *et al.*, 2015; Güreşen *et al.*, 2015; Gerakaris *et al.*, 2017; Rigo *et al.*, 2019)found a decrease in EQR values in urbanized and densely populated sites.

It is also important to consider sites with EQR values near the good/moderate boundary (EQR = 0.550), as recommended by Lopez y Royo et al. (2009b) and Romero et al. (2016). It is the case of the Bou Ismail site, which improves from moderate (with PREI) to good (with MIA-Po), and the Chenoua site, which has an MIAPo EQR of 0.555. If these sites are to be included in monitoring programs, they must be given special attention, and pressures must be precisely identified. The best EQR value obtained using both indices at the Kouali site corresponds to an undisturbed zone with low human pressure. The Kouali meadow has previously been examined and classified as having high primary productivity and vitality (Boumaza & Semroud, 2000; Boumaza et al., 2014). It was classified as having good/high ecological status when using P. oceanica meadow as a BQE (Boumaza et al., 2015; Sengouga, 2017; Sengouga et al., 2019) or the macroalgae community as a BQE for the rocky bottom quality (CFR) index (Anteur et al., 2024). This fact makes the site interesting for classification and protection (PNUE/PAM/CAR PAP, 2006; Mangos & Claudot, 2013; UNEP/MAP-SPA/RAC, 2021).

The results discussed above demonstrate that indices such as MIAPo and PREI, which use a smaller number of metrics, can be valuable and quick tools for monitoring and managing coastal ecosystems, providing critical insights for informed decision-making. However, in this study, some shortcomings were unavoidable. (i) The method for assessing human-induced pressure does provide a quick assessment of human impact, but the level of data aggregation is less precise than that used to collect meadow vitality data. To ensure its effectiveness, it is recommended that it be combined with other precise methods for assessing human pressure. (ii) The relatively small number of sites included in the study may have an impact on the robustness of our findings, especially when establishing the reference conditions. In fact, despite the varied environmental conditions that characterize the study area, no site has a bad ecological status. It appears that the various anthropogenic pressures recorded at these sites had no significant impact on the studied *P. oceanica* meadows.

To confirm the accuracy and robustness of MIAPo in reflecting the ecological status of Algerian water bodies, we aim to apply it to the entire Algerian coastline in sites with diverse environmental conditions (Martínez-Crego et al., 2010). Finally, the P. oceanica meadows must be protected in accordance with national (Law 02-2002; Law 2012: Executive decree no. 12-03) and regional legislation (Barcelona Convention 1976, amended 1995). Thus, non-destructive techniques are recommended to reduce the negative impact of sampling methods on P. oceanica meadows (Montefalcone, 2009). The Non-Destructive Sampling Methods (NDSM) developed by Gobert et al. (2020) have already been tested and approved for the PREI index. Future MIAPo applications will require a similar test because the selected metrics do not require rhizome sampling.

#### Conclusion

The multi-metric index MIAPo offers several advantages. It was built from different types of metrics, and it relates to human pressure and encompasses various time responses to stress. It is simple, quick, effective, and lowcost, meeting managers' and decision-makers needs and inter-calibration requirements. Its inter-calibration with the PREI index yields comparable results with enhanced reliability to human stressors by integrating biochemical and cellular metrics (Martínez-Crego et al., 2010), and it can be assessed using NDSM. However, further investigation is required for the index adoption: (i) a refinement of the approach used to define reference conditions to avoid eventual loss of precision in the classification of the ecological status; (ii) other population-type metrics should also be considered because they provide a more accurate picture of meadow health. Furthermore, it is recommended that the 'lower limit' indicator be calibrated, because most of our meadows exhibit lower limits associated with rocky substrates, controlled by hydrodynamic conditions, as described by Clabaut et al. (2010) in Corsica (France), the northern Gulf of Tunis (Tunisia), and El Kala National Park (Algeria); (iii) integration of environmental parameters measurement when assessing the pressure index, specifically the nitrogen content in interstitial water, in order to establish a relationship with the nitrogen content in tissues (Leoni et al., 2008). In this context, we also propose testing Specific Leaf Area (SLA) as a nutrient enrichment

indicator in P. oceanica meadows, as it has previously been associated with light attenuation and leaf photosynthetic rates (Olesen et al., 2002; Enríquez et al., 2004; Nicastro et al., 2015) and leaf nutrient variation in other magnoliophyte (Ainley et al., 2016); (iv) the tannin cell density in leaves should be compared to other parts of the plant, such as rhizomes, given their tendency to exhibit minor fluctuations in phenolic compounds and a notable response to effluent discharge (Migliore et al., 2007); (v) MIAPo should be applied on a larger scale for comparability and inter-calibration purposes, and its relationship with several human pressure assessment methods should be investigated. Additionally, a comparison with other indices that utilize a greater number of descriptors, such as the POMI or the CS Valencian, is necessary to enhance understanding of its efficacy and facilitate the harmonization of these indices.

# Acknowledgements

We are very grateful to Boukortt Redouane for his significant contributions to our scuba diving work. We also remember Nihal Bouledjouidja's exemplary commitment to laboratory work. It is with great sadness that we mark her passing at such a young age. Her memory will forever be honored.

#### References

- Ainley, L.B., Vergés, A., Bishop, M., 2016. Congruence of intraspecific variability in leaf traits for two co-occurring estuarine angiosperms. *Oecologia*, 181 (4), 1041-1053.
- Anteur, A.C., Bahbah, L., Bensari, B., Seridi, H., 2024. Evaluation of the ecological quality of the macroalgal communities along the Algerian coast (Algeria, Mediterranean Sea). *Regional Studies in Marine Science*, 78, 103767.
- Bakalem, A., Ruellet, T., Dauvin, J.-C., 2009. Benthic indices and ecological quality of shallow Algeria fine sand community. *Ecological indicators*, 9 (3), 395-408.
- Bellissimo, G., Sirchia, B., Ruvolo, V., 2020. Monitoring of *Posidonia oceanica* meadows in the Sicilian coasts under the Water Framework Directive (WFD). p. 510-518.
  In: 8th International Symposium "Monitoring of Mediterranean Coastal Areas. Problems and Measurement Techniques, Livorno, Italy, June 2020. Firenze University Press, Florence. (https://books.fupress.com/doi/capitoli/978-88-5518-147-1 51)
- Bennett, S., Roca, G., Romero, J., Alcoverro, T., 2011. Ecological status of seagrass ecosystems: An uncertainty analysis of the meadow classification based on the *Posidonia oceanica* multivariate index (POMI). *Marine Pollution Bulletin*, 62 (8), 1616-1621.
- Bentaallah, M., 2017. Contribution to the study of Posidonia oceanica through the West Algerian coasts: localisation, characterisation and use for coastal water status assessment. Ph.D. Thesis. University of Milano-Bicocca, Italy, 47 pp.
- Blouet, S., Lenfant, P., Dupuy de la Grandrive, R., Laffon, J-F.,

Chéré, E. et al., 2011. Mise en cohérence des méthodes de suivis des herbiers de posidonies des sites Natura 2000 marins du Languedoc-Roussillon. Rapport ADENACNRS/ EPHE/UPVD-CG66-AAMP, 49 pp.

- Bodéré, J.-C., Cribb, R., Curr, R., Davies, P., Hallégouët, B. et al., 1991. La gestion des milieux dunaires littoraux. Évolution de leur vulnérabilité à partir d'une liste de contrôle. Étude de cas dans le sud du Pays de Galles et en Bretagne occidentale. Norois, 151 (1), 279-298.
- Borja, A., Dauer, D. M., Grémare, A., 2012. The importance of setting targets and reference conditions in assessing marine ecosystem quality. *Ecological Indicators*, 12, 1-7.
- Boudouresque, C. F., Bernard, G., Pergent, G., Shili, A., Verlaque, M., 2009. Regression of Mediterranean seagrasses caused by natural processes and anthropogenic disturbances and stress: a critical review. *Botanica Marina*, 52 (5), 395-418.
- Boumaza, S., Semroud, R., 2000. Evaluation of past and present primary production in *Posidonia oceanica* beds from the cove of Kouali (Tipaza, Algeria) using the lepidochronological method. *Biologia marina mediterranea*, 7, 26-30.
- Boumaza, S., Mokeddem, Z., Hamoul, N., Semroud, R., 2012.
  Effects of disturbances caused by coastal discharges on phenolic compounds in the seagrass *Posidonia oceanica*. p. 34.
  In: 3rd Mediterranean Seagrass Workshop, Essaouira, Morocco, 28 May-1 June, 2012.
- Boumaza, S., Boudefoua, N., Boumaza, R., Semroud, R., 2014. Effects of urban effluents on spatial structure, morphology and total phenols of *Posidonia oceanica*: Comparison with a reference site. *Journal of experimental marine biology* and ecology, 457, 113-119.
- Boumaza, S., Sengouga, A., Misraoui, A., Boumaour, A., Kahlal, M. et al., 2015. Assessment of the ecological status of Algiers coastal waters using two indices based on the seagrass Posidonia oceanica: PREI and BiPo. In: 4th Mediterranean Seagrass Workshop - MSW '15, Oristano, Italy,18-22 May, 2015. (https://mediterranean.seagrassonline.org/sardinia/)
- Boumaza, S., Sengouga, A., Gasmi, D., Misraoui, A., Pergent, G. et al., 2022. Patterns of *Posidonia oceanica* (L.) Delile tannin cells and leaf characteristics according to environmental conditions. *Mediterranean Marine Science*, 23 (4), 879-889.
- Buia, M., Gambi, M. C., Dappiano, M., 2004. Seagrass systems. Biologia Marina Mediterranea, 11 (Suppl 1), 133-183.
- Cancemi, G., Falco, G. D., Pergent, G., 2003. Effects of organic matter input from a fish farming facility on a *Posidonia* oceanica meadow. *Estuarine, Coastal and Shelf Science*, 56 (5-6), 961-968.
- Clabaut, P., Pergent-Martini, C., Pergent, G., Augris, C., Pasqualini, V. et al., 2010. Les relations entre les herbiers de Posidonies, le substrat et l'hydrodynamisme. p. 49-54. In: Proceedings of the 4th Mediterranean Symposium on Marine Vegetation. Yasmine-Hammamet, 2-4 December 2010. RAC/SPA publ,Tunis. (https://www.rac-spa.org/node/670)
- Comeleo, R., Paul, J., August, P., Copeland, J., Baker, C. et al., 1996. Relationships Between Watershed Stressors and Sediment Contamination in Chesapeake Bay Estuaries. Landscape Ecology, 11, 307-319.
- Dallas, H.F., 2021. Rapid Bioassessment Protocols Using Aquatic Macroinvertebrates in Africa-Considerations for

Regional Adaptation of Existing Biotic Indices. *Frontiers in Water*, 3, 628227.

- Dawson, S.P., Dennison, W.C., 1996. Effects of ultraviolet and photosynthetically active radiation on five seagrass species. *Marine Biology*, 125 (4), 629-638.
- Delgado, O., Ruiz, J., Pérez, M., Romero, J., Ballesteros, E., 1999. Effects of fish farming on seagrass (*Posidonia oceanica*) in a Mediterranean bay: seagrass decline after organic loading cessation. *Oceanologica Acta*, 22 (1), 109-117.
- Descamp, P., Holon, F., Ballesta, L., Guilbert, A., Guillot, M. et al., 2011. Fast and easy method for seagrass monitoring: Application of acoustic telemetry to precision mapping of *Posidonia oceanica* beds. *Marine Pollution Bulletin*, 62 (2), 284-292.
- Di Camillo, C., Ponti, M., Storari, A., Scarpa, C., Roveta, C. et al., 2023. Review of the indexes to assess the ecological quality of coralligenous reefs: towards a unified approach. *Frontiers in Marine Science*, 10, 1252969.
- Dimech, M., Borg, J. A., Schembri, P., 2002. Changes in the structure of a *Posidonia oceanica* meadow and in the diversity of associated decapod, mollusc and echinoderm assemblages, resulting from inputs of waste from a marine fish farm (Malta, Central Mediterranean). *Bulletin of Marine Science*, 71, 1309-1321.
- Duarte, C., 1990. Seagrass nutrient content. *Marine Ecology Progress Series*, 67, 201-207.
- Dumay, O., Costa, J., Desjobert, J.-M., Pergent, G., 2004. Variations in the concentration of phenolic compounds in the seagrass *Posidonia oceanica* under conditions of competition. *Phytochemistry*, 65 (24), 3211-3220.
- Dural, B., Aysel, V., Demir, N., Yazici, İ., Erduğan, H., 2012. The status of sensitive ecosystems along the Aegean coast of Turkey: *Posidonia oceanica* (L.) Delile meadows. *Journal of Black Sea / Mediterranean Environment*, 18 (3), 360-379.
- Enríquez, S., Marbà, N., Cebriàn, J., Duarte, C.M., 2004. Annual variation in leaf photosynthesis and leaf nutrient content of four Mediterranean seagrasses. *Botanica marina*, 47 (4), 295-306.
- Fernández Torquemada, Y., Díaz-Valdés, M., Colilla, F., Luna, B., Sánchez Lizaso, J. *et al.*, 2008. Descriptors from *Po-sidonia oceanica* (L.) Delile meadows in coastal waters of Valencia, Spain, in the context of the EU Water Framework Directive. *ICES Journal of Marine Science*, 65 (8), 1492-1497.
- Ferrat, L., Pergent-Martini, C., Roméo, M., 2003. Assessment of the use of biomarkers in aquatic plants for the evaluation of environmental quality: application to seagrasses. *Aquatic Toxicology*, 65 (2), 187-204.
- Fourqurean, J.W., Marbà, N., Duarte, C.M., Diaz-Almela, E., Ruiz-Halpern, S., 2007. Spatial and temporal variation in the elemental and stable isotopic content of the seagrasses *Posidonia oceanica* and *Cymodocea nodosa* from the Illes Balears, Spain. *Marine Biology*, 151, 219-232.
- García-Marín, P., Cabaço, S., Hernández, I., Vergara, J. J., Silva, J. et al., 2013. Multi-metric index based on the seagrass Zostera noltii (ZoNI) for ecological quality assessment of coastal and estuarine systems in SW Iberian Peninsula. Marine Pollution Bulletin, 68 (1), 46-54.
- Gerakaris, V., Panayotidis, P., Vizzini, S., Nicolaidou, A.,

Economou-Amilli, A., 2017. Effectiveness of *Posidonia oceanica* biotic indices for assessing the ecological status of coastal waters in Saronikos Gulf (Aegean Sea, Eastern Mediterranean). *Mediterranean Marine Science*, 18 (1), 161-178.

- Giraud, G., 1979. Sur une méthode de mesure et de comptage des structures foliaires de *Posidonia oceanica* (Linnaeus) Delile. *Bulletin de la Musee d'Histoire Naturelle de Marseille*, 39, 33-39.
- Gobert, S., Lefebvre, L., Boissery, P., Richir, J., 2020. A non-destructive method to assess the status of *Posidonia oceanica* meadows. *Ecological Indicators*, 119, 106838.
- Gobert, S., Sartoretto, S., Rico-Raimondino, V., Andral, B., Chery, A. *et al.*, 2009. Assessment of the ecological status of Mediterranean French coastal waters as required by the Water Framework Directive using the *Posidonia oceanica* Rapid Easy Index: PREI. *Marine Pollution Bulletin*, 58 (11), 1727-1733.
- Güreşen, A., Güreşen, S. O., Aktan, Y., 2020a. Combined synthetic and biotic indices of *Posidonia oceanica* to qualify the status of coastal ecosystems in the North Aegean. *Ecological Indicators*, 113, 106149.
- Güreşen, A., Pergent, G., Güreşen, S. O., Aktan, Y., 2020b. Evaluating the coastal ecosystem status of two Western and Eastern Mediterranean islands using the seagrass *Posidonia oceanica*. *Ecological Indicators*, 108, 105734.
- Hallegouët, B., Hénaff, A., Morel, V., Yoni, C., 1997. Dynamiques économiques et anthropisation des bords de mer : impacts sur les sociétés et les espaces littoraux armoricains / Economie dynamics and anthropization of the sea-side : the impacts upon societies and coastal spaces of Brittany. *Cahiers Nantais*, 47 (1), 377-386.
- Haouchine, A., Haouchine, F. Z., Labadi, A., 2015. Changements climatiques et activités anthropiques: impacts sur les aquifères côtiers en Algérie. *LARHYSS Journal*, (24), 227-241.
- Hawkins, C. P., Olson, J. R., Hill, R. A., 2010. The reference condition: predicting benchmarks for ecological and water-quality assessments. *Journal of the North American Benthological Society*, 29 (1), 312-343.
- Helber, S. B., Procaccini, G., Belshe, E. F., Santillan-Sarmiento, A., Cardini, U. *et al.*, 2021. Unusually Warm Summer Temperatures Exacerbate Population and Plant Level Response of *Posidonia oceanica* to Anthropogenic Nutrient Stress. *Frontiers in Plant Science*, 12, 662682.
- Jiménez-Casero, J., Belando, M.D., Bernardeau-Esteller, J., Marín-Guirao, L., García-Muñoz, R. *et al.*, 2023. A Critical Gap in Seagrass Protection: Impact of Anthropogenic Off-Shore Nutrient Discharges on Deep *Posidonia oceanica* Meadows. *Plants*, 12 (3), 457.
- Jones, B.L., Cullen-Unsworth, L.C., Unsworth, R.K.F., 2018. Tracking Nitrogen Source Using δ15N Reveals Human and Agricultural Drivers of Seagrass Degradation across the British Isles. *Frontiers in Plant Science*, 9, 133.
- Kerninon, F., Payri, C. E., Le Loc'h, F., Alcoverro, T., Maréchal, J.-P. et al., 2021. Selection of parameters for seagrass management: Towards the development of integrated indicators for French Antilles. *Marine Pollution Bulletin*, 170, 112646.
- Kletou, D., Kleitou, P., Savva, I., Attrill, M. J., Charalambous, S. *et al.*, 2020. Seagrass of Vasiliko Bay, Eastern Mediterra-

nean: Lost Cause or Priority Conservation Habitat? *Journal* of Marine Science and Engineering, 8 (9), 717.

- Koçak, F., Uluturhan, E., Gier, G.Y., Önen, S.A., 2011. Impact of environmental conditions on *Posidonia oceanica* meadows in the Eastern Mediterranean Sea. *Indian Journal of Geo-Marine Sciences*, 40 (6), 770-778.
- Larid, M., 2002. Le recul des plages en Algérie : problèmes et perspectives. p. 53-57. In: *CIESM workshp series n°18,* Érosion littorale en Méditerranée occidentale, Tanger, 18-21 September, 2002.
- Legendre, P., Legendre, L. (Ed.), 1998. *Numerical Ecology*. Elsevier, Amsterdam, 853 pp.
- Leoni, V., Pasqualini, V., Pergent-Martini, C., Vela, A., Pergent, G., 2007. Physiological responses of *Posidonia oceanica* to experimental nutrient enrichment of the canopy water. *Journal of experimental marine biology and ecology*, 349 (1), 73-83.
- Leoni, V., Vela, A., Pasqualini, V., Pergent-Martini, C., Pergent, G., 2008. Effects of experimental reduction of light and nutrient enrichments (N and P) on seagrasses: a review. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 18 (2), 202-220.
- Leopold, L.B., 1969. Quantitative comparison of some aesthetic factors among rivers. Vol.620. US Geological Survey, Washington, 16 pp.
- Lopez y Royo, C., Silvestri, C., Pergent, G., Casazza, G., 2009a. Assessing human-induced pressures on coastal areas with publicly available data. *Journal of Environmental Management*, 90 (3), 1494-1501.
- Lopez y Royo, C., Silvestri, C., Salivas-Decaux, M., Pergent, G., Casazza, G., 2009b. Application of an angiosperm-based classification system (BiPo) to Mediterranean coastal waters: Using spatial analysis and data on metal contamination of plants in identifying sources of pressure. *Hydrobiologia*, 633, 169-179.
- Lopez y Royo, C., Casazza, G., Pergent-Martini, C., Pergent, G., 2010a. A biotic index using the seagrass *Posidonia oceanica* (BiPo), to evaluate ecological status of coastal waters. *Ecological Indicators*, 10 (2), 380-389.
- Lopez y Royo, C., Pergent, G., Pergent-Martini, C., Casazza, G., 2010b. Seagrass (*Posidonia oceanica*) monitoring in western Mediterranean: implications for management and conservation. *Environmental Monitoring and Assessment*, 171, 365-380.
- Lopez y Royo, C., Pergent, G., Alcoverro, T., Buia, M., Casazza, G. *et al.*, 2011. The seagrass *Posidonia oceanica* as indicator of coastal water quality: Experimental intercalibration of classification systems. *Ecological Indicators*, 11 (2), 557-563.
- Mancini, I., Rigo, I., Oprandi, A., Montefalcone, M., Morri, C. et al., 2020. What biotic indices tell us about ecosystem change: lessons from the seagrass *Posidonia oceanica*. Vie et Milieu / Life & Environment, 70.
- Mangos, A., Claudot, M.A., 2013. *Economic study of the impacts of marine and coastal protected areas in the Mediterranean*. Plan Bleu, Valbonne, Plan Bleu, No 13, 58 pp.
- Martínez-Crego, B., Alcoverro, T., Romero, J., 2010. Biotic indices for assessing the status of coastal waters: A review of strengths and weaknesses. *Journal of environmental monitoring*, 12, 1013-28.

- Martinez-Haro, M., Beiras, R., Bellas, J., Capela, R., Coelho, J.P. *et al.*, 2015. A review on the ecological quality status assessment in aquatic systems using community based indicators and ecotoxicological tools: what might be the added value of their combination? *Ecological Indicators*, 48, 8-16.
- Mascaró, O., Bennett, S., Marbà, N., Nikolić, V., Romero, J. et al., 2012. Uncertainty analysis along the ecological quality status of water bodies: The response of the *Posidonia* oceanica multivariate index (POMI) in three Mediterranean regions. *Marine Pollution Bulletin*, 64 (5), 926-931.
- Meinesz, A., Laurent, R., 1978. Cartographie et état de la limite inferieure de l'herbier de *Posidonia oceanica* dans les Alpes-maritimes (France) - Campagne Poseïdon 1976. *Botanica marina*, 21 (8), 513-526.
- Migliore, L., Rotini, A., Randazzo, D., Albanese, N. N., Giallongo, A., 2007. Phenols content and 2-D electrophoresis protein pattern: a promising tool to monitor Posidonia meadows health state. *BMC ecology*, 7 (1), 1-8.
- Montefalcone, M., 2009. Ecosystem health assessment using the Mediterranean seagrass *Posidonia oceanica*: A review. *Ecological Indicators*, 9 (4), 595-604.
- Montefalcone, M., Albertelli, G., Nike Bianchi, C., Mariani, M., Morri, C., 2006. A new synthetic index and a protocol for monitoring the status of *Posidonia oceanica* meadows: a case study at Sanremo (Ligurian Sea, NW Mediterranean). *Aquatic Conservation: Marine and Freshwater Ecosystems*, 16 (1), 29-42.
- Nicastro, S., Innocenti, A. M., Passalacqua, N. G., 2015. Histo-anatomical leaf variations related to depth in *Posidonia oceanica*. *Functional Plant Biology*, 42 (4), 418-422.
- Olesen, B., Enríquez, S., Duarte, C., Sand-Jensen, K., 2002. Depth-acclimation of photosynthesis, morphology and demography of *Posidonia oceanica* and *Cymodocea nodosa* in the Spanish Mediterranean Sea. *Marine Ecology Progress Series*, 236, 89-97.
- Oliva, S., Mascaró, O., Llagostera, I., Pérez, M., Romero, J., 2012. Selection of metrics based on the seagrass *Cymodocea nodosa* and development of a biotic index (CYMOX) for assessing ecological status of coastal and transitional waters. *Estuarine, Coastal and Shelf Science*, 114, 7-17.
- Oprandi, A., Bianchi, C.N., Karayali, O., Morri, C., Rigo, I. *et al.*, 2021. RESQUE: A novel comprehensive approach to compare the performance of different indices in evaluating seagrass health. *Ecological Indicators*, 131, 108118.
- Panayotidis, P., Boudouresque, C.F., Marcot-Coqueugniot, J., 1981. Microstructure de l'herbier de *Posidonia oceanica* (Linnaeus) Delile. Microstructure of *Posidonia oceanica* (Linnaeus) beds. *Botanica Marina*, 24 (3), 115-124.
- Parravicini, V., Rovere, A., Vassallo, P., Micheli, F., Montefalcone, M. *et al.*, 2012. Understanding relationships between conflicting human uses and coastal ecosystems status: A geospatial modeling approach. *Ecological Indicators*, 19, 253-263.
- Pérez, M., García, T., Invers, O., Ruiz, J.M., 2008. Physiological responses of the seagrass *Posidonia oceanica* as indicators of fish farm impact. *Marine Pollution Bulletin*, 56 (5), 869-879.
- Pergent, G., Pergent-Martini, C., Boudouresque, C., 1995. Utilisation de l'herbier à *Posidonia oceanica* comme indicateur biologique de la qualité du milieu littoral en Méditer-

ranée: État des connaissances. Mésogée, 54, 3-27.

- Pergent-Martini, C., Leoni, V., Pasqualini, V., Ardizzone, G.D., Balestri, E. *et al.*, 2005. Descriptors of *Posidonia oceanica* meadows: use and application. *Ecological Indicators*, 5 (3), 213-230.
- Pergent, G., Pergent-Martini, C., Bein, A., Dedeken, M., Oberti, P. et al., 2015. Dynamic of *Posidonia oceanica* seagrass meadows in the northwestern Mediterranean: Could climate change be to blame? *Comptes Rendus Biologies*, 338 (7), 484-493.
- Personnic, S., Boudouresque, C.F., Astruch, P., Ballesteros, E., Blouet, S. *et al.*, 2014. An ecosystem-based approach to assess the status of a Mediterranean ecosystem, the *Posidonia oceanica* seagrass meadow. *PloS one*, 9 (6), e98994.
- PNUE/PAM CAR/ASP, 2019. Les protocoles de surveillance des Indicateurs communs 1 et 2 relatifs aux habitats marins. UNEP/MED, Rome, Italie, No WG.474/3, 160 pp.
- PNUE/PAM/CAR PAP., 2006. Programme d'Aménagement Côtier (PAC) de la «zone côtière algéroise». Rapport final intégré. MAP Technical Reports Series, PNUE, Athena, No 164, 190 pp.
- Rende, F., Bacci, T., Penna, M., Trabucco, B., Giovanardi, F. et al., 2011. WFD 2000/60/EC: Posidonia oceanica meadows in the evaluation of ecological status of the Italian coastal waters/WFD 2000/60 CE: Le praterie di Posidonia oceanica nella valutazione dello stato ecologico delle acque costiere Italiane. Biologia Marina Mediterranea, 18 (1), 322-323.
- Rigo, I., Montefalcone, M., Morri, C., 2019. Use of ecological indices to assess the health status of *Posidonia oceanica* meadows in Eastern Liguria. p. 271-280. In: *Planning, nature and ecosystem services*. Gargiulo, C., Zoppi, C. (Eds). Federico II Open Access University Press, Naples.
- Roca, G., Alcoverro, T., de Torres, M., Manzanera, M., Martínez-Crego, B. *et al.*, 2015. Detecting water quality improvement along the Catalan coast (Spain) using stress-specific biochemical seagrass indicators. *Ecological Indicators*, 54, 161-170.
- Rodriguez, W., August, P., Wang, Y., Paul, J., Gold, A. *et al.*, 2007. Empirical relationships between land use/cover and estuarine condition in the Northeastern United States. *Landscape Ecology*, 22, 403-417.
- Romero, J., Martínez-Crego, B., Alcoverro, T., Pérez, M., 2007. A multivariate index based on the seagrass *Posidonia oceanica* (POMI) to assess ecological status of coastal waters under the water framework directive (WFD). *Marine Pollution Bulletin*, 55 (1), 196-204.
- Romero, J., Alcoverro, T., Roca, G., Pérez, M., 2016. Bioindicators, Monitoring, and Management Using Mediterranean Seagrasses: What Have We Learned from the Implementation of the EU Water Framework Directive? p. 161-182. In: *Experiences from Ground, Coastal and Transitional Water Quality Monitoring*. Munné, A., Ginebreda, A., Prat, N. (Eds). Springer International Publishing, Cham.
- Rotini, A., Belmonte, A., Barrote, I., Micheli, C., Peirano, A. et al., 2013. Effectiveness and consistency of a suite of descriptors for assessing the ecological status of seagrass meadows (*Posidonia oceanica* L. Delile). Estuarine, Coastal and Shelf Science, 130, 252-259.
- Scartazza, A., Moscatello, S., Gavrichkova, O., Buia, M.C.,

Lauteri, M. *et al.*, 2017. Carbon and nitrogen allocation strategy in *Posidonia oceanica* is altered by seawater acidification. *Science of The Total Environment*, 607-608, 954-964.

- Sengouga, A., 2017. Evaluation de l'état écologique des eaux côtières algéroises par l'utilisation d'un indice basé sur Posidonia oceanica. Magistère Thesis. École Nationale Supérieure des Sciences de la Mer et de l>Aménagement du Littoral, Algiers, Algeria, 88 pp.
- Sengouga, A., Boumaza, S., Misraoui, A., Semroud, R., 2018. Evaluation de l'état écologique des eaux côtières algéroises par l'utilisation d'indices basés sur *Posidonia oceanica*. In: *3ème Workshop international sur la Biodiversité et la Gestion des Ecosystèmes Exploités BGEE, Algiers, Algeria, 27 October 2018.*
- Sengouga, A., Boumaza, S., Zerrouk, C., Misraoui, A., Semroud, R., 2019. Temporal Evolution of Ecological Status in Algerian Coastal Waters. p. 337-378. In: 14<sup>th</sup>International MEDCOAST Congress on Coastal and Marine Sciences, Engineering, Management and Conservation. Marmaris, Turkey, 22-26, October 2019. Mediterranean Coastal Foundation (MEDCOAST 19), Ortaca, Mugla, Turkey.
- Stoddard, J.L., Herlihy, A.T., Peck, D.V., Hughes, R.M., Whittier, T.R. *et al.*, 2008. A process for creating multimetric indices for large-scale aquatic surveys. *Journal of the North American Benthological Society*, 27 (4), 878-891.
- Terrados, J., Pons, F.J.M., 2008. Epiphyte load on the seagrass Posidonia oceanica (L.) Delile does not indicate anthropogenic nutrient loading in Cabrera Archipelago National Park (Balearic Islands, Western Mediterranean). Scientia

Marina, 72 (3), 503-510.

- Tursi, A., Mastrototaro, F., Bottalico, A., Montesanto, F., Chimienti, G., 2022a. Assessing the conservation status of *Posidonia oceanica* at Tremiti Archipelago (Adriatic Sea): a comparative analysis of different classification systems. pp. 262-266. In: *The 2022 IEEE International Workshop on Metrology for the Sea; Learning to Measure Sea Health Parameters (MetroSea) October 2022.* IEEE.
- Tursi, A., Mastrototaro, F., Montesanto, F., De Giosa, F., Lisco, A. *et al.*, 2022b. The Status of *Posidonia oceanica* at Tremiti Islands Marine Protected Area (Adriatic Sea). *Biology*, 11 (6), 923.
- Udy, J.W., Dennison, W.C., 1997. Growth and physiological responses of three seagrass species to elevated sediment nutrients in Moreton Bay, Australia. *Journal of Experimental Marine Biology and Ecology*, 217 (2), 253-277.
- UNEP/MAP-SPA/RAC., 2021. Conservation de la biodiversité marine et côtière méditerranéenne d'ici 2030 et au-delà By A. Chalabi. SPA/RAC Tunisia, 140 pp.
- Walker, D.I., Campey, M.L., Kendrick, G.A., 2004. Nutrient dynamics in two seagrass species, *Posidonia coriacea* and *Zostera tasmanica*, on Success Bank, Western Australia. *Estuarine, Coastal and Shelf Science*, 60 (2), 251-260.
- Zhang, J.-P., Huang, X.-P., Jiang, Z.-J., Thorhaug, A., 2011. Comparison of the role of the foliar sheath in nutrient (ammonium and phosphate) acquisition by the seagrass *Thalassia hemprichii* (Ehrenb.) Aschers. at two different sites on tropical Hainan Island, China. *Hydrobiologia*, 669 (1), 45-61.