

Mediterranean Marine Science

early view



Non-indigenous and cryptogenic marine species in the Port of Algiers (Western Mediterranean)

BILEL BENSARI, LAMIA BAHBAH, ADEL KACIMI,
RYM BACHETARZI, NOUR EL ISLAM BACHARI

doi: [10.12681/mms.39689](https://doi.org/10.12681/mms.39689)

To cite this article:

BENSARI, B., BAHBAH, L., KACIMI, A., BACHETARZI, R., & BACHARI, N. E. I. (2025). Non-indigenous and cryptogenic marine species in the Port of Algiers (Western Mediterranean). *Mediterranean Marine Science*, early view. <https://doi.org/10.12681/mms.39689>

Non-indigenous and cryptogenic marine species in the Port of Algiers (Western Mediterranean)

Bilel BENSARI¹, Lamia BAHBAH^{1,2}, Adel KACIMF³, Rym BACHETARZI¹, and Nour El Islam BACHARI¹

¹Biological Oceanography and Marine Environment Laboratory, Faculty of Biological Science, University of Science and Technology, Houari Boumediene. B.P 32 El Alia 16111 Bab Ezzouar, Algeria

²National Higher School of Marine Sciences and Coastal Management, 16320 Algiers, Algeria

³Marine and Coastal Ecosystems Laboratory (ECOSYSMarL), Department of Marine and Coastal Environment, National Higher School of Marine Sciences and Coastal Management, 16320 Algiers, Algeria

Corresponding author: Bilel BENSARI; bilel.bensari_fsb@usthb.edu.dz

Contributing Editor: Argyro ZENETOS

Received: 11 December 2024; Accepted: 09 May 2025; Published online: 12 June 2025

Abstract

This study addresses the growing concern of non-indigenous and cryptogenic marine species in the Mediterranean Sea, particularly in high-risk zones such as ports. The Port of Algiers, a significant maritime hub in Algeria, was selected as the study site to document the presence of such species due to its susceptibility to introductions through vessel traffic. Sampling was conducted from 2021 to 2024 across various submerged structures, including mooring buoys and ship fenders. Specimens were collected by scraping surfaces and preserved for identification, which was performed to the lowest taxonomic level possible. The results revealed 18 distinct species, of which 13 were non-indigenous and five cryptogenic. Notably, 11 of these species are recorded for the first time on the Algerian coast. Taxonomically, the species included Ascidiacea (six species), Bryozoa (five species), Crustacea (three species), Polychaeta (three species), and Porifera (one species). This research contributes to the baseline knowledge of non-indigenous and cryptogenic species in Algerian marine environments, providing essential data to support further ecological and biogeographic studies on species introductions in the Mediterranean Sea.

Keywords: Non-indigenous species; Cryptogenic species; Fouling species; Algerian coast; Port.

Introduction

The introduction of non-indigenous marine species (NIS) has become a significant environmental issue worldwide, with profound ecological, economic, and social implications (Ruiz *et al.*, 1997). Non-indigenous species often compete with native species, alter habitats, and disrupt local ecosystems, leading to biodiversity loss and ecosystem imbalances (Katsanevakis *et al.*, 2014). In the Mediterranean Sea, one of the world's most invaded regions, the problem is particularly pronounced due to the basin's high volume of maritime traffic, diverse habitats, and connectivity with the Atlantic Ocean and the Red Sea through the Suez Canal (D'Amen & Azzurro, 2020; Katsanevakis *et al.*, 2014; Zenetos *et al.*, 2010). This connectivity facilitates the spread of species from other regions, making the Mediterranean region a hotspot for biological invasions.

Among the primary pathways of NIS introduction, maritime vessels play a crucial role, mainly through ballast water discharge and hull fouling (Bouda *et al.*, 2016, 2018; Gollasch, 2002). Ports, as key entry points for shipping, are thus critical zones for NIS risk assessment,

as they can serve as initial settlement areas for NIS potentially able to become invasive (Tempesti *et al.*, 2020; Ulman *et al.*, 2017). Monitoring these high-risk areas is essential for managing and mitigating the arrival and establishment of NIS.

Despite the Mediterranean's status as an invasion-prone sea, the study and monitoring of non-indigenous species in Algeria remain limited (Grimes *et al.*, 2018), particularly in ports. To date, only one study has investigated NIS in Algerian ports, highlighting a gap in consistent monitoring and research on this critical issue (Bensari *et al.*, 2020). This lack of comprehensive data presents challenges in developing effective management and conservation strategies to address biological invasions.

The present study aims to fill this knowledge gap by reporting on several NIS observed in the Port of Algiers, contributing valuable data to the limited body of research on this subject. Through this research, we aim to enhance understanding of NIS dynamics in Algerian waters and emphasise the need for regular monitoring to formulate management actions and minimise the impact of future invasions.

Material and Methods

The study was conducted at the Port of Algiers (Fig. 1), one of the oldest ports in Algeria. Various submerged structures, including ropes, ship fender tires, steel dock posts, and the quay surface, were selected as sampling sites. These structures, located at depths not exceeding 2 m, were chosen based on their potential to support a diverse range of marine species. The characteristics of each station, including coordinates, substrate type, and sampling dates, are summarised in Table 1. To optimise species detection, multiple sampling campaigns were conducted. Sampling focused on structures that could be lifted onto the quay, as well as the upper part of the quay. Additionally, an underwater camera was submerged a few centimeters to capture images of species adhering to the quay surface.

Specimens were collected by scraping them off sub-

merged surfaces by using a putty knife, a method that facilitated retrieval while minimizing damage. The sampled organisms were immediately preserved in a 5% formalin solution to maintain their integrity. Although the precise sampling area was not measured, species abundance was estimated qualitatively as low, medium, or high.

Some specimens were identified on-site, which were recognised based on clearly visible morphology. Other specimens were identified using a stereomicroscope and a microscope in the laboratory, while additional identifications were made from photographs. Species identified through photographs or on-site were generally ascidians. Taxonomic keys and expert knowledge were used to identify specimens to the lowest feasible taxonomic level. The nomenclature of identified species was verified according to the World Register of Marine Species (WoRMS Editorial Board, 2024).



Fig. 1: Location of the study site (area outlined with dashed lines) within the Port of Algiers.

Table 1. Location and characteristics of sampling stations.

Station	Coordinates	Substrate	Dates
Station 1	36.78305° N, 3.06427° E	Tire	29/04/2021
Station 2	36.78306° N, 3.06427° E	Rope	
Station 3	36.78329° N, 3.06434° E	Tire	09/05/2022
Station 4	36.78334° N, 3.06436° E	Rope	
Station 5	36.78363° N, 3.06445° E	Tire	26/05/2023
Station 6	36.78362° N, 3.06445° E	Rope	
Station 7	36.78488° N, 3.06549° E	Dock Post	
Station 8	36.78348° N, 3.06441° E	Quay	14/09/2024
Station 9	36.78401° N, 3.06603° E	Tire	

Results

We identified 18 non-indigenous species within the Port of Algiers. Among these, 13 are certainly non-native to the Mediterranean Sea, while five are classified as cryptogenic (Carlton, 1996), meaning their origins remain uncertain (Table 2, Figs. 2 and 3). Remarkably, 11 of these species are documented here for the first time on the Algerian coastline. Ascidiacea comprise the largest group with six species, followed by Bryozoa with five species. Crustacea and Polychaeta each include three species, while Porifera is represented by one species.

Ascidiacea

Botrylloides cf. niger Herdman, 1886

Botrylloides niger is a cosmopolitan colonial tunicate commonly found in tropical and warm water regions. In the Caribbean region, its distribution is nearly uniform, and its genetic diversity is highest, suggesting that it may be native to this region (Sheets *et al.*, 2016). Part of the uncertainty surrounding its origin and distribution arises from its taxonomic confusion with similar species. In the Mediterranean, the species is confirmed in Italy (Della Sala *et al.*, 2022), Turkey (Temiz *et al.*, 2023), and Egypt (Halim & Abdel Messeih, 2016). However, in Spain (Carmona-Rodríguez *et al.*, 2024) and Tunisia (Mnasri-Afifi *et al.*, 2024), it was reported as *B. cf. niger*, indicating

uncertainty in its identification. In Morocco (Mghili *et al.*, 2024), its presence remains questionable. Its distribution in the Mediterranean is likely more extensive, as it is often confused with *Botrylloides leachii* (Savigny). On the iNaturalist platform (consulted on 8 November 2024), this species has been observed 210 times by 30 different observers in the Mediterranean. These observations include sightings in Spain (including Palma), Morocco, Algeria, France, Italy (including Sardinia), Tunisia, Malta, Albania, Greece, and Cyprus.

Almost all specimens photographed closely resemble the “orange-brown” morphotype described by Temiz *et al.* (2023). We identified two morphotypes, the “orange-brown” and the “orange”, with the former being the most abundant (Fig. 3D). Additionally, this species exhibited a very high coverage rate. Since 2021, it has been observed at Alger Plage on the eastern side of the Bay of Algiers (personal observation).

Botrylloides cf. violaceus Oka, 1927

Botrylloides violaceus, first described in Japan in 1927, is native to the Northwest Pacific region, from northern Japan to southern Korea and northern China (Rho *et al.*, 2000). It is now more widely reported in the Northeast Pacific and the North Atlantic (Bock *et al.*, 2011). This colonial tunicate predominantly inhabits artificial substrates such as dock floats, pilings, piers, aquaculture structures, and boat hulls (Simkanin *et al.*, 2012).

Table 2. Non-Indigenous and Cryptogenic Marine Species Identified in the Port of Algiers: First Record, Abundance, Coordinates, and Substrate. **Cat:** Category, **NIS:** Non-Indigenous Species, **DOFF:** date of first found, **Cry:** Cryptogenic **Q:** Quay, **R:** Rope, **T:** Tire, **D:** Dock post.

Taxonomic group	Species	Cat	DOFF	Abun	Coordinates	Substrat			
						Q	R	T	D
Ascidiacea	<i>Botrylloides niger</i>	NIS	29/04/2021	High	36.78305° N, 3.06427° E	+	+	+	+
	<i>Botrylloides violaceus</i>	NIS	26/05/2023	Medium	36.78488° N, 3.06549° E	+	+		+
	<i>Botryllus S chlosseri</i>	Cryp	29/04/2021	High	36.78305° N, 3.06427° E	+	+	+	+
	<i>Polyclinum constellatum</i>	NIS	14/09/2024	Medium	36.78348° N, 3.06441° E	+	-	-	-
	<i>Styela canopus</i>	Cryp	14/09/2024	Low	36.78401° N, 3.06603° E	+	-	+	-
	<i>Styela plicata</i>	NIS	29/04/2021	High	36.78305° N, 3.06427° E	+	+	+	-
Bryozoa	<i>Bugula neritina</i>	Cryp	29/04/2021	High	36.78305° N, 3.06427° E	+	+	+	+
	<i>Bugulina stolonifera</i>	Cryp	29/04/2021	Low	36.78305° N, 3.06427° E	+	-	+	-
	<i>Celleporaria brunnea</i>	NIS	26/05/2023	Low	36.78363° N, 3.06445° E	-	-	+	-
	<i>Schizoporella errata</i>	Cryp	09/05/2022	High	36.78305° N, 3.06427° E	+	-	+	-
	<i>Tricellaria inopinata</i>	NIS	29/04/2021	Medium	36.78305° N, 3.06427° E	-	-	+	+
Crustacea	<i>Balanus trigonus</i>	NIS	14/09/2024	Medium	36.78401° N, 3.06603° E	-	-	+	-
	<i>Jassa slatteryi</i>	Cryp	29/04/2021	Low	36.78305° N, 3.06427° E	-	-	+	-
	<i>Zeuxo coralensis</i>	NIS	29/04/2021	Low	36.78305° N, 3.06427° E	-	-	+	-
Polychaeta	<i>Branchiommma bairdi</i>	NIS	29/04/2021	High	36.78305° N, 3.06427° E	+	+	+	-
	<i>Branchiommma luctuosom</i>	NIS	29/04/2021	High	36.78305° N, 3.06427° E	+	+	+	-
	<i>Hydroides elegans</i>	NIS	29/04/2021	High	36.78305° N, 3.06427° E	+	+	+	-
Porifera	<i>Paraleucilla magna</i>	NIS	29/04/2021	High	36.78305° N, 3.06427° E	+	+	+	+




Fig. 2: NIS and cryptogenic species found in this study. A: *Bugulina stolonifera*, B: *Bugula neritina*, C: *Tricellaria inopinata*, D: *Zeuxo coralensis*, E: *Styela plicata*, F: *Paraleucila magna*, G: *Branchiomma luctuosum*, H: *Schizoporella errata*, I: *Styela canopus*.

In the Mediterranean, this ascidian was first collected in 1993 in the Venetian Lagoon (Zaniolo *et al.*, 1998). Since its first record, it was reported Lake Faro and in Messina, Italy (Palanisamy *et al.*, 2018), as well as in France without a specified location (Massé *et al.*, 2023). This is the first report of this species in Algeria (Fig. 3B).

Botryllus schlosseri (Pallas, 1766)

The tunicate *B. schlosseri*, which is globally distributed and exhibits significant polymorphism in colour pattern and colony shape, is considered cryptogenic in the Mediterranean Sea (Zenetos *et al.*, 2017). According to recent genetic studies, this species has a Mediterranean origin, as the data reveals the presence of five genetically divergent clades in the region. Europe and the Mediterranean are the most diverse regions, where all clades are present, whereas only one clade is found worldwide (Bock *et al.*, 2012; López-Legentil *et al.*, 2006). One possible cause of this complicated taxonomic situation may arise from the fact that most of the findings come from port areas. Ports and marinas are containment zones

where very few fouling species venture beyond. Since they are transported by ships and boats via hulls, there has been greater genetic mixing between Mediterranean ports and marinas due to their higher connectivity compared to other global ports. Moreover, the Mediterranean is a very active region for recreational boating and maritime transport, especially considering its small surface area compared to other oceans (Carreño & Lloret, 2021). This species has been reported in Algeria in the Port of Arzew (Bensari *et al.*, 2020). In our study, *B. schlosseri* was very abundant (Fig. 3C).

Polyclinum constellatum Savigny, 1816

The ascidian *P. constellatum*, first described from specimens collected in Mauritius (Indian Ocean), has a pantropical distribution with a wide range of occurrences across various biogeographic regions (Montesanto *et al.*, 2022). In the Mediterranean, it was first recorded in Egypt (Halim & Abdel Messeih, 2016) and has since been observed in Turkey (Aydin Onen, 2018), Italy, and Greece (Montesanto *et al.*, 2022). This species has likely




Fig. 3: NIS and cryptogenic species found in this study. A: *Balanus trigonus*, B: *Botrylloides* cf. *violaceus*, C: *Botryllus schlosseri*, D: *Botrylloides* cf. *niger*, E: *Polyclinum constellatum*, F: *Jassa slatteryi*, G: *Celleporaria brunnea*, H: *Branchiommma bairdi*, I: *Hydroides elegans*.

spread through a Lessepsian migration route (Halim & Abdel Messeih, 2016). In our study, we report *P. constellatum* for the first time in the western Mediterranean (Fig. 3E). This introduction is most likely due to hull fouling on ships, as there is no clear continuity in its expansion from Egypt to Algeria, further supported by the fact that it is a lecithotrophic larval species. We found this species was abundantly attached to the port dock.

Styela canopus (Savigny, 1816)

The Rough Sea Squirt, *S. canopus*, is broadly distributed across temperate and tropical coastal waters worldwide (Lambert, 2001). This species was first described in the Red Sea in 1816 and was later documented on both sides of the North Atlantic, at Ascension Island in the South Atlantic, in the Northwest Pacific, throughout the tropical Indo-Pacific, and in the temperate waters of Australia (Kott, 1985). It was reported from the Mediterranean [as *Styela canopoides* (Heller)] in 1877 (Heller, 1877). Its long-standing presence in the Mediterranean

and other regions has led to its classification as a cryptogenic species. A genetic analysis of several populations revealed the presence of multiple species (Corrêa de Barros & Moreira da Rocha, 2021). The authors of this study suggested treating *S. canopus* as a species complex. Our study reports this species for the first time in Algeria (Fig. 2I). It was not very frequent in the Port of Algiers.

Styela plicata (Lesueur, 1823)

This species is a well-known, cosmopolitan fouling species found in shallow, sheltered areas of tropical and warm-temperate oceans (Ulman *et al.*, 2017). Genetic analysis indicates that its broad geographic distribution is due to numerous introductions caused by human-mediated hull fouling, leading to multiple introduction events (Locke *et al.*, 2009). Furthermore, most sightings are from artificial substrates or harbours, reinforcing the hypothesis of an ongoing invasion. Consequently, it is challenging to determine the exact origin of this species due to these multiple introductions (Locke *et al.*, 2009). How-

ever, according to Locke *et al.* (2009), the most probable origin of this species is the Western Pacific. It has been found in the Mediterranean since the 19th century (Zenetos & Galanidi, 2020); we report it in Algeria for the first time (Fig. 2E). It is highly likely that this species was established a long time ago along the Algerian coast, but due to a lack of specialists and insufficient investigation, it has not been reported until now. A related species, *Styela plicata* Lesuer, was observed negatively impacting the growth of cultured bivalves in Brazil, Hong Kong, Japan, and Spain (Rocha *et al.*, 2009).

Bryozoa

Bugula neritina (Linnaeus, 1758)

The bryozoan *B. neritina*, a well-known fouling organism, is widely spread along warm-temperate coasts and subtropical regions worldwide (Ryland *et al.*, 2011). Due to its long-standing broad distribution, determining the precise origin of this species remains challenging; therefore, it is considered cryptogenic in the Mediterranean (Zenetos *et al.*, 2017), where it is widely distributed in the eastern and western basins. In Algeria, *B. neritina* was overlooked in bryozoan studies and is absent from the bryozoan inventory compiled by d'Hondt & Ben Ismail (2008). It was first reported in Algeria in 2020 (Bensari *et al.*, 2020), although it is likely that it has been present for a much longer period. We found the species to be quite abundant in the Port of Algiers (Fig. 2B).

Bugulina stolonifera (Ryland, 1960)

The erect bryozoan *B. stolonifera* is commonly found in ports worldwide, including in the Atlantic and Pacific Oceans, as well as in the Mediterranean (Ryland *et al.*, 2011). It was first described by Ryland (1960) based on samples collected from the port at Swansea, Great Britain, and older specimens from an herbarium acquired by the British Museum. These specimens, dating back to before 1875, were labeled “British coast”. The origin of this species remains unknown due to its long-standing confusion with other species such as *Bugulina spicata* (Hincks), *Bugulina avicularia* (Linnaeus), and *Crisularia pulmosa* (Pallas) (Bensari *et al.*, 2020). Additionally, its broad distribution across the globe occurred so long ago that tracking its geographical dispersal is challenging. In Algeria, this species was recorded for the first time in the Port of Arzew (Bensari *et al.*, 2020). In the Port of Algiers, *B. stolonifera* was not frequently observed (Fig. 2A).

Celleporaria brunnea (Hincks, 1884)

This bryozoan was described by Hincks in 1884 as *Cellepora brunnea* from British Columbia, Canada (Canning-Clode *et al.*, 2013). It is naturally found in fouling communities along the northeastern Pacific coasts, rang-

ing from British Columbia to Ecuador (Lezzi *et al.*, 2015). In 2004, *C. brunnea* was found on aquaculture steel cages in South Korea, where it was believed to have been introduced from the eastern Pacific Ocean (Seo & Min, 2009). This species is also found in the northeastern Atlantic in Portugal (Canning-Clode *et al.*, 2013), including the Madeira Archipelago (Souto *et al.*, 2023), France (André *et al.*, 2014), and Norway (Husa *et al.*, 2024). In the Mediterranean, it is found in Lebanon, Italy, Croatia, Turkey, France, Malta, and Greece (Ulman *et al.*, 2017). We document *C. brunnea* in Algeria for the first time (Fig. 3G). It was observed infrequently during the period of this study.

Schizoporella errata (Waters, 1878)

This bryozoan is a widely distributed species found in warm temperate and subtropical regions (Tilbrook *et al.*, 2001). First described from Naples, Italy, it occurs throughout the Mediterranean, West Africa, eastern Canada, the eastern coasts of the Americas from North Carolina through the Caribbean to Brazil, the Pacific coast of North America, the Red Sea, the Persian Gulf, southern Australia, and New Zealand (Tilbrook *et al.*, 2001). Its origin has long been presumed to be the Mediterranean due to its ancient presence in the basin. However, the global distribution of this species challenges the notion of a Mediterranean origin, as it could have been introduced there before the 19th century. It is now considered cryptogenic in the Mediterranean (Katsanevakis *et al.*, 2020). D'Hondt & Ben Ismail (2008) mentioned that the species was reported by Gautier (1955) in Algeria. However, after thoroughly reviewing all of Gautier's references, including the one from 1955, we found no mention of this species. Hence, our observation (Fig. 2H) in the Port of Algiers constitutes the first record of *S. errata* in Algeria.

Tricellaria inopinata d'Hondt & Occhipinti Ambrogi, 1985

This bryozoan was described from the Venetian Lagoon (d'Hondt & Occhipinti Ambrogi, 1985), but its origin is not Mediterranean. It is potentially from the Indo-Pacific Ocean, as the genus *Tricellaria* is typical of this ocean (Dyrynda *et al.*, 2000). Indeed, specimens of *Tricellaria* washed ashore in central Oregon in 2012 on debris from the 2011 Japanese tsunami and were identified as *T. inopinata*, suggesting Japan as its native region (McCuller & Carlton, 2018). This species is present in the northeastern Atlantic, ranging from Germany to Spain, as well as along the British and Irish coasts (Cook *et al.*, 2013) and the coast of Norway (Porter *et al.*, 2015). In the Northwest Atlantic, after its first discovery in Eel Pond, Massachusetts (Johnson *et al.*, 2012), *T. inopinata* expanded its range to the Hampton River in New Hampshire (Wells *et al.*, 2013). In the Mediterranean, it is present in Italy (Lodola *et al.*, 2012), Tunisia (Ben Souissi *et al.*, 2006), France, and Greece (Ulman *et al.*, 2017). The impact of this species is significant, as it dominates

benthic communities and outcompetes other sessile species for space and food (Tsirintanis *et al.*, 2022). Our encounter of it the Port of Algiers marks the first reported occurrence of this species in Algeria (Fig. 2C), where we found it infrequently.

Crustacea

Balanus trigonus Darwin, 1854

Balanus trigonus, commonly called the triangle barnacle, was originally described based on specimens collected from the Pacific region, including Indonesia, Colombia, Peru, Australia, and New Zealand (Darwin, 1854). This barnacle has a global distribution and is considered introduced in the Atlantic Ocean and the Mediterranean Sea (Zullo, 1992). Its first record in the Mediterranean dates to 1927 in the Gulf of Catania, Italy (Zullo, 1992), and it is now widespread across the entire Mediterranean basin (Galanidi *et al.*, 2023). Our observation of it at the Port of Algiers is the first record of this species on the Algerian coast (Fig. 3A), but it is highly likely that was introduced well before. During our sampling, several specimens were found on a tire.

Jassa slatteryi Conlan, 1990

The amphipod *J. slatteryi*, likely native to the eastern Pacific, has an uncertain origin and is considered cryptogenic in other regions, including the Mediterranean, as the species has been misidentified multiple times, leaving its true native and introduced distribution still to be determined (Marchini & Cardeccia, 2017). In a comprehensive study of the genus *Jassa*, Conlan *et al.* (2021) suggested a northeastern Pacific origin for the species, as it is found in artificial and natural habitats in that region, and because the species was never recorded on the European coast before 1900, a zone with over 200 years of natural history collecting. According to Pilgrim & Darling (2010), who found as much genetic variability in *J. slatteryi* as in the native eastern Pacific *Jassa staudei* Conlan, suggesting that *J. slatteryi* is also native to this region. A more recent study based on the genetic analysis of several specimens from various regions suggests that the species has a North Pacific origin (Beermann *et al.*, 2020). Recently, Bakalem *et al.* (2024) classified it as non-indigenous to the Mediterranean, based on the species was not reported in the region before 1997 by Ruffo (1982, 1989, 1993, 1998). Bonifazi *et al.* (2018) cited Conlan (1990) to claim that the species was first observed in Croatia. However, upon consulting Conlan (1990), no reference to Croatia was found. Instead, the species was reported from “Cette”, an old spelling of the city of Sète in the Mediterranean region of France (K. E. Conlan, personal communication). Further examination of her archival notes revealed that the specimen originally identified was not *J. slatteryi* but rather *Jassa morinói* Conlan. Consequently, it is *J. morinói* that was cited from Sète, and there is no evidence to

suggest that *J. slatteryi* has been recorded in the Mediterranean. Based on this evidence, we suggest that the status of this species should be considered non-indigenous. It is worth noting that this species was observed in the port of Ceuta with a very high relative abundance and has been classified as having a high invasion potential (Ros *et al.*, 2020). Our finding of it in the Port of Algiers documents the occurrence of *J. slatteryi* in Algeria for the first time (Fig. 3F). We found the species to be not very frequent in our observations.

Zeuxo coralensis Sieg, 1980

The tanaidacean *Zeuxo coralensis* has a circumtropical distribution (Masunari & Sieg, 1980). In the Mediterranean Sea, it was first discovered in the Bay of Algeciras (southern Spain) between 1992 and 1993 on artificial and natural substrates (Sánchez-Moyano & García-Gómez, 1998). The species has also been encountered in the eastern Mediterranean (Zenetos *et al.*, 2010). *Zeuxo coralensis* was observed for the first time in Algeria at the Port of Arzew (Bensari *et al.*, 2020). We found *Z. coralensis* on a tire (Fig. 2D).

Polychaeta

Branchiomma bairdi (McIntosh, 1885)

The tube-dwelling sabellid *B. bairdi* was first described from Bermuda and is known from central Florida to the Atlantic side of Panama and the Caribbean Islands (Tovar-Hernández *et al.*, 2009). This species was later introduced to the Gulf of California and Hawaii. It has also been introduced to the Mediterranean Sea, the Canary Islands, the Balearic Sea, Madeira (Portugal), Queensland (Australia), the Pacific coast of Mexico, and Coiba National Park in the Gulf of Montijo on the Pacific coast of Panama (Keppel *et al.*, 2018). The initial introduction of this species into the Mediterranean remains uncertain due to frequent misidentification with *Branchiomma boholense* (Grube). However, the first confirmed report of this species was in Cyprus (Çinar, 2005 as *B. boholense*, later re-examined and correctly identified as *B. bairdi* in Çinar, 2009). *Branchiomma bairdi* was also observed in Tunisia in 2012 (Khedhri *et al.*, 2017). We document *B. bairdi* in Algeria for the first time (Fig. 3H), and our observations indicate that the species is highly abundant.

Branchiomma luctuosum (Grube, 1870)

Branchiomma luctuosum, first described in the Indo-Pacific region, was likely introduced to the Mediterranean through the Suez Canal (Licciano & Giangrande, 2008). Following its initial discovery in Italy in 1978 (Knight-Jones *et al.*, 1991), *B. luctuosum* has rapidly spread throughout the Mediterranean and is now widely distributed in the region (Mabrouki & Taybi, 2024), particular-

ly in the northern part. Our encounter of *B. luctuosum* (Fig. 2G) is a new country record for Algeria, where it is well established in the Port of Algiers. Additionally, this species has been observed at Bordj El Bahri to the east of Algiers Bay (personal observation), with a very high density on hard substrates in the infralittoral zone.

Hydroides elegans (Haswell, 1883)

The serpulid *H. elegans* seems to be a circumtropical species that extends into warm-temperate zones and is considered an alien species in the Mediterranean Sea, likely introduced via ship hulls (Zibrowius, 1991). This species is established in the entire Mediterranean basin and is considered long-standing resident (Zenetos *et al.*, 2012). It is considered invasive due to significant impact on biodiversity and ecosystem services (Katsanevakis *et al.*, 2014). *Hydroides elegans* (Fig. 3I) was first reported in Algeria at the Port of Algiers, but its presence along the Algerian coast is likely much older. In the Port of Arzew (Oran), Bensari *et al.* (2020) found the species to be as abundant as in this study.

Porifera

Paraleucilla magna Klautau, Monteiro, & Borojevic, 2004

This calcareous sponge was described from the Brazilian coast where it was considered cryptogenic (Klautau *et al.*, 2004). The origin of this species remains unknown but is likely Indo-Pacific, as is the origin of most species in the genus *Paraleucilla* (Longo *et al.*, 2007). In the Mediterranean, *P. magna* is considered non-indigenous and has rapidly spread to several locations. It is now widespread throughout the Mediterranean (see Katsanevakis *et al.*, 2020 for a more detailed distribution). Its rapid spread is attributed to mussel farming as well as maritime traffic (Longo *et al.*, 2007). Molecular study of samples collected from the Atlantic coast of Brazil and the Adriatic Sea suggests the presence of five genetically distinct populations, likely originating from multiple human-mediated introductions from various sources (Cavalcanti *et al.*, 2020). *Paraleucilla magna* (Fig. 2C) was first observed in Algeria in 2018 simultaneously at several locations, including the Port of Algiers, Pisan Island (Béjaia) (Bachetarzi *et al.*, 2019), and the Port of Arzew (Oran) (Bensari *et al.*, 2020). This suggests that the species may have been introduced some time ago, especially considering its presence on Pisan Island, a location distant from any direct anthropogenic activities likely to lead to an introduction.

Discussion

While several species identified in this study are documented here for the first time in Algeria, their introductions likely date back many years, considering the histor-

ical significance of the Port of Algiers as one of the oldest in the Mediterranean. Notably, *S. plicata*, *H. elegans*, *B. schlosseri*, *C. brunnea*, and *B. bairdi* rank among the most reported non-native species in ports throughout the Mediterranean region (Tempesti *et al.*, 2020).

The 18 species identified in this study underscores our limited understanding of non-indigenous species along the Algerian coast, a reality that is similarly reflected across several Mediterranean countries (Ulman *et al.*, 2017) due to a lack of taxonomic expertise or lack of focused inventories. Other species may be present but were missed because it was not possible to identify all taxa due to limited taxonomic expertise, and sampling was restricted in terms of area, depth, and substrate type.

Maritime transport has played a significant role in the transfer of exotic species between different geographic regions, whether directly from their native ranges or through secondary introductions from other Mediterranean areas. Hull fouling is undoubtedly the most important vector of introduction of the species we observed, as most of these species are biofouling organisms. Moreover, hull fouling is the only viable transport method for species with short lecithotrophic larval stages, such as ascidians. Maritime transport is a vital pathway for secondary introductions, dispersing species across regions through the interconnected network of Mediterranean ports – with even pleasure boats playing a significant role in this process. Ships carry introduced species from one port to another, facilitating their dispersal. For instance, *P. constellatum*, likely a Lessepsian migrant, has established itself in the Mediterranean and was discovered in the Port of Algiers in our study. Its occurrence in the port is most likely due to a secondary introduction originating from the eastern Mediterranean.

The question of the origin of non-indigenous species remains complex for several reasons. On the one hand, these species are often attributed to the Mediterranean or other regions simply because they were first described there – although they were likely widespread long before the 20th century – due, in part, to more intensive sampling of the Northern Hemisphere, especially along European coasts. On the other hand, taxonomic difficulties, particularly for fouling species that are often cosmopolitan and may belong to species complexes, add an additional layer of complexity. This ambiguity highlights the need for historical data, improved taxonomic research, and molecular techniques to more accurately trace the biogeographical history of these organisms and determine their true origins, even though pinpointing an exact origin remains challenging in many cases.

One key question to take away from our study is which species are most likely to spread beyond the port environment. Among the species we encountered, it is evident that *B. cf niger* and *B. luctuosum*, which are widely distributed across Algeria, warrant serious study to assess their impact on native species. For example, based on our own observations outside the port of Algiers, we have noted that *B. luctuosum* is particularly prevalent.

Our study represents a valuable contribution to the enrichment of data and information on non-indigenous

species in port environments in the Mediterranean. It highlights the Algerian coast, a region where information remains scarce, and its results offer insights that bridge knowledge gaps and support future research.

Acknowledgements

We would like to express our sincere gratitude to all those who contributed to the successful completion of this work.

References

- André, F., Corolla, J. P., Lanza, B., Rochefort, G., 2014. *Bryozoaires d'Europe : les carnets du plongeur*. Editions Neptune Plongée.
- Aydin Onen, S., 2018. Distribution of Ascidians with a New Record of the Non-Indigenous Species *Polyclinum constellatum* Savigny, 1816 from the Aegean Coast of Turkey. *Turkish Journal of Fisheries and Aquatic Sciences*, 18 (9), 1077-1089.
- Bachetarzi, R., Rebzani-Zahaf, C., Deudero, S., Vázquez-Luis, M., Uriz, M.J. *et al.*, 2019. The non-indigenous and invasive species *Paraleucilla magna* Klautau, Monteiro & Borojevic, 2004 (Porifera: Calcarea) in the Algerian coast (Southwestern of Mediterranean Sea). *Acta Adriatica*, 60 (1), 41-46.
- Bakalem, A., Dauvin, J.C., Menioui, M., 2024. Diversity of marine Amphipods (Crustacea, Peracarida) from the North African shelf of the Mediterranean Sea (Morocco, Algeria, Tunisia, Libya, Egypt). An updated checklist for 2023. *Mediterranean Marine Science*, 25 (2), 311-373.
- Beermann, J., Hall-Mullen, A.K., Havermans, C., Coolen, J.W., Crooijmans, R.P. *et al.*, 2020. Ancient globetrotters –connectivity and putative native ranges of two cosmopolitan biofouling amphipods. *PeerJ*, 8, e9613.
- Ben Souissi, J., Ben Salem, M., Zaouali, J., 2006. *Tricellaria inopinata* (Bryozoa, Cheilostome), récolté pour la première fois en Tunisie (Lagune sud de Tunis : Tunisie septentrionale) [*Tricellaria inopinata* (Bryozoa, Cheilostome), harvest for the first time in Tunisia (Southern Lagoon of Tunis : northern Tunisia)]. *Actes de La 8^{ème} Conférence Internationale Des Limnologues d'Expression Française 17-21 Mars 2006*.
- Bensari, B., Bahbah, L., Lounaouci, A., Fahci, S.E., Bouda, A. *et al.* 2020. First records of non-indigenous species in port of Arzew (Algeria: southwestern Mediterranean). *Mediterranean Marine Science*, 21 (2), 393-399.
- Bock, D.G., MacIsaac, H.J., Cristescu, M.E., 2012. Multilocus genetic analyses differentiate between widespread and spatially restricted cryptic species in a model ascidian. *Proceedings of the Royal Society B: Biological Sciences*, 279 (1737), 2377-2385.
- Bock, D.G., Zhan, A., Lejeune, C., MacIsaac, H.J., Cristescu, M.E., 2011. Looking at both sides of the invasion: patterns of colonization in the violet tunicate *Botrylloides violaceus*. *Molecular Ecology*, 20 (3), 503-516.
- Bonifazi, A., Mancini, E., Ventura, D., 2018. First record of the invasive and cryptogenic species *Jassa slatteryi* Conlan, 1990 (Crustacea: Amphipoda) in Italian coastal waters. *Journal of Sea Research*, 136, 37-41.
- Bouda, A., Bachari, N.E.I., Bahmed, L., Boubenia, R., 2016. Design of a risk assessment methodology for the introduction of invasive species from ship ballast waters: The case of Arzew port. *Management of Environmental Quality: An International Journal*, 27 (5), 474-490.
- Bouda, A., Bachari, N.I., Nacef, L., Bensari, B., 2018. Risk Analysis of Invasive Species Introduction in the Port of Arzew, by Calculation of Biofouling Surface on Ships' Hulls. *Environmental Modeling and Assessment*, 23 (2), 185-192.
- Canning-Clode, J., Souto, J., McCann, L. 2013. First record of *Celleporaria brunnea* (Bryozoa: Lepraliellidae) in Portugal and in the East Atlantic. *Marine Biodiversity Records*, 6, e108.
- Carlton, J.T., 1996. Biological Invasions and Cryptogenic Species. *Ecology*, 77 (6), 1653-1655.
- Carmona-Rodríguez, A., Antón, C., Climent, M.Á., Garcés, P., Montiel, V. *et al.*, 2024. Development and Succession of Non-Indigenous and Cryptogenic Species over Two Different Substrates in the Port of Alicante (Western Mediterranean). *Journal of Marine Science and Engineering*, 12 (7), 1188.
- Carreño, A., Lloret, J., 2021. Environmental impacts of increasing leisure boating activity in Mediterranean coastal waters. *Ocean & Coastal Management*, 209, 105693.
- Cavalcanti, F.F., Padua, A., Cunha, H., Halasz, M., Nikolić, V. *et al.*, 2020. Population differentiation supports multiple human-mediated introductions of the transatlantic exotic sponge *Paraleucilla magna* (Porifera, Calcarea). *Hydrobiologia*, 847 (17), 3571-3590.
- Çinar, M.E., 2005. Polychaetes from the coast of Northern Cyprus (eastern Mediterranean Sea), with two new records for the Mediterranean Sea. *Cahiers de Biologie Marine*, 46 (2), 143-160.
- Çinar, M.E., 2009. Alien polychaete species (Annelida: Polychaeta) on the southern coast of Turkey (Levantine Sea, eastern Mediterranean), with 13 new records for the Mediterranean Sea. *Journal of Natural History*, 43, (37-38), 2283-2328.
- Conlan, K.E., 1990. Revision of the crustacean amphipod genus *Jassa* Leach (Corophioidea: Ischyroceridae). *Canadian Journal of Zoology*, 68 (10), 2031-2075.
- Conlan, K.E., Desiderato, A., Beermann, J., 2021. *Jassa* (Crustacea: Amphipoda): a new morphological and molecular assessment of the genus. *Zootaxa*, 4939 (1), 1-191.
- Cook, E.J., Stehlíková, J., Beveridge, C.M., Burrows, M.T., de Blauwe, H. *et al.*, 2013. Distribution of the invasive bryozoan *Tricellaria inopinata* in Scotland and a review of its European expansion. *Aquatic Invasions*, 8 (3), 281-288.
- Corrêa de Barros, R., Moreira da Rocha, R., 2021. Genetic analyses reveal cryptic diversity in the widely distributed *Styela canopus* (Asciadiacea: Styelidae). *Invertebrate Systematics*, 35 (3), 298-311.
- D'Amen, M., Azzurro, E., 2020. Lessepsian fish invasion in Mediterranean marine protected areas: a risk assessment under climate change scenarios. *ICES Journal of Marine Science*, 77 (1), 388-397.

- Darwin, C., 1854. *A Monograph on the Sub-Class Cirripedia with Figures of All the Species. The Balanidae, (or Sessile Cirripedia); the Verrucidae, Etc (Vol. 2)*. Ray Society.
- Della Sala, G., Coppola, D., Virgili, R., Vitale, G.A., Tanduo, V. *et al.*, 2022. Untargeted Metabolomics Yields Insights into the Lipidome of *Botrylloides niger* Herdman, 1886, An Ascidian Invading the Mediterranean Sea. *Frontiers in Marine Science*, 9, 865751.
- d'Hondt, J.L., Ben Ismail, D., 2008. Bryozoa from the Algerian coasts. Complementary data on Tunisian Bryozoa. *Bulletin de La Societe Zoologique de France*, 133 (1-3), 55-71.
- d'Hondt, J.L., Occhipinti Ambrogio, A., 1985. *Tricellaria inopinata*, n. sp., un nouveau Bryozoaire Cheilostome de la faune méditerranéenne. *Marine Ecology*, 6 (1), 35-46.
- Dyrynda, P.E.J., Fairall, V.R., Occhipinti Ambrogio, A., D'Hondt, J.L., 2000. The distribution, origins and taxonomy of *Tricellaria inopinata* d'Hondt and Occhipinti Ambrogio, 1985, an invasive bryozoan new to the Atlantic. *Journal of Natural History*, 34 (10), 1993-2006.
- Galanidi, M., Aissi, M., Ali, M., Bakalem, A., Bariche, M. *et al.*, 2023. Validated Inventories of Non-Indigenous Species (NIS) for the Mediterranean Sea as Tools for Regional Policy and Patterns of NIS Spread. *Diversity*, 15 (9), 962.
- Gautier, Y., 1955. Bryozoaires de Castiglione. *Bulletin de La Station d'Aquiculture et de Pêche de Castiglione, N.S.* (7), 227-271.
- Gollasch, S., 2002. The Importance of Ship Hull Fouling as a Vector of Species Introductions into the North Sea. *Biofouling*, 18 (2), 105-121.
- Grimes, S., Benabdi, M., Babali, N., Refes, W., Boudjellal-Kaidi, N. *et al.*, 2018. Biodiversity changes along the Algerian coast (Southwestern Mediterranean basin) from 1834 to 2017. First assessment of introduced species. *Mediterranean Marine Science*, 19 (1), 156-179.
- Halim, Y., Abdel Messeih, M., 2016. Aliens in Egyptian waters. A checklist of ascidians of the Suez Canal and the adjacent Mediterranean waters. *Egyptian Journal of Aquatic Research*, 42 (4), 449-457.
- Heller, C., 1877. Untersuchungen über die Tunicaten des Adriatischen und Mittelmeeres. *Denkschriften Der Kaiserlichen Academie Der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe, Wien*.
- Husa, V., Fossey, F., Davey, M., Agnalt, A.L., Bransegg, H. *et al.*, 2024. *Monitoring marine alien species in Norway: A pilot study for implementing a national program*. Rapport frahvforskningen.
- Johnson, C., Winston, J., Woollacott, R., 2012. Western Atlantic introduction and persistence of the marine bryozoan *Tricellaria inopinata*. *Aquatic Invasions*, 7 (3), 295-303.
- Katsanevakis, S., Wallentinus, I., Zenetos, A., Leppäkoski, E., Çinar, M.E. *et al.*, 2014. Impacts of invasive alien marine species on ecosystem services and biodiversity: A pan-European review. *Aquatic Invasions*, 9 (4), 391-423.
- Katsanevakis, S., Poursanidis, D., Hoffman, R., Rizgalla, J., Rothman, S.B.S. *et al.*, 2020. Unpublished mediterranean records of marine alien and cryptogenic species. *BioInvasions Records*, 9 (2), 165-182.
- Keppel, E., Tovar-Hernández, M.A., Ruiz, G., 2018. New records of the non-indigenous species *Branchiomma bairdi* and *B. conspersum* (Polychaeta: Sabellidae) on the Pacific coast of North America. *BioInvasions Records*, 7 (3), 229-236.
- Khedhri, I., Tovar-Hernández, M., Bonifácio, P., Ahmed, A., Aleya, L., 2017. First report of the invasive species *Branchiomma bairdi* McIntosh, 1885 (Annelida: Sabellidae) along the Tunisian coast (Mediterranean Sea). *BioInvasions Records*, 6 (2), 139-145.
- Klautau, M., Monteiro, L., Borojevic, R., 2004. First occurrence of the genus *Paraleucilla* (Calcarea, Porifera) in the Atlantic Ocean: *P. magna* sp. nov. *Zootaxa*, 710 (1), 1-8.
- Knight-Jones, P., Knight-Jones, W., Ergen, Z., 1991. Sabelliform polychaetes, mostly from Turkey's Aegean coast. *Journal of Natural History*, 25 (4), 837-858.
- Kott, P., 1985. *The Australian Ascidiacea Part 1, Phlebobranchia and Stolidobranchia* (Vol. 23). Memoirs of the Queensland Museum.
- Lambert, G., 2001. A Global Overview of Ascidian Introductions and Their Possible Impact on the Endemic Fauna. In *The Biology of Ascidians* (pp. 249-257). Springer Japan.
- Lezzi, M., Pierri, C., Cardone, F., 2015. Presence of *Celleporaria brunnea* (Bryozoa: Lepraliellidae) in the Central Mediterranean: first occurrence in the Gulf of Taranto. *Marine Biodiversity Records*, 8, e137.
- Licciano, M., Giangrande, A., 2008. The genus *Branchiomma* (Polychaeta: Sabellidae) in the Mediterranean Sea, with the description of *B. maerli* n. sp. *Scientia Marina*, 72 (2), 383-391.
- Locke, A., Carman, M., De Barros, R. C., Da Rocha, R. M., Pie, M. R., 2009. Human-mediated global dispersion of *Styela plicata* (Tunicata, Ascidiacea). *Aquatic Invasions*, 4 (1), 45-57.
- Lodola, A., Savini, D., Occhipinti-Ambrogio, A., 2012. First record of *Tricellaria inopinata* (Bryozoa: Candidae) in the harbours of La Spezia and Olbia, Western Mediterranean Sea (Italy). *Marine Biodiversity Records*, 5, e41.
- Longo, C., Mastrototaro, F., Corriero, G., 2007. Occurrence of *Paraleucilla magna* (Porifera: Calcarea) in the Mediterranean Sea. *Journal of the Marine Biological Association of the United Kingdom*, 87 (6), 1749-1755.
- López-Legentil, S., Turon, X., Planes, S., 2006. Genetic structure of the star sea squirt, *Botryllus schlosseri*, introduced in southern European harbours. *Molecular Ecology*, 15 (13), 3957-3967.
- Mabrouki, Y., Taybi, A. F., 2024. Range expansion of the alien polychaete *Branchiomma luctuosum* (Grube, 1870) (Sabellidae) in Morocco (the southwestern Mediterranean Sea). *Natura Croatica*, 33 (1), 197-202.
- Marchini, A., Cardeccia, A., 2017. Alien amphipods in a sea of troubles: cryptogenic species, unresolved taxonomy and overlooked introductions. *Marine Biology*, 164 (4), 69.
- Massé, C., Viard, F., Humbert, S., Antajan, E., Auby, I. *et al.*, 2023. An Overview of Marine Non-Indigenous Species Found in Three Contrasting Biogeographic Metropolitan French Regions: Insights on Distribution, Origins and Pathways of Introduction. *Diversity*, 15 (2), 161.
- Masunari, S., Sieg, J., 1980. Morphological and ecological notes on *Zeuxo coralensis* Sieg, 1980 from Brazil. *Studies on Neotropical Fauna and Environment*, 15 (1), 1-8.
- McCuller, M., Carlton, J., 2018. Transoceanic rafting of Bryozoa (Cyclostomata, Cheilostomata, and Ctenostomata)

- across the North Pacific Ocean on Japanese tsunami marine debris. *Aquatic Invasions*, 13 (1), 137-162.
- Mghili, B., Lamine, I., Laamraoui, M. R., Aksissou, M., Galanidi, M., 2024. Updating the national list of marine alien species in Morocco. *Mediterranean Marine Science*, 25 (1), 231-249.
- Mnasri-Affifi, I., Zribi, I., Abdelkader, N., Charfi-Cheikhrouha, F., Zakhama-Sraieb, R., 2024. Rapid assessment survey of non-indigenous and cryptogenic species in Tunisian marinas. *BioInvasions Records*, 13 (2), 419-436.
- Montesanto, F., Chimenti, G., Gissi, C., Mastrototaro, F., 2022. *Polyclinum constellatum* (Tunicata, Ascidiacea), an emerging non-indigenous species of the Mediterranean Sea: integrated taxonomy and the importance of reliable DNA barcode data. *Mediterranean Marine Science*, 23 (1), 69-83.
- Palanisamy, S.K., Cosentino, A., Giacobbe, S., 2018. Rapid coastal survey of the distribution of invasive ascidians in Coastal Lake Faro, Messina- Italy. *Poster Presented at the International Invasive Sea Squirt Conference, Woods Hole Oceanographic Institution*, 2018, Massachusetts, United States.
- Pilgrim, E.M., Darling, J.A., 2010. Genetic diversity in two introduced biofouling amphipods (*Ampithoe valida* & *Jassa marmorata*) along the Pacific North American coast: investigation into molecular identification and cryptic diversity. *Diversity and Distributions*, 16 (5), 827-839.
- Porter, J.S., Spencer Jones, M.E., Kuklinski, P., Rouse, S., 2015. First records of marine invasive non-native Bryozoa in Norwegian coastal waters from Bergen to Trondheim. *BioInvasions Records*, 4 (3), 157-169.
- Rho, B.J., Choe, B.L., Song, J.-I., Lee Young J., 2000. Ascidians of Tangsa and its adjacent waters in Korea strait. *Korean Journal of Systematic Zoology*, 16 (1), 39-53.
- Rocha, R.M., Kremer, L.P., Baptista, M.S., Metri, R., 2009. Bi-valve cultures provide habitat for exotic tunicates in Southern Brazil. *Aquatic Invasions*, 4 (1), 195-205.
- Ros, M., Navarro-Barranco, C., González-Sánchez, M., Ostalé-Valriberas, E., Cervera-Currado, L. *et al.*, 2020. Starting the stowaway pathway: the role of dispersal behavior in the invasion success of low-mobile marine species. *Biological Invasions*, 22 (9), 2797-2812.
- Ruiz, G.M., Carlton, J.T., Grosholz, E.D., Hines, A.H., 1997. Global Invasions of Marine and Estuarine Habitats by Non-Indigenous Species: Mechanisms, Extent, and Consequences. *American Zoologist*, 37 (6), 621-632.
- Ryland, J.S., 1960. The British species of *Bugula* (Polyzoa). *Proceedings of the Zoological Society of London*, 134 (1), 65-104.
- Ryland, J., Bishop, J., De Blauwe, H., El Nagar, A., Minchin, D. *et al.*, 2011. Alien species of *Bugula* (Bryozoa) along the Atlantic coasts of Europe. *Aquatic Invasions*, 6 (1), 17-31.
- Sánchez-Moyano, J.E., García-Gómez, J.C., 1998. The arthropod community, especially crustacea, as a bioindicator in Algeciras Bay (Southern Spain) based on a spatial distribution. *Journal of Coastal Research*, 14 (3), 1119-1133.
- Seo, J.E., Min, B.S., 2009. A Faunistic Study on Cheilostomatous Bryozoans from the Shoreline of South Korea, with Two New Species. *Animal Systematics, Evolution and Diversity*, 25 (1), 19-40.
- Sheets, E.A., Cohen, C.S., Ruiz, G.M., da Rocha, R.M., 2016. Investigating the widespread introduction of a tropical marine fouling species. *Ecology and Evolution*, 6 (8), 2453-2471.
- Simkanin, C., Davidson, I.C., Dower, J.F., Jamieson, G., Theriault, T.W., 2012. Anthropogenic structures and the infiltration of natural benthos by invasive ascidians. *Marine Ecology*, 33 (4), 499-511.
- Souto, J., Ramalhosa, P., Ferrario, J., Png-Gonzalez, L., Álvarez, S. *et al.*, 2023. New species and new records of bryozoan species from fouling communities in the Madeira Archipelago (NE Atlantic). *Marine Biodiversity*, 53 (4), 49.
- Ruffo, S., (Ed.), 1982. The Amphipoda of the Mediterranean. Part 1. Gammaridea (Acanthonozomatidae to Gammaridae). *Mémoires de l'Institut Océanographique*, Monaco, 13, 1-364.
- Ruffo, S., (Ed.), 1989. The Amphipoda of the Mediterranean. Part 2. Gammaridea (Haustoriidae to Lysianassidae). *Mémoires de l'Institut Océanographique*, Monaco, 13, 365-576.
- Ruffo S., (Ed.), 1993. The Amphipoda of the Mediterranean. Part 3. Gammaridea (Melphidiidae to Talitridae), Ingolfiellidae, Caprellidae. *Mémoires de l'Institut Océanographique*, Monaco, 13, 577-813.
- Ruffo, S., (Ed.), 1998. The Amphipoda of the Mediterranean. Part 4. Localities and Map. Addenda to Parts –3. Key to Families. Ecology. Faunistics and Zoogeography. Bibliography. Index. *Mémoires de l'Institut Océanographique*, Monaco, 13, 815-959.
- Temiz, B., Öztürk, E., Blanchoud, S., Karahan, A., 2023. Phylogeographic and Morphological Analysis of *Botrylloides niger* Herdman, 1886 from the Northeastern Mediterranean Sea. *Diversity*, 15 (3), 367.
- Tempesti, J., Mangano, M.C., Langeneck, J., Lardicci, C., Maltagliati, F. *et al.*, 2020. Non-indigenous species in Mediterranean ports: A knowledge baseline. *Marine Environmental Research*, 161, 105056.
- Tilbrook, K.J., Hayward, P.J., Gordon, D.P., 2001. Cheilostomatous Bryozoa from Vanuatu. *Zoological Journal of the Linnean Society*, 131 (1), 35-109.
- Tovar-Hernández, M.A., Méndez, N., Salgado-Barragán, J., 2009. *Branchiomma bairdi*: a Caribbean hermaphrodite fan worm in the south-eastern Gulf of California (Polychaeta: Sabellidae). *Marine Biodiversity Records*, 2, e43.
- Tsiritanis, K., Azzuro, E., Crocetta, F., Dimiza, M., Frogila, C. *et al.*, 2022. Bioinvasion impacts on biodiversity, ecosystem services, and human health in the Mediterranean Sea. *Aquatic Invasions*, 17 (3), 308-352.
- Ulman, A., Ferrario, J., Occhpinti-Ambrogi, A., Arvanitidis, C., Bandi, A., 2017. A massive update of non-indigenous species records in Mediterranean marinas. *PeerJ*, 5, e3954.
- Wells, C.D., Pappal, A.L., Cao, Y., Carlton, J.T., Currimjee, Z. *et al.*, 2013. *Report on the 2013 Rapid Assessment Survey of Marine Species at New England Bays and Harbors*. www.mass.gov/czm.
- Zaniolo, G., Manni, L., Brunetti, R., Burighel, P., 1998. Brood pouch differentiation in *Botrylloides violaceus*, a viviparous ascidian (Tunicata). *Invertebrate Reproduction and Development*, 33 (1), 11-23.
- Zenetos, A., Galanidi, M., 2020. Mediterranean non indigenous species at the start of the 2020s: recent changes. *Marine*

Biodiversity Records, 13 (1), 10.

- Zenetos, A., Gofas, S., Verlaque, M., Çinar, M.E., García Raso, J.G. *et al.*, 2010. Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. In *Mediterranean Marine Science*, 11 (2), 381-493.
- Zenetos, A., Gofas, S., Morri, C., Rosso, A., Violanti, D. *et al.*, 2012. Alien species in the Mediterranean Sea by 2012. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part 2. Introduction trends and pathways. *Mediterranean Marine Science*, 13 (2), 328-352.
- Zenetos, A., Çinar, M. E., Crocetta, F., Golani, D., Rosso, A. *et al.*, 2017. Uncertainties and validation of alien species catalogues: The Mediterranean as an example. *Estuarine, Coastal and Shelf Science*, 191, 171-187.
- Zibrowius, H., 1991. Ongoing modification of the Mediterranean marine fauna and flora by the establishment of exotic species. *Mésogée*, 51, 83-107.
- Zullo, V.A., 1992. *Balanus trigonus* Darwin (Cirripedia, Balaninae) in the Atlantic Basin: An Introduced Species? *Bulletin of Marine Science*, 50 (1), 66-74.