

## Mediterranean Marine Science

early view



**Using Local Ecological Knowledge (LEK) to inform the multi-habitat life cycle of wreckfish: A Ligurian Sea case study for Mediterranean conservation**

ELENA DESIDERÀ, DAVIDE DI BLASI, FEDERICA POLI, TREVOR J. WILLIS, SIMONE BAVA, PAOLO GUIDETTI

doi: [10.12681/mms.41040](https://doi.org/10.12681/mms.41040)

### To cite this article:

DESIDERÀ, E., DI BLASI, D., POLI, F., WILLIS, T. J., BAVA, S., & GUIDETTI, P. (2025). Using Local Ecological Knowledge (LEK) to inform the multi-habitat life cycle of wreckfish: A Ligurian Sea case study for Mediterranean conservation. *Mediterranean Marine Science*, early view. <https://doi.org/10.12681/mms.41040>

## Using Local Ecological Knowledge (LEK) to inform the multi-habitat life cycle of wreckfish: A Ligurian Sea case study for Mediterranean conservation

Elena DESIDERÀ<sup>1,2</sup>, Davide Di BLASI<sup>1,3</sup>, Federica POLI<sup>1,4</sup>, Trevor J. WILLIS<sup>2,5</sup>, Simone BAVA<sup>1</sup>, and Paolo GUIDETTI<sup>1,2</sup>

<sup>1</sup> Department of Integrative Marine Ecology (EMI), Stazione Zoologica Anton Dohrn–National Institute of Marine Biology, Ecology and Biotechnology, Genoa Marine Centre (GMC), Villa del Principe, Piazza del Principe 4, 16126, Genoa, Italy

<sup>2</sup> National Biodiversity Future Center (NBFC), Piazza Marina 61, 90133, Palermo, Italy

<sup>3</sup> National Research Council, Institute of Marine Engineering (CNR-INM), Via De Marini 16, 16149, Genoa, Italy

<sup>4</sup> Centre for Ecology and Conservation, University of Exeter, Penryn Campus, Cornwall TR10 9FE, United Kingdom

<sup>5</sup> Department of Integrative Marine Ecology (EMI), Stazione Zoologica Anton Dohrn–National Institute of Marine Biology, Ecology and Biotechnology, Fano Marine Center, Viale Adriatico 1-N, 61032, Fano, Italy

Corresponding author: Elena DESIDERÀ; [elena.desidera@szn.it](mailto:elena.desidera@szn.it)

Contributing Editor: Stelios SOMARAKIS

Received: 08 April 2025; Accepted: 08 July 2025; Published online: 02 September 2025

### Abstract

Local Ecological Knowledge (LEK) from fishers and sea users is invaluable for reconstructing the occurrences of data-deficient and vulnerable fish species, particularly target species. This study explores the distribution patterns of the data-deficient wreckfish *Polyprion americanus* in the Ligurian Sea (NW Mediterranean Sea) using semi-structured interviews with 40 sea users (27 fishers and 13 observational data providers). We identified key habitats associated with five main size classes of wreckfish, from surface waters to 1000 m depth, including vulnerable ecosystems like seamounts and cold-water coral habitats. Smaller individuals (0–1.5 kg) were mainly found in pelagic habitats, frequently near floating objects, while larger individuals (>3 kg) were common in slope and seamount habitats. Abundance declines across all size classes were attributed to overfishing and, according to some respondents, a reduction in floating debris may have specifically affected the presence of smaller individuals. Fishing techniques varied by size class, with technological advances notably impacting larger individuals near seamounts. The study supports LEK as a valuable complementary information source to conventional research, advocating for updated conservation measures, including revised size limits and enhanced protection of vulnerable habitats.

**Keywords:** Data Deficient; overfishing; *Polyprion americanus*; seamount; size limit; vulnerable habitats.

### Introduction

The wreckfish *Polyprion americanus* (Bloch & Schneider, 1801) (Family Polyprionidae), is a marine fish characterized by a worldwide distribution, including the Mediterranean Sea (Ball *et al.*, 2000). Known for its impressive longevity (up to 80 years; Lytton *et al.*, 2015) and maximum size (up to 2 m, 100 kg; Roberts, 1989), it exhibits distinct pelagic juvenile and demersal adult phases (Tortonesi, 1975; Sedberry *et al.*, 1996). Juveniles are commonly found below floating objects, while adults inhabit benthic environments, often near wrecks or areas of underwater relief, up to 1000 m depth (Bini, 1970; Sedberry *et al.*, 1996; Ball *et al.*, 2000). Globally, the wreckfish is an ecologically and economically significant species, serving as a large-sized predator in marine ecosystems and considered a promising candidate for aqua-

culture (Papandroulakis *et al.*, 2004; Pérez *et al.*, 2019). However, its late maturation age (>7 years) (Carbonara *et al.*, 2003; Peres & Klippel, 2003; Wakefield *et al.*, 2013) along with suspected spawning aggregations (Smith, 1949; Menni & López, 1979; Ryall & Hargrave, 1984; Peres & Klippel, 2003; Button *et al.*, 2021) make it highly vulnerable to overfishing. Limited knowledge on its life cycle, biology and ecology (Hardy, 1978; Carbonara *et al.*, 2003) further complicates effective assessment and management. This lack of information is reflected in the wreckfish's classification as Data Deficient in the IUCN Red List global assessment (Sadovy, 2003).

In the Mediterranean Sea, the wreckfish is considered of minor commercial importance due to its low occurrence in commercial catches, with limited data on abundance and population trends (Yokes *et al.*, 2011). Therefore, it is also classified as Data Deficient in the Mediterranean by

the IUCN Red List (Yokes *et al.*, 2011), although recent regional assessments range from Near Threatened (European assessment; Collette *et al.*, 2015) to Vulnerable (Italian assessment; Relini *et al.*, 2017), reflecting potential population declines. Despite these concerns, the only conservation measure currently in place in the Mediterranean Sea is the minimum landing and marketing size, equaling 45 cm, as outlined in Council Regulation (EC) No. 1967/2006 (European Council, 2006). However, this size limit lacks any biological basis, as it was set to match the size established for groupers (family Epinephelidae) based on the wreckfish's grouper-like traits.

In the Mediterranean region, the species may hold historical and cultural significance. Curiously, it is depicted in ancient coinage from Akragas (modern Agrigento, Sicily) dating back to the late 6th-early 5th century BCE (Imhoof-Blumer & Keller, 1889; Ward, 1902; Rizzo, 1946). The fish representation on coins likely reflects its abundance and importance as a food resource. In the Mediterranean, wreckfish are caught by diverse fishing tools across various habitats. The species is primarily targeted by angling and bottom longlines (Bini, 1970; Yokes *et al.*, 2011), but it is also caught by bottom trawlers (Smith, 1949; Bini, 1970) and swordfish longlines (Garibaldi, 2015). Fishing activities may impact wreckfish populations, in the Mediterranean and beyond, as studies have shown that many individuals caught and marketed are below the size of sexual maturity, estimated to be 80-90 cm total length (Carbonara *et al.*, 2003; Peres & Klippel, 2003; Diogo & Pereira, 2013; Mavruk *et al.*, 2018). For instance, most wreckfish marketed in Galicia were found to be immature (Linares *et al.*, 2021), while those caught in the Azores and off Turkey were, on average, below the size of sexual maturity. High demand and rising costs have also led to frequent mislabeling of cheaper fish species (e.g., the Nile perch *Lates niloticus*) as wreckfish (Asensio *et al.*, 2002).

Based on the above, further research is needed to assess wreckfish population status and inform science-based conservation measures. Monitoring wreckfish populations, however, is extremely challenging. In the Mediterranean Sea, fishing experimental surveys (D'Onghia *et al.*, 2012) and deep remotely operated vehicles (ROV) are the most commonly used study methods (Canese & Bava, 2015; D'Onghia *et al.*, 2015; Bo *et al.*, 2021). Yet, the use of ROV has provided only a few occasional records of wreckfish in the region (Canese & Bava, 2015; D'Onghia *et al.*, 2015).

In recent years, Local Ecological Knowledge (LEK) has become a valuable resource for research, particularly for understanding the biology and ecology of commercially important fish species, especially when scientific data is scarce or unavailable (Johannes, 1978; Johannes *et al.*, 2000; Hamilton *et al.*, 2012). LEK is a valuable means of assessing trends in fish abundance, identifying key habitats, and understanding life cycles, often offering information that is difficult to obtain through traditional scientific approaches. The value of LEK has been more and more recognized as crucial for achieving management and conservation goals (Azzurro *et al.*, 2019; Sil-

vano *et al.*, 2023). LEK, in fact, can be a precious complementary source of information integrating scientific knowledge, and a bridge between communities and conservation goals, as it may offer insights into sustainable and customary practices, support dialogue and ultimately promote community-based conservation (Silvano *et al.*, 2017).

This study aimed to collect past and present occurrence data of the wreckfish in the Ligurian Sea (NW Mediterranean Sea) using LEK. The Ligurian Sea was selected as a case study within the Mediterranean basin, described as a “miniature ocean” (Bethoux *et al.*, 1999; Lejeune *et al.*, 2010), due to its unique oceanographic and geomorphological characteristics, which make it an ideal setting for investigating the entire life cycle of the species. These features include a narrow continental shelf (10-20 km) (Relini, 2007), steep slopes with submarine canyons, and a permanent cyclonic circulation driven by the Western Corsica/Mediterranean Current (WCC) and the Tyrrhenian Current (TC) (Cattaneo-Vietti *et al.*, 2010; Canepa *et al.*, 2015), which promotes nutrient upwelling and high primary productivity (Cattaneo-Vietti *et al.*, 2010). Such conditions influence the reproductive biology and recruitment of many species, including commercially important fish (Cattaneo-Vietti *et al.*, 2010). Furthermore, the Ligurian Sea coasts are highly urbanized and host >80 marinas, and numerous harbours and ports serving as key hubs for trade, tourism, and maritime activity (Cattaneo-Vietti *et al.*, 2010). Therefore, these areas increase the likelihood of interaction between local sea users, and marine resources and ecosystems. Historical data (1898) report the wreckfish as a common year-round catch in the Ligurian Sea (Cattaneo-Vietti *et al.*, 1985). However, catches in the area have declined, with no recent records from surveys conducted in habitats where the species was once commonly fished (Bo *et al.*, 2021).

This study aims to utilise LEK to address knowledge gaps concerning wreckfish habitat usage throughout its life cycle, the status of Ligurian wreckfish populations, and the fishing practices impacting the species. Our primary aim is to contribute to the assessment of the species' conservation status and provide relevant information potentially useful for the management and conservation of the wreckfish populations in the Mediterranean.

## Materials and Methods

Records of the past and present occurrence of wreckfish in the Ligurian Sea were collected using a Local Ecological Knowledge (LEK) approach. Individual interviews were conducted with various sea users, employing two types of structured questionnaires (see Annex) to collect two categories of data: 1) fishing data, including information from both recreational and professional fishing activities, and 2) observational data provided by boaters, whale watchers, technical divers, dive photographers, and fish vendors. Informed consent was obtained verbally from all individual participants included in the study, as this was the most appropriate method for the research

context. Participants were assured of their anonymity, with any identifying information removed unless they explicitly chose to disclose it. Formal ethical approval was not required for this study, as per local regulations; however, ethical standards were met through the careful handling of sensitive data and the respect for participants' privacy.

Since the interviewees included people without a scientific background and non-professional fishers, we first ensured they could accurately identify the wreckfish by showing them pictures of grouper species found in the same areas. The interviewees were asked the geographical areas and periods in which wreckfish were caught or observed, and on changes in the species' abundance over the years. Respondents were asked about the size (either weight or total length) and the number of specimens caught or observed per encounter, including size and number ranges when multiple individuals were caught or seen. Additional details collected from fishers included the fishing gear used to catch wreckfish, whether the gonads appeared mature, the co-occurring fish species, and the market value of wreckfish. Photographs of wreckfish specimens were collected whenever available and published with the consent of the interviewees.

Respondents consistently reported wreckfish size in terms of weight rather than length. Accordingly, for the analyses, wreckfish were categorised into size classes by weight. However, to allow comparison with previous studies that report size in terms of length, weight data were occasionally converted to estimated total length (TL) using the wreckfish length-weight relationship from FishBase (Froese & Pauly, 2024).

Key habitat types were identified at geomorphological scale based on the information gathered in the interviews. To infer the distribution and life cycle of the wreckfish, the presence of each size class across the main habitat types was assessed based on the number of respondents reporting their occurrence.

For each size class, the wreckfish conservation status was assessed based on respondent's reports of temporal abundance trends, which were classified as 'decreasing', 'stable', 'increasing' or 'NA' if respondents were unable to determine a trend, and summarised in a bar chart. Additionally, data on the types of fishing gear used to catch wreckfish were illustrated and analysed to determine the use of specific gear for each size class using bar charts.

Chi-square tests were used to determine the dependencies of habitat types, temporal abundance trends and employed fishing gears with the size classes of wreckfish.

## Results

Forty interviews took place between April 2022 and December 2024. Of these, 27 were addressed exclusively to fishers, including both professional and recreational fishers, while 13 were addressed to individuals who provided observational data, some of them being also engaged in recreational fishing. Interviewees were identified through local contacts or approached directly at local

harbours, using a convenience sampling approach (see Bennett *et al.*, 2024).

Reported wreckfish were divided into five size classes, based on those most commonly indicated by the respondents (0-0.4 kg, 0.4-1.5 kg, 1.5-3 kg, 3-10 kg, >10 kg). Five main habitat types were identified - pelagic neritic, pelagic oceanic, continental shelf break, slope, seamount - based on the depth at which specimens were recorded. For pelagic individuals, habitat classification was based on their position relative to the bottom, with pelagic neritic over the continental shelf and pelagic oceanic beyond it, over the continental slope and abyssal plain. Demersal habitats were defined by bottom depth. Specifically, the depth of 200 meters was used as a threshold to distinguish pelagic neritic from pelagic oceanic habitats, and continental shelf from slope habitats (Enrichetti *et al.*, 2019). There was a significant dependency between habitats and size classes ( $\chi^2 = 89.56$ ,  $df = 16$ ,  $P < 0.001$ ). In pelagic neritic and oceanic habitats, smaller size classes (0-0.4 kg and 0.4-1.5 kg ranges) were highly prevalent, with almost the same relative abundances. The occurrences of larger size classes were significantly lower and no records were registered for the >10 kg class. On the continental shelf break, smaller size classes were absent and only mid-to-large size classes were represented, with a dominance of the 3-10 kg class. In slope habitats, including canyons and cold-water coral frameworks of living and dead (thanatocenosis) *Desmophyllum pertusum* and *Madrepora oculata* (Bo *et al.*, 2023), the size distribution was similar to that of the shelf break, with the 3-10 kg class being dominant and a slightly higher frequency of 1.5-3 kg individuals. Seamounts were characterized only by the two largest size classes, with a dominance of the >10 kg class. The presence of each size class per habitat type is illustrated in Table 1.

In terms of the maximum number of specimens reported per encounter or fishing trip, smaller-sized individuals were observed and fished in groups of up to 50 specimens in pelagic habitats (e.g., Fig. 1).

Considering larger-sized individuals (>10 kg), the most remarkable catches were made in previously unexploited seamount habitats. Notably, 20 specimens of about 20 kg were caught in a single event at the Ulisse seamount in June 1973 (Bo *et al.*, 2021), while a maximum of 34 individuals were caught in a single event sometime between 1973 and 1976 off Capo Corso. The interviews revealed a decline in the occurrence of most size classes over time (Fig. 2). Although a decrease in abundance was the most common response for almost all size classes, there was a large incidence of 'NA', especially for the smaller classes. For the 0.4-1.5 kg wreckfish, 'NA' was the most frequent answer rather than a reported decline in abundance (Fig. 2).

There was no dependency between trends in abundance over time and size classes, i.e. the abundance of wreckfish tended to decline independently of size ( $\chi^2 = 8.63$ ,  $df = 12$ ,  $P = 0.734$ ). Respondents suggested that the decline of the smallest class individuals, reported by most interviewees to occur since the early 2000s, was likely due to harvest but also to a general reduction in floating



**Table 1.** Summary table showing the number of respondents reporting the presence of *Polyprion americanus* specimens, categorized by size class and habitat type.

Habitat type	Size class (kg)				
	0-0.4	0.4-1.5	1.5-3	3-10	>10
Pelagic Neritic	10	18	6	3	0
Pelagic Oceanic	9	15	6	2	0
Continental shelf-break	0	0	1	4	1
Slope	0	1	7	13	4
Seamount	0	0	0	5	7



**Fig. 1:** Around 30 juvenile wreckfish (0.4-0.5 kg, centre of the photo) were caught on a single fishing trip near an offshore buoy (pelagic oceanic habitat) in the early 1990s, alongside common dentex *Dentex dentex*, black seabream *Spondyliosoma cantharus*, and rudderfish *Centrolophus niger*. Photograph published with the consent of an interviewee, modified to preserve anonymity.



**Fig. 2:** Grouped bar plot showing the number of respondents reporting the trend (decreasing, stable, or increasing) over time for each identified wreckfish size class (kg) or not providing information (NA).

marine debris, with which small wreckfish are often associated. For larger individuals, respondents emphasized the critical impact of technological advancements in fishing gears. Large individuals, considered fishing trophies, are still caught around seamounts by fishers willing to travel far from the coast and spend significant amounts on fuel when weather conditions are ideal (i.e., flat calm seas and sunny weather).

The analysis of fishing gears used to catch wreckfish revealed a wide range of techniques employed differently across size classes. To clarify our results, we grouped fishing gears into nine categories (Fig. 3).

Bottom longlines and bottom fishing with rods (typically at depths of several hundred meters using electric reels) were reported as the most commonly used techniques, primarily targeting larger size classes. Following these, under-boat lines (baited or with artificial jigs) were frequently used to catch smaller individuals in pelagic habitats near the surface. When lines were not used, other tools such as spearguns, harpoons, and hand nets were employed to catch small pelagic wreckfish. Fishing techniques involving nets were reported less frequently. Both professional and recreational fishers have reported catching wreckfish using “lampara” net fishing that uses a light source, blue strobe lights attached to gillnets, or light baits paired with fishing lines and electric reels as both juveniles (Boero & Guzzardi, 2003) and adults (Ryall & Hargrave, 1984) are known to be attracted to lighted objects.


There was a significant dependency between fishing gear types and size classes ( $\chi^2 = 67.38$ ,  $df = 32$ ,  $P < 0.001$ , Fig. 4). Intermediate-sized individuals were caught using nearly all gear types. The 0.4-1.5 kg size class was primarily harvested at the surface with under-boat lines, while the 1.5-3 kg size class was most commonly caught with longlines. The smallest individuals (0-0.4 kg) were caught exclusively with surface fishing methods (spearfishing, harpooning, hand nets, under-boat lines, and seines), whereas the largest ones were almost exclusively

caught using longlines and bottom fishing. However, the 3-10 kg specimens were occasionally captured with gill-nets or trawling.

Most interviewees did not observe mature gonads in captured specimens, except for one recreational fisher who reported observing them in specimens of >10 kg during summer in slope habitats characterized by cold-water coral frameworks. Another fisher identified a 33 kg individual as female, though with immature gonads.

In terms of species associations, some respondents reported that wreckfish share their habitat with other species, particularly in their juvenile stage. Juvenile wreckfish were mainly observed under floating debris or buoys (Fig. 5), where they were frequently found alongside juveniles of the rudderfish *Centrolophus niger* or the imperial blackfish *Schedophilus ovalis*, occasionally with the pilotfish *Naucrates ductor*, and more rarely with the common dolphinfish *Coryphaena hippurus* or the tripletail *Lobotes surinamensis*. Larger demersal wreckfish were mainly found in areas where the blackspot seabream *Pagellus bogaraveo* was fished. This latter species likely represents a relevant component of the wreckfish diet, as several respondents reported instances of *P. bogaraveo* being preyed upon by wreckfish while hooked in fishing lines.

Interviewees reported that reaching wreckfish fishing grounds, located tens of miles offshore, requires significant time and costs. Nevertheless, interviewees reported also that the wreckfish commercial value remains fairly lower than that of most other species caught and marketed in the study area. For professional fishers, wreckfish is often caught as bycatch on gear targeting other species, such as *Brama brama*, *Lepidopus caudatus*, *Merluccius merluccius*, *P. bogaraveo*, *S. ovalis*, *Xiphias gladius*. Moreover, fishers reported that large adult individuals are not in demand, either by restaurants or customers, as their consumption results in excessive waste due to the high density of the skeletal structure and cranial bones, which contribute to a high percentage of inedible parts.



**Fig. 3:** Stacked bar plot showing the types of fishing gear and the number of respondents who reported using them by wreckfish size class (kg).



**Fig. 4:** Lollipop plots showing the types of fishing gear and the number of respondents who reported using them to target each of the five wreckfish size classes (kg).

## Discussion


In this study, Local Ecological Knowledge (LEK) provided insights into the wreckfish population in the Ligurian Sea. In particular, LEK helped improve the available knowledge on habitat use, assess conservation status, and identify the main types of fishing gear impacting wreckfish in the study area. This region is particularly suitable for studying deep-water species through LEK, as its narrow continental shelf and accessible deep-sea features (e.g., slopes and seamounts) enable frequent interactions between fishers and target species like the wreckfish. Our results highlight a clear ontogenetic shift in habitat use across three broad life stages: pelagic juveniles, demersal subadults, and large adults. These findings contribute to existing knowledge of the species' life cycle (Fig. 6) while also revealing new patterns specific to the Ligurian Sea.

Wreckfish juveniles (0-1.5 kg) were primarily found in pelagic habitats and typically associated with floating objects, as previously reported elsewhere (Sedberry *et al.*, 1999), with no apparent spatial segregation by size


between neritic and oceanic domains. This is likely related to the fact that the wreckfish is known to have a prolonged pelagic phase during which marine currents likely influence juvenile dispersal and population gene flow throughout its geographic range (Ball *et al.*, 2000). Such dynamics help to explain the occasional yet recurrent presence of juveniles in shallow areas such as the Gulf of Trieste in the northern Adriatic Sea (Bettoso *et al.*, 2020). Juveniles were found under various types of objects, including buoys - as previously documented in the literature (Relini *et al.*, 1994; Deudero & Morales-Nin, 2000; Andaloro *et al.*, 2007) - as well as branches, crates, debris, and even live or dead organisms such as the loggerhead sea turtle *Caretta caretta*. In some coastal fishing villages along the Ligurian Sea, the wreckfish is known as '*mangia-morti*' (literally 'dead-people eater') as juveniles were found under floating cadavers, often the remains of crew members from ships torpedoed and sunk during World War II (Gambazza S., personal communication).

The 1.5-3 kg size class marked the onset of the demersal phase, with individuals reported both in pelagic





**Fig. 5:** Objects under which juvenile *Polyprion americanus* were observed. (A) A branch colonized by the barnacle *Lepas anatifera*; (B) A loggerhead sea turtle *Caretta caretta*; (C) A mooring rope for dive boats at the wreck of the Haven oil tanker (a technical diving site) - Close-up of a juvenile in front of two *Thunnus thynnus* specimens; (D) A floating log; (E) A lost fishing gear marker; (F) A mooring buoy at the wreck of the Haven oil tanker; (G) An offshore buoy 2 miles from the coast; (H) An offshore oil platform. Pictures shared with the consent of the photographers: (A, B): Gianni Lucchi; (C): Alessandro Grasso; (D, E): Enrico Maggioni; (F): Marco Panico; (G): Paolo Cappucciati; (H): Vincenzo Paolillo.



**Fig. 6:** Life cycle of the wreckfish in the Ligurian Sea. Image created by Stefano Trainito, modified and used with permission.



waters and near the seabed, particularly along the shelf break and slope. This size class corresponds to individuals measuring 45-57 cm TL, which aligns with reported recruitment sizes in the literature - ranging from 46 to 66 cm TL - in the northern distribution range of the species (Atlantic and Mediterranean waters) (Roberts, 1977, 1989; Relini *et al.*, 1994; Sedberry *et al.*, 1998; Deudero & Morales-Nin, 2000; Machias *et al.*, 2003; Andaloro *et al.*, 2007). In fact, the timing of recruitment seems to be influenced by size, alongside ambient temperature, and is accompanied by morphological changes (Machias *et al.*, 2003).

Subadult wreckfish (3-10 kg) were primarily reported along the shelf break and slope habitats, while adults (>10 kg) were mostly found near seamounts. This distribution pattern suggests an ontogenetic habitat shift culminating in adult residency in offshore areas. In some regions, the slope is known as a relevant habitat for adult wreckfish and for spawning, which is reported to occur in deep waters (>400 m) (Glukhov & Zaferman, 1982; Sedberry *et al.*, 1999, 2006). The wreckfish is a gonochoric species (Roberts, 1989) that likely forms spawning aggregations. In the western North Atlantic, spawning occurs on the Blake Plateau, particularly in an area of high relief and upwelling known as the Charleston Bump, which rises from the continental slope (Brooks & Bane, 1978; Sedberry *et al.*, 1996). In the Southwestern Atlantic, potential spawning aggregations have been suggested along the continental slope (300-500 m) off southern Brazil, based on gonadal analysis and commercial catch data (Peres & Klippel, 2003). In South Africa, remote observations near deep cold-water corals on the Amathole shelf-edge have also indicated possible spawning aggregation behavior (Button *et al.*, 2021). These spawning sites share common environmental conditions, such as high pressure, absence of light, rocky bottoms, the presence of ridges/relief, and upwelling.

However, in the Ligurian Sea, the shelf break and slope do not appear to serve as primary spawning areas. While the 3-10 kg size class partially overlaps with the reported maturation range (8-11 kg, FishBase; Froese & Pauly, 2024), most individuals in this size class are likely still immature. This is further confirmed by fishers, who never observed mature gonads in individuals of this size. In contrast, the presence of larger fish on seamounts supports the hypothesis that spawning may occur in these offshore, structurally complex habitats.

Identifying spawning sites is essential for protecting critical habitats for reproduction, yet the depth range of wreckfish poses a challenge for locating these areas and few data exist on its reproductive activity. While research has been conducted on wreckfish reproductive development (age at maturity, gonadal development, and fecundity), no direct observations on its spawning behaviours exist, neither in captivity, where male breeding behaviours may be impaired (Papadaki *et al.*, 2018), nor in the wild. Wreckfish spawn over a protracted period, typically from late autumn to early spring, though timing varies by region. Reported spawning periods include November-May in the North-Atlantic (Sedberry *et al.*, 2006),

January-April in the Gulf of Naples (Bini, 1970), late autumn-winter in the North-Western Ionian Sea (Carbonara *et al.*, 2003) and February-March (late winter to early spring) in the Eastern Mediterranean (Cretan Sea) (Bini, 1970). However, spawning has also been reported in July (summer) in the Strait of Messina (Bini, 1970). Juvenile wreckfish are most abundant in surface waters during summer in Mallorca (July-August) and the Ligurian Sea (August) (Relini *et al.*, 1994; Deudero & Morales-Nin, 2000) and in autumn-winter (October-March) off Sicily (Andaloro *et al.*, 2007). In our study, most fishers reported never having found mature gonads in captured specimens, except one interviewee. The near absence of reported mature gonads in caught fish is likely due to fishing trips to seamounts occurring only during the “good season” (late spring and summer), a period that does not coincide with the primary spawning season identified in recent literature.

While the IUCN classifies wreckfish as Data Deficient in the Mediterranean and Vulnerable in Italy, our findings indicate a population decline across all size classes or life stages. Overfishing is likely to be a key factor, with numerous fishing techniques directly or indirectly targeting wreckfish. During the pelagic phase, juveniles are caught using hand lines or rod-and-reel setups, often baited with natural or artificial lures. Fishers described juvenile wreckfish as highly voracious, making them easy to catch in large numbers. Other fishing methods also impact this stage, contributing to its vulnerability. Respondents reported catching dozens of juvenile wreckfish in single events, suggesting that this species may be at risk of overexploitation at early life stages. Uncontrolled catches of juveniles could lead to ‘growth overfishing’, where excessive capture of juveniles prevents individuals from reaching reproductive maturity. This is particularly concerning given the minimum legal size of 45 cm, which is below the species’ size at maturity (~80-90 cm total length). The fishing of juvenile wreckfish is likely to have occurred since historical times, as their presence in surface waters makes them more accessible than adults. This is supported by an artwork from Santorini, Greece (1500 B.C.), which depicts a man carrying multiple common dolphinfish *C. hippurus* (Begossi & Caires, 2015), a species that shares the pelagic habitat with wreckfish. Juvenile wreckfish were reported to share “shaded environments” with the juveniles of other species known to be attracted to floating objects (Castro *et al.*, 2001), including *C. niger* and *S. ovalis*, both common in the Ligurian Sea, and the rare *L. surinamensis*. Similarly, *N. ductor* and *C. hippurus* are frequently associated with floating objects and may be vulnerable to the same fishing techniques during juvenile stages. Respondents also suggested that juvenile wreckfish encounters have declined due to reduced marine debris. Legislative actions (EU, 2000), mild negative trends in summer rainfall in the northern Ligurian Sea (1998-2017) (Kalimeris & Kolios, 2019), and a general decrease in waste may have reduced the number of floating objects serving as shelters for juveniles, while also offering fewer opportunities for people traveling at sea to encounter wreckfish near float-

ing wreckage.

As wreckfish recruit on the continental shelf, slope, or seamounts, they become susceptible to longline and bottom fishing. Longlines are used by both professional and recreational fishers, while bottom fishing is mostly practiced by advanced recreational fishers using highly technological equipment (e.g., electric reel). Unlike in other regions, like the Southern Atlantic (Gauvin *et al.*, 1994), wreckfish is rarely the primary target of fishing operations in the study area. Instead, it is usually caught incidentally during multi-species fishing for *P. bogaraveo*, *Brama brama*, *Lepidopus caudatus*, *Merluccius merluccius*, *S. ovalis*, *Xiphias gladius*, and other less common species. In shelf and slope habitats, medium-sized wreckfish (1.5–10 kg) are also caught as bycatch in gillnets or trawling operations.

The main reason for the decline of medium- and large-sized wreckfish is thus likely overfishing. Furthermore, the deep-water habitats inhabited by wreckfish are home to fragile structural organisms, such as cold-water corals, which can be easily damaged by fishing gear (Bo *et al.*, 2021). Cold-water coral habitats are recognised as vulnerable habitats (Bo *et al.*, 2021) that serve as nurseries and spawning grounds for many commercially valuable species (Baillon *et al.*, 2012).

Our study, in conclusion, highlights the importance of protecting habitats critical to different ontogenetic stages of wreckfish, from pelagic zones to offshore, deep-water spawning habitats, and suggests the need for enhanced protection measures in these critical areas. These findings also emphasize the need to update the minimum legal size for this species in the Mediterranean Sea to prevent overfishing, as both juveniles and adults are particularly vulnerable to fishing pressure. Although further biological and ecological data are needed to advance our knowledge on the species and assess its conservation status, the LEK-based information presented here provides new insights into the wreckfish life cycle and its fishery in the Ligurian Sea. Notably, the consistency of responses across different sea users (i.e., fishers and other categories of observers) reinforces the reliability of the patterns identified. Nonetheless, some limitations of LEK must be acknowledged, such as the lack of information on sex or maturity. Despite these gaps, the results provide valuable data for local population assessments and the development of management and conservation measures at the Mediterranean scale.

## Acknowledgements

We thank all the interviewees, Giulia Calogero and Elia Biasissi from the Menkab Association, and Simonpietro Canese for generously sharing their knowledge. We also acknowledge the photographers (Giorgio Barsotti, Paolo Cappucciati, Roberto Croce, Alessandro Grasso, Gianni Lucchi, Enrico Maggioni, Gabriella Motta, Marco Panico, Vincenzo Paolillo) for sharing their valuable observations. We dedicate this work to Anni Croce, in memory of his remarkable discoveries and contributions.

This work was supported by the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 - Call for tender No. 3138 of 16 December 2021, rectified by Decree n.3175 of 18 December 2021 of Italian Ministry of University and Research funded by the European Union - NextGenerationEU; Award Number: Project code CN\_00000033, Concession Decree No. 1034 of 17 June 2022 adopted by the Italian Ministry of University and Research, CUP C63C22000520001, Project title “National Biodiversity Future Center - NBFC”.

## References

- Andaloro, F., Campo, D., Castriota, L., Sinopoli, M., 2007. Annual trend of fish assemblages associated with FADs in the southern Tyrrhenian Sea. *Journal of Applied Ichthyology*, 23, 258–263.
- Asensio, L., Gonzalez, I., Fernandez, A., Rodriguez, M. A., Lobo, E. *et al.*, 2002. Application of Random Amplified Polymorphic DNA (RAPD) Analysis for Identification of Grouper (*Epinephelus guaza*), Wreck Fish (*Polyprion americanus*), and Nile Perch (*Lates niloticus*) Fillets. *Journal of Food Protection*, 65 (2), 432–435.
- Azzurro, E., Sbragaglia, V., Cerri, J., Bariche, M., Bolognini, L. *et al.*, 2019. Climate change, biological invasions, and the shifting distribution of Mediterranean fishes: A large-scale survey based on local ecological knowledge. *Global Change Biology* 25 (8), 2779–2792.
- Baillon, S., Hamel, J.-F., Wareham, V.E., Mercier, A., 2012. Deep cold-water corals as nurseries for fish larvae. *Frontiers in Ecology and the Environment*, 10 (7), 351–356.
- Ball, A.O., Sedberry, G.R., Zatzoff, M.S., Chapman, R.W., Carlin, J.L., 2000. Population structure of the wreckfish *Polyprion americanus* determined with microsatellite genetic markers. *Marine Biology*, 137, 1077–1090.
- Begossi, A., Caires, R., 2015. Art, fisheries and ethnobiology. *Journal of Ethnobiology and Ethnomedicine*, 11, 16.
- Bennett, N.J., Calò, A., Guidetti, P., Milazzo, M., Prato, G. *et al.*, 2024. Access and well-being in small-scale fisheries. *Marine Policy*, 169, 106328.
- Bethoux, J.P., Gentili, B., Morin, P., Nicolas, E., Pierre, C. *et al.*, 1999. The Mediterranean Sea: a miniature ocean for climatic and environmental studies and a key for the climatic functioning of the North Atlantic. *Progress in Oceanography*, 44 (1–3), 131–146.
- Bettoso, N., Faresi, L., Felluga, A., Lipej, L., 2020. On the occurrence of the wreckfish *Polyprion americanus* in the Gulf of Trieste (northern Adriatic Sea). *Acta Adriatica*, 61 (1), 107–112.
- Bini, G. (Ed.), 1970. *Polyprion americanum*. p.63–64. In: *Atlante Dei Pesci Delle Coste Italiane. Volume VII-Osteitti*. Mondo sommerso, Rome.
- Boero, M., Guzzardi, R., 2003. *Pesci salati: La pesca delle acciughe e la loro salagione*. Associazione culturale Storie di barche, 93 pp.
- Bo, M., Coppari, M., Betti, F., Enrichetti, F., Bertolino, M. *et al.*, 2021. The high biodiversity and vulnerability of two Mediterranean bathyal seamounts support the need for creating offshore protected areas. *Aquatic Conservation: Ma-*

- rine and Freshwater Ecosystems*, 31 (3), 543-566.
- Bo, M., Enrichetti, F., Betti, F., Gay, G., Quarta, G. *et al.*, 2023. The cold-water coral province of the eastern Ligurian Sea (NW Mediterranean Sea): historical and novel evidences. *Frontiers in Marine Science*, 10, 1114417.
- Brooks, D.A., Bane, J.M., Jr, 1978. Gulf Stream deflection by a bottom feature off Charleston, South Carolina. *Science*, 201 (4326), 1225-1226.
- Button, R.E., Parker, D., Coetzee, V., Samaai, T., Palmer, R.M. *et al.*, 2021. ROV assessment of mesophotic fish and associated habitats across the continental shelf of the Amathole region. *Scientific Reports*, 11, 18171.
- Canepa, E., Pensieri, S., Bozzano, R., Faimali, M., Traverso, P. *et al.*, 2015. The ODAS Italia 1 buoy: More than forty years of activity in the Ligurian Sea. *Progress in Oceanography*, 135, 48-63.
- Canese, S., Bava, S., 2015. The decline of top predators in deep coral reefs. p.67–68. In: *1st Mediterranean Symposium on the Conservation of Dark Habitats (Portorož, Slovenia, 31 October 2014)*, Langar, H., Bouafif, C., A., O. (Eds.), RAC/SPA publ., Tunis.
- Carbonara, P., Costantino, G., Giovine, G., Lembo, G., Spedicato, M.T. *et al.*, 2003. Some aspects of the life history of *Polyprión americanus* (Schneider, 1801) along the coasts of the north western Ionian Sea. *Biologia Marina Mediterranea*, 10 (2), 102-112.
- Castro, J.J., Santiago, J.A., Santana-Ortega, A.T., 2001. A general theory on fish aggregation to floating objects: An alternative to the meeting point hypothesis. *Reviews in Fish Biology and Fisheries*, 11 (3), 255–277.
- Cattaneo-Vietti, R., Orsi Relini, L., Wurtz, M., 1985. *La pesca in Liguria*. Centro Studi Unioncamere liguri, Genova, 190 pp.
- Cattaneo-Vietti, R., Albertelli, G., Aliani, S., Bava, S., Bavestrello, G. *et al.*, 2010. The Ligurian Sea: present status, problems and perspectives. *Chemistry in Ecology*, 26, 319–340.
- Collette, B., Fernandes, P., Heessen, H., Smith-Vaniz, W.F., 2015. *Polyprión americanus* (European assessment). The IUCN Red List of Threatened Species 2015: e.T43972A45795607. (Accessed April 21, 2022).
- Deudero, S., Morales-Nin, B., 2000. Occurrence of *Polyprión americanus* under floating objects in western Mediterranean oceanic waters, inference from stomach contents analysis. *Journal of the Marine Biological Association of the United Kingdom*, 80 (4), 751-752.
- Diogo, H., Pereira, J.G., 2013. Recreational boat fishing pressure on fish communities of the shelf and shelf break of Faial and Pico Islands (Azores Archipelago): implications for coastal resource management. *Acta Ichthyologica et Piscatoria*, 43 (4), 267-276.
- D'Onghia, G., Maiorano, P., Carlucci, R., Capezzuto, F., Carluccio, A. *et al.*, 2012. Comparing deep-sea fish fauna between coral and non-coral “megahabitats” in the Santa Maria di Leuca cold-water coral province (Mediterranean Sea). *PLoS One*, 7 (9), e44509.
- D'Onghia, G., Capezzuto, F., Carluccio, A., Carlucci, R., Giove, A. *et al.*, 2015. Exploring composition and behaviour of fish fauna by *in situ* observations in the Bari Canyon (Southern Adriatic Sea, Central Mediterranean). *Marine Ecology*, 36 (3), 541-556.
- Enrichetti, F., Dominguez-Carrió, C., Toma, M., Bavestrello, G., Betti, F. *et al.*, 2019. Megabenthic communities of the Ligurian deep continental shelf and shelf break (NW Mediterranean Sea). *PLoS One*, 14 (10), e0223949.
- EU, 2000. *Directive 2000/59/EC of the European Parliament and of the Council of 27 November 2000 on port reception facilities for ship-generated waste and cargo residues - Commission declaration*. Official Journal of the European Communities L332. 81-90.
- European Council, 2006. *Council Regulation (EC) No. 1967/2006 concerning management measures for the sustainable exploitation of fishery resources in the Mediterranean Sea, amending Regulation (EEC) No. 2847/93 and repealing Regulation (EC) No. 1626/94*. Official Journal of the European Union L 409.
- Froese, R., Pauly, D., 2024. Fish Base. <http://www.fishbase.org>. (Accessed February 13, 2024).
- Garibaldi, F., 2015. By-catch in the mesopelagic swordfish longline fishery in the Ligurian Sea (Western Mediterranean). *Collective Volume of Scientific Papers ICCAT*, 71 (3), 1495-1498.
- Gauvin, J.R., Ward, J.M., Burgess, E.E., 1994. Description and evaluation of the wreckfish (*Polyprión americanus*) fishery under individual transferable quotas. *Marine Resource Economics*, 9, 99–118.
- Glukhov, A.A., Zaferman, M.L., 1982. Observations on the behaviour of *Polyprión americanus* (Schneider) (Serranidae). *Voprosy Ikhtologii*, 22 (2), 334-336.
- Hamilton, R., de Mitcheson, Y.S., Aguilar-Perera, A., 2012. The role of local ecological knowledge in the conservation and management of reef fish spawning aggregations. p.331-369. In: *Reef Fish Spawning Aggregations: Biology, Research and Management*. Sadovy de Mitcheson, Y., Colin, P.L. (Eds). Springer Netherlands, Dordrecht.
- Hardy, J.D., 1978. *Development of fishes of the Mid-Atlantic Bight. An atlas of egg, larval and juvenile stages. Vol. 3: Aphredoderidae through Rachycentridae*. Fish and Wildlife Service, U.S. Department of the Interior, 402 pp.
- Imhoof-Blumer, F., Keller, O., 1889. *Tier-und Pflanzenbilder auf Münzen und Gemmen des klassischen Altertums*. Teubner, Leipzig, 240 pp.
- Johannes, R.E., 1978. Traditional marine conservation methods in Oceania and their demise. *Annual Review of Ecology and Systematics*, 9 (1), 349-364.
- Johannes, R.E., Freeman, M.M.R., Hamilton, R.J., 2000. Ignore fishers' knowledge and miss the boat. *Fish and Fisheries*, 1 (3), 257-271.
- Kalimeris, A., Kolios, S., 2019. TRMM-based rainfall variability over the Central Mediterranean and its relationships with atmospheric and oceanic climatic modes. *Atmospheric Research*, 230, 104649.
- Lejeusne, C., Chevaldonné, P., Pergent-Martini, C., Boudouresque, C.F., Pérez, T., 2010. Climate change effects on a miniature ocean: the highly diverse, highly impacted Mediterranean Sea. *Trends in Ecology & Evolution*, 25 (4), 250-260.
- Linares, F., Pérez Rial, E., Rodríguez, J.L., Pazos, G., Álvarez-Blázquez, B., 2021. Biometric parameters and biochemical composition of wild wreckfish (*Polyprión ameri-*



- canus). *Marine Biology Research*, 17 (3), 234-246.
- Lytton, A.R., Ballenger, J.C., Reichert, M.J., Smart, T.I., 2015. Age validation of the North Atlantic stock of wreckfish (*Polyprion americanus*), based on bomb radiocarbon ( $^{14}\text{C}$ ), and new estimates of life history parameters. *Fishery Bulletin*, 114 (1), 77-88.
- Machias, A., Somarakis, S., Papadroulakis, N., Spedicato, M.-T., Suquet, M. *et al.*, 2003. Settlement of the wreckfish (*Polyprion americanus*). *Marine Biology*, 142 (1), 45-52.
- Mavruk, S., Saygu, İ., Bengil, F., Alan, V., Azzurro, E., 2018. Grouper fishery in the Northeastern Mediterranean: An assessment based on interviews on resource users. *Marine Policy*, 87, 141-148.
- Menni, R.C., López, H.L., 1979. Biological data and otolith (sagitta) morphology of *Polyprion americanus* and *Schedophilus griseolineatus* (Osteichthyes, Serranidae and Centrolophidae). *Studies on Neotropical Fauna and Environment*, 14 (1), 17-32.
- Papadaki, M., Peleteiro, J.B., Alvarez-Blázquez, B., Rodríguez Villanueva, J.L., Linares, F. *et al.*, 2018. Description of the annual reproductive cycle of wreckfish *Polyprion americanus* in captivity. *Fishes*, 3 (4), 43.
- Papandroulakis, N., Suquet, M., Spedicato, M., Machias, A., Fauvel, C. *et al.*, 2004. Feeding Rates, Growth Performance and Gametogenesis of Wreckfish (*Polyprion americanus*) kept in Captivity. *Aquaculture International: Journal of the European Aquaculture Society*, 12, 395-407.
- Peres, M.B., Klippel, S., 2003. Reproductive biology of southwestern Atlantic wreckfish, *Polyprion americanus* (teleostei: Polyprionidae). *Environmental Biology of Fishes* 68 (2), 163-173.
- Pérez, E., Linares, F., Rodríguez Villanueva, J.L., Vilar, A., Mylonas, C.C. *et al.*, 2019. Wreckfish (*Polyprion americanus*). New knowledge about reproduction, larval husbandry, and nutrition. Promise as a new species for aquaculture. *Fishes*, 4, 14.
- Relini, G., 2007. La pesca batiale in Liguria. *Biologia Marina Mediterranea*, 14 (2), 190-244.
- Relini, M., Orsi Relini, L., Relini, G., 1994. An offshore buoy as a FAD in the Mediterranean. *Bulletin of Marine Science*, 55 (2-3), 1099-1105.
- Relini, G., Tunesi, L., Vacchi, M., Andaloro, F., D'Onghia, G. *et al.*, 2017. Lista Rossa IUCN dei Pesci ossei marini Italiani. Comitato Italiano IUCN e Ministero dell'Ambiente e della Tutela del Territorio e del Mare, Roma, 52 pp.
- Rizzo, G.E. (Ed.), 1946. *Monete greche della Sicilia: Descritte e illustrate da Giulio Emanuele Rizzo. Vol. 1,2.* Arnaldo Forni Editore, Bologna.
- Roberts, C.D., 1977. The wreckfish *Polyprion americanus* (Schneider, 1801) in Irish waters: An underwater sighting and review of the Irish records. *The Irish Naturalists' Journal*, 19 (4), 108-112.
- Roberts, C.D., 1989. Reproductive mode in the percomorph fish genus *Polyprion* Oken. *Journal of Fish Biology*, 34 (1), 1-9.
- Ryall, P.J.C., Hargrave, B.T., 1984. Attraction of the Atlantic wreckfish (*Polyprion americanus*) to an unbaited camera on the Mid-Atlantic Ridge. *Deep-Sea Research. Part A, Oceanographic Research Papers*, 31 (1), 79-83.
- Sadovy, Y., 2003. *Polyprion americanus*. The IUCN Red List of Threatened Species 2003: e.T43972A10845280. (Accessed October 2, 2023).
- Sedberry, G.R., Carlin, J.L., Menezes, G.M., 1998. Movements of a pelagic-phase wreckfish, *Polyprion americanus* (Schneider, 1801) as indicated by tag and recapture. *Arguipelago. Life and Marine Sciences*, 69-72.
- Sedberry, G.R., Carlin, J.L., Chapman, R.W., Eleby, B., 1996. Population structure in the pan-oceanic wreckfish, *Polyprion americanus* (Teleostei: Polyprionidae), as indicated by mtDNA variation. *Journal of Fish Biology*, 49, 318-329.
- Sedberry, G.R., Andrade, C.A.P., Carlin, J.L., Chapman, R.W., Luckhurst, B.E., *et al.*, 1999. Wreckfish *Polyprion americanus* in the North Atlantic: fisheries, biology, and management of a widely distributed and long-lived fish. p.27-50. In: *American Fisheries Society Symposium*, 23, 27-50.
- Sedberry, G.R., Pashuk, O., Wyanski, D.M., Stephen, J., Weinbach, P., 2006. Spawning locations for Atlantic reef fishes off the southeastern U.S. *Gulf and Caribbean Fisheries Institute*, 57, 464-514.
- Silvano, R.A.M., Nora, V., Andreoli, T.B., Lopes, P.F.M., Begossi, A., 2017. The "ghost of past fishing": Small-scale fisheries and conservation of threatened groupers in sub-tropical islands. *Marine Policy*, 75, 125-132.
- Silvano, R.A.M., Baird, I.G., Begossi, A., Hallwass, G., Huntington, H.P. *et al.*, 2023. Fishers' multidimensional knowledge advances fisheries and aquatic science. *Trends in Ecology & Evolution*, 38 (1), 8-12.
- Smith, J.L.B. (Ed.), 1949. *The sea fishes of Southern Africa*. Central News Agency, Ltd, Cape Town, 580 pp.
- Tortonese, E., 1975. *Polyprion americanus*. p.60-62. In: *Fauna d'Italia, "Osteichthyes": Pesci Ossei*, Tortonese, E. (Ed.). Calderini, Bologna.
- Wakefield, C.B., Newman, S.J., Boddington, D.K., 2013. Exceptional longevity, slow growth and late maturation infer high inherent vulnerability to exploitation for bass groper *Polyprion americanus* (Teleostei: Polyprionidae). *Aquatic Biology*, 18 (2), 161-174.
- Ward, J., 1902. *Greek coins and their parent cities*. Hill, G.F. (Ed). J. Murray, London.
- Yokes, B., Kara, M.H., Bizsel, C., Kada, O., Bariche, M. *et al.*, 2011. *Polyprion americanus* (Mediterranean assessment). The IUCN Red List of Threatened Species 2011: e.T43972A10845635. (Accessed October 2, 2023).

## Appendix

This document contains the two types of questionnaires used in the study: the Fishing Data questionnaire and the Observation Data questionnaire.

### FISHING DATA QUESTIONNAIRE

#### INFORMATION ON CATCHES

1. Have you ever caught wreckfish?
2. The first time you caught it, how could you be sure it was this species?
3. In which years and seasons did you catch it?
4. How many specimens (approximately) have you caught, and where? Please indicate the locations.
5. What gear did you use? Has it remained the same over the years? (Please specify the exact years, if possible)
6. What is the size range (length/weight) of the specimens you have caught?
  - Minimum size:
  - Most common size:
  - Maximum size:
7. Have you ever opened (gutted) the captured specimens?  
If so:
  - Have you identified male/female specimens and/or observed eggs/milt?
  - Have you recognized what the specimen has eaten?
8. In general, where is the wreckfish most likely to be caught?

Wreckfish size	Season	Habitat	Depth	Site (whenever possible)

9. Over time, how has the trend in catches changed?

From what year to what year		Size
<input type="checkbox"/>	Stable	
<input type="checkbox"/>	Increasing	
<input type="checkbox"/>	Decreasing	
<input type="checkbox"/>	NA	

10. Do you remember any exceptional catches of other fish species (very large ones and/or with eggs/milt)?

#### SOCIO-ECONOMIC INFORMATION

11. Have you ever sold wreckfish, and at what price?
12. During the same period, at what price did you sell the following:
  - Sea bass \_\_\_\_\_
  - Dusky grouper \_\_\_\_\_
  - Common dentex \_\_\_\_\_
13. Who did you mainly sell your catch to?
  - Private individuals \_\_\_\_\_ %
  - Restaurateurs/Hoteliers \_\_\_\_\_ %
  - Wholesalers \_\_\_\_\_ %
  - Fishers \_\_\_\_\_ %

14. Who do you currently sell the specimens to, and how much do you earn from selling them? Is there demand for them?
15. Do you know if there is a minimum catch size?

## OBSERVATION DATA QUESTIONNAIRE

### INFORMATION ON OBSERVATIONS

1. Have you ever seen a wreckfish?
2. The first time you saw it, how could you be sure it was this species?
3. In which years and seasons did you observe it?
4. Total number of trips (from year \_\_\_\_ to year \_\_\_\_):
5. Number of trips/observation frequency where floating wreckage was recorded **without** wreckfish:
6. Number of trips/observation frequency where floating wreckage was recorded **with** wreckfish:
7. How many specimens (approximate number) have you seen in total, and in which areas? Please indicate the sites, whenever possible.
8. What is the minimum, maximum, and most frequently observed number of specimens per floating wreckage/site?
  - Minimum n°/range:
  - Most common n°/range:
  - Maximum n°/range:
9. What is the size range (length/weight) of the observed specimens?
  - Minimum size:
  - Most common size:
  - Maximum size:
10. What type of floating wreckage were the wreckfish found under?
11. Were other species present as well?
12. In general, where is it most likely to see wreckfish?

Wreckfish size	Season	Habitat	Depth	Site (whenever possible)

13. Over time, how has the frequency of observations changed?

	From what year to what year	Size
<input type="checkbox"/>	Stable	
<input type="checkbox"/>	Increasing	
<input type="checkbox"/>	Decreasing	
<input type="checkbox"/>	NA	



**Specific record information of *Polyprion americanus***  
*especially if more than one individual was observed*

	Observation #1	Observation #2	Observation #3	Observation #4	Observation #5
Photo ID (if available)					
Size (total length or weight)					
N. of specimens					
Wreckage type					
Area or depth					
Weather conditions					
Date or season					
Co-occurring species					