DOI: https://doi.org/10.12681/mms.41614

Westernmost record of *Caulerpa taxifolia* var. *distichophylla* (Chlorophyta, Caulerpaceae) in the Mediterranean: a 20-year journey

Jose MARTÍNEZ-GARRIDO^{1*}, Diego MORENO^{2*}, Irene ROJO¹, Manuel MANCHADO³, Agustín BARRAJÓN², and Jaime BERNARDEAU-ESTELLER¹

¹ Grupo de Ecología de Angiospermas Marinas. Instituto Español de Oceanografía,
 Centro Oceanográfico de Murcia (IEO-CSIC), c/ El Varadero, 1, 30740 - Lo Pagan, Murcia, Spain
 ² Agencia de Medio Ambiente y Agua de Andalucía, M.P./ Consejería de Sostenibilidad y Medio Ambiente,
 Junta de Andalucía, c/Johann G. Gutenberg, 1, 41092, Sevilla, Spain
 ³ Instituto de Formación Agraria y Pesquera de Andalucía, El Toruño,
 Camino del Tiro Pichón, 11500 - El Puerto de Santa María, Cádiz, Spain

Corresponding author: Jose MARTÍNEZ-GARRIDO; jose.martinez@ieo.csic.es

*These authors contributed equally to this paper

Contributing Editor: Vasillis PAPATHANASIOU

Received: 28 May 2025; Accepted: 10 September 2025; Published online: 15 October 2025

Abstract

The invasive alga *Caulerpa taxifolia* has been detected for the first time in the Cabo de Gata-Níjar Natural Park (Andalusia, southern Spain), after more than 30 years of active monitoring for invasive species. Morphological and phylogenetic analyses identified the taxon as *C. taxifolia* var. *distichophylla* (Sonder) Verlaque, Huisman & Procaccini 2013, a variety native to Australia and distinct from the aquarium strain introduced in Monaco in 1984. This variety is part of an independent invasion originating in the eastern Mediterranean, first reported in Syria in 2003, which since then has spread westward through the southern Mediterranean Sea. This record from Andalusia represents a significant westward range extension from previous occurrences in Tunisia (2015) and Sardinia, Italy (2016). In just two decades, the species has crossed the Mediterranean from the eastern to the westernmost coasts, demonstrating a remarkable capacity for long-distance dispersal and adaptation to a wide range of habitats. However, compared to the aquarium-released *C. taxifolia* strain, it differs in size, thermal tolerance, and potentially in reproductive traits and secondary metabolite profiles. These differences may determine its potential for spreading and its ecological impact during the ongoing invasion.

Keywords: alien species; Andalusia; Caulerpa taxifolia var. distichophylla; Mediterranean Sea.

Introduction

The genus *Caulerpa* is represented in the Mediterranean Sea by a single native species, *C. prolifera* (Forsskål) J.V. Lamouroux, and seven non-indigenous taxa (in alphabetical order): *C. chemnitzia* (Esper) J.V. Lamouroux, *C. cylindracea* Sonder, *C. mexicana* Sonder *ex* Kützing, *C. racemosa* f. *requienii* (Montagne) Weber Bosse, *C. scalpelliformis* (R. Brown *ex* Turner) C. Agardh, and *C. taxifolia* (M. Vahl) C. Agardh, *C. taxifolia* var. *distichophylla* (Sonder) Verlaque, Huisman and Procaccini (Verlaque *et al.*, 2015). Among them, *C. taxifolia* has been included in the list of the 100 "Worst Invasive" species in the Mediterranean Sea (Streftaris & Zenetos, 2006).

Caulerpa taxifolia, a tropical marine alga, was accidentally released from the aquariums of the Musée Océanographique in Monaco in 1984 (Meinesz & Hesse,

1991). Following its introduction, it has spread rapidly across the Mediterranean Sea, reaching the French coast, Italy (extending to Messina), and Spain (Balearic Islands) in 1992 (Pou *et al.*, 1993), Croatia in 1994 (Span *et al.*, 1998), with Tunisia in 2000 (Langar *et al.*, 2000) and Turkey (Turan *et al.*, 2011).

The area covered by *C. taxifolia* increased dramatically, from just 1 hectare in 1989 (limited to France and Monaco) to over 13,000 hectares by 2000 across multiple Mediterranean countries (Meinesz *et al.*, 2001), causing a significant impact on native communities. However, after this phase of exponential expansion, its dispersal and colonisation capacity declined markedly, even leading to its disappearance in many of the previously invaded areas.

The invasive alga *Caulerpa taxifolia* var. *disticho-phylla*, native to the Pacific, was first recorded in the Mediterranean Sea along the Syrian coast in 2003. Ini-

tially, it was misidentified as a dwarf form of C. mexicana or C. taxifolia (RAC/SPA-UNEP/MAP, 2003; Bitar et al., 2010, 2017). After that, a population detected in the Gulf of Iskenderun (Turkey) in 2006 was analysed in detail, revealing that it had thinner thalli compared to the more robust *C. taxifolia* released from the aquariums already present in the region. Its morphology closely resembled that of the Australian species C. distichophylla (Sonder, 1845), but due to insufficient molecular distinction, it was initially classified as C. taxifolia (Famà et al., 2002; Cevik et al., 2007). Subsequent molecular studies revealed genetic differences, leading to the designation of C. taxifolia var. distichophylla, characterised by narrow, elongated thalli and fine rhizoids (Jongma et al., 2013). The variety was reported in the south of Sicily (Punta Bracetto and Marina di Ragusa) in 2007 (Jongma et al., 2013), and later in northern Sicily (Sant'Ambrogio) in 2013 (Musco et al., 2014). Its spread has been rapid throughout the central and eastern Mediterranean basins, reaching Cyprus in 2009 (Çiçek et al., 2013), Rhodes (Greece) in 2010 (Aplikioti et al., 2016), Malta in 2013 (Schembri et al., 2015), Tunisia in 2015 (Chartosia et al., 2018), Calabria and Sardinia (Italy) in 2016 (Di Martino et al., 2018), Lebanon in 2016 (Bitar et al., 2017), and Libya in 2017 (Shakman et al., 2017). In Sicily, due to the extensive proliferation recorded in recent years (Picciotto et al., 2016; Mannino & Balistreri, 2017; Mannino et al., 2019), it is now considered fully naturalised. Recently, it has also expanded along the Turkish Aegean coast (Mutlu et al., 2025), reaching the Marmara Sea in 2022 (Taşkın et al., 2023; Zenetos et al., 2025). In contrast to the strain of C. taxifolia from the Monaco aquarium, the variety distichophylla is currently in an expansion phase. Since its initial detection in 2003 on the easternmost coasts of the Mediterranean, it has shown a continuous pattern of westward spread.

The region of Andalusia (southern Spain) is located at the southwestern limit of the Mediterranean Sea. Owing to its considerable distance from the Suez Canal, the primary entry source of exotic marine species into the Mediterranean Sea, the region remained relatively unaffected by invasive algae for years. In response to growing concern caused by the rapid expansion of C. taxifolia in the western Mediterranean, early detection efforts were carried out between 1994 and 1996 along the Andalusian coastline, particularly focusing on the coast of Almería and Alborán Island. These surveys, funded by the European LIFE Programme, did not detect any presence of C. taxifolia in the area (Ballesteros et al., 1996; Moreno, 2009, 2010). However, this situation has changed in recent decades with the arrival and establishment of several non-indigenous macroalgae with highly invasive potential in the region, such as Asparagopsis taxiformis (Báez et al., 2001), Caulerpa cylindracea (Moreno, 2010), and Rugulopteryx okamurae (Altamirano et al., 2016), which have been included with C. taxifolia in the Spanish Catalogue of Invasive Alien Species (Royal Decree 630/2013 and subsequent updates). Despite this increasing trend of macroalgal invasions, C. taxifolia var. distichophylla has only been recorded in the eastern Mediterranean Sea, with its westernmost occurrence reported in the Sardinia Channel. In this study, we provide the first evidence of its recent expansion to the westernmost part of the Mediterranean Sea, along with a detailed analysis of its distribution and habitat preferences.

Material and Methods

Sampling sites

The study was conducted in the coastal waters of Almería, Granada and Málaga (Alborán Sea, Andalusia). This area is strongly influenced by Atlantic waters, resulting in relatively cooler and eutrophic conditions that support high algal productivity and diverse macroalgal communities. The monitoring and assessment of seagrass habitats along the coast of Andalusia was carried out in June and July 2023 by the Spanish Institute of Oceanography of the Spanish National Research Council (IEO-CSIC) in the context of the Marine Strategy Framework Directive (MSFD). A total of 46 linear km of seabed were recorded using georeferenced underwater video, 74 ground-truth points were surveyed, and 49 stations were sampled via scuba diving, covering approximately 300 km of coastline from Terreros (Almería) to Nerja (Málaga). On 5 July, 2023, a putative specimen of *Caulerpa taxifolia* var. distichophylla was collected at Punta de la Media Naranja (Cabo de Gata-Níjar Natural Park, Almería), with its presence also detected at two nearby sites (<250 m).

Additional sampling was conducted for monitoring of seagrass meadows and exotic species in Andalusia under the Sustainable Management Programme for the Andalusian Marine Environment (Department of Sustainability and Environment, Junta de Andalucía). Several scuba dives were performed from 18 July, 2023 to 6 August, 2024 to assess the distribution and abundance of the invasive taxa. A total area of 2 km² between Punta de la Media Naranja and El Plomo was prospected and 10 locations with a putative *Caulerpa taxifolia* var. *distichophylla* were detected. For each site, the geographic coordinates, depth, type of substrate, abundance and any other relevant characteristics such as pigmentation and size were recorded (Table 1).

Morphological identification

Specimens were morphologically identified in the laboratory based on diagnostic traits (stolon, fronds, pinnules, rhizoidal pillars, midrib) following the criteria used by Jongma *et al.*, (2013). Morphological characterisations were conducted on 10 individuals from each site. Furthermore, the thalli were photographed both *in situ* and in the laboratory. Six herbarium voucher specimens were deposited at the Research Centre for Scientific Collections of the University of Almería (CECOUAL), with collection numbers: HUAL-ALGA259 to HUAL-ALGA264.

Table 1. Records of Caulerpa taxifolia var. distichophylla in the Cabo de Gata-Níjar Natural Park (Andalusia, Spain).

Site	Date	Coordinates*	Depth (m)	Substrata	Abundance	Observations	Source
Punta de la Media Naranja	2023-07-05	36.93280° N 01.91555° W	10	C. nodosa	Scarce	First observation and samples	IEO-CSIC
Punta de la Media Naranja	2023-07-05	36.93182° N 01.91557° W	14	C. nodosa	Scarce	Record	IEO-CSIC
Punta de la Media Naranja	2023-07-05	36.93137° N 01.91535° W	18	C. nodosa	Scarce	Record	IEO-CSIC
Punta de la Media Naranja	2023-07-18	36.93559° N 01.92257° W	6	C. nodosa	Frequent	Samples and photographs	SMPAME (JA)
Punta de la Media Naranja	2023-07-18	36.93304° N 01.91552° W	7	C. nodosa	Scarce	Record	SMPAME (JA)
Agua Amarga	2023-07-18	36.93789° N 01.93381° W	3	P. oceanica & C. nodosa	Frequent	Also dead matte. Samples and photographs	SMPAME (JA)
El Plomo	2023-07-18	36.92446° N 01.95063° W	6	P. oceanica	Scarce	Record	SMPAME (JA)
Agua Amarga (POSIMED St.)	2023-11-29	36.93615° N 01.93082° W	12	C. nodosa	Scarce	Photographs	SMPAME (JA)
Agua Amarga	2024-03-17	36.93882° N 01.93348° W	3	C. nodosa	Scarce	Molecular analysis and photographs	SMPAME (JA)
Punta de la Media Naranja	2024-08-06	36.93333° N 01.91458° W	9.5	C. nodosa	Frequent	Photographs	SMPAME (JA)
Punta de la Media Naranja	2024-08-06	36.93306° N 01.91478° W	10	C. nodosa	Abundant	Samples and photographs	SMPAME (JA)
Punta de la Media Naranja	2024-08-06	36.93290° N 01.91507° W	10.5	C. nodosa	Frequent	Samples and photographs	SMPAME (JA)
Punta de la Media Naranja	2024-08-06	36.93253° N 01.91557° W	11.5	C. nodosa	Abundant	Samples and photographs	SMPAME (JA)
Agua Amarga (POSIMED St.)	2024-12-11	36.93615° N 01.93082° W	12	C. nodosa	Absent		SMPAME (JA)

^{*} ETRS89 Datum. IEO-CSIC: Instituto Español de Oceanografía- Centro Superior de Investigaciones Científicas. SMPAME (JA): Sustainable Management Programme for the Andalusian Marine Environment (Junta de Andalusía)

Molecular identification

Specimens were also identified through molecular analyses of the rDNA 18S-5.8S-28S region, conducted at the Andalusian Institute of Agricultural and Fisheries Research and Training (IFAPA) (Department of Agriculture, Fisheries, Water and Rural Development, Regional Government of Andalusia) at El Puerto de Santa María (Cádiz). Two algal samples collected from Agua Amarga, 17 March, 2024 (Table 1) were homogenized using a TissueRuptor® tissue homogenizer (Qiagen) in five vol of lysis buffer from the NucleoSpinTM Tissue Mini Kit (Macherey-Nagel). Thereafter, total genomic DNA was extracted following the manufacturer's protocol. DNA

concentration was determined using a Nanodrop™ ND-8000, and quality was assessed via agarose gel electrophoresis.

The 18S-5.8S-28S region was amplified in two separate PCR reactions using the primer pairs 18S-ITS1F-new/TW13 (Omelchenko *et al.*, 2019; Fang *et al.*, 2023) and AB28/T13N (De Clerck *et al.*, 2006). Each PCR reaction was performed in a final volume of 12.5 µl, containing 40 ng of template DNA, 300 nM of each specific forward and reverse primer, and 6.25 µl of Platinum Multiplex PCR Master Mix (2x, Thermo Fisher Scientific). The thermal cycling conditions included an initial denaturation at 94 °C for 4 min, followed by 30 cycles of 94 °C for 40 seconds, annealing at 46 °C for 1 min, and

extension at 72 °C for 2 min, with a final step at 72 °C for 6 min. PCR products were cloned into a TOPO-TA vector and subsequently sequenced using the BigDye® Terminator v3.1 kit (Life Technologies) when necessary. Sequences obtained were assembled using SeqMan v8.2 (DNASTAR) and analysed for similarity using BLAST (blastn) at NCBI. The final sequences were deposited in GenBank/EMBL/DDBJ under accession numbers PV037361 and PV037362.

Phylogenetic analyses were conducted using the maximum likelihood method in IQ-TREE 2 v2.1.2 (Trifinopoulos et al., 2016), with 1,000 ultrafast bootstrap replications and a minimum correlation threshold of 0.99 as the convergence criterion. The TPM2+F+G4 substitution models were selected based on the Bayesian information criterion (BIC) score. Phylogenetic trees were visualised with FigTree1.4.4 with Caulerpa serrulata designated as the outgroup. The accession numbers of the species included in the phylogenetic analysis were as follows: Caulerpa taxifolia var. distichophylla, LN559619, LN559620, EF190263; Caulerpa taxifolia, AF316358, AJ299754, AF316355, AY034869, AJ299772, AJ299757, AJ316268, AJ316269; Caulerpa sertularioides, JQ745715; Caulerpa racemosa, AJ297652, AJ297653, JQ745722; Caulerpa mexicana, JQ745720; Caulerpa scalpelliformis, JQ745714; Caulerpa serrulata, JQ745711.

Results

Distribution, morphological and molecular identifica-

Specimens morphologically consistent with Caulerpa taxifolia var. distichophylla were found at 13 locations within the Cabo de Gata-Níjar Natural Park (Almería, southern Spain) along approximately 4 km of coastline between Punta de la Media Naranja and El Plomo (Table 1; Fig. S1). The macroalgae showed highly variable abundances in the invaded sites, ranging from a presence that is especially abundant at the edges of Cymodocea nodosa meadows and in the understorey of these formations, where it is generally hidden by the leaves of this angiosperm, to small colonies comprising a few isolated individuals. No monospecific patches have been observed in the area. No clear pattern associated with depth was observed, with a presence between 3 and 18 metres. The species identified in this study predominantly colonise C. nodosa meadows, occurring both within the meadows and along their edges. Additionally, the alga was observed at the edges of Posidonia oceanica meadows and on P. oceanica dead matte, but it was not detected within the meadows themselves or on loose sandy substrates.

All collected specimens were characterised by a generally thin and delicate thallus. The stolons were cylindrical and thin, approximately 1 mm in diameter, very light whitish-green in colour (Fig. 1A), with thin and




Fig. 1: Caulerpa taxifolia var. distichophylla from Cabo de Gata-Níjar Natural Park (Andalusia, Spain). A: Specimen within a Cymodocea nodosa meadow (-9.5 m; 6 August, 2024). B: Specimens in a gap of a Posidonia oceanica meadow (-3 m; 18 July, 2023). C: Middle portion of distichous fronds (-3 m; 17 March, 2024). D: Detail of the upper portion of a distichous frond (-3 m; 18 July, 2023). E: Detail of the upper portion of an irregular frond (-3 m; 18 July, 2023). F: Depigmented specimen in a dense Cymodocea nodosa meadow (-9.5 m; 6 August, 2024). In C, D and E, each interval in the scale bar corresponds to 1 mm. Photographs by Agustin Barrajón (A, F) and Diego Moreno (B, C, D, E)/ Sustainable Management Programme for the Andalusian Marine Environment/ Junta de Andalucía).

branched rhizoidal pillars. The fronds were green, generally simple and more rarely divided (Fig. 1B), with lengths ranging from 3 to 10 cm (maximum 12 cm), and 2-3 mm wide (Fig. 1C). Along the stolon, the fronds were often grouped in sets of 3-10 fronds (Fig. 1B). The rachis was cylindrical, less than 1 mm in diameter, with numerous flattened pinnules on each side, very close to each other, with a wider shape in the centre, and pointed and somewhat curved at the end (Fig. 1D). The pinnules were generally all located on the same plane with respect to the rachis, giving the frond a characteristic distichous appearance (Fig. 1D), although occasionally thalli with pinnules arranged irregularly around the rachis were also observed (Fig. 1E). Numerous depigmented thalli were observed in August 2024 within a dense C. nodosa meadows at Punta de la Media Naranja, between depths of 10 and 11 metres (Fig. 1F). However, most thalli exhibited the characteristic light green pigmentation.

The nucleotide sequences of the analysed samples were 1090 and 1093 bp in length and showed a 100% match with previously published sequences classified as *C. taxifolia* var. *distichophylla* at NCBI/EMBL/DDBJ. Phylogenetic analysis confirmed that our samples clustered in a group highly supported by a bootstrap value of 94 with *C. taxifolia* var. *distichophylla* specimens (Fig. 2).

Discussion

Morphological and molecular analyses confirmed the presence of Caulerpa taxifolia var. distichophylla for the first time in 13 sites of the Cabo de Gata-Níjar Natural Park (Almería, southern Spain). This finding provides new evidence of the species' westward expansion, with the nearest known population located over 1000 km away in the Sardinia Channel. The species is now established from the easternmost to the westernmost coasts of the Mediterranean Sea. Nucleotide sequences of rDNA grouped the Almería samples within a strongly supported clade that includes specimens of C. taxifolia var. distichophylla from Sicily (Jongma et al., 2013), Greece and Cyprus (Aplikioti et al., 2016). Furthermore, the thalli observed in the samples studied are morphologically similar to those described in other Mediterranean populations of this species. The fronds of the specimens inhabiting C. nodosa meadows reached lengths of 10–12 cm. In Sicily, established populations show frond lengths ranging from 8–10 cm on sand (sparse fronds), up to 15 cm in sheltered and shaded breakwaters, and 20-35 cm in P. oceanica meadows, in areas shaded by the seagrass canopy (Musco et al., 2014). In contrast, the fronds of the aquarium strain of C. taxifolia in Cap Martin (France) can exceed 60 cm (Meinesz & Hesse, 1991),




Fig. 2: Phylogenetic tree of *Caulerpa* species using the Maximum Likelihood method. Sequences from this study are labelled as Seq1 and Seq2. Accession numbers for each species are provided. Clades corresponding to *C. taxifolia* var. *distichophylla* and *C. taxifolia* are shaded in blue and green, respectively. Bootstrap values higher than 70 are shown in each node.

reaching up to 80 cm in autumn (Boudouresque *et al.*, 1995). The modest size observed in Almería may reflect an early stage of colonization, with potential for increasing growth as populations become more established. One plausible vector for the arrival of this invasive species is maritime traffic, particularly large commercial vessels operating in the vicinity. The nearby port of Carboneras (<6 km), which hosts intensive industrial activity including a cement plant and a thermal power station, could be a potential point of introduction.

The bathymetric range of Caulerpa taxifolia var. distichophylla in Almería (3-18 m) is consistent with previous records from Turkey, Cyprus, Malta, and Italy (0-20 m) (Cevik et al., 2007; Çiçek et al., 2013; Schembri et al., 2015; Mannino & Balistreri, 2017; Mannino et al., 2019). However, deeper occurrences have also been reported in the eastern Mediterranean Sea, reaching depths of 28 m in Malta (Ellul et al., 2019), 48 m in Cyprus and Lebanon (Aplikioti et al., 2016; Bitar et al., 2017), and up to 100 m in Greece (Aplikioti et al., 2016). In comparison, C. taxifolia has been observed from shallow waters to depths of 100 m in Monaco (Belsher & Meinesz, 1995), and C. cylindracea typically ranges from 0 to 70 m (Klein & Verlaque, 2008). The shallow distribution of the Almería populations suggests a preference for warmer habitats, possibly due to thermal constraints, but further studies are necessary to confirm this hypothesis. In this study, C. taxifolia var. distichophylla was mainly found in C. nodosa meadows with lower occurrences near P. oceanica edges, dead matte, and occasionally near C. cylindracea. In contrast, in Sicily, where the species is widespread, it shows high coverage (>50%) in P. oceanica, especially on dead matte and meadow edges, with lower abundance in C. nodosa (<10%). It has also been observed on sand, rocks, artificial substrates, and mooring lines in Sicily (Musco et al., 2014). Across the Mediterranean, the species has been recorded on a wide variety of substrates (e.g. mud, sand, gravel, rock, biogenic hard bottoms, shipwrecks) and habitats, including bioconstructions (e.g., Sabellaria spp.), macroalgal communities (Cystoseira spp., Codium bursa, Caulerpa spp., Penicillus capitatus), and seagrass meadows (C. nodosa, P. oceanica, Halophila stipulacea) (Mannino et al., 2019). This ecological plasticity, coupled with the availability of suitable habitats (e.g., C. nodosa and P. oceanica meadows and some areas of dead matte) suggests a high potential for further spread in Almería unless constrained by other environmental factors (e.g., temperature).

The aquarium-derived strain of Caulerpa taxifolia expanded along the cooler northern Mediterranean coasts in the late 20th century—from Monaco, France, Italy, Croatia, and Spain (Balearic Islands)—but remained rare in warmer southern regions, with the exception of Tunisia (Meinesz et al., 2001; 2010). Its spread is limited by a lower thermal tolerance (lethal temperature below 7 °C) (Boudouresque et al., 1995). In contrast, C. taxifolia var. distichophylla has primarily colonized warmer regions of the eastern and southern Mediterranean Sea, including Syria, Lebanon, Cyprus, southern Turkey, Greece, Libya, Malta, Tunisia, Sicily, Calabria, and southern Sar-

dinia (Mannino et al., 2019). Its distribution appears to be constrained by the temperature of the 15 °C February isotherm, which may explain its absence from colder northern coasts (Jongma et al., 2013). Recent westward records in Tunisia (2015) (Chartosia et al., 2018) and southern Sardinia (2016) (Di Martino et al., 2018) remain within thermally mild zones. Similarly, in Panarea (north of Sicily, Italy), its presence is restricted to hydrothermal vent areas characterized by warmer waters, where it coexists with C. cylindracea and H. stipulacea (Gaglioti & Gambi, 2018). These patterns suggest that the biogeographic range of this variety is largely determined by thermal limitations, although the reproductive potential and other ecological traits probably also constrain its distribution.

Depigmented fronds observed in some C. taxifolia var. distichophylla samples from Almería may indicate a reproductive process, potentially linked to gamete release. However, their uniformly whitish appearance contrasts with the partial depigmentation observed in fertile thalli of other Caulerpa species (e.g., C. taxifolia, C. cylindracea, C. chemnitzia) (Panayotidis & Žuljević, 2001; Phillips 2009; Caronni et al., 2021). Unlike the aquarium-derived strain of C. taxifolia, which shows no evidence of sexual reproduction in the Mediterranean, likely due to its hybrid origin (Meinesz, 1992; Boudouresque et al., 1995), the sexual reproductive capacity of C. taxifolia var. distichophylla remains unknown. The decline in C. taxifolia expansion since the early 2000s across the Mediterranean Sea may be partially explained by the lack of sexual reproduction and the emergence of algal disease, possibly caused by pathogenic agents (Meinesz et al., 2010; Žuljević et al., 2019; Kleitou et al., 2025). Therefore, investigating the reproductive strategy of C. taxifolia var. distichophylla is critical to assess its potential for long-term persistence and expansion.

Although no ecological impacts have been observed yet in Almería, the presence of C. taxifolia var. distichophylla raises concerns due to its invasive potential. Its size and morphology, distinct from the aquarium-derived C. taxifolia, may result in different ecological interactions (Musco et al., 2014). While the impacts of C. taxifolia on benthic communities and seagrass meadows are well documented (Boudouresque et al., 1995), the effects of C. taxifolia var. distichophylla remain poorly understood. Preliminary studies suggest possible impacts on polychaete assemblages in muddy sand (Cevik et al., 2012) and P. oceanica dead matte (Musco et al., 2014). The absence of caulerpine in native populations of this variety (Schwede et al., 1986) could imply lower toxicity compared to other Caulerpa species (Lemée et al., 1996), potentially reducing its biotic impact. However, its ability to monopolize space, accumulate as wrack, and cause socioeconomic impacts such as interference with fishing activities (Musco et al., 2014) reinforce the importance of ongoing monitoring in the western Mediterranean Sea. Additionally, seasonal variability may modulate its ecological impact. In Malta, fronds seem to regress during winter (Schembri et al., 2015), potentially reducing ecological impacts on native assemblages (Ellul *et al.*, 2019).

A similar seasonal pattern has been observed in Almería (southeastern Spain), where the species is abundant in summer but becomes scarce or absent in November. This suggests temperature-dependent dynamics, but long-term systematic monitoring is required to confirm seasonal trends and their ecological significance.

The future expansion of C. taxifolia var. distichophylla in southeastern Spain remains uncertain. In contrast to C. taxifolia in the Mediterranean Sea, C. cylindracea has exhibited a continuous and rapid spread since its first detection in Almería in 2008 (Moreno, 2010), reaching multiple locations in Andalusia within a few years (Junta de Andalucía, 2010, 2013, 2014; Arroyo et al., 2015). After more than 15 years of presence in Andalusia, and despite its wide distribution from the shallow photophilous algae communities to depths exceeding 30 m, C. cylindracea has shown limited impact on well-structured *P. oceanica* meadows. It predominantly colonises sandy patches within meadows, dead matte, and maerl beds at depths from 20 to over 30 m (Carreto et al., 2021). Similar patterns have been described around the Balearic Islands (Cebrian & Ballesteros, 2009). Interestingly, recent data indicate a decline or disappearance of C. cylindracea in some previously dense populations (Junta de Andalucía, 2024). The available evidence suggests that C. taxifolia var. distichophylla may expand along southeastern Spain and further north, especially under continued sea surface warming, which may benefit this thermophilic, invasive taxon. However, its potential expansion into the Alborán Sea could be constrained by the Almería-Oran front, a well-known biogeographic barrier for many species (Real et al., 2021), and by the influence of colder Atlantic waters. Ongoing monitoring will be key to tracking these dynamics.

Acknowledgements

The authors are very grateful to the Department of Sustainability and Environment (Regional Government of Andalusia) for initiating the seabed monitoring work in 1994–1996 and the "Plan de detección del alga *Caulerpa taxifolia*". In particular, we thank José Guirado Romero and the recently deceased Hermelindo Castro Nogueira. We also thank the current heads of the Sustainable Management Programme for the Andalusian Marine Environment (Junta de Andalucía), specifically José Eugenio Montes Gómez, and the coordinator of the programme, Soledad Vivas Navarro, as well as all the members of the team, including the crew of the *Arca* vessel, Pascual Hernández and Martín Carrillo. This study has been partially funded by the project FEM.DIP2019.001, programme FEMP 2014-2020 Andalucía.

The IEO-CSIC MSFD monitoring programme was developed within the framework of the ESMARES2 projects, as part of the scientific programme "Asesoramiento científico-técnico para la protección del medio marino: Evaluación y seguimiento de las Estrategias Marinas, Seguimiento de los espacios marinos protegidos de competencia estatal (2018–2021)", funded by the Spanish Ministry of Ecological Transition and the Demographic

Challenge (MITECO) and the BIODIV project: "Asesoramiento científico-técnico para el seguimiento de la biodiversidad marina: espacios y especies marinos protegidos de competencia estatal (2022–2025)", funded by the European Union – NextGenerationEU through the Recovery, Transformation and Resilience Plan; and promoted by the Directorate General for Biodiversity, Forests and Desertification of the Ministry for Ecological Transition and the Demographic Challenge and CSIC, through the Spanish Institute of Oceanography (IEO). We are especially grateful to the members of the GEAM team –Juan Manuel Ruiz, Lázaro Marín, María Dolores Belando, Marta García, Mari Carmen Mompeán, Fernando García, and Rocío García– for their invaluable collaboration in sampling and helpful comments on this article.

References

Altamirano, M., De La Rosa, J., Martínez, F.J., 2016. Arribazones de la especie exótica *Rugulopteryx okamurae* (E.Y. Dawson) I.K. Hwang, W.J. Lee and H.S. Kim (Dictyotales, Orchrophyta) en el Estrecho de Gibraltar: primera cita para el Atlántico y España. *Algas*, 52, 20.

Aplikioti, M., Louizidou, P., Mystikou, A., Marcou, M., Stavrou, P. et al., 2016. Further expansion of the alien seaweed Caulerpa taxifolia var. distichophylla (Sonder) Verlaque, Huisman & Procacini (Ulvophyceae, Bryopsidales) in the Eastern Mediterranean Sea. Aquatic Invasions, 11 (1), 11-20.

Arroyo, M.C., Barrajón, A., Brun, F.G., del Castillo, F., De la Rosa, J. *et al.*, 2015. Praderas de angiospermas marinas de Andalucía. p. 312-397. In: *Atlas de las praderas marinas de España*. Ruiz, J.M., Guillén, J.E., Ramos Segura, A., Otero, M.M. (Eds). IEO/IEL/UICN, Murcia-Alicante-Málaga.

Báez, J.C., Conde, F., Flores-Moya, A., 2001. Notas corológicas del macrofitobentos de Andalucía (España). V. Acta Botanica Malacitana, 26, 193-196.

Ballesteros, E., Aranda, A., Pou, S., Grau, A.M., Riera, F. et al., 1996. Valoración de la campaña de sensibilización y vigilancia sobre la invasión de Caulerpa taxifolia (Vahl) C. Agardh en las costas mediterráneas españolas. Programa Life (1993-1994). Publicacions Universitat Barcelona. Second International Workshop on Caulerpa taxifolia, 63-66.

Belsher, T., Meinesz, A., 1995. Deep-water dispersal of the tropical alga *Caulerpa taxifolia* introduced into the Mediterranean. *Aquatic Botany*, 51, 163-169.

Bitar, G., 2010. La flore marine benthique introduite de la côte libanaise. Etat actuel de trois espèces envahissantes. p. 107-114. In: *INOC*, *International conference on Biodiversity of the Aquatic Environment*. Tishreen University, Syria.

Bitar, G., Ramos-Esplá, A.A., Ocaña, O., Sghaier, Y.R., Forcada A. *et al.*, 2017. Introduced marine macroflora of Lebanon and its distribution on the Levantine coast. *Mediterranean Marine Science*, 18 (1), 138-155.

Boudouresque, C.F., Meinesz, A., Ribera, M.A., Ballesteros, E., 1995. Spread of the green alga *Caulerpa taxifolia* (Caulerpales, Chlorophyta) in the Mediterranean possible consequences of a major ecological event. *Scientia Marina*, 59 (1), 21-29.

- Caronni, S., Delaria, M.A., Gentili, R., Montagnani, C., Navone, A. *et al.*, 2021. First report of gametogenesis and spawning for the invasive alga *Caulerpa cylindracea* in the Tyrrhenian Sea: the key role of water motion and temperature. *Frontiers in Marine Science*, 8, 774274.
- Carreto, M.T., Barrajón Domenech, A., Moreno, D., Remón, J.M., De la Rosa, J. et al., 2021. Cartografía temática de praderas de angiospermas marinas para la obtención de información de los stocks de carbono azul en Andalucía. Chronica naturae, 8, 57-110.
- Cebrian, E., Ballesteros, E., 2009. Temporal and spatial variability in shallow- and deep-water populations of the invasive *Caulerpa racemosa* var. *cylindracea* in the Western Mediterranean. *Estuarine Coastal and Shelf Science*, 83, 469-474.
- Cevik, C., Yokes, B., Cavas, L., Erkol, L., Derici, O.B. et al., 2007. First report of *Caulerpa taxifolia* (Bryopsidales, Chlorophyta) on the Levantine coast (Turkey, Eastern Mediterranean). *Estuarine, Coastal and Shelf Science*, 74 (3), 549-556.
- Chartosia, N., Anastasiadis, D., Bazairi, H., Crocetta, F., Deidun, A. et al., 2018. New Mediterranean Biodiversity Records (July 2018). Mediterranean Marine Science, 19 (2), 398-415.
- Çiçek, B.A., Kurt, O., Taskin, E., Öztürk, M., 2013. First report of Caulerpa taxifolia var. distichophylla (Sonder) Verlaque, Huisman and Procaccini (Caulerpaceae, Chlorophyta) from Northern Cyprus. Rapport Commission International Mer Méditerranée, 40, 600.
- De Clerck, O., Leliaert, F., Verbruggen, H., Lane, C.E., De Paula, J.C. et al., 2006. A revised classification of the dictyoteae (dictyotales, phaeophyceae) based on rbcl and 26s ribosomal dna sequence analyses 1. Journal of Phycology, 42, 1271-1288.
- Di Martino, V., Stancanelli, B., Cantasano, N., 2018. The alien alga *Caulerpa taxifolia* (Vahl) C. Agardh var. distichophylla (Sonder) Verlaque, Huisman and Procaccini move their northern and western limits. *Journal of the Black Sea/Mediterranean Environment*, 24 (2), 140-148.
- Ellul, T., Evans, J., Schembri, P.J., 2019. Invasion alert: rapid range expansion of *Caulerpa taxifolia* var. *distichophylla* in Maltese waters (central Mediterranean). *BioInvasions Records* 8 (2), 208-217.
- Famà, P., Wysor, B., Kooistra, W.H.C.F., Zuccarello, G.C., 2002. Molecular phylogeny of the genus *Caulerpa* (Caulerpales, Chlorophyta) inferred from chloroplast *tufA* gene. *Journal of Phycology*, 38, 1040-1050.
- Fang, C., Sun, X., Fan, F., Zhang, X., Wang, O. et al., 2023. High-resolution single-molecule long-fragment rRNA gene amplicon sequencing of bacterial and eukaryotic microbial communities. Cell Reports Methods, 3, 100437.
- Gaglioti, M., Gambi, M.C., 2018. The natural laboratory of the CO₂ vents off Panarea Aeolian Islands, Italy: A special ecological setting and a further stepping stone for some alien macrophytes. *Notiziario S.I.B.M.*, 74, 111-117.
- Jongma, D.N., Campo, D., Dattolo, E., D'Esposito, D., Duchi, A. et al., 2013. Identity and origin of a slender Caulerpa taxifolia strain introduced into the Mediterranean Sea. Botanica Marina, 56 (1), 2739.
- Junta de Andalucía, 2008-2024. Informes Regionales del Pro-

- grama de Gestión Sostenible del Medio Marino Andaluz. *Consejería de sostenibilidad y Medio Ambiente. Junta de Andalucía.* https://www.junta-andalucia.es/medioambiente/portal/home. (Accessed 18 March 2025).
- Klein, J., Verlaque, M., 2008. The *Caulerpa racemosa* invasion: A critical review. *Marine Pollution Bulletin*, 56, 205-225.
- Kleitou, P., Agius, D., Akalin, S., Albano, M., Ammar, I. A. et al., 2025. New records of introduced species in the Mediterranean Sea (February 2025). Mediterranean Marine Science, 26 (1), 175-198.
- Langar, H., Djellouli, A., Ben Mustapha, K., El Abed, A.,
 2000. Première signalisation de *Caulerpa taxifolia* (Vahl)
 C. Agardh en Tunisie. *Bulletin de l'Institut National des Sciences et Techniques de la Mer (Tunisia)*, 27 (1), 1-8.
- Lemée, R., Boudouresque, C.F., Gobert, J., Malestroit, P., Mari, X. et al., 1996. Feeding behaviour of *Paracentrotus lividus* in the presence of *Caulerpa taxifolia* introduced in the Mediterranean Sea. *Oceanologica Acta*, 19, 245-253.
- Mannino, A.M., Balistreri, P., 2017. An updated overview of invasive *Caulerpa* taxa in Sicily and circum-Sicilian Islands, strategic zones within the NW Mediterranean Sea. *Flora Mediterranea*, 27, 221-240.
- Mannino, A.M., Cicero, F., Toccaceli, M., Pinna, M., Balistreri, P., 2019. Distribution of *Caulerpa taxifolia* var. *disticho-phylla* (Sonder) Verlaque, Huisman & Procaccini in the Mediterranean Sea. *Nature Conservation*, 37, 17-29.
- Meinesz, A., 1992. Modes de dissémination de l'algue *Cauler-pa taxifolia* introduite en Méditerranée. *Rapport de la Com-mission Internationale sur la Mer Méditerranée*, 33, 44.
- Meinesz, A., 2001. *Killer algae*. The University of Chicago, Chicago, 360 pp.
- Meinesz, A., Chancollon, O., Cottalorda, J.M., 2010. Observatoire sur l'expansion de Caulerpa taxifolia et Caulerpa racemosa en Méditerranée: campagne janvier 2008 juin 2010. Université Nice Sophia Antipolis, E.A. 4228 ECOMERS publ., 50 p.
- Meinesz, A., Hesse, B., 1991. Introduction et invasion de l'algue tropicale *Caulerpa taxifolia* en Mediterranée nord-occidentale. *Oceanologia Acta*, 14 (4), 415-426.
- Meinesz, A., Belsher, T., Thibaut, T., Antolic, B., Ben Mustapha, K. et al., 2001. The introduced green alga Caulerpa taxifolia continues to spread in the Mediterranean. Biological Invasions, 3, 201-210.
- Moreno, D., 2009. Gestión sostenible del medio marino en Andalucía: las fanerógamas marinas. p. 114-128. In: Posidonia oceanica. Redes de seguimiento y estado de conservación en el Mediterráneo español. Instituto de Ecología Litoral, Diputación de Alicante, Ministerio de Medio Ambiente y Medio Rural y Marino (Eds), Alicante.
- Moreno, D., 2010. Flora y fauna alóctona del medio marino andaluz. p. 214-229. In: Especies exóticas invasoras en Andalucía. Talleres provinciales 2004-2006. Cobos, F.J., Ortega, F. (Eds). Consejería de Medio Ambiente, Junta de Andalucía, Sevilla.
- Musco, L., Andaloro, F., Mikac, B., Mirto, S., Vega Fernández, T. et al. 2014. Concern about the spread of the invader seaweed Caulerpa taxifolia var. distichophylla (Chlorophyta: Caulerpales) towards the West Mediterranean. Mediterranean Marine Science, 15 (3), 532-538.
- Mutlu, E., Akçalı, B., Özvarol, Y., Narlı, Z., Aslan, B.E. et al.,

- 2025. Recent plant traits of *Caulerpa taxifolia* var. *distichophylla* in the Turkish Aegean Sea. *Aquatic Sciences and Engineering*, 40 (2), 63-73.
- Omelchenko, D.O., Speranskaya, A.S., Ayginin, A.A., Khafizov, K., Krinitsina, A.A. *et al.*, 2019. Improved Protocols of ITS1-Based Metabarcoding and Their Application in the Analysis of Plant-Containing Products. *Genes*, 10 (2), 122.
- Panayotidis, P., Žuljević A., 2001. Sexual reproduction of the invasive green alga *Caulerpa racemosa* var. *occidentalis* in the Mediterranean Sea. *Oceanologica Acta*, 24 (2), 199-203.
- Phillips, J.A., 2009. Reproductive ecology of *Caulerpa taxi-folia* (Caulerpaceae, Bryopsidales) in subtropical eastern Australia. *European Journal of Phycology*, 44 (1), 81-88.
- Pou, S., Ballesteros, E., Delgado, O., Grau, A.M., Riera, F. et al., 1993. Sobre la presencia del alga Caulerpa taxifolia (Vahl) C. Agardh (Caulerpales, Chlorophyta) en aguas costeras de Mallorca. Bolletí de la Societat d'Història Natural de les Balears, 36, 83-89.
- RAC/SPA-UNEP/MAP, 2003. Report of second mission relating to the development of Marine Protected Areas on Syrian coasts, 1-18 August 2003. In: *Regional Project for the Development of Marine and Coastal Protected Areas in the Mediterranean Region*. Bitar, G., Dupuy de la Grandrive, R., Foulquié, M. (Eds)., RAC/SPA-MedMPAnet Project, Tunis.
- Real, R., Gofas, S., Altamirano, M., Salas, C., Báez, J.C. et al., 2021. Biogeographical and Macroecological Context of the Alboran Sea. p. 431-457. In: Alboran Sea-Ecosystems and Marine Resources. Báez, J.C. (Ed). Springer Nature Switzerland AG.
- Royal Decree 630/2013, of August 2, regulating the Spanish Catalogue of Invasive Alien Species. Ministry of Agriculture, Food and Environment "BOE" No. 185, of August 3, 2013: 56764-56786 pp. Reference: BOE-A-2013-8565. (See update in Order TED/1126/2020, of November 20).
- Schembri, P.J., Barbara, J., Deidun, A., Lanfranco, E., Lanfranco, S., 2015. It was only a matter of time: Occurrence of *Caulerpa taxifolia* (Vahl) C. Agardh var. *distichophylla* (Sonder) Verlaque, Huisman and Procaccini in the Maltese Islands (Chlorophyta, Ulvophyceae, Caulerpaceae). *BioInvasions Records*, 4 (1), 9-16.

- Schwede, J.G., Cardellina, J.H., Grode, S.H., James Jr, T.R., Blackman, A.J., 1986. Distribution of the pigment caulerpin in species of the green alga *Caulerpa*. *Phytochemistry*, 26 (1), 155-158.
- Shakman, E.A., Abdalha, A.B., Thala, F., Al-Faturi, A., Bariche M., 2017. First records of seven marine organisms of different origins from Lybia (Mediterranea Sea). *BioInvasions Records*, 6 (4), 377-382.
- Span, A., Antolic, B., Žuljević, A., 1998. The genus Caulerpa (Caulerpales, Chlorophyta) in Adriatic Sea. Rapport de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée, 35, 584-585.
- Streftaris, N., Zenetos, A., 2006. Alien marine species in the Mediterranean the 100 'Worst Invasives' and their impact. *Mediterranean Marine Science*, 7 (1), 87-118.
- Taşkın, E., Evcen, A., Bilgiç, F., 2023. Further expansion of the alien marine green macroalga *Caulerpa taxifolia* var. *distichophylla* (Sonder) Verlaque, Huisman & Procacini in Türkiye. *Journal of Black Sea/Mediterranean Environment*, 29 (1), 121-126.
- Trifinopoulos, J., Nguyen, L.T., von Haeseler, A., Minh, B.Q., 2016. W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Research*, 44, 232-235.
- Turan, G., Tekogul, H., Cirik, S., Meinesz, A., 2011. First record of the invasive green seaweed *Caulerpa taxifolia* (Bryopsidales) on the coast of Turkey. *Cryptogamie, Algologie*, 32 (4), 379-382.
- Verlaque, M., Ruitton, S., Mineur, F., Boudouresque, C.F., 2015. CIESM Atlas of exotic macrophytes in the Mediterranean Sea. *Rapport de la Commission de l'Exploration de la mer Méditerranée*, 38, 14.
- Zenetos, A., Doğan, A., Bakir, A.K., Chatzigeorgiou, G., Corsini-Foka, M. et al., 2025. Non-Indigenous Species (NIS) Know No Geopolitical Borders -An Update of NIS in the Aegean Sea. Diversity, 17, 12.
- Žuljević A., Cvitkovic, I., Despalatovic, M., Lucic, P., Zunec, A. et al., 2019. Decline of Caulerpa taxifolia in the Adriatic Sea. p. 82-87. In: Langar H. y Ouerghi A. (Eds). Proceedings of the 1st Mediterranean Symposium on the Non-Indigenous Species. SPA/RAC, Antalya, Turkey.

Supplementary Data

The following supplementary information is available online for the article:

S1: Distribution map of Caulerpa taxifolia var. distichophylla in the Mediterranean Sea (red dots, bibliography records) and in Andalusia, Spain (yellow dots, present study). A: First records in the eastern Mediterranean basin and distribution of the species in the Mediterranean Sea. B: Location of Agua Amarga (Andalusia, Spain). C: New records of the species (yellow star, first observation) between Punta de la Media Naranja and El Plomo (Cabo de Gata-Níjar Natural Park, Andalusia).