

Mediterranean Marine Science

Vol 7, No 2 (2006)



Decapod crustaceans on the Gökçeada (Imbros) island continental shelf (north-eastern Aegean Sea)

A.S. ATES, T. KATAGAN, A. KOCATAS, M. SEZGIN

doi: [10.12681/mms.169](https://doi.org/10.12681/mms.169)

To cite this article:

ATES, A., KATAGAN, T., KOCATAS, A., & SEZGIN, M. (2006). Decapod crustaceans on the Gökçeada (Imbros) island continental shelf (north-eastern Aegean Sea). *Mediterranean Marine Science*, 7(2), 55–60.
<https://doi.org/10.12681/mms.169>

Mediterranean Marine Science
Volume 7/2, 2006, 55-60

Decapod crustaceans on the Gökçeada (Imbros) island continental shelf (north-eastern Aegean Sea)

A.S. ATEŞ¹, T.KATAĞAN², A. KOCATAŞ² and M. SEZGİN³

¹Çanakkale Onsekiz Mart University,
Fisheries Faculty, 17100 Çanakkale, Turkey

²Ege University, Fisheries Faculty 35100 Bornova - İzmir, Turkey

³Ondokuz Mayıs University, Sinop Fisheries Faculty 57000 Sinop, Turkey

e-mail: asuatates@yahoo.com

Abstract

The present composition of decapod crustaceans found at the sublittoral depths (5-104 m) off the coast of the island of Gökçeada (north-eastern Aegean Sea) is presented. A total of 28 species (11 caridean shrimps, 1 thalassinid ghost crab, 7 anomurans and 9 brachyuran crabs) and 277 specimens were recorded. The caridean shrimp, *Athanas nitescens* had the highest abundance with a dominance value of 20.94% in samples. The dominant group is caridean, represented by a total of 11 species and an occurrence frequency of 39.29%.

Keywords: Decapoda Crustacea; Gökçeada (Imbros) island; North-eastern Aegean Sea; Turkey.

Introduction

Faunistic and ecological studies on the decapod crustaceans of the sublittoral coast of Gökçeada (Imbros) are quite scarce, and are reported in the study by BALKIS *et al.*, 2001. Lately, KOCATAŞ & KATAĞAN (2003) have published a checklist of decapods in Turkish Seas that included 220 species, among which 181 species were found in the Turkish Aegean Sea. Recently, ATEŞ *et al.* (2004) have increased to 186 the number of species known for the area with five new records i.e. the caridean shrimp, *Processa macrodactyla* Holthuis, 1952, the thalassinid, *Callianassa trryhena* (Petagna, 1792), the brachyurans, *Ebalia tumefacta* (Montagu, 1808), *Liocarcinus maculatus* (Risso, 1827), *Palicus caronii* (Roux, 1830). BALKIS *et al.* (2001) found 32 species of brachyuran crabs

on the littoral bottoms of Gökçeada island. ATEŞ *et al.* (2005) reported *Munida rugosa* (J.C. Fabricius, 1775) as a new record for the Aegean coast of Turkey. Thus, the number of decapod species known for the Turkish Aegean Sea increased to 187. The present paper aims to describe the decapod fauna occurring on the continental shelf of Gökçeada Island.

Material and Methods

The study area is located on the coast of the Gökçeada continental shelf (the north-eastern Aegean Sea) (Fig. 1). The bottoms surveyed are situated between 5 and 104 m depth. Samples were collected during daytime hours in August 2000 using either beam-trawl or dredge at 5 sampling sites. The bottom at the shallowest stations was




Fig. 1: Locations of the sampling stations off Gökçeada Island.

mostly covered with a *Posidonia oceanica* (L.) Delile meadow. The details on date, GPS coordinates, sampling gear, depth and substratum are presented in Table 1.

After collection, all decapods in the samples were preserved in 4 % formalin. In the laboratory, preserved samples were rinsed over a 0.5 mm sieve. Decapods were counted and identified to species level whenever possible, based on the work of ZARIQUIEY AL-

VAREZ (1968), INGLE (1993) and FALCIAI & MINERVINI (1996). The higher classification follows MARTIN & DAVIS (2001).

Dominance was calculated as $D_i = (n_i/N) \times 100$, where D_i is the mean dominance index for species i ; n_i , the number of individuals belonging to species i ; N , the total number of individuals of all species (BELLAN-SANTINI, 1969). SOYER's (1970) frequency index was used to determine the frequency of spe-

Table 1

List of sampling stations. Substrates: P, *Posidonia oceanica* meadow; M, mud, sampling gears: D, dredge; B, beam-trawl.

Station	Date	Coordinates	Sampling gear	Depth (m)	Type of substratum
1	17.08.2000	40°05'45" N 25°50'45" E	D	27	P
2	13.08.2000	40°07'22" N 25°39'50" E	B	5	P
3	13.08.2000	40°07'30" N 25°40'39" E	D	10	P
4	13.08.2000	40°10'40" N 25°40'50" E	D	104	M
5	14.08.2000	40°13'20" N 26°03'00" E	D	96	M

cies at the stations, and in biotopes. The frq/x 100, where, m = number of stations where the species occurred and M = number of all stations.

Results

A total of 277 decapod specimens belonging to 28 species was captured (Table 2). Caridea was the most diversified group with 11 species followed by Brachyurans (9),

Anomurans (7) and Thalassinideas (1). The decapod taxocoenosis is dominated by few species, *Athanas nitescens* being the most abundant (Di 20.94%), followed by *Lysmata seticaudata*, and *Pisidia longimana*: dominance values of 19.86% and 15.16%, respectively. *Galathea squamifera*, *Macropodia rostrata*, and *Parthenope massena* have the lowest value of 0.36% (Table 2).

Total abundance of higher taxa was 39.29% for Caridea, 32.14% for Brachyura

Table 2
Species composition, abundance of individuals at stations, values of frequency (f%) and dominance (Di%).

Species	1	2	3	4	5	
Total specimens	94	127	33	3	20	
Total species	18	9	9	3	7	
CARIDEA						Di%
<i>Alpheus macrocheles</i> (Hailstone, 1885)	0	0	21	0	0	7.58
<i>Athanas nitescens</i> (Leach, 1814)	16	42	0	0	0	20.94
<i>Hippolyte inermis</i> Leach, 1815	7	3	0	0	0	3.61
<i>Lysmata seticaudata</i> (Risso, 1816)	27	28	0	0	0	19.86
<i>Palaemon elegans</i> Rathke, 1837	3	0	0	0	0	1.08
<i>Philocheas bispinosus</i> (Hailstone, 1835)	1	0	0	0	2	1.08
<i>Philocheas sculptus</i> (Bell, 1847)	2	0	0	0	0	0.72
<i>Processa macrodactyla</i> Holthuis, 1952	3	0	0	0	5	2.89
<i>Processa macrophthalma</i> Nouvel & Holthuis, 1957	5	0	0	0	0	1.81
<i>Processa modica</i> Williamson & Rochanburanon, 1979	0	2	0	0	3	1.81
<i>Processa nouveli</i> Al-Adhub & Williamson, 1975	0	0	0	1	2	1.08
<i>Upogebia pusilla</i> (Petagna, 1792)	3	0	1	0	0	1.44
<i>Pagurus anachoretus</i> Risso, 1827	0	0	1	0	3	1.44
<i>Pagurus cuanensis</i> Bell, 1846	11	0	0	0	0	3.97
<i>Pagurus forbesii</i> Bell, 1845	3	1	0	0	0	1.44
<i>Galathea bolivari</i> Zariquiey Alvarez, 1950	5	0	2	0	0	2.53
<i>Galathea intermedia</i> Lilljeborg, 1851	0	5	0	0	2	2.53
<i>Galathea squamifera</i> Leach, 1814	0	0	1	0	0	0.36
<i>Pisidia longimana</i> (Risso, 1816)	0	42	0	0	0	15.16
<i>Achaeus cranchii</i> Leach, 1817	0	2	3	0	0	1.81
<i>Achaeus gracilis</i> O. G. Costa, 1839	0	2	0	0	0	0.72
<i>Ethusa mascarone</i> (Herbst, 1785)	1	0	1	0	0	0.72
<i>Goneplax rhomboides</i> (Linnaeus, 1758)	0	0	0	1	3	1.44
<i>Inachus dorsettensis</i> (Pennant, 1777)	1	0	1	0	0	0.72
<i>Liocarcinus maculatus</i> (Risso, 1827)	2	0	2	0	0	1.44
<i>Macropodia rostrata</i> (Linnaeus, 1761)	1	0	0	0	0	0.36
<i>Parthenope massena</i> (Roux, 1830)	1	0	0	0	0	0.36
<i>Pilumnus hirtellus</i> (Linnaeus, 1758)	2	0	0	1	0	1.08

Table 3
Dominance (Di%) values of species found on *Posidonia oceanica* beds and muddy bottoms.

Species	<i>Posidonia oceanica</i> beds	Muddy bottom
CARIDEA	Dominance (Di%)	Dominance (Di%)
<i>Alpheus macrocheles</i>	8.26	-
<i>Athanas nitescens</i>	22.83	-
<i>Hippolyte inermis</i>	3.93	-
<i>Lysmata seticaudata</i>	21.65	-
<i>Palaemon elegans</i>	1.18	-
<i>Philocheras bispinosus</i>	0.39	8.69
<i>Philocheras sculptus</i>	0.78	-
<i>Processa macrodactyla</i>	1.18	21.73
<i>Processa macrophthalma</i>	1.96	-
<i>Processa modica</i>	0.78	13.04
<i>Processa nouveli</i>	-	13.04
THALASSINIDEA		
<i>Upogebia pusilla</i>	1.57	-
ANOMURA		
<i>Pagurus anachoretus</i>	0.39	13.04
<i>Pagurus cuanensis</i>	4.33	-
<i>Pagurus forbesii</i>	1.57	-
<i>Galathea bolivari</i>	2.75	-
<i>Galathea intermedia</i>	1.96	8.69
<i>Galathea squamifera</i>	0.39	-
<i>Pisidia longimana</i>	16.53	-
BRACHYURA		
<i>Achaeus cranchii</i>	1.96	-
<i>Achaeus gracilis</i>	0.78	-
<i>Ethusa mascarone</i>	0.78	-
<i>Goneplax rhomboides</i>	-	17.39
<i>Inachus dorsettensis</i>	0.78	-
<i>Liocarcinus maculatus</i>	1.57	-
<i>Macropodia rostrata</i>	0.39	-
<i>Parthenope massena</i>	0.39	-
<i>Pilumnus hirtellus</i>	0.78	4.34

25% for Anomura and 3.5% for Thalassinidea. The number of species and specimens found at the sampling stations is presented in Table 2.

The *P. oceanica* stations had the richest decapoda fauna with 254 individuals (f%=92.86) and 26 species, followed by the mud, with 23 specimens (f%=28.57) and 8 species. The commonest decapod of the *P. oceanica* beds was *A. nitescens* with a

dominance value of 22.83%, followed by *L. seticaudata* (Di=21.65%), and the *P. longimana* (Di=16.53%). *Processa macrodactyla* showed the highest abundance (Di=21.73%) in mud stations, followed by *Goneplax rhomboides* (Di=17.39%), *Processa modica*, *Processa nouveli* and *Pagurus anachoretus* (each with Di=13.04) (Table 3). As to the frequency of occurrence, only 2% of the species collected in *P. oceanica* meadows can

be considered as rare, and 98% are continuous. 28% of species recorded from mud sediments are common and 72% rare.

Discussion

P. oceanica meadows exhibit a complex biotic community (PÉRÈS & PICARD, 1964) and have a very high productivity and population density (GALLMETZER *et al.*, 2005). Decapod crustaceans are one of the most remarkable taxa inhabiting these meadows. In recent years, many studies on decapods associated with *P. oceanica* beds have been published (SCIPIONE *et al.*, 1983; MAZZELLA *et al.*, 1989; GARCÍA RASO, 1990; ZUPO, 1990; BORG, 1991; GARCÍA RASO *et al.*, 1996; BORG & SCHEMBRI, 2000). In our study, 26 decapod species were found on *P. oceanica* meadows and only 8 in muddy bottoms.

GARCÍA RASO (1990), stated that only 9 out of 50 decapod species captured in *P. oceanica* beds in southern Spain have a dominance value higher than 1%, and the commonest species were *Cestopagurus timidus* (31.6%), *Calcinus tubularis* (25.3%), and *A. nitescens* (13.6%). Recently, BORG & SCHEMBRI (2000) reported a total of 41 decapods from the same habitat along the Malta coasts, and the same authors indicated that *C. timidus* was the most abundant species with a dominance value of 51.2%, followed *A. nitescens* (Di=10.97%). According to GARCÍA RASO *et al.* (1996), the *P. oceanica* beds occurring in the Alboran Sea are characterized by two hermit crabs, *C. tubularis* and *C. timidus*. Overall GARCÍA RASO (1990), GARCÍA RASO *et al.* (1996) and BORG & SCHEMBRI (2000) exposed relatively similar results regarding the dominant decapod species of *P. oceanica* beds in the western and central Mediterranean. In addition, according to GALLMETZER *et al.* (2005), the most abundant decapods occurring in the *P. oceanica* meadows of Corsica were *A. nitescens*, *G. intemedia*, *L. navigator*, and *Pisa tetraodon*. Nevertheless, *A.*

nitescens, *L. seticaudata*, and *P. longimana* are commonest in the *P. oceanica* sites studied here.

The species such as *L. seticaudata*, *Palaeomon elegans*, *Philocheras bispinosus*, *Philocheras sculptus*, *P. macrodactyla*, *P. macrophthalma*, *P. modica*, *Upogebia pusilla* (Peltagna, 1792), *Pagurus forbesii*, *G. intermedia*, *Inachus dorsettensis*, *Liocarcinus maculatus*, *Macropodia rostrata*, and *Parthenope massena* which are cited here in *Posidonia oceanica* meadows are not present in the study of BORG & SCHEMBRI (2000).

Alpheus dentipes, *Thorulus cranchii*, *C. tubularis*, *C. timidus*, and *P. longicornis* which are noted as the most abundant species of *P. oceanica* beds in littoral zones off the coast of southern Spain by GARCÍA RASO, could not be found off the Gökçeada coast (1990). During our study, while *I. dorsettensis*, *M. rostrata*, *P. massena*, and *Pilumnus hirtellus* were observed in *P. oceanica* beds, on the contrary, BALKIS *et al.* (2001) previously recorded these species on the rocky, sandy and muddy bottoms of the Gökçeada coast.

Finally, the structure of the decapod community mainly depends on the physical characteristics of the biotopes. Differences in the species diversity obtained here may be related to different depth, biotope variations and hydrodynamic conditions. Further faunistic surveys which should be carried out in deeper zones, covering more sampling stations in the region will undoubtedly lead to better information regarding the decapod fauna.

References

- ATEŞ, A. S., İŞMEN, A., ÖZEKİNCİ, U. & YIĞIN, C. Ç., 2005. A new record of *Munida rugosa* (J. C. Fabricius, 1775) (Decapoda, Anomura, Galatheididae) from the Eastern Aegean Sea, Turkey. *Crustaceana*, 78 (10): 1265-1267.
- ATEŞ, A. S., KATAĞAN, T. & KOCATAŞ, A., 2004. New decapod species for the Turkish seas. *Crustaceana*, 77 (4): 507-512.

- BALKIS, H., BALKIS, N. & ALTINSAÇLI, S., 2001. The crab species found on the coasts of Gökçeada (Imbros) island in the Aegean Sea. *Hydrobiologia*, 449: 99-103.
- BELLAN-SANTINI, D., 1969. Contribution à l'étude des peuplement infralittoraux sur substrat rocheux (Étude qualitative et quantitative de la frange supérieure). *Recherche Travaux Station Marine Endoume*, France, 63 (47): 9-294.
- BORG, J.A., 1991. Species richness and abundance of decapod crustaceans associated with a Maltese *Posidonia oceanica* (L.) Delile meadow, Unpublished MSc dissertation, vi + 144 pp. Malta: University of Malta.
- BORG, S. A. & SCHEMBRI, P. J., 2000. Bathymetric distribution of decapods associated with a *Posidonia oceanica* meadow in Malta (central Mediterranean). *Crustaceana*, 12: 119-130.
- FALCIAI, L. & MINERVINI, R., 1996. Guide des homards, crabes, langoustes, crevettes et autres crustacés décapodes d'Europe: 1-287. (Delachaux et Niestlé SA, Lausanne-Paris).
- GALLMETZER, I., PFLUGFELDER, B., ZEKELY, J. & OTT, J. A., 2005. Macrofauna diversity in *Posidonia oceanica* detritus: distribution and diversity of mobile macrofauna in shallow sublittoral accumulations of *Posidonia oceanica* detritus. *Marine Ecology*, 147: 517-523.
- GARCÍA RASO, J. E., 1990. Study of a crustacea decapoda taxocoenosis of *Posidonia oceanica* beds from the southeast of Spain. *Marine Ecology*, 11(4): 309-326.
- GARCÍA RASO, J. E., LÓPEZ DE LA ROSA, I. & ROSALES, J. M., 1996. Decapod crustacean communities from calcareous seaweed and *Posidonia oceanica* (Rhizome stratum) in shallow waters. *Opheelia*, 45 (2): 143-158.
- INGLE, R., 1993. Hermit crabs of the North-eastern Atlantic Ocean and Mediterranean Sea. *National History Museum Publications*, 495 p.
- KOCATAŞ, A. & KATAĞAN., T., 2003. The decapod crustacean fauna of the Turkish Seas. *Zoology in the Middle East*, 29: 63-74.
- MARTIN, J. W. & DAVIS, G. E., 2001. An updated classification of the recent crustacea. *National History Museum Science Series*, 39: 1-124.
- MAZZELLA, L., SCIPIONE, M. B. & BUIA, M. C., 1989. Spatio-temporal distribution of algal and animal communities in a *Posidonia oceanica* meadow. *P. S. Z. N. I. Marine Ecology*, 10 (2): 107-129.
- PÉRÈS, J. M. & PICARD, J., 1964. Nouveau manuel de bionomie benthique de la Méditerranée. *Recl. Trav. Stn. Mar. Endoume*, 31 (47): 1-137.
- SCIPIONE, M. B., FRESI, E. & WITTMAN, K. J., 1983. The vagile fauna of *Posidonia oceanica* (L.) Delile foliar stratum: a community approach. *Rapport de la Commission Internationale Exploration de la Mer Méditerranée*, 28 (3): 141-142.
- SOYER, J., 1970. Bionomie benthique du plateau continental de la côte catalane française. III. Les peuplements de Copepodes harpacticoides (Crustacea). *Vie et Milieu*, 21: 337-511.
- ZARIQUIEY-ÁLVAREZ, R., 1968. Crustáceos Decápodos Ibéricos. *Investigación Pesquera*, Barcelona, 32: 1-510.
- ZUPO, V., 1990. I Decapodi delle praterie di *Posidonia oceanica*: confronto tra metodiche di campionamento in rapporto alla zonazione del taxon. *Oebalia* vol. XVI: 817-822.

Accepted in July 2007