

Journal of Humanitarian Cardiovascular Medicine

Vol 1, No 3 (2024)

Current Publications



IVUS and iFR: A Powerful duo for uncovering intermediate lesions in coronary artery disease

Antonios Samaras, Thrasivoulos Konstantinou, Evangelos Sdogkos, Ioannis Vogiatzis

doi: [10.12681/jhcv.36986](https://doi.org/10.12681/jhcv.36986)

Copyright © 2024, Antonios Samaras, Thrasivoulos Konstantinou, Evangelos Sdogkos, Ioannis Vogiatzis



This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/).

To cite this article:

Samaras, A., Konstantinou, T., Sdogkos, E., & Vogiatzis, I. (2024). IVUS and iFR: A Powerful duo for uncovering intermediate lesions in coronary artery disease. *Journal of Humanitarian Cardiovascular Medicine*, 1(3). <https://doi.org/10.12681/jhcv.36986>

IVUS and iFR: A Powerful duo for uncovering intermediate lesions in coronary artery disease

Antonios Samaras, Thrasimvoulos Konstantinou, Evangelos Sdogkos, Ioannis Vogiatzis


Department of Cardiology, General Hospital of Veroia

Corresponding author


Dr. Ioannis Vogiatzis, Department of Cardiology, General Hospital of Veroia, Veroia 59100, Greece
Tel. +302331351253 / +306944276230
e-mail: ivogia@hotmail.gr



A



B



C

A 55-year-old patient with a non-ST elevation myocardial infarction (NSTEMI).

A: Cranial view of the left main coronary artery with an apparent 50% lumen stenosis near the orifice.

B: The dominant right coronary artery that appears to be 70% stenotic near its orifice.

C: Intravascular ultrasound depiction of the vessel lumen, where the minimal lumen area appears to be 7.4 mm². iFR calculation uses a ratio of proximal and distal coronary pressures over the wave-free period in diastole to assess the limitation of blood flow.