

## Journal of the Hellenic Veterinary Medical Society

Vol 62, No 3 (2011)



### First identification of *Bartonella coopersplainsensis* in wild rodents (*Rattus norvegicus*) in Greece

E. PAPADOGIANNAKIS (Ε. ΠΑΠΑΔΟΓΙΑΝΝΑΚΗΣ), G. SPANAKOS (Γ. ΣΠΑΝΑΚΟΣ), I. KONTOU (Ι. ΚΟΝΤΟΥ), V. KONTOS (Β. ΚΟΝΤΟΣ), E. VELONAKIS (Ε. ΒΕΛΟΝΑΚΗΣ), D. SKLIVAS (Δ. ΣΚΛΗΒΑΣ), C. KOUTIS (Χ. ΚΟΥΤΗΣ)

doi: [10.12681/jhvms.14851](https://doi.org/10.12681/jhvms.14851)

#### To cite this article:

PAPADOGIANNAKIS (Ε. ΠΑΠΑΔΟΓΙΑΝΝΑΚΗΣ) E., SPANAKOS (Γ. ΣΠΑΝΑΚΟΣ) G., KONTOU (Ι. ΚΟΝΤΟΥ) I., KONTOS (Β. ΚΟΝΤΟΣ) V., VELONAKIS (Ε. ΒΕΛΟΝΑΚΗΣ) E., SKLIVAS (Δ. ΣΚΛΗΒΑΣ) D., & KOUTIS (Χ. ΚΟΥΤΗΣ) C. (2017). First identification of *Bartonella coopersplainsensis* in wild rodents (*Rattus norvegicus*) in Greece. *Journal of the Hellenic Veterinary Medical Society*, 62(3), 212–215. <https://doi.org/10.12681/jhvms.14851>

## ■ First identification of *Bartonella coopersplainsensis* in wild rodents (*Rattus norvegicus*) in Greece.

**Papadogiannakis E.<sup>a</sup> DVM, CertSAD, PhD, Spanakos G.<sup>b</sup> PhD, Kontou I.<sup>a</sup> MD, MPH,  
Kontos V.<sup>a</sup> DVM, PhD, Velonakis E.<sup>c</sup> MD, PhD, Sklivas D.<sup>a</sup> MPH, Koutis C.<sup>d</sup> MD, PhD**

<sup>a</sup> Department of Veterinary Public Health, National School of Public Health, Athens, Greece

<sup>b</sup> Department of Parasitology, Entomology and Tropical Diseases, National School of Public Health, Athens, Greece

<sup>c</sup> Department of Microbiology and Applied Immunology, National School of Public Health, Athens, Greece

<sup>d</sup> Department of Public Health, Technological Educational Institution of Athens, Athens, Greece

## ■ Η ταυτοποίηση της *Bartonella coopersplainsensis* σε άγρια τρωκτικά (*Rattus norvegicus*) για πρώτη φορά στην Ελλάδα.

**Ε. Παπαδογιαννάκης<sup>a</sup> DVM, CertSAD, PhD, Γ. Σπανάκος<sup>b</sup> PhD, Ι. Κοντού<sup>a</sup> MD, MPH,  
Β. Κοντός<sup>a</sup> DVM, PhD, Ε. Βελονάκης<sup>c</sup> MD, PhD, Δ. Σκλήβας<sup>a</sup> MPH, Χ. Κουτής<sup>d</sup> MD, PhD**

<sup>a</sup> Τομέας Κτηνιατρικής Δημόσιας Υγείας, Εθνική Σχολή Δημόσιας Υγείας, Αθήνα

<sup>b</sup> Τομέας Παρασιτολογίας, Εντομολογίας και Τροπικών Νοσημάτων, Εθνική Σχολή Δημόσιας Υγείας, Αθήνα

<sup>c</sup> Τομέας Μικροβιολογίας και Εφαρμοσμένης Ανοσολογίας, Εθνική Σχολή Δημόσιας Υγείας, Αθήνα

<sup>d</sup> Τομέας Δημόσιας Υγιεινής, Τ.Ε.Ι. Αθήνας

**ABSTRACT.** This study was a preliminary attempt to detect *Bartonella* sp. in tissues (liver and heart) obtained from wild rodents (*Rattus norvegicus*) in central Greece. Eighteen (18) liver and eighteen (18) heart samples were examined, which were obtained from twenty three (23) animals. Two (2) liver samples were found positive, while the heart samples examined were found negative for *Bartonella* sp. 16S-23S intergenic spacer rDNA gene by PCR. Sequencing of the positive PCR products and comparison with those available in GenBank using the BLAST program revealed the same species of *Bartonella* in both positive samples with 100% sequence homology to *Bartonella coopersplainsensis*.

**Keywords:** *Bartonella coopersplainsensis*, *Rattus norvegicus*, Greece, PCR

**ΠΕΡΙΛΗΨΗ.** Η μελέτη αυτή ήταν μία πρώτη προσπάθεια για ανίχνευση της *Bartonella* sp. σε ιστούς (ήπαρ και καρδιά) που πάρθηκαν από άγρια τρωκτικά (*Rattus norvegicus*) στην κεντρική Ελλάδα. Εξετάστηκαν δεκαοκτώ (18) δείγματα ήπατος και δεκαοκτώ (18) δείγματα καρδιάς, τα οποία προέρχονταν από είκοσι τρία (23) ζώα. Δύο (2) δείγματα ήπατος βρέθηκαν θετικά ενώ όλα τα δείγματα καρδιάς ήταν αρνητικά για το γονίδιο 16S-23S intergenic spacer rDNA της *Bartonella* sp. Η αλληλουχία των βάσεων του DNA των θετικών δειγμάτων και η σύγκριση με τα διαθέσιμα δεδομένα της GenBank με τη χρησιμοποίηση του προγράμματος BLAST, αποκάλυψε το ίδιο είδος *Bartonella* και στα δύο θετικά δείγματα με 100% ομοιότητα με την αλληλουχία των βάσεων της *Bartonella coopersplainsensis*.

**Λέξεις ενδετηρίασης:** *Bartonella coopersplainsensis*, *Rattus norvegicus*, Ελλάδα, PCR

**Correspondence:** Papadogiannakis E.  
Department of Veterinary Public Health, National School of Public Health, 115 21 Athens, Greece  
Tel.: +030 2132010217, Fax.: +030 2109818444, E-mail: dermpap1@otenet.gr

**Αλληλογραφία:** E. Παπαδογιαννάκης  
Τομέας Κτηνιατρικής Δημόσιας Υγείας, Εθνική Σχολή Δημόσιας Υγείας,  
Δ. Αλεξανδρας 196, 115 21 Αθήνα  
Τηλ.: +0302132010217, Fax.: +0302109818444, E-mail: dermpap1@otenet.gr

**Submission date:** 10.05.2011  
**Approval date:** 03.10.2011

**Ημερομηνία υποβολής:** 10.05.2011  
**Ημερομηνία εγκρίσεως:** 03.10.2011

The genus *Bartonella* includes fastidious, Gram-negative, aerobic, facultatively intracellular bacilli of erythrocytes and endothelial cells of various mammals. A significant number of *Bartonella* species is considered zoonotic pathogens and, apart from humans, they can infect several animal species including rodents, ruminants, cats and dogs (Breitschwerdt and Kordick 2000). In addition, *Bartonella* species have been detected in several arthropod vectors like fleas, lice and ticks (Billeter et al. 2008). It has been reported that *Bartonella* species are widely distributed among rodents in southern China (Ying et al. 2002), Thailand (Castle et al. 2004), Bangladesh (Bai et al. 2007), Europe (Birtles and Harrison 1994, Ellis et al. 1999, Engbaek and Lawson 2004) and America (Kosoy et al. 1997, Ellis et al. 1999). There has been only one report in which three *Bartonella* strains have been detected in wild rodents in northern Greece (Tea et al. 2004).

This study was a preliminary attempt to detect *Bartonella* sp. in tissues (liver and heart) obtained from wild rodents (*Rattus norvegicus*) in Greece.

All animals were live trapped at randomly selected suburban areas in the cities of Athens and Piraeus (it is the greater Athens area) in central Greece. Wire mesh traps were put from 6 p.m to 7 a.m.

Upon arrival at the Molecular Biology Laboratory of the National School of Public Health, the rodents were given general anaesthesia using a combination of xylazine and ketamine intramuscularly. Under aseptic conditions, the liver and heart were removed and the animal was then euthanized. A small part of the excised tissues was placed in sterile eppendorf tubes and stored at -80°C until DNA isolation. Samples were handled under sterile conditions to avoid the risk of cross-contamination. Liver and heart are among the internal organs affected following experimental inoculation of *Bartonella* sp. in immunocompetent mice (Colton et al. 2010).

In total, 18 heart and 18 liver samples were examined, which were obtained from 23 animals. All rodents were adult and belonged to the species *Rattus norvegicus*. Thirteen were males (56.52%) and 10 were females (43.48%). In 13 animals, the heart and liver samples were obtained from the same animal.

DNA was isolated from approximately 10 mg of liver or heart tissue, using the QIAamp DNA Mini kit (Qiagen, Valencia, CA, USA). DNA isolation was

performed according to the manufacturer's recommendations and DNA was eluted in 100 µl elution buffer (EB buffer) for increased concentration. Detection of *Bartonella* DNA by PCR was performed at 25 µl total reaction volume containing 1x PCR buffer, 2 mM Magnesium Chloride, 0.2 mM of each dNTP 1 µM of each primer and 5 µl of template DNA. Thermal cycling was performed with the following conditions: an initial denaturation step at 94°C for 2 min was followed by 45 cycles of 94°C for 15 sec, 66°C for 15 sec and 72°C for 15 sec and a final elongation step at 72°C for 10 min. Primers used were

325s:

(5CTTCAGATGATGATCCCAAGCCTCTGGCG) and

1100as:

(5GAACCGACGACCCCTGCTTGCAAAGCA3),

obtained from Syntezza (Israel), as previously described (Maggi et al. 2006). These primers amplify a region of 16S-23S ribosomal RNA intergenic spacer (ITS) (Cadenas et al. 2008, Maggi et al. 2006). One positive (DNA extracted from *Bartonella henselae* from culture) and one negative control were included in each reaction. PCR products were electrophorised in 2% agarose gel containing ethidium bromide, visualized under UV light and photographed with a digital camera. The band of interest was excised and DNA was isolated using the QIAquick gel extraction kit (Qiagen, Valencia, CA, USA) according the manufacturer's instructions. Both strands of the DNA were directly sequenced at Biogenomica (Vrillisia, Athens, Greece) using the primers used for PCR. Nucleotide sequences were manually corrected using finchtv 1.3.1 for linux (downloaded from <http://www.geospiza.com/Products/finchtv.shtml>). Sequences obtained were compared to those available in GenBank using the web interface of BLAST 2.2.23+ program (Zhang et al. 2000).

Two (2) liver samples were found positive for *Bartonella* sp. by PCR, representing a prevalence of 11.11% of the total number of liver samples and 8.69% of the total number of animals examined. In contrast, all heart samples examined were found to be negative. A heart sample was available and examined in only one out of the two positive animals. Sequencing of the PCR products obtained in two independent reactions and comparison with those available in GenBank using the

BLAST program revealed sequences identical to each other (GenBank accession number HM749323), with 100% sequence homology to *Bartonella coopersplainsensis* strain AUST/NH20 (EU111770), *Bartonella* sp. strain Rt222sm (AY277896) and *Bartonella* sp. strain Lao/Nh1 (EU714976).

Rodents seem to play an important role in *Bartonella* infection, since several species of *Bartonella* have been detected in rodents by isolation and/or molecular detection. It has been shown that a) rodents constitute a natural “reservoir” of various species of *Bartonella*, b) the bacterium is commonly isolated from rodents living in suburban areas and c) following experimental infection congenital transmission has been observed (Ellis et al. 1999). Accordingly, the determination of the *Bartonella* species in wild rodents in a specific area (or country) greatly contributes to the better understanding of the epidemiology and prevention of *Bartonella* spreading to animals and humans.

There has been only one published report in which *Bartonella taylorii*, *Bartonella birtlesii* and *Bartonella grahamii* were isolated from the blood of wild rodents (*Apodemus flavicollis*, *Microtus agresti* and *Dryomys nitedula*) in northern Greece (Tea et al. 2004). However, to the best of the authors' knowledge, *Bartonella coopersplainsensis* identified in the present study has never been reported either in Greece or in Europe until today. It seems that different species of rodents host different *Bartonella* species, although different geographical distribution of the bacterium cannot be excluded.

*Bartonella coopersplainsensis* isolated from the blood of wild rats of the species *Rattus leucopus* in Australia has recently been identified as a novel *Bartonella* species (Gundi et al. 2009).

It has been shown that *Bartonella* sp. has been detected in the liver and heart of experimentally inoculated mice (Colton et al. 2010). This probably explains the detection of *Bartonella* in the liver samples in our study, although the heart sample was not

available from one positive animal. In a recent study carried out in Japan, bacteria belonging to *Bartonella* species were not detected in the blood obtained from *Rattus norvegicus* (Inoue et al. 2008), whereas a similar study in southern France revealed a 16.6% infection in the blood and heart samples obtained from *Rattus norvegicus* (Gundi et al. 2004).

The zoonotic potential of rodent-borne *Bartonella* spp. is emerging, as suggested by a novel rodent *Bartonella*-associated febrile disease in the rural southwestern USA, based on serological evidence (Iralu et al. 2006). At present, only a limited number of human infections caused by rodent-borne *Bartonella* species has been reported. However, previous experience has shown that *Bartonella* species firstly detected or isolated in animals could be later involved in humans, such as *Bartonella alsatica*, which was firstly isolated from wild rabbits (Heller et al. 1999) and later identified in cases of endocarditis (Raoult et al. 2006) and cat scratch disease (Angelakis et al. 2008) in humans. Accordingly, although the public health significance of *Bartonella coopersplainsensis* identified in our study remains to be clarified, it should be prudent for people working in conditions bringing them in contact with wild rodents to take the appropriate measures to protect themselves from direct contact with these animals, as well as their ectoparasites (e.g. fleas). Arthropods (ectoparasites) constitute one of the main ways of *Bartonella* transmission from one animal to another or to humans (Morick et al. 2009).

Further studies are needed to clarify in which way the wild rodents *Rattus norvegicus* contaminate the environment with *Bartonella coopersplainsensis* and possible mode of transmission to humans and other animals.

#### Conflict of interest statement

None of the authors of this paper has a financial or personal relationship with other people or organizations that could inappropriately influence or bias the content of the paper. 

## REFERENCES

Angelakis, E., Lepidi, H., Cane, A., Rispal, P., Perraudeau, F., Barre, I., Rolain, J., Raoult, D., (2008). Human case of *Bartonella alsatica* lymphadenitis. *Emerging Infectious Diseases* 14(12): 1951-1953.

Bai, Y., Montgomery, S., Sheff Kelly, W., Chowdhury, M., Breiman, R., Kabeya, H., Kosoy, M., (2007). *Bartonella* Strains in Small Mammals from Dhaka, Bangladesh, Related to *Bartonella* in America and Europe. *American Journal of Tropical Medicine and Hygiene* 77(3): 567-570.

Billeter, S., Levy, M., Chomel, B., Breitschwerdt, E., (2008). Vector transmission of *Bartonella* species with emphasis on the potential for tick transmission. *Medical and Veterinary Entomology*, 22(1): 1-15.

Birtles, R., Harrison, T., (1994). *Grahamella* in small woodland mammals in the U.K.: isolation, prevalence and host specificity. *Annals of Tropical Medicine and Parasitology*, 88: 317-327.

Breitschwerdt, E., Kordick, D., (2000). *Bartonella* infection in animals: carriership, reservoir potential, pathogenicity and zoonotic potential for human infection. *Clinical Microbiology Reviews*, 13: 428-38.

Cadenas, M., Bradley, J., Maggi Ricardo, G., Takara, M., Hegarty, B., Breitschwerdt, E., (2008). Molecular Characterization of *Bartonella vinsonii* subsp. *berkhoffii* Genotype III. *Journal of Clinical Microbiology*, 46(5): 1858-1860.

Castle, K., Kosoy, M., Lerdthusnee, K., Phelan, L., Bai, Y., Gage, K., Leepitakrat, W., Monkanna, T., Khlaimanee, N., Chandranoj, K., Jones, J., Coleman, R., (2004). Prevalence and diversity of *Bartonella* in rodents of northern Thailand: A comparison with *Bartonella* in rodents from southern China. *American Journal of Tropical Medicine and Hygiene*, 70(4): 429-433.

Colton L., Zeidner N., Lynch T., and Kosoy M. (2010). Human isolates of *Bartonella tamiae* induce pathology in experimentally inoculated immunocompetent mice. *BMC Infectious Diseases*, 10: 229-237.

Ellis, B., Regnery, R., Beati, L., (1999). Rats of the genus *Rattus* are reservoir hosts for pathogenic *Bartonella* species: an old world origin for a new world disease? *Journal of Infectious Diseases*, 180: 220-224.

Engbaek, K., Lawson, P., (2004). Identification of *Bartonella* species in rodents, shrews and cats in Denmark: detection of two *B. henselae* variants, one in cats and the other in the long tailed field mouse. *Acta Pathologica Microbiologica et Immunologica*, 112: 336-341.

Gundi, V., Davoust, B., Khamis, A., Boni, M., Raoult, D., La Scola, B., (2004). Isolation of *Bartonella ratti* sp. nov. and *Bartonella phoceensis* sp. nov. from European *Rattus norvegicus*. *Journal of Clinical Microbiology*, 42(8): 3816-3818.

Gundi, V., Taylor, C., Raoult, D., La Scola, B., (2009). *Bartonella ratta* sp. nov., *Bartonella queenslandensis* sp. nov. and *Bartonella coopersplainensis* sp. nov., identified in Australian rats. *International Journal of Systematic and Evolutionary Microbiology*, 59(12): 2956-2961.

Heller, R., Kubina, M., Mariet, P., Riegel, P., Delacour, G., Dehio, C., Lamarque, F., Kasten, R., Boulouis, H., Monteil, H., Chomel, B., Piemont, Y., (1999). *Bartonella alsatica* sp. nov., a new *Bartonella* species isolated from the blood of wild rabbits. *International Journal of Systematic Bacteriology*, 49(1): 283-288.

Inoue, K., Maruyama, S., Kabeya, H., Yamada, N., Ohashi, N., Sato, Y., Yukawa, M., Masuzawa, T., Kawamori, F., Kadosaka, T., Takada, N., Fujita, H., Kawabata, H., (2008). Prevalence and Genetic Diversity of *Bartonella* Species Isolated from Wild Rodents in Japan. *Applied and Environmental Microbiology*, 74(16): 5086-5092.

Iralu, J., Bai, Y., Crook, L., Tempest, B., Simpson, G., Mckenzie, T., Koster, F., (2006). Rodent-associated *Bartonella* febrile illness, South-western United States. *Emerging Infectious Diseases*, 12(7): 1081-1086.

Kosoy, M., Regnery, R., Tzianabos, T., (1997). Distribution, diversity and host specificity of *Bartonella* in rodents from the southeastern United States. *American Journal of Tropical Medicine and Hygiene*, 57: 578-588.

Maggi, R., Diniz, M., Cadenas, B., Breitschwerdt, E., (2006). The use of molecular diagnostic techniques to detect *Anaplasma*, *Bartonella* and *Ehrlichia* species in arthropods or patients. *Proceedings of the First International Canine Vector-Borne Diseases Symposium* organized by Bayer, Birmingham, United Kingdom, pp: 9-13.

Morick, D., Baneth, G., Avidor, B., Kosoy, M., Mumcuoglu, K., Mintz, D., Eyal, O., Goethe, R., Mietze, A., Shpigel, N., Harrus, S., (2009). Detection of *Bartonella* spp. in wild rodents in Israel using HRM real-time PCR. *Veterinary Microbiology*, 139: 293-297.

Raoult, D., Roblot, F., Rolain, J., Besnier, J., Loulergue, J., Bastides, F., Choutet, P., (2006). First isolation of *Bartonella alsatica* from a valve of a patient with endocarditis. *Journal of Clinical Microbiology*, 44(1): 278-279.

Tea, A., Alexiou-Daniel, S., Papoutsis, A., Papa, A., Antoniadis, A., (2004). *Bartonella* species isolated from rodents, Greece. *Emerging Infectious Diseases*, 10(5): 963-964.

Ying, B., Kosoy, M., Maupin, G., (2002). Genetic and ecological characteristics of *Bartonella* communities in rodents in southern China. *American Journal of Tropical Medicine and Hygiene*, 66: 622-627.

Zhang, Z., Schwartz, S., Wagner, L., Miller, W., (2000). A greedy algorithm for aligning DNA sequences. *Journal of Computerized Biology*, 7(1-2): 203-14.