

Journal of the Hellenic Veterinary Medical Society

Vol 68, No 3 (2017)



Genetic Polymorphisms of Cyp19 and Myostatin Genes in Turkish Indigenous Sheep Breeds

I. AKIŞ, F. ESEN GÜRSEL, N. HACIHASANOĞLU ÇAKMAK, G. ATMACA, H. YARDIBI, A. ATEŞ, M. H. DURAK, K. ÖZTABAK

doi: [10.12681/jhvms.15475](https://doi.org/10.12681/jhvms.15475)

Copyright © 2018, I AKIŞ, F ESEN GÜRSEL, N HACIHASANOĞLU ÇAKMAK, G ATMACA, H YARDIBI, A ATEŞ, MH DURAK, K ÖZTABAK



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

To cite this article:

AKIŞ, I., ESEN GÜRSEL, F., HACIHASANOĞLU ÇAKMAK, N., ATMACA, G., YARDIBI, H., ATEŞ, A., DURAK, M. H., & ÖZTABAK, K. (2018). Genetic Polymorphisms of Cyp19 and Myostatin Genes in Turkish Indigenous Sheep Breeds. *Journal of the Hellenic Veterinary Medical Society*, 68(3), 313–318. <https://doi.org/10.12681/jhvms.15475>

Genetic Polymorphisms of Cyp19 and Myostatin Genes in Turkish Indigenous Sheep Breeds

Akış, I.¹, Esen Gürsel, F.¹, Hacıhasanoğlu Çakmak, N.², Atmaca,
G.¹, Yardibi, H.¹, Ateş, A.¹, Durak M.H.³, Öztürk, K.¹

¹ Istanbul University, Faculty of Veterinary Medicine, Department of Biochemistry, Avcılar, 34310, Istanbul, Turkey

² Istanbul Medipol University, Vocational School of Health Services, Medical Laboratory Techniques Program,
Istanbul, Turkey

³ Dicle University, Faculty of Veterinary Medicine, Department of Biochemistry, 21280, Diyarbakır, Turkey

ABSTRACT. Growth and meat production traits are very important in sheep breeding. *Cyp19* gene has a major role in reproductive activity and growth due to its function in estrogen synthesis. Another gene affecting growth traits is Myostatin (*MSTN*) gene, which mainly regulates skeletal muscle growth. In this study allele frequencies of genetic polymorphism in *Cyp19* and Myostatin genes were identified by PCR-RFLP method in five indigenous Turkish sheep breeds, Chiose, Imroz, Kivircik, Zom and Morkaraman. Digestion of *Cyp19* gene with *Hae*III only revealed uncut AA genotype and digestion of *MSTN* with *Dra*I also revealed only uncut AA genotype. Both loci analyzed in this study were found to be monomorphic in five Turkish indigenous sheep breeds. These highly conserved parts of the two genes can be useful for molecular evolutionary studies in sheep. Further studies regarding association analysis of *Cyp19* and *MSTN* in sheep should be conducted.

Key words: Cyp19, myostatin, sheep, SNP, PCR-RFLP

Corresponding Author:

Assos. Prof. Feraye Esen Gürsel
Istanbul University, Faculty of Veterinary Medicine,
Department of Biochemistry, Avcılar-Istanbul, Turkey
E-mail: fesen@istanbul.edu.tr

Date of initial submission: 4.7.2016
Date of revised submission: 8.8.2016
Date of acceptance: 8.8.2016

INTRODUCTION

Sheep is one of the most important livestock animals bred in Turkey. Turkey has 31 million sheep and 10% of total meat production in the country consists of sheep meat (www.tuik.gov.tr). There are several indigenous sheep breeds in Turkey, which are bred for different production traits like meat, milk and wool (Yalcin, 1986).

Growth and meat production traits are significant economic traits in sheep. In the past years many single nucleotide polymorphisms (SNPs) and in/del polymorphisms in livestock have been identified by genome-wide association studies in order to facilitate the potential utilization of genes involved in growth and meat production traits. But the number of quantitative trait loci (QTL) identified in sheep still remains limited (Zhang et al., 2013).

Oestrogen is an important endocrine, paracrine and autocrine acting hormone, which has a major role in the regulation of reproduction system. It is also involved in fat deposition and growth (Heine et al., 2000; Jones et al., 2000, Simpson et al., 2000). The aromatase cytochrome P450 enzyme encoded by the *Cyp19* gene, is responsible for estrogen biosynthesis by conversion or aromatization of androgens into estrogens (Simpson et al., 1994).

In sheep, the *Cyp19* gene that encodes the aromatase enzyme is located in chromosome 7 (Payen et al., 1995). It is transcribed by four different promoter regions that have organ-specific activities. The P2 promoter region is mostly active in granulosa cells, P1.5 and P1.1 in the placenta and P1.4 is active in the brain (Vanselow et al., 1999; Vanselow et al., 2001).

Myostatin (growth and differentiation factor - GDF8) is the major regulator of myogenesis in mammals. It is a member of the transforming growth factor- β superfamily and acts as a negative regulator of muscle growth. It directly affects muscular hypertrophy and carcass conformation (Nakev et al., 2013; Zhang et al., 2013). It has been reported, that mutations in *MSTN* gene located on OAR2, which encodes myostatin, are associated with increased skeletal muscle mass in sheep (Kijas et al. 2007). The *MSTN* gene has 3 exons and 2 introns (Bellinge et al., 2005) and encodes a glycoprotein which is widely expressed in skeletal muscle. The differentiations

in this protein may be responsible for changing the composition of muscle fibers and causing a variation of muscle weight (Chen, 2008). Increased levels of muscle myostatin protein may also have a role in mediating effects of estrogen on growth in skeletal muscle (Jung et al., 2007).

The aim of the present study is to determine allele frequencies of genetic polymorphisms for *Cyp19* and *MSTN* genes in indigenous Turkish sheep breeds, namely Chios, İmroz, Kivircik, Zom and Morkaraman.

MATERIALS AND METHODS

Animals and DNA isolation

In this study unrelated animals from Chiose (n=35), İmroz (n=35), Kivircik (n=38), Zom (n=35) and Morkaraman (n=33) sheep breeds were investigated. Blood samples were collected into 2 ml sterilized tubes with EDTA from *V. jugularis*. Genomic DNA was isolated by standard salt-out method (Miller et al., 1998).

The study had an approval from the Ethical Committee of Istanbul University, Faculty of Veterinary Medicine with number 30/04/2008- 39.

PCR-RFLP analysis

PCR was performed in a reaction volume of 25 μ l using 1 U Taq DNA polymerase (Fermantas Life Sciences, Canada), 2-2.5 μ l 10X PCR buffer, 1.5mM MgCl₂, 50-100 ng genomic DNA, 100 μ M dNTP (Takara, Biotechnology Co, Ltd, Japan) and 10 pmol of each primer. Amplification was carried out with an initial denaturation at 94 °C for 5 min; 35 cycles of 94 °C for 1min, primer specific annealing temperature for 1 min, 72 °C for 1 min; and a final extension at 72 °C for 10 min. Primer sequences used to amplify intron 9 of *Cyp19* gene and 5'UTR of *MSTN* gene, their annealing temperatures (T_m) and product sizes were given in Table 1.

For RFLP analysis 10 μ l of the PCR products were digested with 10 units of *Hae*III and *Dra*I restriction enzymes at 37°C overnight in order to genotype *Cyp19* and *MSTN* polymorphisms, respectively. The digested DNA fragments were separated by electrophoresis in 2% agarose gel including ethidium bromide and visualized under UV light.

Table 1. Primer sequences, annealing temperatures and PCR products' sizes

Gene symbol	Primer sequence	Tm (°C)	Product size (bp)
Cyp19	F: 5' - CCA AAG CCT AAT GAA TTT ACT C - 3'	56	266
	R: 5' - CTC TCG TGT GCC CTC CAT GAA G - 3'		
MSTN	F: 5' - TGC CGT TAC TCA AAA GCA AA - 3'	54	497
	R: 5' - AAC AGC AGT CAG CAG AGT CG - 3'		

Statistical analysis

Direct counting was used to estimate genotype and allele frequencies of the genetic variants for all loci. Genetic relationships regarding *Cyp19* gene among populations analyzed in this study and seven sheep breeds from Europe (Zsolnai et al., 2002) were visualized in a dendrogram constructed by unweighted paired group cluster analysis (UPGMA), from a modified NEIGHBOR procedure implemented in PHYLIP version 3.5 software also using PopGene32 (Yeh et al., 2000). The UPGMA dendrogram of population was constructed based on Nei's genetic distance (Nei, 1972). No further analysis was carried out, since the two loci analyzed in five sheep breeds were found to be monomorphic.

RESULTS

The SNP in intron 9 of *Cyp19* gene could be assayed by RFLP analysis with *Hae*III restriction enzyme. Digestion of the 266 bp PCR product reveals allele G (266 bp undigested fragment) and allele C (190 and 76 bp fragments). All of the breeds analyzed in this study were found to be monomorphic for this locus (Table 2). Only GG genotype was observed (Figure 1).

The digestion of 5'UTR of *MSTN* gene by *Dra*I restriction enzyme results in two alleles, A (497 bp undigested fragment) and B (427 and 70 bp fragments). In this study only AA genotype was observed (Figure 1). The indigenous Turkish sheep breeds were found to be monomorphic for this locus (Table 2).

The phylogenetic tree based on G/C SNP in ovine *Cyp19* gene presented two clusters among the sheep breeds analysed in this study and different European

breeds (Figure 2). All the Turkish sheep breeds were clustered in the same branch with the Racka breed from Hungary.

DISCUSSION


Due to its function in reproductive activity and growth, many studies have been conducted on ovine *Cyp19* gene in different sheep breeds (Vanselow et al., 1999; Lobo et al., 2009; Tanamati et al., 2013).

The G/C transversion in intron 9 of ovine *Cyp19* gene was firstly identified in Spanish breeds Carranzana, Latxa Black Face and Latxa White Face, the Merino and British Milk Sheep breeds and indigenous Hungarian breeds, Cikta, Racka and

Figure 1. PCR-RFLP fragments of two SNPs. M: 100 bp marker, Lane 1, 2 and 3: Cyp19/*Hae*III GG genotype (266 bp), Lane 4, 5 and 6: MSTN/*Dra*I AA genotype (497 bp)



Figure 2. Genetic distance between Turkish and European sheep breeds according to *Cyp19*/HaeIII polymorphism. Dendrogram based on Nei's genetic distance.



Tsigai. Allele C were observed more frequently in all the breeds with frequencies ranged between 0.650 and 0.988 (Zsolnai et al., 2002). In the present study the five indigenous Turkish sheep breeds, Chiose, Imroz, Kivircik, Zom and Morkaraman were found to be GG homozygous. Based on the analysis of this genetic locus the breeds from Turkey seem to be genetically apart from the breeds originated from Europe. The phylogenetic tree (Figure 2) obtained from the *Cyp19* G/C polymorphism analysis of the five Turkish sheep breeds analyzed in the present study and seven European breeds mentioned above, presented two clusters. The Turkish breeds clustered with Racka breed from Hungary and the rest European breeds were grouped in the second cluster.

Absence of allele C in indigenous Turkish breeds may be due to a mutation exists in sheep from warmer regions. The geographic origin of the animals is an important factor regarding the sheep reproductive activity, as it determines the photoperiods (Sa and Sa, 2006), which affects melatonin synthesis. Melatonin

is an indoleamine derived from the serotonin secreted during darkness that acts as a natural inhibitor of the aromatases (Hafez and Hafez, 2003). The climatic and seasonal conditions of Turkey differ from the ones in Europe. Especially, three breeds in this study Kivircik, Zom and Morkaraman are raised mainly in central and south-eastern parts of Turkey, where the climate is semi-arid (Bilgili et al., 2013). We may suggest a relationship between *Cyp19* G/C polymorphism and the geographical origin of sheep breeds.

Another gene, which is largely investigated in different sheep breeds is *MSTN*, which is associated with meat-related traits, especially muscle growth (Nakev et al., 2013).

A deletion of a small DNA fragment (TTTTA) in the 5'UTR of myostatin gene was identified in several goat breeds (Xianglong et al., 2008; Singh et al., 2014). Xianglong et al. (2008) suggested a relationship of this deletion with body weight and size in goats and that the heterozygote populations present

better growth traits than homozygotes. In a study on Indian goat breeds researchers observed that the allele A which presents TTTA deletion, is almost fixed with a frequency of 0.98 (Singh et al., 2014). The results of the study on Egyptian sheep breeds showed that this deletion is not unique for goats. The researchers could only observe AB genotype. Allele A had frequencies ranged between 0.07 and 0.23 in three Indian sheep breeds (Shafey et al., 2014). In the present study, five indigenous Turkish sheep breeds were found to be monomorphic for this deletion. All individuals analyzed had AA genotype. No association analysis of this deletion has been performed in sheep breeds until now. Regarding to its effects in goats, we may suggest that the Turkish breeds analyzed in the present study may have a disadvantage in terms of growth traits.

In addition, Khani et al. (2014) reported a significant association of TTTTA deletion in 5'UTR of *MSTN* gene with twinning in goats. Although Chiose sheep breed has a higher twinning rate compared to the other indigenous breeds in this study, our

results did not present any difference among the five sheep breeds. Association analysis of this in/del polymorphism in sheep may provide results different than these found in goats.

CONCLUSION

Both loci analysed in this study were found to be monomorphic in five Turkish indigenous sheep breeds. *Cyp19* and *MSTN* genes have highly conserved parts, which can be helpful for molecular evolutionary studies in sheep. Further analysis which will analyze the association between these polymorphisms and production traits in sheep should be conducted.

ACKNOWLEDGEMENT

This study was supported by Istanbul University, Faculty of Veterinary Medicine with project numbers VET-TB-16-1, VET-TB-16-2, BEK-2016-22550 and BEK-2017-24900

REFERENCES

- Bilgili AV, Güzel Y, İkinci A (2013) Modeling temporal variability of soil CO₂ emissions from an apple orchard in the Harran Plain of southeastern Turkey. *Turk J Agric For* 37(6):744-761.
- Zsolnai A, Anton I, Fésüs L, Estonba A, Schwerin M, Vanselow J (2002) Allele distributions of two novel SNPs within the sheep CYP19 gene. *J Anim Breed Genet* 119(6): 402-405
- Bellinge RH, Liberles DA, Iaschi SP, O'Brien PA, Tay GK (2005). Myostatin and its implications on animal breeding: a review. *Anim Genet* 36:1-6.
- Chen TT (2008) Polymorphisms of MSTN, IGFBP-3 gene and the related research with growth performance of Tianfu goat. Sichuan Agricultural University, Ya'an.
- Hafez B, Hafez ESE (2003) Ciclos reprodutivos. In: WAHID H, HAFEZ ESE. *Reprodução animal*. 7th ed. São Paulo: Manole: pp 55-67.
- Heine PA, Taylor JA, Iwamoto GA, Lubahn DB, Cooke PS (2000) Increased adipose tissue in male and female estrogen receptor-alpha knockout mice. *P Natl Acad Sci USA* 97(23):12729-12734.
- Jones ME, Thorburn AW, Britt KL, Hewitt KN, Wreford NG, Proietto J, OZ OK, Leury BJ, Robertson KM, Yao S., Simpson ER (2000) Aromatase-deficient (ArKO) mice have a phenotype of increased adiposity. *P Natl Acad Sci USA* 97(23):12735-12740
- Kijas JW, McCulloch R, Edwards JE, Oddy VH, Lee S.H, Van Der Werf J (2007) Evidence for multiple alleles effecting muscling and fatness at the ovine GDF8 locus. *BMC Genet* 8: 80-90.
- Zhang L, Jiasen L, Fuping Z, Hangxing R, Lingyang X, Jian L, Shifang Z, Xiaoning Z, Caihong W, Guobin L, Youmin Z, Lixin D (2013) Genome-wide association studies for growth and meat production traits in sheep. *Plos One* 8(6): e66569.
- Lôbo AMO, Lôbo RN, Paiva SR (2009) Aromatase gene and its effects on growth, reproductive and maternal ability traits in a multibreed sheep population from Brazil. *Genet Mol Biol* 32(3): 484-490.
- Miller M, Dykes DD, Polesky HF (1998) A simple salting out procedure for extracting DNA from human nucleated cells. *Nucl Acids Res* 16: 1215.
- Nakev G, Stamberov P, Dimitrova I, Stancheva N, Georgieva S, Hristova D, Angelov G, Mehmedov T, Genova K, Teneva A (2013) Growth and development of skeletal muscle in connection with the expression of the Myostatin Gene (MSTN). In: *Proceedings of the 10th International Symposium Modern Trends in Livestock Production*, Belgrade, Serbia: pp 640-647.
- Payen E, Saidi-Mehtar N, Pailhoux E, Cotinot C (1995) Sheep gene mapping: Assignment of ALDOB, CYP19, WT and SOX2 by somatic cell hybrid analysis. *Anim Genet* 26: 331-333.
- Sa CO, Sa JL (2006) Estacionalidade reprodutiva em ovinos Piracicaba. <http://www.farmpoint.com.br/radares-tecnicos/reproducao/estacionalidade-reprodutiva-em-ovinos-155n.aspx> (accessed 14 February 2016)
- Simpson ER, Mahendroo MS, Means GD, Kilgore MW, Hinshelwood MM, Graham-Lorence S, Amarnah B, Yuji I, Fisher CR, Michael MD, Mendelson CR, Bulun SE (1994) Aromatase cytochrome P450, the enzyme responsible for estrogen biosynthesis. *Endocr Rev* 15:342-355.
- Simpson E, Rubin G, Clyne C, Robertson K, O'Donnell L, Jones M, Davis S. (2000) The role of local estrogen biosynthesis in males and females. *Trends Endocrin Met* 11(5):184-188.
- Singh SP, Kumar R, Kumari P, Kumar S, Mitra A (2014) Characterization of 5' upstream region and investigation of TTTTA deletion in 5' UTR of myostatin (MSTN) gene in Indian goat breeds. *Anim Biotech* 25(1):55-68.
- Tanamati F, Silva SCC, Mora NHAP, Schuroff GP, Macedo FAF, Gasparino E (2013) Efeito do polimorfismo no gene da aromatase sobre a característica peso vivo ao abate em ovinos Santa Inês. In: *Anais do X Simpósio Brasileiro de Melhoramento Animal* [online], Uberaba, MG. <http://sbmaonline.org.br/anais/x/trabalhos/pdf/6NKA.pdf> (accessed 20 May 2016)
- Vanselow J, Fürbass R, Zsolnai A, Kalbe C, Said HM, Schwerin M (2001) Expression of the aromatase cytochrome P450 encoding gene in cattle and sheep. *J Steroid Biochem Mol Biol* 79:279-288.
- Vanselow J, Zsolnai A, Fésüs L, Schmidt P, Schwerin MA (1999) Bsp1431 PCR-RFLP in exon 3 of the ovine aromatase gene (CYP19). *Anim Genet* 30:382-405.
- Li, X, Liu, Z, Zhou R, Zheng G, Gong Y, Li L (2008) Deletion of TTTTA in 5' UTR of goat MSTN gene and its distribution and genetic effect on body weight. *Front Agric China* 2(1):103-109.
- Yalçın BC (1986) Sheep and goats in Turkey, Animal production and protection paper 60, Food and Agriculture Organisation of the United Nations, Rome <http://www.fao.org/3/a-ah224e.pdf> (accessed 20 May 2016)
- Yeh F, Yang RC, Boyle T: Popgene (v. 1.32), Microsoft Windows based freeware for Population Genetic Analysis. Retrieved from <http://www.ualberta.ca/~fyeh/Pop32.exe>. 2000.
- Zhang ZJ, Ling YH, Wang LJ, Hang YF, Guo XF, Zhang YH, Ding JP, Zhang XR (2013) polymorphisms of the myostatin gene (MSTN) and its relationship with growth traits in goat breeds. *Genet Mol Res* 12(2): 965-971.