

Journal of the Hellenic Veterinary Medical Society

Vol 66, No 1 (2015)



Small Ruminant Lenti-virus infection in imported sheep in Greece

N. D. GIADINIS (Ν.Δ. ΓΙΑΔΙΝΗΣ), C. I. DOVAS (Χ.Ι. ΔΟΒΑΣ), E. CHATZINASIΟΥ (Ε. ΧΑΤΖΗΝΑΣΙΟΥ), I. BOUZALAS (Η. ΜΠΟΥΖΑΛΑΣ), G. ARSENOS (Γ. ΑΡΣΕΝΟΣ), M. PAPANASTASSOPOYLOU (Μ. ΠΑΠΑΝΑΣΤΑΣΣΟΠΟΥΛΟΥ)

doi: [10.12681/jhvms.15580](https://doi.org/10.12681/jhvms.15580)

To cite this article:

GIADINIS (Ν.Δ. ΓΙΑΔΙΝΗΣ) N. D., DOVAS (Χ.Ι. ΔΟΒΑΣ) C. I., CHATZINASIΟΥ (Ε. ΧΑΤΖΗΝΑΣΙΟΥ) E., BOUZALAS (Η. ΜΠΟΥΖΑΛΑΣ) I., ARSENOS (Γ. ΑΡΣΕΝΟΣ) G., & PAPANASTASSOPOYLOU (Μ. ΠΑΠΑΝΑΣΤΑΣΣΟΠΟΥΛΟΥ) M. (2017). Small Ruminant Lenti-virus infection in imported sheep in Greece. *Journal of the Hellenic Veterinary Medical Society*, 66(1), 27–30. <https://doi.org/10.12681/jhvms.15580>

Small Ruminant Lenti-virus infection in imported sheep in Greece

Giadinis N.D.^{1*}, Dovas C.I.², Chatzinasiou E.³, Bouzalas I.³,
Arsenos G.⁴, Papanastassopoulou M.³

¹Clinic of Farm Animals, Faculty of Veterinary Medicine, Aristotle University of Thessaloniki, Greece

²Diagnostic Laboratory, Faculty of Veterinary Medicine, Aristotle University of Thessaloniki, Greece

³Laboratory of Microbiology and Infectious Diseases, Faculty of Veterinary Medicine, Aristotle University of Thessaloniki, Greece

⁴Laboratory of Animal Husbandry, Faculty of Veterinary Medicine, Aristotle University of Thessaloniki, Greece

Μόλυνση από Lenti-ιούς σε εισαγόμενα πρόβατα στην Ελλάδα

Γιαδίνης Ν.Δ.^{1*}, Δόβας Χ.Ι.², Χατζηνάσιου Ε.³, Μπουζαλάς Η.³,
Αρσένος Γ.⁴, Παπαναστασσοπούλου Μ.³

¹Κλινική Παραγωγικών Ζώων, Τμήμα Κτηνιατρικής, Σχολή Επιστημών Υγείας, Α.Π.Θ.

²Διαγνωστικό Εργαστήριο, Τμήμα Κτηνιατρικής, Σχολή Επιστημών Υγείας, Α.Π.Θ.

³Εργαστήριο Μικροβιολογίας και Λοιμωδών Νοσημάτων, Τμήμα Κτηνιατρικής, Σχολή Επιστημών Υγείας, Α.Π.Θ.

⁴Εργαστήριο Ζωοτεχνίας, Τμήμα Κτηνιατρικής, Σχολή Επιστημών Υγείας, Α.Π.Θ.

ABSTRACT. During the last years, a large number of Assaf-E sheep have been brought into Greece from Spain. In total, 57 randomly selected Assaf-E sheep purchased from Spain to five 5 different farms in Greece were examined serologically on arrival for *Small Ruminant Lentiviruses*. Of these, 25 animals were found to be seropositive against *Small Ruminant Lentiviruses*. Taking into account the above findings, we suggest a more strict legislation at the sheep sales.

Keywords: Assaf-E, Greece, sheep, *Small Ruminant Lentiviruses*, Spain

ΠΕΡΙΛΗΨΗ. Πρόσφατα, μεγάλος αριθμός προβάτων φυλής Assaf-E έχουν εισαχθεί στην Ελλάδα από διάφορες ισπανικές εκτροφές. Από αυτά, 57 πρόβατα επιλεγμένα τυχαία από πέντε διαφορετικές Ελληνικές εκτροφές, εξετάστηκαν ορολογικά για Lenti-ιούς των μικρών μηρυκαστικών κατά την άφιξή τους. Συνολικά, 25 ζώα βρέθηκαν οροθετικά στους παραπάνω ιούς. Λαμβάνοντας υπόψη τα ευρήματα, προτείνουμε την καθιέρωση πιο αυστηρής νομοθεσίας για τις αγοραπωλησίες προβάτων.

Λέξεις ευρετηρίασης: Ελλάδα, Ισπανία, πρόβατα, Assaf-E, Lenti-οί μικρών μηρυκαστικών

Correspondence: N.D. Giadinis,
Clinic of Farm Animals, Faculty of Veterinary Medicine,
Aristotle University of Thessaloniki, 11 St. Voutyra str., 54627, Thessaloniki, Greece.
E-mail: ngiadini@vet.auth.gr

Αλληλογραφία: Ν.Δ. Γιαδίνης,
Κλινική Παραγωγικών Ζώων, Τμήμα Κτηνιατρικής
Α.Π.Θ., Στ. Βουτυρά 11, 54627, Θεσσαλονίκη.
E-mail: ngiadini@vet.auth.gr

Date of initial submission: 10 February 2014
Date of acceptance: 21 February 2014

Ημερομηνία αρχικής υποβολής: 10 Φεβρουαρίου 2014
Ημερομηνία αποδοχής: 21 Φεβρουαρίου 2014

INTRODUCTION

Small Ruminant Lentiviruses (SRLVs) can cause a significant chronic disease with detrimental consequences for animal health and economic viability of sheep flocks, while hosts are infected for life. In adult sheep, the disease shows various clinical manifestations, e.g., chronic interstitial pneumonia (Maedi) (Pritchard and McConnell, 2007), chronic encephalomyelitis (Visna) (Glaria et al., 2012), chronic arthritis (Glaria et al., 2009) and chronic indurative mastitis (hard udder) (Giadinis et al., 2012). Also, reproductive disorders may possibly be caused by the same virus group (Pérez et al., 2010). However, often the infection remains asymptomatic for years and infected sheep can transmit the infection to other animals (Álvarez et al., 2006).

SRLVs can be transmitted by colostrum and milk consumption, by direct contact through respiratory secretions, while trans-placental infection also seems possible (Pritchard and McConnell, 2007). A factor that significantly contributed to the spread of the disease in many flocks seems to be the extensive uncontrolled trading of animals with unknown SRLV status (Peterhans et al., 2004). In Greece, national legislation does not require testing for SRLV. Hence, infection rate remains increased (Eltahir et al., 2006).

The present study describes the trans-boundary introduction of SRLVs from Spain to Greece and aims to alert veterinarians and farmers for the possibility of introduction of this disease from other countries.

MATERIALS AND METHODS

The study was conducted from February 2011 to September 2012 in five sheep farms (F_1 - F_5), located in various regions of Greece. Farm 1 was located in Epirus, farm 2 in Macedonia and farms 3-5 in Peloponnesus. All farms had purchased Assaf-E sheep directly from Spain, with no intermediate contact with other animals. On the day of arrival of the imported sheep at the farm, a number of animals was selected at random for blood sampling for subsequent serological testing for SRLV infection. In total,

57 sheep were selected for blood sampling: 18 in F_1 , 14 in F_2 , 5 in F_3 , 10 in F_4 and 10 in F_5 .

Blood samples were then transferred to the laboratory; serum was obtained by centrifugation of blood at 2500 g for 15 min. and stored at -20 °C until examination by using a commercially available ELISA test, specifically the CHEKIT-CAEV/MVV ELISA kit (IDEXX, Switzerland). Descriptive statistics were employed to present the results.

RESULTS

Seropositive animals were found in all examined flocks; proportion of flocks with seropositive animals was 1.0000 (95% confidence intervals: 0.4629-1.0000). Prevalence of SRLVs seropositive animals was 0.4386 (25/57) (95% confidence intervals: 0.3098-0.5757). Within farm seroprevalence varied from 20% to 50% (9/18 in F_1 , 7/14 in F_2 , 2/5 in F_3 , 5/10 in F_4 , 2/10 in F_5 ; 95% confidence intervals: 0.0354-0.7986).

DISCUSSION

The national flock of sheep in Greece includes approx. 9 million animals. The country has a strong tradition in dairy sheep farming, with Chios-breed sheep being the most representative high producing indigenous breed (Gelasakis et al., 2013). In recent years, intensively or semi-intensively reared flocks are fast growing, with their farmers choosing to import foreign breeds, as most Greek indigenous breeds have been neglected from genetic improvement programs and have low productivities. Popular imported sheep breeds are Lacaune, East Friesian, Awassi and, in recent times, Assaf-E. These animals are becoming increasingly popular, although no consideration is given to their health status.

There is strong evidence in the available literature that Spain has a high SRLV prevalence in sheep flocks. In recent studies, 77 % of intensively managed sheep flocks in Northern Spain were found seropositive and in over 50 % of Assaf-E flocks the seropositivity being over 60 % (Leginakicoica et al., 2006; Polledo et al., 2013). Those findings pro-

vided an interest to examine a sample of imported Assaf-E animals.

Our results are in accord to those in the literature. The findings do not provide evidence that the sheep will develop clinical disease, but highlight its potential. There is possibly an increased susceptibility of the Assaf-E breed for seroconversion of infected animals. In the past, previous studies have also suggested increased susceptibility of some breeds to SRLVs (Sarafidou et al., 2013).

Taking into account, that eradication of SRLV infections is difficult (Peterhans et al., 2004; Pritchard and McConnell 2007), we suggest that national legislation regarding sheep imports into Greece would be revised. Further, we recommend that, in the interim, sheep farmers considering to purchase Assaf-E or

other sheep breeds request an examination of sheep for possible SRLV infection. Attending veterinarians should guide flock owners for performing specific examinations of sheep that are brought from other flocks.

CONCLUDING REMARKS

Everybody concerned should be alerted about the possibility of SRLVs introduction into Greek flocks with the import of sheep from other countries, as it has been documented with Assaf-E sheep bought from Spain, in which increased seroprevalence of SRLVs has been detected. ■

REFERENCES

- Álvarez V, Daltabuit-Test M, Arranz J, Leginagoikoa I, Juste RA, Amorena B, de Andrés D, Luján LL, Badiola JJ, Berriatua E (2006) PCR detection of colostrums-associated Maedi-Visna virus (MVV) infection and relationship with ELISA-antibody status in lambs. *Res Vet Sci* 80:226-234.
- Eltahir YM, Dovas CI, Papanastassopoulou M, Koumbati M, Giadinis N, Verghese-Nikolakaki S, Koptopoulos G (2006) Development of a semi-nested PCR using degenerate primers for the generic detection of small ruminant lentivirus proviral DNA. *J Virol Meth* 135:240-246.
- Gelasakis AI, Oikonomou G, Bicalho RC, Valergakis GE, Fthenakis GC, Arsenos G (2013) Clinical characteristics of lameness and potential risk factors in intensive and semi-intensive dairy sheep flocks in Greece. *J Hell Vet Med Soc* 64:123-130.
- Giadinis ND, Arsenos G, Tsakos P, Psychas V, Dovas CI, Papadopoulos E, Karatzias H, Fthenakis GC (2012) "Milk-drop syndrome of ewes": Investigation of the causes in dairy sheep in Greece. *Small Rumin Res* 106:33-35.
- Glaria I, Reina R, Crespo H, de Andrés X, Ramírez H, Biescas E, Pérez MM, Badiola J, Luján L, Amorena B, de Andrés D (2009) Phylogenetic analysis of SRLV sequences from an arthritic sheep outbreak demonstrates the introduction of CAEV-like viruses among Spanish sheep. *Vet Microbiol* 138:156-162.
- Glaria I, Reina R, Ramírez H, de Andrés X, Crespo H, Jauregui P, Salazar E, Luján L, Pérez MM, Benavides J, Pérez V, Polledo L, García-Marín JF, Riezu JI, Borrás F, Amorena B, de Andrés D (2012) Visna/Maedi virus genetic characterization and serological diagnosis of infection in sheep from a neurological outbreak. *Vet Microbiol* 155:137-146.
- Leginagoikoa I, Daltabuit-Test M, Álvarez V, Arranz J, Juste RA, Amorena B, de Andrés D, Luján LL, Badiola JJ, Berriatua E (2006) Horizontal Maedi-Visna Virus (MVV) infection in adult dairy-sheep raised under varying MVV-infection pressures investigated by ELISA and PCR. *Res Vet Sci* 80:235-241.
- Pérez M, Biescas E, de Andrés X, Leginagoikoa I, Salazar E, Berriatua E, Reina R, Bolea R, de Andrés D, Juste RA, Cancr J, Gracia J, Amorena B, Badiola JJ, Luján L (2010) Visna/Maedi virus serology in sheep: Survey, risk factors and implementation of a stressful control programme in Aragón (Spain). *Vet J* 186:221-225.
- Peterhans E, Greenland T, Badiola J, Harkiss G, Bertoni G, Amorena B, Eliasiewicz M, Juste RA, Krassnig R, Lafont JP, Lenihan P, Pétursson G, Pritchard G, Thorley J, Vitu C, Mornex JF, Pépin M (2004) Routes of transmission and consequences of small ruminant lentiviruses (SRLVs) infection and eradication schemes. *Vet Res* 35:257-274.
- Polledo L, González J, Fernández C, Miguélez J, Martínez- Fernández B, Morales S, Ferreras MC, García-Marín JF (2013) Simple control strategy to reduce the Maedi-Visna infection in sheep flocks with high prevalence values (>90 %). *Small Rumin Res* 112:224-229.
- Pritchard GC, McConnell I (2007) Maedi-Visna. In: (ed.: Aitken ID), *Diseases of Sheep*, 4th ed., Blackwell, Oxford.
- Sarafidou T, Stamatis C, Kalozoumi G, Spyrou V, Fthenakis GC, Billinis C, Mamuris Z (2013) Toll like receptor 9 (TLR9) polymorphism G520R in sheep is associated with seropositivity for Small Ruminant Lentivirus. *Plos One* 8:e63901