



Journal of the Hellenic Veterinary Medical Society

Vol 69, No 1 (2018)



To cite this article:

SYMEONIDOU, I., BONOS, E., MOUSTAKIDIS, K., FLOROU-PANERI, P., CHRISTAKI, E., & PAPAZAHARIADOU, M. (2018). Botanicals: a natural approach to control ascaridiosis in poultry. *Journal of the Hellenic Veterinary Medical Society*, *69*(1), 711–722. https://doi.org/10.12681/jhvms.16383

Botanicals: a natural approach to control ascaridiosis in poultry

Symeonidou I.^{1*}, Bonos E.², Moustakidis K.¹, Florou-Paneri P.³, Christaki E.³, Papazahariadou M.¹

¹ Laboratory of Parasitology and Parasitic Diseases, School of Veterinary Medicine, Faculty of Health Sciences,

Aristotle University of Thessaloniki, Greece.

² Research Institute of Animal Science / HAO-DEMETER, Paralimni Giannitsa, Greece.

³ Laboratory of Nutrition, School of Veterinary Medicine, Faculty of Health Sciences,

Aristotle University of Thessaloniki, Greece.

Φυτοβιοτικά: Νέα φυσική προσέγγιση αντιμετώπισης της ασκαριδίωσης των πτηνών

Συμεωνίδου Η.^{1*}, Μπόνος Ε.², Μουστακίδης Κ.¹, Φλώρου-Πανέρη Π.³, Χρηστάκη Ε.³, Παπαζαχαριάδου Μ.¹

ABSTRACT. Parasites (protozoa, helminthes, arthropods) represent a main threat for poultry worldwide. Among helminthes, nematodes constitute the most important group of parasites of poultry. The nematode *Ascaridia galli*, the cause of ascaridiosis in poultry, is one of the most important and prevalent parasites, resulting in serious economic losses, associated with the treatment cost, the decreased feed efficiency, and the poor egg and meat production. During the past few decades the indiscriminate use of anthelmintic drugs has generated several cases of resistance in helminthes in poultry, situation which is coupled with the severity of residues in poultry products. For this reason, nowadays attention has been drawn to the use of botanicals in poultry diet, due to their anthelmintic properties. Furthermore, the dietary use eco-friend-

Corresponding Author: isaia@vet.auth.gr Laboratory of Parasitology and Parasitic Diseases, School of Veterinary Medicine, Faculty of Health Sciences, Aristotle University of Thessaloniki, 54124, Thessaloniki, Greece

Date of initial submission: 29-4-2017 Date of acceptance: 21-5-2017 ly of these plant derived substances compared to conventional synthetic anthelmintic drugs is considered as a natural and ecofriendly approach by the consumers. The focus of the present review is to recapitulate the studies, both *in vivo* and *in vitro*, that have demonstrated the anthelmintic efficacy of various dietary botanicals in controlling poultry ascaridiosis.

Keywords: Helminthes, Ascaridia galli, Botanicals, Poultry.

ΠΕΡΙΛΗΨΗ. Τα παράσιτα (πρωτόζωα, έλμινθες, αρθρόποδα) αποτελούν σημαντική απειλή για τα πτηνά παγκοσμίως. Μεταξύ των ελμίνθων, τα νηματώδη συνιστούν τη σπουδαιότερη ομάδα παρασίτων των πτηνών. Το νηματώδες Ascaridia galli, αίτιο της ασκαριδίωσης των πτηνών, είναι ένα από τα πλέον σημαντικά και συχνά παράσιτα, προκαλώντας σοβαρές οικονομικές απώλειες, που συνδέονται με το κόστος θεραπείας, τη μειωμένη αξιοποίηση της τροφής και τη μείωση της παραγωγής αυγών και κρέατος. Κατά τη διάρκεια των τελευταίων δεκαετιών, η αλόγιστη χρήση ανθελμινθικών φαρμάκων οδήγησε στην ανάπτυξη ανθεκτικότητας σε κάποια ανθελμινθικά που χρησιμοποιήθηκαν σε πτηνά, γεγονός που σχετίζεται με την παρουσία καταλοίπων στα προϊόντα των πτηνών. Για το λόγο αυτό, τα τελευταία χρόνια το ενδιαφέρον των ερευνητών έχει στραφεί στη χρήση φυτοβιοτικών στη διατροφή των πτηνών εξαιτίας των ανθελμινθικώ ιδιοτήτων τους. Επιπλέον, η χρήση αυτοβιοτικών προελευσης ουσιών σε σύγκριση με τα συμβατικά συνθετικά ανθελμινθικά φάρμακα θεωρείται ως μια φυσική και φιλική προς το περιβάλλον προσέγγιση από τους καταναλωτές. Σκοπός της παρούσας ανασκόπησης είναι η παράθεση των ερευνών, τόσο *in vivo* όσο και *in vitro*, που παρουσιάζουν την ανθελμινθική αποτελεσματικότητα διαφόρων φυτοβιοτικών στον έλεγχο της ασκαριδίωσης των πτηνών.

Λέξεις ευρετηρίασης: Έλμινθες, Ascaridia galli, Φυτοβιοτικά, Πτηνά.

INTRODUCTION

Darasitism with protozoa, helminthes and arthropods remains a main threat for poultry worldwide inducing heavy production losses in animals (Ahmad et al., 2013). Between helminthes, like Roundworms (Nematodes), Tapeworms (Cestodes) and Flukes (Trematodes), nematodes are the most important ones (Rafi et al., 2011). Ascaridia galli (A. galli), Heterakis gallinarum (H. gallinarum) and Capillaria spp. are the most common roundworms of poultry, with A. galli being the most prevalent (Kaufman et al., 2011). Researchers' interest in relation to alternative control measures has been focused mainly on A. galli. This parasite is responsible for clinical and subclinical parasitism. In heavily infected poultry the clinical signs include droopiness, diarrhea and hemorrhages. Also, during heavy infestation birds may show signs of decreased weight gain and retarded growth, due to damaged integrity of the intestinal mucosa and subsequent impaired nutrient utilization (Das et al., 2010). In more severe cases and especially in young birds, intestinal blockage may occur, leading to

death (Abdelqader et al., 2008). *A. galli* infections result in serious economic losses, usually associated with treatment cost, decreased feed efficiency and poor egg and meat production (Martin-Pacho et al., 2005). Another very debilitating factor resulting in economic losses is the ability of *A. galli* eggs to act as vectors for transmission of fatal bacterial infectious organisms, such as *Salmonella enterica* and *Escherichia coli* (Permin et al., 2006).

During the past few decades, novel research on the transmission of helminthes has enabled scientists to develop efficient products for their control. However, the indiscriminate use of antiparasitic drugs in poultry has generated cases of resistance to conventional drugs, especially piperazine and benzimidazoles, such as fenbendazole and albendazole (Abdelqader et al., 2012; Yazwinski et al., 2013). This situation, coupled with the severity of the associated risks of chemical residues in poultry products and the high cost of treatment compliance in endemic regions, necessitates further efforts into the discovery of novel drugs from either natural or synthetic sources (Anthony et al., 2005).

For sustainable control of *A. galli* different approaches have been employed such as, nutrition of host animal (Das et al., 2010), utilization of genetic resistance (Kaufman et al., 2011), biological control (Braga et al., 2011), and the use of plants with promising anthelmintic activity (Anthony et al., 2005).

Attention has been drawn to the screening of botanicals for their anthelmintic properties (Anthony et al., 2005). Accordingly, the knowledge of traditional herbal remedies is scientifically examined in order to find new plants having potent broad spectrum anthelmintic activity with less toxicity (Mali and Mehta, 2008). Compared to conventional synthetic anthelmintic drugs, plant derived products are considered natural and eco-friendly. Moreover, many such products are certified as GRAS (Generally Recognized As Safe) by the FDA (Food and Drug Administration) and therefore could be ideal candidates as feed additives (Brenes and Roura, 2010; Christaki et al., 2012). The use of phytogenic bioactive compounds for poultry nematode control is increasing in different commercial production systems and it has a reduced impact on environment (George et al., 2009). Furthermore, the active components of plants are compounds with great structural diversity and low molecular weight. These components can be active against many biological processes of the parasites and this diversity can preclude the development of anthelmintic resistance (Tariq et al., 2009).

Aim of this review is to summarize the current knowledge regarding the use of plant derived substances to control *A. galli* parasitism in poultry. Under this effort their anthelmintic activities and various proposed modes of action are discussed.

BOTANICALS

Recently, strong research interest is focused on botanicals (or phytogenics or phytobiotics). Botanicals are made from plants, algae, fungi or lichens (European Food Safety Authority, 2009b). Currently, they are under examination for their various bioactive activities: improving feed intake and flavour; stimulating the secretion of digestive enzymes; increasing gastric and intestinal motility; endocrine stimulation; anticoccidial and other antiparasitic activities; antimicrobial, anti-viral, immune-stimulating, anti-inflammatory and antioxidative activity. The main active substances of botanicals are considered to be the plant secondary metabolites. Botanicals vary greatly due to the enormous variety of different plants used, the different methods used in their preparation, and their formulations (Christaki et al., 2012; Bozkurt et al., 2013).

Traditionally, the easiest way to prepare botanicals is to separate the plant part that contains the active substances (seeds, leaves, bark, etc) and then dry and grind it to powder (Christaki et al., 2012; Bozkurt et al., 2013). Nevertheless, the research focus is now on the separation and identification of their active ingredients, using gas chromatography and mass spectrometry (Brenes and Roura, 2010).

Some of the main botanical extracts are essential oils and oleoresins. Essential oils (volatile oils; ethereal oils; aetherolea) are aromatic oily liquids that originate from plants. Essential oils can be complex mixtures of many secondary plant metabolites, mainly low boiling terpenes (examples: linalool, geraniol, borneol, menthol, thujanol, citronnillol, α -terpineol), phenols (examples: thymol, carvacrol, eugenol, gaiacol), aromatic aldehydes (examples: cinnamaldehyde, cuminal and phellandral), and their alcohol, aldehyde or ester derivatives. (Brenes and Roura, 2010; Christaki et al., 2012).

Oleoresins are naturally occurring mixtures of essential oils and resins. They can be obtained from plants by extraction with a nonaqueous solvent (alcohol, ether, or acetone), followed by the removal of the solvent through evaporation. They contain volatile and non-volatile plant constituents (McCloud, 2010).

Botanicals can be produced from single plants or they can be mixtures from different plants. The idea of using mixtures is to exploit possible synergistic effects, so as to maximize the bioactive effects of different secondary metabolites and to minimize the concentrations required to achieve a particular effect (Kirkpinar et al., 2011).

The solvent of the extract also plays an important role in the activity of the botanical. Different solvents have variable physical properties such as polarity, which can affect the solubility and the activity of the plant metabolites, when the extract is ingested by the parasite or comes in contact with its surface (transcuticular absorption), especially under *in vitro* experiments (Ahmad et al., 2013; Kaingu et al., 2013).

ACTIVE SUBSTANCES OF BOTANICALS WITH ANTHELMINTIC ACTIVITIES

As already mentioned, the active substances of botanicals are plant metabolites that are synthesized by plants throughout their life cycle. They are distinguished in primary and secondary metabolites (Hrcova and Velebny, 2013; Marin et al., 2015).

Plant primary metabolites such as carbohydrates, lipids, proteins and nucleic acids, are the main compounds of basic metabolic pathways and also precursors for the synthesis of the plant secondary metabolites (Hrcova and Velebny, 2013; Marin et al., 2015).

Plant secondary metabolites are organic compounds synthesized by plants that have important functions for the plant, mediating interaction with other plants or organisms; for example protection against microbial or insect attack or attraction of pollinators and seed-dispersing. Plant secondary metabolites are often colored, fragrant or flavorful compounds. Based on their biosynthetic origin and chemical structure they be divided into three wide groups: terpenes (or terpenoids), phenolics (or phenols or phenolic compounds) and nitrogen-containing compounds (Hrcova and Velebny, 2013).

MODES OF ACTION OF BOTANICALS AGAINST HELMINTHES

Various modes of action have been suggested for the botanicals although it is possible that other mechanisms are not sufficiently identified yet.

1. Antelminthic activity of plant secondary metabolites

An overview of plant secondary metabolites effects against helminthes is presented in Figure 1.

1.1. Starvation

Some monoterpenes, like ascaridole (from plant

Fig 1. Plant secondary metabolites and their modes of action against helminthes



J HELLENIC VET MED SOC 2018, 69(1) ΠΕΚΕ 2018, 69(1) *Chenopodium ambrosioides*), can disrupt the tubulin polymerization in the intestinal cells of the parasites, which leads to their degeneration and death (Wink, 2012; Jain et al., 2013). Ascaridole is effective against helminthes and has been used since the 1900s, but today its use has been limited, as it is considered mutagenic and poisonous. Also, some plant secondary metabolites separately or jointly can block glucose uptake by the parasite (Jain et al., 2013).

Tannins have the capacity to bind free protein in the digestive tract, thus limiting the nutrient availability and possibly resulting in larval starvation (Jain et al., 2013). Another possible action of tannins is the decrease in gastrointestinal metabolism directly through inhibition of oxidative phosphorylation, again leading to larval death (Athanasiadou et al., 2001; Kateregga et al., 2014). Tannins and flavonoids could inhibit energy production in the parasite cells by blocking phosphorylation reactions (Sharma and Prasad, 2014). Tannins may also bind to free protein in the gastrointestinal tract, as well as to proteins of the cuticle of parasite, thus limiting the ability to absorb nutrients. Condensed tannin ingested by the larvae of nematodes can bind to their intestinal mucosa and cause autolysis (Schultz, 1989; Athanasiadou et al., 2001).

Steroidal alkaloids and oligoglycosides can limit the amount of sugars, such as sucrose that reach the small intestine from the stomach, as well as affect the generation of nitrates (Borba et al., 2010). These effects modify the local conditions in the intestine, making them unfavorable for the development of intestinal parasites (Borba et al., 2010).

Likewise, saponins can restrict feed intake, limiting the available nutrients for the helminthes and possibly causing their death (Kateregga et al., 2014).

Also, it has been hypothesized that some plant extracts induce an inflammatory response in the gastric and intestinal mucosal of the host, which disrupts the local homeostasis that is necessary for the development of the helminthes (Borba et al., 2010). In addition, it has been suggested that some plant secondary metabolites play an important role in the regulation of the carbohydrate metabolism (Bazh and El-Bahy, 2013). In this case, it is speculated that the branch point of phosphoenolpyruvate carboxykinase/ pyruvate kinase forms the basis of the anthelmintic attack by the plant derived components (Bazh and El-Bahy, 2013).

1.2. Damage to the helminth cuticle

Any damage to the mucopolysaccharide membrane of the parasite results in movement restriction and possibly in paralysis (Chandrashekhar et al., 2008; Jain et al., 2013). The binding effect of tannins on the cuticle results in the loss of its flexibility due to hydrogen bonding. This type of reactivity may lead to increased toughness of the cuticle and hence the helminthes become immobile and nonfunctional. Larvae are then unable to burrow into the mucosal lining of the small intestine of the host, and are subsequently eliminated from the host (Salhan et al., 2011; Jain et al., 2013).

1.3. Effects on mobility

Alkaloids may act on central nervous system of the parasite and cause paralysis of helminthes (Roy et al., 2010; Jain et al., 2013). Alkaloids pelletierine from *Punica granatum* (Lythraceae) and arecoline from *Areca catechu* (Arecaceae), target acetylcholine receptors as competitive antagonists and can cause paralysis of the helminthes (Wink, 2012). The anthelmintic activity of alkaloids has been demonstrated in two rat nematodes; *Strongyloides ratti* and *Strongyloides venezuelensis* (Kateregga et al., 2014). Paralyzed parasites in the digestive tract are unable to remain adhered on the intestinal wall of the host and are removed through peristalsis. It is also possible that these parasites starve to death (Wink, 2012).

Some terpenes, such as thymol and carvacrol, are neurotoxic to nematodes and interact with Ser-2, a *Caenorhabditis elegans* tyramine receptor (Lei et al., 2010; Kaplan et al., 2014). In nematodes, tyramine is believed to play a role in foraging behavior and pharyngeal pumping (Rex et al., 2004). Tyramine is a nonpeptidic hormone that can only be found in invertebrates and thus it is considered an important substance in antiparasitic research (Klowden, 2007).

Phloroglucinols, such as aspidin, deaspidin, and filixic acid, which can be found in *Dryopteris filix-mas* (*Dryopteridaceae*) can paralyze helminthes, and this effect is more pronounced on cestodes (Murthy et al., 2011; Wink, 2012).

1.4. Impact on growth and reproduction

It is possible that, tannins directly or indirectly reduce the number of hatching eggs, the rate of larvae development and the number of eggs produced from adult parasites (Athanasiadou et al., 2001; Van Krimpen et al., 2010), via not sufficiently identified mechanisms. It has been proposed that substances with hormonal effect, such as triterpenes, disrupt the reproductive cycle of the parasite. These effects have been investigated for some plants, for example for the genus *Artemisia*, against the nematodes *Ascaris suum* of pigs, as well as *Toxocara* spp. of carnivores and the cestodes *Moniezia* spp. of ruminants (Githiori, 2004; Van Krimpen et al., 2010; Acton, 2012).

It is possible that the secondary metabolites affect helminthes variously depending on different stages of their development. Also, a main factor that contributes to the anthelmintic activity of the secondary metabolites is the conditions in the digestive tract. For example, the formation and disassociation of complexes between proteins and tannins is greatly affected by the *pH*. Complexes between condensed tannins and protein remain stable in *pH* between 5 and 7, but they disassociate in *pH* higher or lower than the above (Athanasiadou et al., 2001). Also, the presence of surfactants, such as bile acids, has been reported to be important for the disassociation of tannin-protein complexes.

Further research is required in order to identify additional modes of action of botanicals against helminthes of poultry, as well as the possible interactions between major and minor bioactive components.

2. Additional beneficial activities of plant secondary metabolites

Botanicals can have additional effects that benefit poultry health and performance such as antimicrobial activity (Christaki et al., 2012), immunomodulatory effects (Anthony et al., 2005), antioxidant activity (Christaki et al., 2012), anti-inflammatory properties (Borba et al., 2010) and appetite and digestion enhancing effects (Borba et al., 2010).

IN VIVO AND IN VITRO STUDIES

Several up-to-date studies have demonstrated the anthelmintic efficacy of different botanicals and most

of them evaluate the effect of plants against A. galli in chicken (Gallus gallus domesticus). It has been reported that botanicals exert similar anthelmintic activity both in vivo and in vitro, usually in a concentration and time-dependent manner (Alawa et al., 2003; Adedapo et al., 2007). In a number of studies, the anthelmintic effects of botanicals against A. galli in poultry were compared to conventional antiparasitic drugs, such as albendazole, levamisole, piperazine, etc, suggesting that botanicals can partially or totally substitute those reference drugs (Akhtar and Riffat 1985, Al-Harbi 2011, Bazh and El-Bahy 2013). It is notable that in most cases the efficacy of botanicals was adequate, although not up to par, with the anthelmintic drugs. Since testing biological activity under in vivo conditions has several difficulties, such as the inherent features of animals and self-cure phenomenon, most of the studies refer to in vitro screenings of the anthelmintic efficacy of different botanicals (Sandoval-Castro et al., 2012).

1. In vivo studies

Under *in vivo* studies when botanicals were supplemented, either in the water or in the feed, they depressed the fecal egg count of *Ascaridia galli* as well as reduced the adult worm burden in parasitized poultry (Table 1).

In a study with Lohmann Leghorn chicks a mixture of ethanol extracts from orange (Citrus x sinensis), lemon (Citrus x lemon), and mandarin (Citrus reticulata), was added in the feed (at 300, 600 or 1200 mg/ kg of b.w.) and a significant dose dependent reduction in fecal egg output and parasitic worm burden was recorded (Abdelgader et al., 2012). Moreover, Melia azedarach fruit (powder at 20 mg/kg of b.w. or equivalent amounts of water extract, methanol extract or ethanol extract) were found to inhibit A. galli egg development in chickens infected with the parasite (Akhtar and Riffat, 1985). Likewise, it was shown that A. galli challenged cockerels exhibited a dose dependent reduction in fecal egg count, when treated with graded doses of ethanolic extract (100, 200 and 400 mg / kg b.w.) from the bark of *Piliostigma* thonningii (Asuzu and Onu, 1994). In addition, it has been demonstrated that Caesalpinia crista, known as karanjwa, when administered to broilers as seed powder (at 30, 40, and 50 mg/kg of b.w. or as equivalent

				5	
Plant name	Plant part used	Examined form and dosage	Source of <i>A</i> . <i>galli</i> worms	Effect against <i>A. galli</i>	References
Citrus x sinensis Citrus x lemon Citrus reticulata	Peels Peels Peels	Dietary addition of mix- tures at 300, 600 and 1200 mg/kg of body weight	Chickens	Fecal egg reduction; Worm motility inhibition	(Abdelqader et al., 2012)
Melia azedarach	Fruit	Fruit powder at 20 mg/kg; Equivalent water extract, methanol extract, ethanol extract	Chickens	Egg development inhibi- tion	(Akhtar and Riffat, 1985)
Piliostigma thonningii	Bark	Ethanolic extracts at 100, 200 and 400 mg / kg body weight	Chickens	Fecal egg reduction	(Asuzu and Onu, 1994)
Caesalpinia crista	Seeds	Powder and methanolic extracts at 30, 40 and 50 mg/kg body weight	Chickens	Fecal egg reduction	(Javed et al., 1994)
Tephrosia vogelli Vernonia amygdalina	Leaves Leaves	Water extracts. Doses not mentioned	Chickens	Fecal egg reduction; Reduction of adult worms population	(Siamba et al., 2007)
Punica granatum	Peel	Dry peel at 0.5, 1.0 and 1.5 g/kg of body weight	Laying hens	Fecal egg reduction Increased packed cell vol- ume, total serum proteins, body weight	(Sabri, 2013)
Azadirachta indica	Leaves	Aqueous extract at 200 mg / kg of body weight	Chickens	Increased bird body weight; Parasite death	(Khokon et al., 2014)

Table 1	1. In vivo	anthelmintic	activities	of various	botanicals	against.	Ascaridia galli	
							····· ··· · · · · · · · · · · · · · ·	

amounts of water and methanol extracts) reduced A. galli egg numbers in chicken faeces, analogously to piperazine (Javed et al., 1994). Also, the extracts of Tephrosia vogelli and Vernonia amygdalina (doses not mentioned) not only significantly depressed fecal egg output, but also reduced the adult worm population in A. galli parasitized poultry (Siamba et al., 2007). Another study reported that Punica granatum dry peel, orally administered to infected hens at 0.5, 1.0 and 1.5 g/kg of b.w., reduced fecal egg count, analogously to levamisole, while increasing hematocrit (packed cell volume, PCV), total serum proteins and body weight (Sabri, 2013). Furthermore, neem (Azadirachta indica) leaves extract was supplemented to chickens at 200 mg/kg of b.w., leading to a significant increase in body weight (Khokon et al., 2014).

2. In vitro studies

Researchers have evaluated different plant extracts under *in vitro* conditions against *A. galli* collected from freshly slaughtered poultry. The commonly observed anthelmintic effects included paralysis and death, as well as inhibition of egg and larvae development (Table 2).

Bazh and El-Bahy (2013) revealed that when living worms were incubated at 37 °C in media containing ginger (*Zingiber officinale*) and curcumin (*Curcuma longa* L) methanolic extracts at three concentration levels (25, 50 and 100 mg/ml), their physical activity (spontaneous movement) as well as their survival was reduced in a concentration and time dependent manner. Likewise, Lal et al. (1976) demonstrated that extracts from Carica papaya seeds (alcohol extract at 25 mg/ml), Sapindus trifoliatum fruit pericarp (alcohol extract at 10 mg/ml), Butea frondosa seeds (alcohol extract at 200 mg/ml) and Momordica charantia fresh juice (alcohol extract at 100 mg/ ml) caused paralysis and death of A. galli procured from fowls. Additionally, crude alcohol and aqueous extracts of seeds of Cleome viscosa exhibited considerable dose-dependent antiparasitic results (10, 50, 100 mg/ml) against A. galli worms (Mali et al., 2007). Kosalge and Fursule (2009) recorded the paralysis and death of A. galli and Raillietina spp. when administering the aqueous extract of Thespesia lampas roots at concentrations of 10, 20 and 50 mg/ ml, and thus proposed that this extract can be used effectively as an anthelmintic. Another similar study showed that the methanolic extract (10, 25, 50 mg/ ml) of Cymbopogon citratus leaves displayed better antihelmintic efficacy against A. galli in terms of paralysis and death, than the aqueous extract (10, 25, 50 mg/ml) of the same plant (Gore et al., 2010). Al-Harbi (2011) comparing aqueous solution of dried Artemisia abssinthium leaves and powder suspension of dried Lepidium sativium seeds with levamisole against A. galli, found that all of them have the same effectiveness against this parasite, causing paralysis and subsequently death after a period of exposure.

Kundu et al. (2012) revealed the broad wormicidal (paralysis and death) in vitro activity of Cassia alata, Cassia angustifolia and Cassia occidentalis (crude ethanol extracts at 10, 20 and 40 mg/ml) against various parasites (trematode Catatropis spp., cestode Raillietina tetragona and nematode H. gallinarum) collected from domestic fowl. The observed effects could be attributed to the large amount of alkaloids, flavonoids, glycosides, tannins that these plants are known to contain (Hossain et al., 2012). Another study revealed similar effects for the ethanolic extract of the leaves of Eupatorium triplinerve (50 and 100 mg/ml) and of the rhisome of Alpinia galangal (100 mg/ml) against A. galli. The activity of these plants was comparable to albendazole and can be attributed for A. galangal to the many flavonoids that it contains, such as kaempferide, kaemperol, galangin and alpinin, whereas for the crude extracts of E. triplinerve to the presence of phenolic compounds and coumarins (Charles et al., 1992;

Subash et al., 2012). Likewise, it was demonstrated that various concentrations of crude hydroalcoholic extracts (25 and 50 mg/ml) and aqueous extract (50 mg/ml) from *Mentha longifolia* leaves resulted in concentration and time dependent paralysis and death of *A. galli* (Ahmad et al., 2013). Also, aqueous (25, 50, 100 mg/ml) and ethanol extracts (10, 25, 50 mg/ml) of *Azadirachta indica* leaves, *Carica papaya* seeds and *Momordica charantia* bark, caused cessation of motility and increased mortality of *A. galli* (Shah Alam et al., 2014), comparable to the effects of levamisole and piperazine. Moreover, it has been reported that the extracts of *Tephrosia vogelli* and *Vernonia amygdalina* (doses not mentioned) can inhibit larvae mobility (Siamba et al., 2007).

In another study, A. galli treated with different concentrations (5, 10, 20 mg/ml) of Acacia oxyphylla methanol extract demonstrated extensive structural alterations, such as rupture of the ovaries and deformity on the egg membranes, detachment of the cuticle, disintegration of the muscular layers of the nematode and subsequent death (Lalchhandama, 2008). The wormicidal in vitro potential of leaves and fruits of aqueous extracts of Sesbania grandiflora and Solanum torvum (10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, and 130 mg/ 100 ml) was proven against A. galli from laying hens (Jothi Karumari et al., 2014). Furthermore, Kateregga et al. (2014) revealed that the use of Cassia occidentalis methanolic leaf extract (8, 12, 16, 20 and 24 mg/ ml) caused the death of H. gallinarum and A. galli. The main phytochemicals of C. occidentalis leaves are saponins, flavonoids, terpenes, sterols, alkaloids and tannins (Muyibi et al., 2000). Likewise, another study demonstrated that a mixture of ethanol extracts from orange (Citrus x sinensis), lemon (Citrus x lemon), and mandarin (Citrus reticulata) at 50 mg/ml has potential wormicidal properties against A. galli in vitro (Abdelqader et al., 2012). Similarly, neem (Azadirachta indica) leaves aqueous extract, in a range of concentrations (1, 2, 4 and 20 mg/ml), presented sufficient anthelmintic effect, causing death of A.galli worms (Khokon et al., 2014). In addition, Kaushik et al. (1981) evaluated 11 plants extracts (Amomum aromaticum root and rhisome, Ammora wallichii stem, Anthocephalus indicus stem and bark, Calamintha umberosa plant, Dalbergia latifolia stem

		• • •		- ·
Plant name	Plant part used	Examined form and dosage	Effect against <i>A</i> . <i>galli</i>	References
Zingiber officinale Curcuma longa L.	Root Root	Methanolic extracts at 25, 50 and 100 mg/ml.	Paralysis; Death	(Bazh and El-Bahy, 2013)
Carica papaya, Sapindus trifoliatum Butea frondosa Momordica charantia	Seeds Fruit pericarp Seeds Fresh juice	Alcohol extract at 25 mg/ml Alcohol extract at 10 mg/ml Alcohol extract at 200 mg/ml Alcohol extract at 100 mg/ml	Paralysis; Death	(Lal et al., 1976)
Cleome viscosa	Seeds	Crude alcohol and aqueous extracts at 10, 50 and 100 mg/ml	Paralysis; Death	(Mali et al., 2007)
Thespesia lampas	Roots	Aqueous extracts at 10, 20 and 50 mg/ml.	Paralysis; Death	(Kosalge and Fursule, 2009)
Cymbopogon citrates	Leaves	Methanolic and aqueous extracts 10, 25 and 50 mg/ml	Paralysis; Death	(Gore et al., 2010)
Artemisia abssinthium Lepidium satirium	Leaves Seeds	Aqueous solution and powdered suspension. Dose not mentioned	Paralysis; Death	(Al-Harbi, 2011)
Cassia alata Cassia angustifolia Cassia occidentalis	Leaves	Crude ethanol extracts at 10, 20 and 40 mg/ml	Paralysis; Death	(Kundu et al., 2012)
Eupatorium triplinerve Alpinia galangal	Leaves Rhisome	Ethanolic extracts at 50 and 100 mg/ml	Paralysis; Death	(Subash et al., 2012)
Mentha longifolia	Leaves	Crude hydroalcoholic extracts at 25 and 50 mg/ml; Crude aqueous extract at 50 mg/ml	Paralysis; Death	(Ahmad et al., 2013)
Azadirachta indica Carica papaya Momordica charantia	Leaves Seeds Bark	Aqueous extracts at 25, 50 and 100 mg/ml; Ethanol extracts at 10, 25 and 50 mg/ml	Paralysis; Death	(Shah Alam et al., 2014)
Tephrosia vogelli Vernonia amygdalina	Leaves Leaves	Water extracts. Doses not mentioned	Paralysis	(Siamba et al., 2007)
Acacia oxyphylla	Bark	Ethanol extracts at 5, 10 and 20 mg/ml	Body structural alterations; Death	(Lalchhandama, 2008)
Sesbania grandiflora Solanium torrum	Leaves, fruit Leaves, fruit	Aqueous extracts at 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120 and 130 mg/ 100 ml	Death	(Jothi Karumari et al., 2014)
Cassia occidentalis	Leaves	Methanolic extracts at 8, 12, 16, 20 and 24 mg/ml	Death	(Kateregga et al., 2014)
Citrus x sinensis Citrus x lemon Citrus reticulata	Peels Peels Peels	Mixture of ethanol extracts of the three plants at 50 mg/ml	Death	(Abdelqader et al., 2012)
Azadirachta indica	Leaves	Aqueous extracts at 1, 2, 4 and 20 mg/ml.	Death	(Khokon et al., 2014)
Amomum aromaticum Ammora wallichii Anthocephalus indicus Calamintha umberosa, Dalbergia latifolia Datura quercifolia Datura metel Ficus religiosa Sentia myrtina Sumplocos crataegoides	Root, rhizome Stem, bark Plant Stem, bark Fruit Plant Stem, bark Plant Leaves	_	Death	(Kaushik et al., 1981)
Azadirachta indica Carica papaya Momordica charantia Polygonum hydropiper Swietenia macrophylla	Leaves Leaves Leaves Leaves Leaves	Fresh juice at 5%, 10% and 20%; Aqueous extract at 1%, 2% and 4%; Ethanol extract at 1%, 2% and 4%; Methanol extract at 1%, 2% and 4%; Powder at 10% and 20%	Inhibition of egg development	(Islam et al., 2008)
Aloe secundiflora	Leaves	Hexane, ethylacetate, acetone, methanol, and chlo- roform extracts at 5, 10, 20, 40 and 50 mg/ml	Inhibition of larvae development	(Kaingu et al., 2013)

Table 2. In vitro anthelmintic activity of various botanicals against Ascaridia galli from freshly slat	ughtered poultry.

J HELLENIC VET MED SOC 2018, 69(1) ПЕКЕ 2018, 69(1) and bark, *Datura quercifolia* fruit, *Datura metel* plant, *Ficus religiosa* stem and bark, *Sentia myrtina* plant, and *Sumplocos crataegoides* leaves) which were all proven lethal to *A. galli*.

Some in vitro studies reported other anthelmintic effects of botanicals against A. galli, such as inhibition of egg and larvae development. Particularly, fresh juice (at 5, 10 and 20%), aqueous extract (at 1, 2 and 4%), ethanol extract (at 1, 2 and 4%), methanol extract (at 1, 2 and 4%), and powder (at 10 and 20%) of leaves of Azadirachta indica, Carica papaya, Momordica charantia, Polygonum hydropiper, and Swietenia macrophylla inhibited the development of A. galli eggs, with C. papaya showing the highest efficacy 92% when the concentration was 4% (Islam et al., 2008). Another experiment using larvae development assays of A. galli revealed that different types of extracts (hexane; ethylacetate; acetone; methanol; chloroform) of Aloe secundiflora at 5, 10, 20, 40 and 50 mg/ml have inhibitory effects on the parasite's development (Kaingu et al., 2013).

CONSTRAINTS OF USING BOTANICALS AS ANTHELMINTIC AGENTS

One important problem with botanicals is the difficulty to characterize and standardize their ingredients and composition. Many factors can influence the chemical composition of the plants, such as species, subspecies, geographical location, harvesting and the collected part, such as seeds, leaf, root or bark (Christaki et al., 2012; Bozkurt et al., 2013). Also, the processing technique (cold expression, steam or alcohol distillation, extraction with non-aqueous solvents, etc.) can modify the active substances and associated compounds in the final product (Tariq et al., 2009; Windisch et al., 2009).

Another important consideration is that botanicals may also have adverse or toxic side effects for the treated animals. It has been demonstrated that plant substances, which interfere with parasite development, such as steroidal alkaloids, may also exhibit toxic effects on animal tissues. These effects include mutagenicity, embryotoxicity, hepatotoxicity, central nervous system symptoms, cardiac arrhythmia, etc (European Food Safety Authority, 2009a; Borba et al., 2010). Therefore, further studies are required to evaluate plant substances with possible detrimental effects for the animal, as well as to carefully quantify the optimal beneficial doses, versus the potential harmful ones (Wu et al., 2004; Bozkurt et al., 2013). Additionally, residue studies should be required before botanicals can be safely integrated in poultry management system.

CONCLUSION

The diversity of botanicals provides an important source of bioactive compounds, which may lead to potential new candidates remedies of natural origin against ascaridiosis of chickens. Keeping in view the economic importance of the parasitic infections in the development of profitable poultry industry, the anthelmintic properties of botanicals represent a very promising alternative solution to overcome current treatment inadequacies.

J HELLENIC VET MED SOC 2018, 69(1) ПЕКЕ 2018, 69(1)

REFERENCES

- Abdelqader A, Gauly M, Wollny CB, Abo-Shehdada MN (2008) Prevalence and burden of gastro-intestinal helminths among local chickens in northern Jordan. Prev Vet Med 85: 17-22.
- Abdelqader A, Qarallah B, Al-Ramamneh D, Das G (2012) Antelmintic effects of citrus peels ethanolic extracts against Ascaridia galli. Vet Parasitol 188: 78-84.
- Acton AQ (2012) Cestode infections Advances in research and treatment. Scholarly Editions, Atlanta, USA: pp. 8-9.
- Adedapo AA, Otesile AT, Soetan KO (2007) Assessment of the anthelmintic efficacy of an aqueous crude extract of Vernonia amygdalina. Pharm Biol 45: 564-568.
- Ahmad J, Tanveer S, Zargar BA (2013) In vitro anthelmintic activity of Mentha longfolia (L.) leaves against Ascaridia galli. Global Vet 11: 112-117.
- Akhtar MS, Riffat S (1985) Evaluation of Melia azedarach Linn. seeds (Bahain) and piperazine against Ascaridia galli infection in chickens. Pak Vet J 5: 34-37.
- Al-Harbi LG (2011) Comparative study in vitro on the efficacy of some plants extracts and Levamisole drug against Round worms of chickens natural infected. King Abdulaziz University, AL Faisaliah Campus, Saudi Arabia.
- Alawa CBI, Adamu AM, Gefu JO, Ajanusi OJ, Abdu PA, Chiezey NP, Alawa JN, Bowman DD (2003) In vitro screening of two Nigerian medicinal plants (Vermonia amygdalina and Annona senegalensis) for anthelmintic activity. Vet Parasitol 113: 73-81.
- Anthony JP, Fyfe L, Smith H (2005) Plant active components a resource for antiparasitic agents? Trends Biotechnol 21: 462-468.
- Asuzu IU, Onu UO (1994) Anthelmintic activity of the ethanolic extract of Piliostigma thonningii bark in Ascaridia galli infected chickens. Fitoterapia 65: 291-297.
- Athanasiadou S, Kyriazakis I, Jackson F, Coop RL (2001) Direct anthelmintic effects of condensed tannins towards different gastrointestinal nematodes of sheep: in vitro and in vivo studies. Vet Parasitol 99: 205-219.
- Bazh EKA, El-Bahy NM (2013) In vitro and in vivo screening of anthelmintic activity of ginger and curcumin on Ascaridia galli. Parasitol Res 112: 3679-3686.
- Borba HR, Freire RB, Albuquerque AC, Cardoso MEO, Braga IG, Almeida STP, Ferreira MJC, Fernandes GLT, camacho ACLF, Lima RC, Almeida ACC, Mattos DMM, Duarte RM, Nascimento SF, Frami RA, Dire GF (2010) Anthelmintic comparative study of Solanum lycocarpum St. Hill extracts in mice naturally infected with Aspiculuris tetraptera. Nat Sci 8: 95-100.
- Bozkurt M, Giannenas I, Kucukyilmaz K, Christaki E, Florou-Paneri P (2013) An update on approaches to controlling coccidia in poultry using botanical extracts. Br Poult Sci 54: 713-727.
- Braga FR, Araujo JV, Araujo JM, Frassy LN, Tavela AO, Soares FEF, Carvalho RO, Queiroz LM, Queiroz JH (2011) Pochonia chlamydosporia fungal activity in a solid medium and its crude extract against eggs of Ascaridia galli. J Helminthol 85: 1-5.
- Brenes A, Roura E (2010) Essential oils in poultry nutrition: Main effects and modes of action. Animal Feed Sci Technol 158: 1-14.
- Chandrashekhar CH, Latha KP, Vagdevi HM, Vaidya VP (2008) Anthelmintic activity of the crude extracts of Ficus racemosa. Int J Green Pharm 2: 100-103.
- Charles DJR, Simon JE, Singh NK (1992) Bio active principles/Chemical constituents: The essential oil of Alpinia galanga willd. J. Essential Oil Res 4: 81-82.
- Christaki E, Bonos E, Giannenas I, Florou-Paneri P (2012) Aromatic plants as a source of bioactive compounds. Agriculture 2: 228-243.
- Das G, Kaufmann F, Abel H, Gauly M (2010) Effect of extra dietary lysine

in Ascaridia galli infected grower layers. Vet Parasitol 170: 238-243.

- European Food Safety Authority (2009a) EFSA Compedium of botanicals that have been reported to contain toxic, addictive, phychotropic or other substances of concern. EFSA J 7: 289.
- European Food Safety Authority (2009b) Guidance on safety assessment of botanicals and botanical preparations intended for use as ingredients in food supplements. EFSA J 7: 1249.
- George DR, Smith TJ, Shiel RS, Sparagano OAE, Guy JH (2009) Mode of action and variability in efficacy of plant essential oils showing toxicity against the poultry red mite, Dermanyssus gallinae. Vet Parasitol 161: 276-282.
- Githiori JB (2004) Evaluation of anthelmintic properties of ethnoveterinary plant preparations used as livestock dewormers by pastoralists and small holder farmers in Kenya. Swedish University of Agricultural Science, Uppsala.
- Gore MS, Tare HL, Deore SR, Bidkar JS, Dama GY (2010) Hemintholytic potential of Cymbopogon citratus leaves extact and its formulation as an emulsion. Int J Pharmac Sci Res 1: 174-177.
- Hossain E, Chandra G, Nandy AP, Mandal SC, Gupta JK (2012) Anthelmintic effect of a methanol extract of Bombax malabaricum leaves on Paramphistomum explanatum. Parasitol Res 110: 1097-1102.
- Hrcova G, Velebny S (2013) Pharmacological potential of selected natural compounds in the control of parasitic diseases. Springer, NY.
- Islam KR, Farjana T, Begum N, Mondal MMH (2008) In vitro efficacy of some indigenous plants on the inhibition of development of eggs of Ascaridia galli (Digenia: Nematoda). Bangl J Vet Med 6: 159-167.
- Jain P, Singh S, Singh SK, Verma SK, Kharya MD, Solanki S (2013) Anthelmintic potential of herbal drugs. Int J Pharmac Sci Res 2: 412-427.
- Javed I, Akhtar MS, Rahman ZU, Khaliq T, Ahmad M (1994) Comparative anthelmintic efficacy and safety of Caesalpinia crista seed and piperazine adipate in chickens with artificially induced Ascaridia galli infection. Acta Vet Hung 42: 103-109.
- Jothi Karumari R, Sumathi S, Vijayalakshmi K, Ezhilarasi Balasubramanian S (2014) Anthelmintic efficacy of Sesbania grandiflora leaves and Solanum torvum fruits against the nematode parasite Ascaridia galli. Am J Ethnomed 1: 326-333.
- Kaingu F, Kibor A, Waihenya R, Shivairo R, Mungai L (2013) Efficacy of Aloe secundiflora crude extracts on Ascaridia galli in vitro. Sustain Agric Res 2: 49-53.
- Kaplan RM, Storey BE, Vidyashankar AN, Bissinger BW, Mitchell SM, Howell SB, Mason ME, Lee MD, Pedroso AA, Akashe A, Skrypec DJ (2014) Antiparasitic efficacy of a novel plant-based functional food using an Ascaris suum model in pigs. Acta Trop 139: 15-22.
- Kateregga JN, Nabayunga M, Vudriko P, Ndukui JG (2014) Anthelmintic activity of Cassia occidentalis L. methanolic leaf extract on Ascaridia galli and Heterakis gallinarum and its acute toxicity. Int J Basic Clin Pharmacol 3: 114-119.
- Kaufman F, Das G, Sohnrey B, Gauly M (2011) Helminth infections in laying hens kept in organic free range systems in Germany. Livestock Sci 141: 182-187.
- Kaushik RK, Katiyar JC, Sen AB (1981) A new in vitro screening technique for anthelmintic activity using Ascaridia galli as a test parasite. Indian J Animal Sci 51: 869-872.
- Khokon JU, Sharifuzzaman A, Sarker EH, Rahman MA, Kisku JJ, Mostofa M (2014) Efficacy of neem leaf extract against ascariasis in indigenous chicken. Int J Nat Soc Sci 1: 25-30.
- Kirkpinar F, Unlu HB, Ozdemir G (2011) Effects of oregano and garlic essential oils on performance, carcase, organ and blood characteristics and intestinal microflora of broilers. Livestock Sci 137: 219-225.

J HELLENIC VET MED SOC 2018, 69(1) ПЕКЕ 2018, 69(1)

- Klowden MJ (2007) Physiological systems in insects. Academic Press, Burlington.
- Kosalge SB, Fursule RA (2009) Investigation of in vitro anthelmintic activity of Thespesia lampas (Cav.). Asian J Pharmac Clin Res 2: 69-71.
- Kundu S, Roy S, Lyndem LM (2012) Broad spectrum anthelminitic potential of Cassia plants. Asian Pacific J Clin Nutr 4: S436-S441.
- Lal J, Chandra S, Prakash VR, Sabir M (1976) In vitro anthelmintic action of some indigenous medicinal plants on A. galli worms. Indian J Physiol Pharmacol 20: 64-68.
- Lalchhandama K (2008) Nematocidal effects of piperazine and the extract of Acacia oxyphylla stem bark on the poultry nematode Ascaridia galli. Pharmac Onl 3: 864-869.
- Lei J, Leser M, Enan E (2010) Nematicidal activity of two monterpenoids and SER-2 tyramine receptor of Caenorhabditis elegans. Biochem Pharmacol 79: 1062-1071.
- Mali RG, Mahajan SG, Mehta AA (2007) In vitro screening of Cleome viscosa extract for anthelmintic activity. Pharm Biol 45: 766-768.
- Mali RG, Mehta AA (2008) A review on anthelmintic plants. Nat Prod Radiance 7: 466-475.
- Marin L, Miguelez M, Villar CJ, Lombo F (2015) Bioavailability of dietary polyphenols and gut microbiota metabolism: Antimicrobial properties. BioMed Res Int ID905215: 1-18.
- Martin-Pacho JR, Montoya MN, Aranguena T, Toro C, Morchon R, Marcos-Atxutegi C, Simon F (2005) A coprological and serological survey for the prevalence of Ascaridia spp. in laying hens. J Vet Med B Infect Dis Vet Public Health 82: 238-242.
- McCloud TG (2010) High throughput extraction of plant, marine and fungal specimens for preservation of biologically active molecules a review. Molecules 15: 4526-4563.
- Murthy PK, Joseph SK, Murthy PSR (2011) Plant products in the treatment and control of filariasis and other helminth infections and assay systems for antifilarial/anthelmintic activity. Planta Med 77: 647-661.
- Muyibi SA, Olorode BR, Onyeyili PA, Osunkwo UA, Muhammad BY, Ajagbonna OP (2000) Haematological and histopathological changes of Cassia occidentalis leaf extract in rats. Nig J Nat Prod Med 4: 48-51.
- Permin A, Christensen JP, Bisgaard M (2006) Consequences of concurrent A. galli and Escherichia coli infections in chickens. Acta Vet Scand 47: 43-54.
- Rafi KP, Karthikeyan M, Kannan M, Rajasekar S (2011) Anthelmintic activity of Nerium olender flower extract in Indian adult earthworm. J Nat Prod Plant Res 1: 40-46.
- Rex E, Molitor SC, Hapiak V, Xiao H, Hernderson M, Komuniecki R (2004) Tyramine receptor (SER-2) isoforms are involved in regulation of pharyngeal pumping and foraging behavior in Caenorhabditis elegans. J Neurochem 91: 1104-1115.
- Roy H, Chakraborty A, Bhanja S, Nayak BS, Mishra SR, Ellaiah P (2010) Preliminary phytochemical investigation and anthelmintic activity of Acanthospermum hispidum DC. J Pharmaceut Sci Technol 2: 217-221.

- Sabri MH (2013) Comparison of the effeciancy of levamisole and dry Punica granatum peel against Ascaridia galli in layer hens that naturally infested Euphrates J Agric Sci 5: 1-13.
- Salhan M, Kumar B, Tiwari P, Sharma P, Sandhar HK, Gautam M (2011) Comparative anthelmintic activity of aqueous and ethanolic leaf extracts Of Clitoria ternatea. Int J Drug Dev Res 3: 62-69.
- Sandoval-Castro CA, Torres-Acosta JFJ, Hoste H, Salem AZM, Chan-Perez JI (2012) Using plant bioactive materials to control gastrointestinal tract helminths in livestock. Animal Feed Sci Technol 176: 192-201.
- Schultz JC (1989) Tannin-Insect interactions. In: Hemingway RW, Karchesy JJ (eds.), Chemistry and significance of condensed tannins, Plenum Press, New York: pp. 417-433.
- Shah Alam M, Alam KJ, Begum N, Amin MR (2014) Comparative efficacy of different herbal and modern anthelmintics against gastrointestinal nematodiasis in fowl. Int J Biol Res 2: 145-148.
- Sharma M, Prasad SB (2014) Evaluation of anthelmintic activity of leaves extracts of Guazuma Tomentosa. Int J Pharmacol Pharmaceut Sci 1: 1-5.
- Siamba DN, Okitoi LO, Watai MK, Wachira AM, Lukibisi FB, Mukisira EA (2007) Efficacy of Tephrosia vogelli and Vernonia amygdalina as anthelmintics against Ascaridia galli in indigenous chicken Livestock Res Rural Dev 19: Article 176.
- Subash KR, Jagan Rao N, Cheriyan BV, Bhaarati GM, Kumar KS (2012) The anthelmintic activity of Eupatorium triplinerve and Alpinia galanga in Pheritima posthuma and Ascardia galli: A comparative Study. J Clin Diagn Res 6: 947-950.
- Tariq KA, Chishti MZ, Ahmad F, Shawl AS (2009) Anthelmintic activity of extracts of Artemisia absinthium against ovine nematodes. Vet Parasitol 160: 83-88.
- Van Krimpen MM, Binnendijk GP, Borgsteede FHM, Gaasenbeek CPH (2010) Anthelmintic effects of phytogenic feed additives in Ascaris suum inoculated pigs. Vet Parasitol 168: 269-277.
- Windisch W, Rohrer E, Schedle K (2009) Phytogenic feed additives to young piglets and poultry: Mechanisms and application. In: Steiner T (ed.), Phytogenics in animal nutrition. Natural concepts to optimize gut health and performance, Nottingham University Press, Nottingham: pp. 19-38.
- Wink M (2012) Medicinal plants: A source of anti-parasitic secondary metabolites. Molecules 17: 12771-12791.
- Wu KM, Farrelly J, Birnkrant D, Chen S, Dou J, Atrakchi A, Bigger A, Chen C, Chen Z, Freed L, Ghantous H, Goheer A, Hausner E, Osterberg R, Rhee H, Zhang K (2004) Regulatory toxicology perspectives on the development of botanical drug products in the United States. Am J Ther 11: 213-217.
- Yazwinski TA, Tucker CA, Wray E, Jones L, Clark FD (2013) Observations of benzimidazole efficacies against Ascaridia dissimilis, Ascaridia galli, and Heterakis gallinarum in naturally infected poultry. J Appl Poultry Res 22: 75-79.