

## Journal of the Hellenic Veterinary Medical Society

Vol 69, No 1 (2018)



**Effects of yeast based toxin binder supplementation on growth performance and intestinal microarchitecture in male buffalo calves (*Bubalus bubalis*) exposed to different concentrations of Aflatoxin B1.**

O. NASEER, M. U. SALEEM, M. L. SOHAIL, J. NASEER, J. KHAN, M. O. OMER, K. M. ANJUM, A. S. AHMAD

doi: [10.12681/jhvms.16440](https://doi.org/10.12681/jhvms.16440)

Copyright © 2018, O NASEER, MU SALEEM, ML SOHAIL, J NASEER, JA KHAN, MO OMER, KM ANJUM, AS AHMAD



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

### To cite this article:

NASEER, O., SALEEM, M. U., SOHAIL, M. L., NASEER, J., KHAN, J., OMER, M. O., ANJUM, K. M., & AHMAD, A. S. (2018). Effects of yeast based toxin binder supplementation on growth performance and intestinal microarchitecture in male buffalo calves (*Bubalus bubalis*) exposed to different concentrations of Aflatoxin B1. *Journal of the Hellenic Veterinary Medical Society*, 69(1), 853–862. <https://doi.org/10.12681/jhvms.16440>

**Effects of yeast based toxin binder supplementation on growth performance and intestinal microarchitecture in male buffalo calves (*Bubalus bubalis*) exposed to different concentrations of Aflatoxin B1.**

**Naseer O.<sup>1\*</sup>, Saleem M.U.<sup>2</sup>, Sohail M. L.<sup>1</sup>, Naseer J.<sup>4</sup>, Khan J.A. <sup>6</sup>, Omer M.O.<sup>5</sup>,  
Anjum K.M.<sup>4</sup>, Ahmad A.S.<sup>3</sup>**

<sup>1</sup>University College of Veterinary and Animal sciences, The Islamia University of Bahawalpur, Pakistan.

<sup>2</sup>Department of Biosciences, Faculty of Veterinary Sciences, Bahauddin Zakariya University, Multan, Pakistan

<sup>3</sup>Department of Parasitology, University of Veterinary and Animal Sciences, Lahore, Pakistan

<sup>4</sup>Department of Wildlife & Ecology, University of Veterinary and Animal Sciences, Lahore, Pakistan

<sup>5</sup>Department of Pharmacology and Toxicology, Faculty of Biosciences, UVAS Lahore, Pakistan

<sup>6</sup>Department of Clinical Medicine and Surgery, University of Veterinary and Animal Sciences, Lahore, Pakistan

**ABSTRACT.** The purpose of study was to analyze the effects of yeast based toxin binder (YTB) on growth performance and intestinal microarchitecture of buffalo calves exposed to different concentrations of aflatoxin B1 (AFB1). Male buffalo calves (n=72) having no disease exposure in the last four months, aged  $10 \pm 1$  months and weighing  $150 \pm 12.22$  kg (Mean  $\pm$  standard deviation) were purchased from a buffalo calf fattening farm and divided in 8 equal groups (n=9 animals in each group). Animals that were fed common basal diet and basal diet along with yeast cell wall based toxin binder (YTB) and had no exposure to AFB1 were served as controls. Different concentrations of AFB1 (6mg, 8mg and 10mg per kg of feed) were given to 27 animals, similarly 2mg of YTB (per kg of contaminated feed) was added to concentrate with different AFB1 concentrations and fed to another 27 animals. Average daily gain of each animal was recorded. Animals were slaughtered after 27 days and intestinal segments from duodenum, jejunum, ileum and colon were collected which were stained with hematoxylin/eosin or combined alcian blue and periodic acid schiff stain.

*Corresponding Author:*

Omer Naseer

Email: dromersheikh@gmail.com

Postal Address: University College of Veterinary and Animal sciences,  
The Islamia University of Bahawalpur, Pakistan.

*Date of initial submission: 2-3-2017*

*Date of revised submission: 15-4-2017*

*Date of acceptance: 21-5-2017*

Average daily gain and intestinal morphometric parameters (villus height, villus width, crypt depth and surface area) in all the selected intestinal segments decreased ( $P<0.05$ ) whereas, intraepithelial lymphocytes, goblet cells (having acidic and neutral mucins) and proliferative cells increased ( $P<0.05$ ) with AFB1 exposure. It was also observed that YTB neutralized ( $P<0.05$ ) the harmful effects of AFB1 on intestinal morphology, boosted activity of goblet cells and decreased ( $P<0.05$ ) number of proliferative cells. It was concluded that YTB supplementation is advantageous in eliminating the negative effects of AFB1 on gut of buffalo calves.

**Keywords:** Buffaloes, aflatoxin B1, yeast based toxin binder supplementation, growth performance, intestinal histology.

## INTRODUCTION

Aflatoxins (AF), a cluster of intoxicating mycotoxins are secondary metabolites from certain strains of filamentous fungi specially *Aspergillus Flavus* and *Aspergillus Parasiticus* (Rashid et al., 2012). These fungi colonize on feed and food products and produce AF whose presence may lead to economic losses and raise public health concerns which at times can prove to be fatal (Mendoza et al., 2011). Harmful outcomes of AF which include teratogenic, mutagenic, growth inhibitory and carcinogenic effects have been widely studied (Oguz et al., 2000, Sur and Celik, 2003). Aflatoxins suppress the immune system and cause both macro and micro pathological alterations in the normal structure of liver, kidney and spleen making animal more susceptible to infectious agents (Magnoli et al., 2011).

To decrease the toxic effects of AF toxin binders (TB) are being used (Schatzmayr et al., 2003). Inclusion of TB in feed contaminated with AF has been the most promising approach for reducing the detrimental effects of AF (Galvano et al., 2001). In-vivo and in-vitro trails have been conducted over the last decade with TB for estimating their efficacy against toxins but results from both the trials do not correlate very well (Doll et al., 2004; Diaz et al., 2004). Many complex indigestible carbohydrates (polysaccharides in yeast cell wall and cellulose) and bacteria (glucomannans and peptidoglycans) have been reported to absorb AF among these yeast cell wall based toxin binder (YTB) is being preferred for its better nutritional properties and superior binding activity than other commercially available toxin binders (Huwig et al., 2001).

Among several types of AF aflatoxin B1 (AFB1) is most commonly encountered and is considered most potent than any other type (Yunus et al., 2011),

causing stern health concerns for animal populations (Sirajudeen et al., 2009). Negative effects of AF have been an active area of research (Yunus et al., 2011), and studies have revealed that the primary site for absorption of AF is small intestine (predominantly duodenum) (Mendoza et al., 2011). To the best of our knowledge influence of YTB supplementation on the quantification and differentiation of goblet cells (GC), role of intraepithelial lymphocytes (IEL) as an indicator of mucosal damage and immunohistochemistry of proliferative cells in intestine of buffalo calves exposed to different concentrations of AFB1 has not been assed. This study will be pivotal in recognizing the structural alterations in buffalo gut associated with AFB1 and YBT supplementation as it aims to evaluate the effects of different concentrations of AFB1 on intestinal microarchitecture (morphometric variables, IEL count, GC quantification and differentiation and immunohistochemistry of proliferative cells) in gut of buffalo calves and efficacy of YTB to counter toxic effects of AFB1 on intestinal morphology.

## MATERIALS AND METHODS

### Experimental design and grouping of animals

A total of 72 having no disease exposure in the last four months, aged  $10 \pm 1$  months and weighing  $150 \pm 12.22$  kg (Mean  $\pm$  standard deviation) were purchased from a buffalo calf fattening farm and divided in 8 equal groups ( $n=9$  animals in each group). These animals were then acclimatized for 15 days in an environmentally control shed having an optimum temperature of  $22 \pm 2$  °C and a relative humidity of  $55 \pm 10$  % before the start of trial during which they were given ad-libitum corn silage (Table 1). Silage was tested for the presence of AFB1 on weekly basis by high performance liquid chromatog-

raphy. Animals were dewormed and vaccinated for foot and mouth disease and hemorrhagic septicemia (UVAS-FMD+HS-VAC, Pakistan). At the start of trail all animals were individually housed in separate pens made of stainless steel and were offered corn silage (free of AFB1, 15 kg per animal) and concentrate (1 kg per animal) (Table 1) per day and there refusal were weighed daily. Animals were divided into eight groups with each group having (n = 9) animals. Basal diet (BD) and BD plus YTB (BD-YTB) were not exposed to AFB1. For investigating the harmful effects of AFB1 different concentrations of AFB1 were given to animals of 3 groups which included AFB1 6mg per kg of concentrate (AFB1 0.6), AFB1 8mg per kg of concentrate (AFB1 0.8) and AFB1 10mg per kg of concentrate (AFB1 1.0) after being formulated in the concentrate (Table 1) for 27 days. Furthermore for analysis the effects of YTB during aflatoxicosis in buffalo calves, 2mg per kg of YTB (Fixar viva dry yeast based, Bentoli Agrinutrition Asia Pte. Ltd, USA) was mixed in feed having different concentrations of AFB1 and given to another 3 groups of buffalo calves (AFB1 0.6-YTB, AFB1 0.8-YTB, and AFB1 1.0-YTB). Throughout the trail animals had open access to water. This study was conducted according to the guidelines of Animal Care and Use Committee, University of Veterinary and Animal Sciences (UVAS), Lahore.

#### Production of AFB1, preparation of diet and safety measures

Aflatoxin B1 being used in the trial was produced in Microbiology laboratory UVAS using toxic strains of *Aspergillus Flavus* IMI-90, obtained from International Mycology Institute, London as described by Tessari et al., (2006). Briefly, chloroform (30 ml chloroform per 10 grams of culture) was used to extract coconut culture after shaking for 30 minutes. Contents post filtration via

filter paper (Whatman#1) was evaporated to dryness. Densitometry was applied for quantification and water bath (WNE 14 water bath, Memmert, USA) was used for evaporation of chloroform solution having AFB1 at 60 °C. Contents were resuspended in fish oil that had been proven negative for AF presence. This fish oil with different concentrations

**Table 1.** Ingredients (as fed basis) and nutrient composition (% dry matter basis) of concentrate and corn silage fed to male buffalo calves.

Ingredients (Concentrate)	% unless indicated
Corn Meal	22.8
Sugarcane molasses	2
Wheat midds	65.8
Soyabean meal	3.3
Limestone	3.6
Fish oil	1
Salt	1
Premix*	0.4
<b>Chemical composition (Concentrate)</b>	
Dry matter	86
Crude protein	16.8
Crude fat	4.2
Crude fiber	12.3
Neutral detergent fiber	38.6
Metabolizable energy	2.6 M cal/kg
<b>Chemical composition (Corn Silage)</b>	
Dry matter	35.7
Crude protein	7.3
Crude fat	1.3
Crude fiber	26.4
Neutral detergent fiber	66.2
Metabolizable energy	2.2 M cal/kg

*M cal, Mega calories.*

\* Each kg of premix contained; vitamin A, 495 IU; vitamin E, 0.33 IU; vitamin D3, 165 IU; zinc, 16mg; copper, 4mg; selenium, 0.1mg; iodine, 0.6mg; and cobalt, 0.6mg.

(6mg, 8mg and 10mg) of AFB1 was used in the concentrate formulation (Table 1). Final concentrations of AFB1 in concentrate to be offered were confirmed using procedure used by Shephard et al., (1990).

Aflatoxin B1 is a toxic substance therefore it was manipulated as solutions to avoid formation of aerosol and dust. Aprons, masks and nitrile gloves were used at every step of manufacturing and manipulation (Corcuera et al., 2011).

### **Growth performance, sampling and histological morphometry**

Animals were weighed on weekly basis and after completion of trial were slaughtered by Halal slaughter method as described by Gregory et al., (2008), and 3cm intestinal segments at midpoints of duodenum, jejunum, ileum and colon were collected. Samples were washed with physiological saline solution, opened longitudinally and immediately fixed in 10% buffered formaldehyde solution for 48 hours. Intestinal segments were then rinsed with water and after dehydration with graded series of absolute alcohol (50%, 60%, 70%, 80%, 90% and 100%) were cleared twice with benzene. Sections were embedded with paraffin and 4 µm thin sections were obtained which were stained with hematoxylin/eosin (Wang et al., 2009). Slides were observed under light microscope (Labomed CXL, New York Microscope Co, USA) fitted with camera (Moticam CMOS Digital Camera, New York Microscope Co, USA) and measurements were made with commercial morphometric program (Pixel Pro, Labomed, USA). Double blind analysis was done in triplicate on 5 well oriented villi that were selected on the basis of intact lamina propria. Variables that were measured included villus height (from tip of villus to villus crypt junction), villus width, crypt depth (from base to the transition region between villus and crypt), surface area ( $2 \times \text{villus width}/2 \times \text{villus length}$ ) (Solis de los Santos et al., 2007), and IEL (rounded cell that had central or slightly eccentric nucleus with scant cytoplasm) number.

### **Histo-chemistry of goblet cells**

Slides obtained from paraffin embedding technique were subjected to alcian blue and periodic acid Schiff staining methods (Bancroft and Stevens, 2007), to

evaluate goblet cells containing acidic and neutral mucin types respectively. Acidic mucins were stained blue by AB whereas neutral mucins were stained magenta by PAS (Leknes, 2010).

### **Immunohistochemistry of proliferative cells**

Proliferative cells were counted in all the groups with antibody MIB-1 (Sigma-Aldrich, St. Louis, MO., USA) directed against proliferation marker Ki-67. Regarding antigen accessibility sections were incubated in boiling 0.01 M citrate buffer having pH 6 and washing was carried out in phosphate buffered saline (PBS) with 0.05% Tween (PBS-T). For blocking the endogenous peroxidase activity slides were treated with 1% hydrogen peroxide and to avoid binding of unspecific antibody slides were treated with 10% goat serum (Life technologies, Scientific supplies Ltd. Pakistan). At 4 °C binding of MIB-1 (1:50 in PBS) was done overnight. Sections were then incubated with polyclonal goat anti-mouse antibody which had been labeled with horse red-dish peroxidase. Visualization was done with 3,3'-diaminobenzidine solution (KPL Inc., Gaithersburg, Maryland USA). Quantification of proliferative cells positive to MIB-1 antibody was done as described by Masanatz et al., (2010) in the five consecutive crypts of selected intestinal segments and the average of the results were reported.

### **STATISTICAL ANALYSIS**

Statistical differences were analyzed by using Statistical Package for Social Sciences (SPSS 13.3, SPSS, Chicago, USA). For estimating the normal distribution of data Shapiro- Wilk test was applied, data was found to be normally distributed. Data for comparison among groups were subjected to one way ANOVA keeping BD, different concentrations of AFB1, treatment of AFB1 contaminated groups with YTB and inclusion of YTB to BD as constant factors. Statistical differences among means were considered significant at  $P < 0.05$  and calculated using Duncan's multiple range test.

### **RESULTS**

Effects of feeding different levels of AFB1 on average daily gain and morphometric variables (villus




height, villus width, crypt depth, villus: crypt ratio and surface area) of selected intestinal segments are shown in Table 2. Exposure of buffalo calves to AFB1 resulted in lower ( $P<0.05$ ) average daily gain and decreased ( $P<0.05$ ) morphometric variables except for crypt depth which increased ( $P<0.05$ ) with AFB1 for all the selected intestinal segments. Villus height, villus width, villus: crypt ratio and surface area decreased ( $P<0.05$ ) whereas crypt depth increased ( $P<0.05$ ) as AFB1 concentrations were increased. It was observed that by addition of YTB harmful effects of AFB1 on intestinal microarchitecture were neutralized as no difference was observed for morphometric variables in AFB1 0.6-YTB, AFB1 0.8-YTB and AFB1 1.0-YTB compared to BD. Moreover inclusion of YTB in BD not contaminated with AFB1 increased ( $P<0.05$ ) villus height, villus

width, and surface area.


A higher ( $P<0.05$ ) IEL count was observed in all selected intestinal segments of male buffalo calves exposed to different concentrations of AFB1 compared to BD animals. Inclusion of YTB to contaminated diet decreased ( $P<0.05$ ) IEL count of animals compared to those exposed to AFB1. Number of GC, GC having acidic mucin and neutral mucin increased ( $P<0.05$ ) with exposure to AFB1 compared to BD and gradually increased ( $P<0.05$ ) with increasing AFB1 concentrations in feed. Supplementation of YTB caused an increase ( $P<0.05$ ) in GC having acidic mucin and neutral mucin in selected intestinal segments of animals exposed to AFB1 and BD-YTB animals.

For all the selected intestinal segments number of proliferative cells increased ( $P<0.05$ ) with increas-


**Fig 1.** Acidic type of goblet cells in in the duodenum of calves fed AFB1 0.8 -YTB.




**Fig 2.** Mixed type of goblet cells in in the duodenum of calves fed AFB1 0.8 -YTB.



**Fig 3.** Acidic and mixed mucin having types of goblet cells in in the duodenum of calves fed AFB1 -0.8.



**Fig 4.** Intra epithelial lymphocytes in the Duodenum of calves fed AFB1- 0.8.



**Table 2.** Comparison of intestinal morphometric parameters and intraepithelial lymphocyte count in intestinal segments of male buffalo calves.

Intestinal segments	Variables	BD	Contaminated			Treated			BD-YTB	SEM	P-Value
			AFB <sub>1</sub> 0.6	AFB <sub>1</sub> 0.8	AFB <sub>1</sub> 1.0	AFB <sub>1</sub> 0.6-YTB	AFB <sub>1</sub> 0.8-YTB	AFB <sub>1</sub> 1.0-YTB			
Duodenum	Average daily gain (grams)	502 <sup>d</sup>	411 <sup>c</sup>	372 <sup>b</sup>	310 <sup>a</sup>	491 <sup>d</sup>	505 <sup>d</sup>	511 <sup>d</sup>	557 <sup>c</sup>	31.75	< 0.01
	Villus height (mm)	0.73 <sup>d</sup>	0.51 <sup>c</sup>	0.45 <sup>b</sup>	0.39 <sup>a</sup>	0.72 <sup>d</sup>	0.71 <sup>d</sup>	0.68 <sup>d</sup>	0.78 <sup>c</sup>	0.04	< 0.01
	Villus width (mm)	0.15 <sup>d</sup>	0.11 <sup>c</sup>	0.09 <sup>b</sup>	0.06 <sup>a</sup>	0.12 <sup>d</sup>	0.11 <sup>d</sup>	0.12 <sup>d</sup>	0.18 <sup>c</sup>	0.01	< 0.01
	Crypt depth (mm)	0.23 <sup>a</sup>	0.35 <sup>c</sup>	0.41 <sup>d</sup>	0.50 <sup>e</sup>	0.24 <sup>a</sup>	0.21 <sup>a</sup>	0.25 <sup>a</sup>	0.27 <sup>ab</sup>	0.04	< 0.01
	Villus: crypt ratio	3.17 <sup>e</sup>	1.45 <sup>c</sup>	1.09 <sup>b</sup>	0.78 <sup>a</sup>	3.01 <sup>e</sup>	3.38 <sup>f</sup>	2.72 <sup>d</sup>	2.88 <sup>de</sup>	0.39	< 0.01
	Surface area (mm <sup>2</sup> )	0.37 <sup>e</sup>	0.18 <sup>c</sup>	0.15 <sup>b</sup>	0.08 <sup>a</sup>	0.27 <sup>d</sup>	0.25 <sup>d</sup>	0.26 <sup>d</sup>	0.45 <sup>c</sup>	0.04	< 0.01
	Intraepithelial lymphocytes	24.61 <sup>a</sup>	38.14 <sup>d</sup>	42.73 <sup>e</sup>	49.55 <sup>f</sup>	27.32 <sup>ab</sup>	30.24 <sup>b</sup>	33.17 <sup>b</sup>	24.18 <sup>a</sup>	3.38	< 0.01
Jejunum	Villus height (mm)	0.69 <sup>d</sup>	0.44 <sup>c</sup>	0.36 <sup>b</sup>	0.25 <sup>a</sup>	0.68 <sup>d</sup>	0.71 <sup>d</sup>	0.73 <sup>d</sup>	0.79 <sup>e</sup>	0.07	< 0.01
	Villus width (mm)	0.14 <sup>c</sup>	0.08 <sup>b</sup>	0.07 <sup>b</sup>	0.05 <sup>a</sup>	0.13 <sup>c</sup>	0.15 <sup>c</sup>	0.15 <sup>c</sup>	0.19 <sup>d</sup>	0.01	< 0.01
	Crypt depth (mm)	0.26 <sup>a</sup>	0.38 <sup>c</sup>	0.44 <sup>d</sup>	0.48 <sup>e</sup>	0.25 <sup>a</sup>	0.27 <sup>a</sup>	0.29 <sup>a</sup>	0.31 <sup>ab</sup>	0.03	< 0.01
	Villus: crypt ratio	2.65 <sup>d</sup>	1.15 <sup>c</sup>	0.81 <sup>b</sup>	0.52 <sup>a</sup>	2.72 <sup>d</sup>	2.62 <sup>d</sup>	2.51 <sup>d</sup>	2.54 <sup>d</sup>	0.35	< 0.01
	Surface area (mm <sup>2</sup> )	0.32 <sup>d</sup>	0.17 <sup>c</sup>	0.08 <sup>b</sup>	0.04 <sup>a</sup>	0.28 <sup>d</sup>	0.31 <sup>d</sup>	0.34 <sup>d</sup>	0.45 <sup>c</sup>	0.05	< 0.01
	Intraepithelial lymphocytes	22.61 <sup>a</sup>	31.25 <sup>d</sup>	35.43 <sup>e</sup>	42.72 <sup>f</sup>	23.69 <sup>ab</sup>	25.57 <sup>b</sup>	24.91 <sup>ab</sup>	21.89 <sup>a</sup>	2.64	< 0.01
	Villus height (mm)	0.64 <sup>e</sup>	0.41 <sup>c</sup>	0.35 <sup>b</sup>	0.33 <sup>a</sup>	0.61 <sup>de</sup>	0.58 <sup>d</sup>	0.53 <sup>d</sup>	0.71 <sup>f</sup>	0.05	< 0.01
Ileum	Villus width (mm)	0.13 <sup>d</sup>	0.09 <sup>c</sup>	0.06 <sup>b</sup>	0.04 <sup>a</sup>	0.11 <sup>d</sup>	0.12 <sup>d</sup>	0.11 <sup>d</sup>	0.15 <sup>e</sup>	0.01	< 0.01
	Crypt depth (mm)	0.33 <sup>a</sup>	0.39 <sup>c</sup>	0.47 <sup>d</sup>	0.53 <sup>e</sup>	0.29 <sup>a</sup>	0.31 <sup>a</sup>	0.34 <sup>a</sup>	0.36 <sup>ab</sup>	0.03	< 0.01
	Villus: crypt ratio	1.93 <sup>ef</sup>	1.05 <sup>c</sup>	0.74 <sup>b</sup>	0.62 <sup>a</sup>	2.10 <sup>e</sup>	1.87 <sup>c</sup>	1.55 <sup>d</sup>	1.97 <sup>ef</sup>	0.22	< 0.01
	Surface area (mm <sup>2</sup> )	0.28 <sup>e</sup>	0.12 <sup>c</sup>	0.07 <sup>b</sup>	0.04 <sup>a</sup>	0.21 <sup>d</sup>	0.22 <sup>d</sup>	0.19 <sup>d</sup>	0.34 <sup>f</sup>	0.03	< 0.01
	Intraepithelial lymphocytes	27.19 <sup>a</sup>	37.27 <sup>d</sup>	44.31 <sup>e</sup>	46.22 <sup>f</sup>	29.64 <sup>b</sup>	30.43 <sup>b</sup>	28.88 <sup>ab</sup>	26.91 <sup>a</sup>	2.81	< 0.01
	Villus height (mm)	0.59 <sup>d</sup>	0.45 <sup>c</sup>	0.39 <sup>b</sup>	0.32 <sup>a</sup>	0.57 <sup>d</sup>	0.55 <sup>d</sup>	0.58 <sup>d</sup>	0.67 <sup>e</sup>	0.04	< 0.01
	Villus width (mm)	0.11 <sup>d</sup>	0.07 <sup>c</sup>	0.05 <sup>b</sup>	0.04 <sup>a</sup>	0.11 <sup>d</sup>	0.10 <sup>d</sup>	0.12 <sup>d</sup>	0.16 <sup>e</sup>	0.01	< 0.01
Colon	Crypt depth (mm)	0.37 <sup>a</sup>	0.43 <sup>c</sup>	0.49 <sup>d</sup>	0.57 <sup>e</sup>	0.36 <sup>a</sup>	0.39 <sup>a</sup>	0.41 <sup>a</sup>	0.43 <sup>ab</sup>	0.02	< 0.01
	Villus: crypt ratio	1.59 <sup>e</sup>	1.04 <sup>c</sup>	0.79 <sup>b</sup>	0.56 <sup>a</sup>	1.58 <sup>c</sup>	1.41 <sup>d</sup>	1.42 <sup>d</sup>	1.55 <sup>e</sup>	0.15	< 0.01
	Surface area (mm <sup>2</sup> )	0.22 <sup>d</sup>	0.09 <sup>c</sup>	0.06 <sup>b</sup>	0.04 <sup>a</sup>	0.19 <sup>d</sup>	0.17 <sup>d</sup>	0.21 <sup>d</sup>	0.28 <sup>e</sup>	0.03	< 0.01
	Intraepithelial lymphocytes	31.11 <sup>a</sup>	47.02 <sup>e</sup>	54.55 <sup>d</sup>	59.88 <sup>c</sup>	35.62 <sup>a</sup>	34.47 <sup>a</sup>	35.23 <sup>a</sup>	31.97 <sup>a</sup>	3.92	< 0.01

Results are demonstrated as mean  $\pm$  s.e.m.

Difference of superscripts (a-f) within a row indicates significance difference  $P < 0.05$ .

BD, basal diet (negative for AFB<sub>1</sub>); AFB<sub>1</sub> 0.6, aflatoxin B<sub>1</sub> 6mg per kg of concentrate; AFB<sub>1</sub> 0.8, aflatoxin B<sub>1</sub> 8mg per kg of concentrate; AFB<sub>1</sub> 1.0, aflatoxin B<sub>1</sub> 10 mg per kg of concentrate; AFB<sub>1</sub> 0.6- YTB, aflatoxin B<sub>1</sub> 6mg per kg of concentrate plus 2mg of yeast based toxin binder per kg of feed; AFB<sub>1</sub> 0.8- YTB, aflatoxin B<sub>1</sub> 8mg per kg of concentrate plus 2mg of yeast based toxin binder per kg of feed; AFB<sub>1</sub> 1.0- YTB, aflatoxin B<sub>1</sub> 10mg per kg of concentrate plus 2mg of yeast based toxin binder per kg of feed; BD-YTB, basal diet plus 2mg per kg of yeast based toxin binder; SEM, standard error of mean.

**Table 3.** Goblet cells, histochemistry of goblet cells and immunohistochemistry of proliferative cells in intestinal segments of male buffalo calves.

Variables	Intestinal segments	BD	Contaminated			Treated			BD-YTB	SEM	P-Value
			AFB <sub>1</sub> 0.6	AFB <sub>1</sub> 0.8	AFB <sub>1</sub> 1.0	AFB <sub>1</sub> 0.6-YTB	AFB <sub>1</sub> 0.8-YTB	AFB <sub>1</sub> 1.0-YTB			
Goblet cells (per villus)	Duodenum	49.21 <sup>a</sup>	59.24 <sup>b</sup>	62.34 <sup>bc</sup>	65.37 <sup>c</sup>	67.52 <sup>d</sup>	69.98 <sup>d</sup>	73.29 <sup>e</sup>	99.86 <sup>f</sup>	5.56	< 0.01
	Jejunum	55.82 <sup>a</sup>	68.13 <sup>b</sup>	71.14 <sup>b</sup>	75.02 <sup>bc</sup>	78.09 <sup>d</sup>	81.27 <sup>de</sup>	85.12 <sup>e</sup>	116.71 <sup>f</sup>	6.69	< 0.01
	Ileum	71.65 <sup>a</sup>	79.84 <sup>b</sup>	80.23 <sup>b</sup>	83.26 <sup>b</sup>	89.17 <sup>c</sup>	93.45 <sup>c</sup>	96.15 <sup>c</sup>	141.34 <sup>d</sup>	8.12	< 0.01
	Colon	88.91 <sup>a</sup>	93.23 <sup>b</sup>	95.77 <sup>b</sup>	98.65 <sup>c</sup>	101.24 <sup>d</sup>	105.36 <sup>d</sup>	108.91 <sup>d</sup>	168.79 <sup>e</sup>	9.65	< 0.01
Goblet cells having acidic mucin (per villus)	Duodenum	18.14 <sup>a</sup>	31.86 <sup>b</sup>	34.25 <sup>c</sup>	36.97 <sup>c</sup>	41.69 <sup>d</sup>	44.26 <sup>d</sup>	49.87 <sup>d</sup>	51.27 <sup>e</sup>	3.59	< 0.01
	Jejunum	26.61 <sup>a</sup>	30.74 <sup>b</sup>	33.37 <sup>b</sup>	35.42 <sup>c</sup>	39.19 <sup>d</sup>	43.84 <sup>e</sup>	44.36 <sup>e</sup>	58.17 <sup>f</sup>	3.43	< 0.01
	Ileum	29.22 <sup>a</sup>	36.11 <sup>b</sup>	40.42 <sup>bc</sup>	41.24 <sup>c</sup>	45.13 <sup>d</sup>	48.27 <sup>d</sup>	51.86 <sup>d</sup>	62.88 <sup>e</sup>	3.87	< 0.01
	Colon	33.15 <sup>a</sup>	38.21 <sup>b</sup>	40.07 <sup>b</sup>	41.14 <sup>c</sup>	49.11 <sup>d</sup>	51.24 <sup>d</sup>	53.62 <sup>d</sup>	74.16 <sup>e</sup>	4.84	< 0.01
Goblet cells having neutral mucin (per villus)	Duodenum	15.26 <sup>a</sup>	22.17 <sup>b</sup>	23.79 <sup>b</sup>	25.32 <sup>c</sup>	21.35 <sup>d</sup>	22.13 <sup>d</sup>	23.03 <sup>d</sup>	48.13 <sup>e</sup>	3.71	< 0.01
	Jejunum	24.27 <sup>a</sup>	27.21 <sup>b</sup>	28.53 <sup>b</sup>	30.79 <sup>b</sup>	32.84 <sup>c</sup>	34.21 <sup>c</sup>	36.18 <sup>c</sup>	49.27 <sup>d</sup>	2.97	< 0.01
	Ileum	25.04 <sup>a</sup>	30.18 <sup>b</sup>	33.29 <sup>c</sup>	34.96 <sup>c</sup>	38.21 <sup>d</sup>	40.39 <sup>d</sup>	42.58 <sup>d</sup>	57.83 <sup>e</sup>	3.72	< 0.01
	Colon	29.04 <sup>a</sup>	34.55 <sup>b</sup>	36.11 <sup>b</sup>	37.25 <sup>c</sup>	42.24 <sup>d</sup>	44.95 <sup>d</sup>	47.14 <sup>e</sup>	65.39 <sup>f</sup>	4.19	< 0.01
Proliferative cells (per mm of crypt)	Duodenum	38.41 <sup>b</sup>	42.91 <sup>c</sup>	43.74 <sup>c</sup>	46.19 <sup>d</sup>	36.29 <sup>b</sup>	39.57 <sup>b</sup>	39.86 <sup>b</sup>	34.79 <sup>a</sup>	1.45	< 0.01
	Jejunum	32.13 <sup>b</sup>	35.62 <sup>c</sup>	37.89 <sup>cd</sup>	39.27 <sup>d</sup>	30.61 <sup>b</sup>	31.46 <sup>b</sup>	32.87 <sup>b</sup>	29.28 <sup>a</sup>	1.36	< 0.01
	Ileum	27.57 <sup>b</sup>	31.73 <sup>c</sup>	33.54 <sup>d</sup>	37.28 <sup>e</sup>	26.12 <sup>b</sup>	28.99 <sup>b</sup>	29.15 <sup>b</sup>	24.85 <sup>a</sup>	1.54	< 0.01
	Colon	25.38 <sup>b</sup>	28.25 <sup>d</sup>	30.42 <sup>d</sup>	34.67 <sup>e</sup>	24.11 <sup>bc</sup>	26.43 <sup>bc</sup>	27.49 <sup>c</sup>	21.09 <sup>a</sup>	1.53	< 0.01

Results are demonstrated as mean  $\pm$  s.e.m.

Difference of superscripts (a-f) within a row indicates significance difference  $P < 0.05$ .

BD, basal diet (negative for AFB<sub>1</sub>); AFB<sub>1</sub> 0.6, aflatoxin B<sub>1</sub> 6mg per kg of concentrate; AFB<sub>1</sub> 0.8, aflatoxin B<sub>1</sub> 8mg per kg of concentrate; AFB<sub>1</sub> 1.0, aflatoxin B<sub>1</sub> 10 mg per kg of concentrate; AFB<sub>1</sub> 0.6- YTB, aflatoxin B<sub>1</sub> 6mg per kg of concentrate plus 2mg of yeast based toxin binder per kg of feed; AFB<sub>1</sub> 0.8- YTB, aflatoxin B<sub>1</sub> 8mg per kg of concentrate plus 2mg of yeast based toxin binder per kg of feed; AFB<sub>1</sub> 1.0- YTB, aflatoxin B<sub>1</sub> 10mg per kg of concentrate plus 2mg of yeast based toxin binder per kg of feed; BD-YTB, basal diet plus 2mg per kg of yeast based toxin binder; SEM, standard error of mean.

ing AFB<sub>1</sub> concentrations but YTB supplementation decreased ( $P < 0.05$ ) there number and brought it back to normal levels as no difference for proliferative cells was observed between animals of AFB<sub>1</sub> 0.6-YTB, AFB<sub>1</sub> 0.8-YTB, AFB<sub>1</sub> 1.0-YTB and BD groups. However, BD-YTB animals had the least number ( $P < 0.05$ ) of proliferative cells in the selected intestinal segments compared to animals of other groups.

## DISCUSSION

Outcomes of YTB supplementation in BD and feed contaminated with different concentrations of AFB<sub>1</sub> on intestinal morphology were studied in male buffalo calves. Impairments in the intestinal mucosa caused by AFB<sub>1</sub> decreases its nutrient absorbing ability (Liu et al., 2011). Villus surface area depends upon the dimensions of villus (Hou et al., 2012), and higher growth rates are associated with greater villus




surface area (Awad et al., 2009), and increased villus: crypt ratio (Wu et al., 2004), which is in agreement with our results. Aflatoxin B1 induces alterations in intestinal microarchitecture (Yang et al., 2012), such as short villi, deep crypts, reduced villus: crypt ratio and surface area (Wan et al., 2013). This study confirms previous reports by Dogi et al., (2011), and Motawe et al., (2014), that YTB supplementation nullifies the harmful effects of AFB1 on intestinal morphology. Yeast cell wall (YCW) binds with AFB1, reduces its absorption in intestine and thus is useful in protecting ruminants from detrimental effects of AFB1 (Firmin et al., 2011). Yeast improves intestinal microarchitecture by increasing concentration of useful microbes present in the intestine and suppressing the concentration of pathogenic bacteria thus increasing growth performance of animal (Gao et al., 2008). It was also observed that YTB supplementation improved intestinal microarchitecture in BD-YTB animals compared to BD, similar results were seen in duodenum and jejunum of broilers by Gao et al., (2008), and in duodenum of pigs by Shen et al., (2009), who reported that YCW supplementation increases villus height and villus: crypt ratio.

The IELs play an important role in the regulations of immune response as they form specialized lymphoid compartments and are the first cells to encounter antigens in intestinal lumen (Finamore et al., 2008). Increased number of IEL in all the selected intestinal segments are results of intestinal inflammatory response (Quinteiro-Filho et al., 2010), towards AFB1. This increase in IEL count may also


be associated with lesions (O,Handley et al., 2001), caused by AFB1. Beta-glucans ( $\beta$ -glucans) are carbohydrates having linked glucose molecules which are major components of yeast cell wall (Volman et al., 2008). These components have the ability to enhance mucosal immunity of intestine (Battilana et al., 2001), as after oral administration  $\beta$ -glucans regulate immune cells of Peyer's patches and IELs (Suzuki et al., 1990; Tsukada et al., 2003). Supplementation of YTB decreased number of IEL in intestinal mucosa compared to AFB1. Intestinal motility is due to stimulation of parasympathetic nerves, since lymphocytes carry cholinergic receptors it can be assumed that stimulation of parasympathetic nerves then activate the mucosal immune system in the intestine leading to an increase in IEL number.

Highly viscous mucus layer that covers intestinal mucosa, helps in lubrication of ingested food and


**Fig 5.** Intra epithelial lymphocytes in the Duodenum of calves fed AFB1 0.8- YTB.



**Fig 6.** Histological section of duodenum showing villus height of calves fed AFB1 0.8- YTB.



**Fig 6.** Histological section of duodenum showing villus height of calves fed AFB1 0.8.



protects intestinal epithelium by acting as a physical barrier to pathogens is secreted by GC (Kim and Ho, 2010). Goblet cells are responsible for secretion of mucin which after hydration becomes mucus (Cone, 2009). Number of GC in intestinal epithelium may vary with challenging substances and diet (Machado-Neto et al., 2013). Number of intestinal GC on exposure to AFB1 increased in our study, similar results were observed by Kenawy et al., (2009), who observed an increased GC number in intestine after exposure to AFB1 contaminated diet. Moreover, addition of YCW increases GC number in intestine (Chee et al., 2010; Morales-Lopez et al., 2010; Muthusamy et al., 2012), but there is no consensus on whether increase in GC number is considered an improvement in animal health or not (Lea et al., 2013).

To our knowledge no data is present regarding histochemistry of GC and immunohistochemistry of proliferative cells in male buffalo calves exposed to AFB1 or supplemented with YTB. Mucins secreted by goblet cells are either acidic, neutral or mixed in nature. In fish GC having acidic mucins protect intestinal epithelium against chemical agents while GC having neutral mucins provide protection against chemical agents (Cruz et al., 2014). Factors that lead to an increase in GC having acidic and neutral mucin under the influence of AFB1 or YTB supplementation are yet to be investigated. However increased number of GC having acidic or neutral mucin in response to

YTB supplementation indicates greater intestinal protection against pathogens in male buffalo calves.

Amplified villus length is associated with an increase in number of proliferative cells as longer villi indicate faster proliferation in intestinal crypts (Wu et al., 2013). Decline in proliferative cells number reduces the amount of energy required for maintaining microarchitecture of gut (Masanetz et al., 2010). Yet the exact mechanisms that caused a diminution in proliferative cells number after YTB supplementation are needed to be explored.

## CONCLUSIONS

Collectively, results of the current study confirm that feeding aflatoxin B1 (AFB1) contaminated diet exerts deleterious effects on intestinal microarchitecture. Supplementation of 2mg/kg yeast cell wall based toxin binder (YTB) not only alleviates harmful effects on intestinal microarchitecture induced by AFB1 but also supports modulations in defense system of male buffalo calves. Therefore, use of YTB in AFB1 contaminated feed is beneficial and recommended.

## ACKNOWLEDGEMENTS

This work was financially supported by the Higher Education Commission (HEC), Islamabad, Pakistan under the Indigenous PhD fellowship program, PIN no. 112-22340-2AV1-034.

## REFERENCES

- Awad WA, Ghareeb K, Abdel-Raheem S, Bohm J (2009) Effects of dietary inclusion of probiotic and symbiotic on growth performance, organ weights and intestinal histomorphology of broiler chickens. *PoultSci* 88:49-55.
- Bancroft JD, Stevens A (2007) *Theory and practice of histological techniques*. 4th ed., Churchill, Livingstone, USA, pp. 180-201.
- Battilana P, Ornstein K, Minehira K, Schwarz JM, Acheson K, Schneider P, Burri J, Jequier E, Tappy L (2001) Mechanisms of actions of  $\beta$ -glucan in postprandial glucose metabolism in healthy men. *Eur J Clin Nutr* 55:327-333.
- Chee S, Iji P, Choct M, Mikkelsen L, Kocher A (2010) Functional interactions of mannan-oligosaccharide with dietary threonine in chicken gastrointestinal tract. I. growth performance and mucin dynamics. *Brit PoultSci* 51:677-685.
- Cone RA (2009) Barrier properties of mucus. *Adv Drug Deliver Rev* 61:75-85.
- Corcuera LA, Vea MI, Vettorazzi A, Penas EG (2011) Validation of a UHPLC-FLD analytical method for the simultaneous quantification of aflatoxin B1 and ochratoxin A in rat plasma, liver and kidney. *J Chromatogr B Analyt Technol Biomed Life Sci* 879: 2733-2740.
- Cruz TMP, Moretti DB, Nordin WM, Cyrino JEP, Machado-Neto R (2014) Intestinal epithelium of juvenile *Salminus brasiliensis* (Cuvier, 1816) fed diet with lyophilized bovine colostrum. *Aquacult Res* 1: 1-7.
- Diaz DE, Hagler WM, Blackwelder JT, Eve JA, Hopkins BA, Anderson KL, Jones FT, Whitlow LW (2004) Aflatoxin binders II: reduction of aflatoxin M1 in milk by sequestering agents of cows consuming aflatoxin in feed. *Mycopathologia* 57:233-241.
- Dogi CA, Armando R, Luduena R, de Moreno de Leblanc A, Rosa CAR, Dalcero A, Cavaglieri L (2011) *Saccharomyces cerevisiae* strains retain their viability and aflatoxin B1 binding ability under gastrointestinal conditions and improve ruminal fermentation. *Food Addit Contam A* 28: 1705-1711.
- Doll S, Danicke S, Valenta H, Flachowsky G (2004) In vitro studies on the evaluation of mycotoxins detoxifying agents for their efficacy on deoxynivalenol/zearalenone. *Arch Anim Nutr* 58:311-324.

- Finamore A, Roselli M, Britti S, Monastra G, Ambra R, Turrini A, Mengheri E (2008) Intestinal and peripheral immune response to MON810 maize ingestion in weaning and old mice. *J Agric Food Chem* 56: 11533-11539.
- Firmin R, Morgani DP, Yiannikouris A, Boudra H (2011) Effectiveness of modified yeast cell wall extracts to reduce aflatoxin B1 absorption in dairy ewes. *J Dairy Sci* 94: 5611-5619.
- Galvano F, Piva A, Ritieni A, Galvano G (2001) Dietary strategies to counteract the effects of mycotoxins: A review. *J Food Prod* 64:120-131.
- Gao J, Zhang HJ, Yu SH, Wu SG, Yoon I, Quigley J, Gao YP, Qi GH (2008) Effects of yeast culture in broiler diets on performance and immunomodulatory functions. *PoultSci* 87:1377-1384.
- Gregory NG, Wenzlawowicz MV, Alam RM, Anil HM, Yesildere T, Fletcher AS (2008) False aneurysms in carotid arteries of cattle and water buffalo during shechita and halal slaughter. *Meat Sci* 79:285-288.
- Hou Y, Wang I, Zhang W, Yang Z, Ding B, Zhu H, Liu Y, Qui Y, Yin Y, Wu G (2012) Protective effects of N- acetylcysteine on intestinal functions of piglets challenged with lipopolysaccharide. *Amino Acids* 43:1233-1242.
- Huwig A, Freimund S, Kappeli O, Dutler H (2011) Mycotoxin detoxication of animal feed by different adsorbents. *Toxicol Lett* 122:179-188.
- Kenawy AM, El-Genaidy HM, Authman MMN, Abdel-Wahab MA (2009) Pathological studies on effects of aflatoxin on *Oreochromis niloticus* with application of different trials of control. *Egypt J Comp Path & Clinic Path* 22:175-193.
- Kim YS, Ho SB (2010) Intestinal goblet cells and mucins in health and disease: recent insights and progress. *CurrGastroenterol Rep* 12:319-330.
- Lea H, Spring P, Taylor-Pickard J, Burton E (2013) A natural carbohydrate fraction Actigen™ from *saccharomyces cerevisiae* cell wall: effects on goblet cells, gut morphology and performance of broiler chickens. *J ApplAnimNutr* 1:1-7.
- Leknes LI (2010) Histochemical study on the intestinal goblet cells in cichlid and poeciliid species (Teleostei). *Tissue Cell* 42:61-64.
- Liu YL, Meng GQ, Wang HR, Zhu HL, Hou YQ, Wang WJ, Ding BY (2011) Effect of three mycotoxin adsorbents on growth performance, nutrient retention and meat quality in broilers fed on mould-contaminated feed. *Brit. PoultSci* 52:255-263.
- Machado-Neto R, Pontin MCF, Nordi WM, Lima AL, Moretti DB (2013) Goblet cell mucin distribution in the small intestine of newborn goat kids fed lyophilized bovine colostrum. *LivestSci* 157:125-131.
- Magnoli AP, Monge MP, Miazzo RD, Cavaglieri LR, Magnoli CE, Merkis CE, Cristofolini AL, Dalcero AM, Chiacchiera SM (2011) Effect of low levels of aflatoxin B1 on performance, biochemical parameters, and aflatoxin B1 in broiler liver tissues in the presence of monensin and sodium bentonite. *PoultSci* 90:48-58.
- Masanetz S, Wimmer N, Plitzner C, Limbeck E, Preibinger W, Pfaffl MW (2010) Effect of inulin and lactulose on the intestinal morphology of calves. *Animal* 4:739-744.
- Mendoza AH, Cordova AFG, Cordoba BV, Gracia HS (2011) Effect of oral supplementation of *Lactobacillus reuteri* in reduction of intestinal absorption of aflatoxin B1 in rats. *J. Basic Microbiol* 51:263-268.
- Morales-Lopez R, Auclair E, Van Immerseel F, Ducatelle R, Garcia F, Brufau J (2010) Effects of different yeast cell wall supplements added to maize- or wheat-based diets for broiler chickens. *Brit PoultSci* 51:399-408.
- Motawe HFA, Abdel Salam AF, El Meleigy KM (2014) Reducing the toxicity of aflatoxin in broiler chickens diet by using probiotic and feed. *Int J PoultSci* 13:397-407.
- Muthusamy N, Halder S, Ghosh T, Bedford M (2012) Effects of hydrolysed *saccharomyces cerevisiae* yeast and yeast cell wall components on live performance, intestinal histo-morphology and humoral response of broilers. *Brit PoultSci* 52:694-703.
- O,Handley RM, Buret AG, McAllister TA, Jelinski M, Olson ME (2001) Giardiasis in dairy calves: effects of fenbendazole treatment on intestinal structure and function. *Int J Parasitol* 31:73-79.
- Oguz H, Kurtoglu V, Coskun B (2000) Preventive efficacy of clinoptilolite in broiler during chronic aflatoxin (50 and 100 ppb) exposure. *Res Vet Sci* 69:197-201.
- Quinteiro-Filho WM, Ribeiro A, Ferraz-de-Paula V, Pinheiro ML, Sakai M, Sa LR, Ferreira AJ, Palermo-Neto J (2010) Heat stress impairs performance parameters, induces intestinal injury and decreases macrophage activity in broiler chickens. *PoultSci* 89:1905-1914.
- Rashid N, Bajwa MA, Rafeeq M, Khan MA, Ahmad Z, Tariq MM, Wadood A, Abbas F (2012) Prevalence of aflatoxin B1 in finished commercial broiler feed from west central Pakistan. *J Anim Plant Sci* 22:6-10.
- Schatzmayer G, Heidler D, Fuchs E, Nitsh S, Mohni M, Taubel M, Loibner AP, Braun R, Binder EM (2003) Investigation of different yeast strains for the detoxification of ochratoxin A. *Mycotoxin Res* 19:124-128.
- Shen YB, Piao XS, Kim SW, Wang L, Liu P, Yoon I, ZhenYG (2009) Effects of yeast culture supplementation on growth performance, intestinal health and immune response of nursery pigs. *J AnimSci* 87:2614-2624.
- Shepherd GS, Sydenham EW, Thiel PG, Gelderblom WGA (1990) Quantitative determination of fumonisins B1 and B2 by high performance liquid chromatography with fluorescence detection. *J LiqChromatogr* 13:2077-2087.
- Sirajudeen M, Gopi K, Tyagi JS, Moudgal RP, Mohan J, Singh R (2009) Protective effects of melatonin in reduction of oxidative damage and immunosuppression induced by aflatoxin B1 contaminated diets in young chicks. *Environ Toxicol* 26:153-160.
- Solis de los Santos F, Donoghue AM, Farnell MB, Huff GR, Huff WE, Donoghue DJ (2007) Gastrointestinal maturation is accelerated in turkey poults supplemented with a mannan-oligosaccharide yeast extract (alphanune). *PoultSci* 86:921-930.
- Sur E, Celik I (2003) Effects of aflatoxin B1 on the development of bursa of fabricius and blood lymphocyte acid phosphatase of the chicken. *Brit PoultSci* 44:558-566.
- Suzuki I, Tanaka H, Kinoshita A, Oikawa S, Osawa M, Yadamoe T (1990) Effect of orally administered beta-glucan on macrophage function in mice. *Int J Immunopharmacol* 12:675-684.
- Tessari ENC, Oliveria, CAF, Cardoso ALSP, Ledoux, DR, Rottinghaus GE (2006). Effects of aflatoxin B1 and fumonisin B1 on body weight, antibody titers and histology of broiler chicks. *Brit PoultSci* 47:357-364.
- Tsukada C, Yokoyama H, Miyaji C, Ishimoto Y, Kawamura H, Abo T (2003). Immunopotential of intraepithelial lymphocytes in the intestine by oral administrations of  $\beta$ -glucan. *Cell Immunol* 221:1-5.
- Volman JJ, Ramakers JD, Plat J (2008) Dietary modulation of immune function by  $\beta$ -glucans. *PhysiolBehav* 94:276-284.
- Wan XL, Yang ZB, Yang WR, Jiang SZ, Zhang GG, Johnston SL, Chi F (2013) Toxicity of increasing aflatoxin B1 concentrations from contaminated corn with or without clay adsorbent supplementation in ducklings. *PoultSci* 92:1244-1253.
- Wang YH, Xu M, Wang FN, Yu ZP, Yao JH, Zan LS, Yang FX (2009) Effect of dietary starch on rumen and small intestine morphology and digesta pH in goats. *LivestSci* 122:48-52.
- Wu QJ, Wang LC, Zhou YM, Zhang JF, Wang T (2013) Effects of clinoptilolite and modified clinoptilolite on the growth performance, intestinal microflora and gut parameters of broilers. *PoultSci* 92:684-692.
- Wu YB, Ravindran V, Thomas DG, Britles MJ, Hendricks WH (2004) Influence of method of whole wheat inclusion and xylanase supplementation on the performance, apparent metabolizable energy, digestive tract measurements and gut morphology of broilers. *Brit PoultSci* 45:385-394.
- Yang J, Bai F, Zhang K, Lv X, Bai S, Zhao L, Peng, X, Ding X, Li Y, Zhang J (2012) Effects of feeding corn naturally contaminated with AFB1 and AFB2 on performance and aflatoxin residue in broilers. *Czech J AnimSci* 57:506-515.
- Yunus AW, Fazeli ER (2011) Aflatoxin B1 in affecting broiler's performance, immunity, and gastrointestinal tract: a review of history and contemporary issues. *Toxins* 3:566-590.