Journal of the Hellenic Veterinary Medical Society

Vol 69, No 2 (2018)

Granulosa Cell Tumor In A Spayed Young Queen

Z. GÜNEY UÇMAK, M. UÇMAK, Ç. TEK, L. KOENHEMSI, Ö. ERDOĞAN BAMAÇ, A. GÜREL, E. YILDAR

doi: [10.12681/jhvms.18022](https://doi.org/10.12681/jhvms.18022)

Copyright © 2018, Z. GÜNEY UÇMAK, M. UÇMAK, Ç. TEK, L. KOENHEMSI, Ö. ERDOĞAN BAMAÇ, A. GÜREL, E. YILDAR

This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

To cite this article:

GÜNEY UÇMAK, Z., UÇMAK, M., TEK, Ç., KOENHEMSI, L., ERDOĞAN BAMAÇ, Ö., GÜREL, A., & YILDAR, E. (2018). Granulosa Cell Tumor In A Spayed Young Queen. *Journal of the Hellenic Veterinary Medical Society*, 69(2), 1010-1015. <https://doi.org/10.12681/jhvms.18022>

Granulosa Cell Tumor In A Spayed Young Queen

Z. Günay Uçmak^{1,*}, M. Uçmak¹, Ç. Tek¹, L. Koenhems²,
Ö. Erdoğan Bamaç³, A. Gürel³, E. Yıldar⁴

¹ Department of Obstetrics and Gynaecology, Faculty of Veterinary Medicine, Istanbul University, Istanbul, Turkey

² Department of Internal Medicine, Faculty of Veterinary Medicine, Istanbul University, Istanbul, Turkey

³ Department of Pathology, Faculty of Veterinary Medicine, Istanbul University, Istanbul, Turkey

⁴ Institute of Health Sciences, Istanbul University, Istanbul, Turkey

ABSTRACT. A four years old cat was presented to Clinic of Obstetrics and Gynaecology with complaints of weakness, inappetency, vomiting and estrus signs although it was spayed. Blood tests, radiography and ultrasonography revealed abdominal mass and uterine stump which were then removed surgically. Multilobular mass was defined as solid granulosa cell tumor (GCT). Increase of estrogen (E2) and insulin-like growth factor-1 (IGF-1) values were detected on the 10th postoperative day. On the 40th postoperative day, the cat was brought to Internal Medicine Clinic with the complaints of weakness, inappetency and cachexia. Anemia, leucocytosis, uremia, hyperglycemia, sensitiveness and pain in the right abdomen were determined. A tumor was detected in the liver by radiography and ultrasonography and was suspected to be GCT metastasis. Despite medical therapy, the cat died after four months.

In conclusion; retained ovarian tissue after erroneous ovariectomy may cause, regular estrus signs and GCT development. Even if GCTs are removed by surgical approach, they have metastatic potential that deteriorates the prognosis. Evaluating IGF-1 and E2 in the short postoperative term are beneficial for determining the metastatic potential of GCTs.

Keywords: Granulosa cell tumor, IGF-1, estrogen, ovary, queen.

Corresponding Author:

Z. Günay Uçmak, PhD. DVM. Doğum ve Jinekoloji Anabilim Dalı,
Veteriner Fakültesi, İstanbul Üniversitesi, 34320,
Avcılar, İstanbul, Türkiye.
vethekezeynep86@gmail.com

Date of initial submission: 4-4-2017
Date of revised submission: 2-8-2017
Date of acceptance: 21-8-2017

CASE HISTORY

A four-years-old female cat, that was spayed when was one year old at a private clinic, was brought to Obstetrics and Gynaecology Clinic with regular estrus signs for three years. The cat had received a treatment because of vomiting, anorexia and weakness for 15 days beforehand in a proprietary clinic.

Complete blood cell count (CBC), ALT (Alanine aminotransferase), AST (Aspartate aminotransferase), ALP (Alkaline phosphatase), GGT (Gamma glutamyl transpeptidase), urea, creatinine, glucose, cholesterol, IGF-1, E2, progesterone (P4) analysis, thoracoabdominal radiography and abdominal ultrasonography were performed. Blood parameter results on operation day are notified in Table 1.

On ultrasonographic examination, a mass was viewed behind the left kidney (Figure 1a.,1b.) and uterine stump was determined. On radiography there was no visible metastasis in thorax; however the abdominal mass was clearly seen (Figure 2). A surgical approach was decided.

The anesthesia protocol consisted of sc atropine (Atropin, Vetaş, Turkey) as premedication (0.005 mg/kg) followed by the induction of the anesthesia 15 min. later with iv propofol (Pofol 1%, Dongkook Pharm, Korea) (6 mg/kg), and the intubation and maintenance with 3% isoflurane (Forane likid, Abbott Laboratories, England). Uterine stump (Figure 3) and a mass sized 12x10x5 cm which was located behind left kidney, around the left ovary, were removed by

Table 1. Blood parameter results on operation day and postoperative 10th, 40th and 45th day.

Blood Parameters	Reference Ranges In Cats	Operation Day	Postoperative 10th Day	Postoperative 40th Day	Postoperative 45th Day
RBC	5.8-11 uL*	4.5	6.84	4.5	3.6
HGB	8.6-16 g/dl*	6.5	9.6	6.3	4.6
HCT	28-47%*	20	31	18	14
WBC	(3.7-20.5) x1000/uL*	51.5	14.7	55.8	51.6
PLT	(160-660) x1000/uL*	494	386	95	157
MCV	37.7-50 fL*	45	46	39	40
MCH	12.3-17.2 pg*	15	14	14	13
MCHC	(31.1-36) x10 g/L*	32	31	36	32
GLU.	56-153 mg/dL*	224	164	170	190
UREA	18-36 mg/dL*	71	57	56	61
CREA.	0.6-2 mg/dL*	0.7	0.6	0.6	0.4
AST	14-54 U/L*	17	34	22	19
ALT	26-128 U/L*	26	51	28	20
ALP	14-102 U/L*	90	41		
GGT	0-5 U/L*	1	1		
Cholesterol	71-218 mg/dL*	134	165		
IGF-1	5-70 nmol/L**	88	118		
E2 Base level	5-14 pg/ml***	56.2	67.1		
In estrus	50-70pg/ml***				
P4 Luteal phase	1-4 ng/ml***	0.48	0.37		
In estrus	< 1ng/ml***				

RBC= Red Blood Cell Count; HCT= Hematocrit; HGB= Hemoglobin; MCV= Mean Corpuscular Volume; MCH= Mean Corpuscular Hemoglobin; MCHC= Mean Corpuscular Hemoglobin Concentration; WBC= White Blood Cell Count; PLT = Platelet Count; CREA= Creatinine; GLU= Glucose; AST= Aspartate Aminotransferase; GGT= Gamma Glutamyltransferase; ALP= Alkaline Phosphatase; ALT= Alanine aminotransferase; IGF-1= Insulin-like growth factor-1; E2= Estrogen; P4= Progesterone. Reference values: *Plumb (2008), **Maden and Çuhadar (2013), ***Shille et al., (1979).

Figure 1a. The mass behind left kidney on cranoabdominal regio, **1b** Tumoral mass on ultrasonographic examination.

Figure 2. Tumoral area on thoraco-abdominal radiography of the cat.

Figure 3. Uterine stump

median laparotomy. Before laparotomy incision was closed, we were sure that there was no residual ovarian tissue in the abdomen. Monofilament and absorbable suture material (Monocryl No: 2/0, Medeks, Turkey) was used for ligations, muscular, subdermal and dermal sutures. The mass and stump uterus were delivered to Pathology Department for histopathologic examination. Twenty ml/kg iv 0.9 % isotonic sodium chloride twice a day (b.i.d), 20 ml/kg iv Lactated Ringer (Laktatlı Ringer, Polifarma, Turkey) b.i.d., 40 µg/kg intramuscular (im) vitamin B12 (Dodex, Deva, Turkey) once daily (s.i.d.), B and C complex vitamin (Hepargriseovim, Deva, Turkey), 20 mg/kg im ampicillin sodium (Ampisina, Mustafa Nevzat, Turkey) b.i.d. and 15 mg/kg im clindamycin (Klindan, Bilim, Turkey) b.i.d. were administrated to the cat for ten days postoperatively. Also inactivated paravoxvirus ovis (Zylexis, Pfizer, USA) was injected once a week for four weeks by subcutaneous way.

The first control was made on the 10th postoperative day. On anamnesis; having appetite, absence of vomiting and vitality were informed. Blood parameter results on the 10th postoperative day are notified in Table1. After isotonic serum infusion and low protein diet for seven days, blood profile was as follows; hematocrit 28%, hemoglobin 8.6 g/dl, glucose 166 mg/dl, urea 45 mg/dl and creatinine 0.7 mg/dl. According to this blood profile, 0.9% isotonic sodium chloride (Izotonik, Eczacıbaşı, Turkey), Lactated Ringer (Laktatlı Ringer, Polifarma, Turkey), B and C vitamin complex (Hepargriseovim, Deva, Turkey) and vitamin B12 (Dodex, Deva, Turkey) were administrated. Three weeks after the prescrip-

Figure 4. Abdominal radiography of the tumor in liver.

Figure 5a,b Hyperechoic areas within liver on ultrasonography.

tion applied, the cat was brought again to Internal Medicine Clinic with the complaint of weakness, anorexia and cachexia. Anemia, leucocytosis, uremia and hyperglycemia were detected on the 40th postoperative day (Table 1). As a result, 20 mg/kg ampicillin sodium and sulbactam combination (Sulcid, İ.E. Ulagay, Turkey), 15 mg/kg clindamycin (Klindan,

Figure 6. Tumoral mass in macroscopic examination.

Figure 7a Cystic structures with diffuse, atypical granulosa cells. **7b, 7c** Density of distinctive tubular pattern in stroma. **7d** Cystic glandular metritis.

Bilim, Turkey) and vitamin B12 (Dodex, Deva, Turkey) were prescribed for 5 days. At the end of this therapy, blood tests (Table 1) were repeated. Because signs of pain were detected on palpation of the right abdominal region, abdominal radiography (Figure 4) and ultrasonography (Figure 5a,b) were performed. According to ultrasonographic examination, a tumor on the liver was determined. Because of the poor prognosis, surgery was not performed. Medical therapy with 25 mg/kg im Seftriaxon (Novosef, Zentiva, Turkey), 15 mg/kg im clindamycin (Klindan, Bilim, Turkey), 0.9% iv isotonic sodium chloride (Izotonik, Eczacıbaşı, Turkey) and 5% iv dextrose (Dekstroz, Eczacıbaşı, Turkey) serum infusion, im vitamin B12 (Dodex, Deva, Turkey) and 16 mg im cortisone

(Prednol, Mustafa Nevzat, Turkey) was applied for 7 days. After that last therapy, survival time was only four months.

Histopathologic examination was performed at Department of Pathology. In the macroscopic examination, the tumoral mass of the ovary was measured 12x10x5 cm in size and had multinodular structure (Figure 6). Tumor samples were fixed in 10% neutral buffered formalin for at least 24 h and 5 µm thick serial sections were cut, deparaffinized, rehydrated with water in descending concentrations of ethanol and used for hematoxylin and eosin (HE) staining. Tissues were observed under a light microscope. The histological evaluation revealed diffuse, atypical, granulosa-like cells with spherical to oval, hyperchromatic nuclei, distinct nucleoli, and scant eosinophilic cytoplasm. In the diffuse areas, there were cystic structures surrounded by granulosa cells (Fig 7a). Also, a distinctive tubular pattern in which a dense stroma was present (Fig 7b, c). In some slides, areas of luteinisation and hemorrhage were seen. Cystic glandular metritis was also observed in the same patient (Fig 7d).

DISCUSSION

GCTs are the most common sex-cord stromal tumor in all animal species (Nielsen and Kennedy, 1990). Holzworth (1987) reported that mean age of GCT occurrence in cats is 11 years. Contrarily, reported cat which affected with GCT was four years old.

In our case, GCT was detected in a spayed cat as Spoor et al. (2014) detected GCT in spayed Bulldog. In accordance with Gelberg and McEntee (1985), the cat had also estrus signs although spayed in early age. In line with Ball et al. (2010), it is thought to be as a result of incomplete excised ovarian or ectopic ovarian tissues.

Gelberg and McEntee (1985) reported that GCTs are the most common ovarian tumor type in cats which have malignant character. The appearance of the tumor in liver 45 days after the ovarian tumor resection supports GCTs' malignancy as Gelberg and McEntee (1985) reported.

In accordance with Gündüz et al. (2010) ultrasonographic imaging of the ovaries is used for diagnosis of GCT. In line with Hayes and Harvey (1979),

we performed surgery for treatment and also after surgery immunotherapeutic agent was used for four weeks.

Noakes (2009) reported that ovariectomy cause elimination of E2, P4, inhibin, activin, and follistatin in ovaries. Shille et al. (1979) notified that E2 in plasma elevate from a base level of 5-14 pg/ml to peaks of 50-70 pg/ml in estrus. Contradicting to Noakes (2009), in this report, we expected base level in plasma E2 after the GCT resection but E2, which was measured 56 pg/ml in the operation day, increased to 62 pg/ml on the 10th postoperative day.

Fontbonne et al. (2007) reported that ovarian remnant syndrome (ORS) leads to elevated levels of ovarian hormones. Shanbhogue et al. (2010) notified that hyperandrogenism and hyperestrogenism are the most common causes of ovarian tumors. Although we had no data about the E2 levels of the erroneous ovariohysterectomized cat before the GCT diagnosis, in line with the researchers (Fontbonne et al. 2007, Shanbhogue et al. 2010) this ORS case was followed by GCT formation.

Buijtelts et al. (2010) reported that GCTs produce E2 and small amount of P4 which causes endometrial growth and glandular secretion. In accordance with Buijtelts et al (2010), the cat with GCT had high E2 level and also cystic glandular metritis in the uterine stump.

Boormann (2002) notified that increase of E2 levels after surgery happen if a residual hormone producing tumor may exist. Increased E2 levels detected on the 10th postoperative day and presence of tumor on liver 45 days after surgery are in accordance with Boormann (2002).

In accordance with Spicer and Echternkamp (1995), that reported a positive correlation between follicular growth and IGF-1 concentration, IGF-1 concentration was measured high on the operation day in this case.

Despite the resection of GCT and uterine stump, the increased IGF-1 concentration on the 10th postoperative day may be related to IGF-1 synthesis by the metastatic liver as reported (Maden and Çuhadar, 2013).

The histopathological findings of this granulosa tumour did not differ from the literature and they

are in accordance with the aggressive behaviour described for this tumour in cats (Giacóia et al., 1999).

A second operation for tumor resection in liver was not performed due to the poor body condition. It is in accordance with Boormann's (2002) report which defends that second-look operations are unnecessary unless significant masses and complications such as obstruction of the bowel occur.

In conclusion; retained ovarian tissue after erro-

neous ovariectomy may cause regular estrus signs and GCT development. Even if GCTs are removed by surgical approach, they have metastatic potential that deteriorates the prognosis. Evaluating IGF-1 and E2 in the short postoperative term are beneficial for determining the GCT metastasis.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

- Ball RL, Birchard SJ, May JR, Threlfall WR, Young GS (2010) Ovarian remnant syndrome in dogs and cats: 21 cases (2000-2007). *J Am Vet Med Assoc* 236:548-553.
- Boormann EV (2002) Granulosa cell tumors. In: *Proceedings of UCLA Health Care* 6:1-6. <http://www.med.ucla.edu/modules/xfsection/article.php?articleid=177>
- Buijts JJCWM, De Gier J, Kooistra HS, Veldhuis Kroeze EJB, Okkens AC (2010) Alterations of the pituitary-ovarian axis in dogs with a functional granulosa cell tumor. *Theriogenology* 73:11-19.
- Fontbonne A, Levy X, Fontaine E, Gilson C (2007) Guide pratique de reproduction clinique canine et feline. Med'com.
- Gelberg HB, Mc Entee K (1985) Feline ovarian neoplasms. *Vet Pathol* 22:572-576.
- Giacóia MR, Maiorka PC, Oliveira CM, Sinhorini IL, Daglı MLZ (1999) Granulosa cell tumour with metastasis in a cat. *Braz J Vet Res Anim Sci* 36(5).
- Gündüz MC, Kaşıkçı G, Kılıçarslan MR, Uçmak M, Düzgün O, Tek Ç (2010) Reproductive performance following unilateral ovariectomy for treatment of ovarian tumors in 7 mares. *Turk J Vet Anim Sci* 34(3):283-287.
- Hayes A, Harvey HJ (1979) Treatment of metastatic granulosa cell tumor in a dog. *JAVMA* 174(12):1304-1306.
- Holzworth J (1987) Tumours of the female reproductive system In: *Diseases of the cat: medicine and surgery*. 1st. ed., W.B. Saunders, Philadelphia: pp 520-521.
- Maden M, Çuhadar F (2013) Kedi ve köpeklerde endokrin hastalıkların laboratuvar tanısı. *Türkiye Klinikleri J Vet Sci* 4(3): 35-60.
- Nielsen SW, Kennedy PC (1990) Tumours of the genital systems In: *Tumours in domestic animals*. 3rd. ed., Berkeley University of California Press, California: pp 458-475.
- Noakes D. (2009). Endogenous and exogenous control of ovarian cyclicity In: *Veterinary reproduction and obstetrics*. 9th ed., Pa: Saunders, Philadelphia: pp 6-9.
- Shanbhogue AKP, Shanbhogue DKP, Prasad SR, Surabhi VR, Fasih N, Menias CO (2010) Clinical Syndromes Associated with Ovarian Neoplasms: A Comprehensive Review. *RadioGraphics* 30(4): 903-919
- Shille VM, Lundström KE, Stabenfeldt GH (1979) Follicular function in domestic cat as determined by estradiol-17 β concentrations in plasma: relation to estrous behaviour and cornification of exfoliated vaginal epithelium. *Biology of Reproduction* 23: 953-963.
- Spicer LJ, Echternkamp SE (1995) The ovarian insulin and insulin-like growth factor system with an emphasis on domestic animals. *Domestic Animal Endocrinology* 12: 223-245.
- Spoor MS, Flesner BK, Trzil JE, Whitney MS, Shaw DP, Selting KA (2014) What is your diagnosis? Intra-abdominal mass in female spayed dog. *Vet Clin Pathol* 43(1): 109-110.