



Journal of the Hellenic Veterinary Medical Society

Vol 70, No 1 (2019)



To cite this article:

SHEIKHANI-SHAHIN, H., MEHRABANI, D., ASHRAF, M. J., RAJABI, H., NOROUZIAN, M., RAHMANIFAR, F., DEHGHANI NAZHVANI, S., & ZARE, S. (2019). The healing effect of bone marrow-derived stem cells and aquatic activity in Achilles tendon injury. *Journal of the Hellenic Veterinary Medical Society*, *70*(1), 1373–1380. https://doi.org/10.12681/jhvms.20342

The healing effect of bone marrow-derived stem cells and aquatic activity in Achilles tendon injury

H. Sheikhani-Shahin¹, D. Mehrabani^{2,3,4,5*}, M.J. Ashraf⁶, H. Rajabi⁷, M. Norouzian⁷,

F. Rahmanifar⁸, S. Dehghani Nazhvani⁹, S. Zare³

¹Department of Physical Education, Zand Institute of Higher Education, Shiraz, Iran

²Burn and Wound Healing Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

³Stem Cell Technology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

⁴Comparative and Experimental Medicine Center, Shiraz University of Medical Sciences, Shiraz, Iran

5Rohan Gene Cell Tech, Shiraz, Iran

⁶Department of Pathology, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

⁷Department of Exercise Physiology, Faculty of Physical Education and Sport Sciences, Kharazmei University, Tehran, Iran

⁸Department of Anatomy, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

⁹Department of Surgery, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

ABSTRACT. The common treatment recommended for Achilles tendon rupture as the most common tendon injury during exercise is surgical intervention, while it eventually causes various clinical problems. This study assessed the healing effect of bone marrow-derived stem cells (BMSCs) and aquatic activities in Achilles tendon injury. Forty rats were randomly divided into 5 equal groups. Group 1 underwent aquatic activity, 72 h after a crush lesion formed on Achilles tendon, group 2 received 1×106 intra-articular BMSCs post-tendon injury, group 3 had aquatic activity together with BMSCs transplantation after tendon damage, group 4 just experienced tendon injury without any treatment intervention and group 5 was considered as the control group while did not undergo any tendon injury and did not receive any treatment measure. After 8 weeks, the animals were sacrificed and the tendons were transferred in 10% formalin for histological evaluation. There was a significant increase in fibroblast number in group 3 in comparison to other groups. However, there was a significant increase in collagen deposition in groups 2, 3 and 5 in comparison to group 1 and 4. A significant decrease was noted for cellularity in group 2 when compared to groups 1 and 4. Regarding tendon diameter in group 3; a significant healing was observed when compared to groups 2, 4 and 5. It was shown that aquatic activity together with cell transplantation was an effective therapeutic measure enhancing the healing in tendon injuries.

Keywords: Aquatic activity, Bone marrow, Mesenchymal stem cells, Healing, Tendon.

Corresponding Author:

Date of initial submission: 25-03-2018 Date of revised submission: 25-03-2018 Date of acceptance: 30-02-2018

Davood Mehrabani, Assistant Professor of Stem Cell Technology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran. Email address: mehrabad@sums.ac.ir

INTRODUCTION

The common treatment recommended for Achilles tendon rupture as the most common tendon injury during exercise is surgical intervention, while it eventually causes various clinical problems (Rodeo et al., 2010). During the recent decade, it was shown that biological factors may accelerate tendon healing in damaged tendons (Rajabi et al., 2015). Although many studies have indicated that exercising and physical activity is among effective factors in increasing tendon strength, the question whether physical activity after the surgery could accelerate the recovery of tendons biomechanical properties and lead to fundamental changes in healing process have still remained controversial (Rajabi et al., 2015).

Several researchers have mentioned the significant role of mesenchymal stem cells (MSCs) in tendon healing (McGoldrick et al., 2017; Romero et al., 2017). MSCs have been isolated from different tissues including bone marrow (BM) (Aliborzi et al., 2015), adipose tissue (Mehrabani et al., 2015a), menstrual blood (Faramarzi et al., 2016), dental pulp (Mehrabani et al., 2017), and endometrium (Shamosi et al., 2017). They have osteogenic (Shaterzadeh Yazdi et al., 2015), adipogenic (Mehrabani et al., 2015b) and neurogenic (Razeghian Jahromi et al., 2015) differentiation properties with similar features such as favorable proliferative capability, selfrenewal, and differentiation potential (Mehrabani et al., 2013). Bone marrow-derived stem cells BMSCs) have opened a new window in regenerative medicine and cell transplantation purposes getting help from tissue engineering too (Hosseinkhani et al., 2014).

The beneficial effects of MSCs in healing of Achilles tendon injuries were previously reported demonstrating that use of MSCs could improve the biological and mechanical parameters of the tissue (Chong et al., 2007). Therefore, this study was conducted to determine the healing effect of a combination of BMSCs and aquatic activities in Achilles tendon injuries in experimental rat model.

MATERIAL AND METHODS

Forty male Sprague-Dawley rats weighing 200±20 g and aging 3 months were provided from Laboratory Animal Center of Shiraz University of Medical Sciences, Shiraz, Iran and housed in groups of 4 in transparent polycarbonate cages with dimensions of $54 \times 18 \times 18$ cm at an environment of $21\pm 2^{\circ}$ C with the lighting of 12:12 light-to-dark ratio; light at 7:30 AM in 50 ± 5 percent humidity. All procedures were done based on laws of Animal Care by Iran Veterinary Organization. The rats were randomly divided into 5 equal groups.

Group 1 underwent aquatic activity, 72 h after a crush lesion formed on Achilles tendon using a mosquito hemostat (Akinbo et al., 2008), group 2 received 1×106 intra-articular BMSCs post-tendon injury, group 3 had aquatic activity together with BMSCs transplantation after tendon damage, group 4 just experienced tendon injury without any treatment intervention and group 5 was considered as the control group while did not undergo any tendon injury and did not receive any treatment measure.

In order to prepare BMSCs, 5 rats were euthanized and under sterile conditions, femoral and tibial bones were removed while, the muscular and connective tissues were later detached. Under sterile conditions, both ends of the bones were cut and the BM was flushed out using a 10 ml syringe filled with Dulbecco's Modified Eagle Medium (DMEM; Biovet, Bulgaria) and 1% penicillin streptomycin (Sigma, USA). BM was then transferred on ice to the stem cell laboratory, Shiraz University of Medical Sciences, Shiraz, Iran.

The BM was diluted in an equal volume of DMEM, and centrifugation was undertaken at 1200 rpm for 7 minutes. After removal of the supernatant, the precipitate was transferred in 25 cm2 flasks containing DMEM supplemented with 10% fetal bovine serum (FBS; Biovet, Bulgaria), 1% L-glutamine (Sigma, USA) and 1% penicillin and streptomycin and the culture flasks were placed in CO2 incubator at 37°C with 5% CO2 and saturated humidity while the medium was replaced every 3 days. The adherent cells were subcultured at 80% confluency, by washing twice with PBS (Gibco, USA) and adding 0.25% trypsin (Gibco, USA) for 3 minutes. To inactivate the trypsin activity, equal volume of DMEM was added. Cell passage was continued until passage 5 and at each passage, the cells were counted too. Cell morphology was assessed by inverted microscope (Olympus, USA) as described by Gashmardi et al. (2017).

To assess the osteogenic differentiation property, BMSCs from passage 5 were seeded into 6 well plates and at 80% confluency, the cells were cultured for 21 days with low glucose DMEM containing 100 nM dexamethasone (Sigma, USA), 0.051 M ascorbate-2-phosphate (Wako Chemicals, USA), 10 mM b-glycerophosphate (Sigma, USA), 1% penicillin/streptomycin and 10% FBS. The medium was changed every 3 days. After 3 weeks, osteogenic differentiation was evaluated with Alizarin Red (Sigma, USA) staining (Gashmardi et al., 2017).

RT-PCR was done to determine the expression of MSC markers. So after extraction of the total RNA using the column RNA isolation kit (Denazist-Asia, Iran) based on manufacturer's instructions, it was evaluated by spectrophotometry. The complementary DNA (cDNA) was provided by AccuPower Cycle Script (RT PreMix Kit Bioneer, Korea) upon manufacturer's instruction. For each reaction; 15 μ L of total RNA was used to reach a volume of 20 μ L with the DEPC water. Twelve thermal cycles were conducted including 30 s at 20°C for primer annealing, 4 min at 42°C for cDNA synthesis, and 30 s at 55°C for melting secondary structure and cDNA synthesis and 5 min at 95°C for inactivation (Rahmanifar et al., 2016).

Then, 1 µL of template cDNA and PCR buffer, H2O, dNTPs, MgCl2, Taq DNA polymerase, and forward and reverse primers were mixed. The microtubules containing 20 µL of the mentioned mixture were transferred to a thermocycler (Eppendorf Mastercycler Gradient, Eppendorf, Hamburg, Germany) and 30 amplification cycles were performed (30 s denaturation at 95°C, 30 s annealing at 64°C, 62°C, and 61°C and 30 s extension at 72°C with the 5 min at 95°C for primary denaturation and 5 min at 72°C for final extension). PCR products were assessed for defined bands by gel electrophoresis by DNA safe stain in 1.5% agarose gel medium. The bands were seen by UV radiation and a gel documentation system (UVtec, Cambridge, UK) and photography was done.

Aquatic activity was undertaken for 8 weeks (five sessions per week). For adaptation, rats were put into a glass aquarium with the dimensions of $50 \times 40 \times 50$ cm filled with 34°C water for 5 minutes for 3 days. Then the aquatic activity was extended for 8 weeks

5 days per week (5-30 minutes each session, Table 1) as described by Hart et al. (2001) and McVeigh et al. (2010). After 8 weeks, the animals were euthanized and sacrificed and the Achilles tendon was removed and

Table 1. Eight-week incremental aquatic activity pro-gram of studied rats.

Week	Stage				
	Introduction stage: In this stage, to have				
	optional activity, rats were put into a glass				
	aquarium				
1	With the dimensions of $50 \times 40 \times 50$ cm filled				
2	with 34°C water for 5 minutes for 3 days.				
3	Aquatic activity: 5 min: Overload Stage:				
4	days 1 to 5				
5	days 1 to 5: Aquatic activity: 10 min				
6	days 1 to 5: Aquatic activity: 15 min				
7	days 1 to 5: Aquatic activity: 20 min				
8	days 1 to 5: Aquatic activity: 25 min				
	days 1 to 5: Aquatic activity: 25 min				
	days 1 to 5: Aquatic activity: 30 min				
	days 1 to 5: Aquatic activity: 30 min				

transferred into 10% formalin buffer for histological evaluation using hematoxylin and eosin (H&E) staining. The studied factors in histological evaluation were the number of fibroblasts, collagen deposition, cellularity and tendon diameter. For statistical analysis, SPSS software (version 16, Chicago, IL, USA) was used applying Kolmogorov Smirnov, one-way ANOVA and Tukey post hoc tests. The significance level was considered a p value less than 0.05.

RESULTS

BMSCs were plastic adherent and fibroblast-like throughout all passages (Figure 1A-C). Culture of BMSCs in osteogenic media resulted into osteogenic differentiation of these cells due to presence

Figure 1. BMSCs were fibroblast-like throughout all passages. A: Passage 1, B: Passage 2 and C: Passage 3.



of calcium deposits after three weeks verified by Alizarin Red staining (Figure 2A, 2B). They showed

Figure 2. The osteogenic differentiation of BMSCs by Alizarin Red staining. A: Control, B: Osteogenic induction). C: The positive expression of CD73 marker confirming the mesenchymal stem cell property and negative expression of CD45 for markers of hematopoietic stem cells.



positive expression of CD73 marker confirming the mesenchymal stem cells and absence of CD34 and CD45 as markers for hematopoietic stem cells (Figure 2C). Table 2 shows the histological findings denoting to a significant increase in the number of fibroblast in group 3 in comparison to other groups (Figure 3A). A significant increase was visible in collagen deposition in groups 2, 3 and 5 in comparison to group 1 (p=0.001) and 4 (p=0.001,

Figure 3. The mean and standard deviation of Fibroblast numbers (A), Collagen deposition (B), Cellularity (C) and Tendon diameter (D) were shown in different groups. *P<0.05.



Figure 3B). There was a significant decrease in cellularity in group 2 compared to groups 1 (p=0.009) and 4 (p=0.04, Figure 3C).

Regarding tendon diameter revealing healing process, in the group receiving BMSCs and undergoing aquatic activity, an increase in tendon diameter was seen compared to the group just underwent cell transplantation (p=0.04) and the group with tendon injury and any treatment measure (p=0.01) and the control group with no tendon injury (p=0.04, Figure 3D). Figure 4 shows a moderate healing in Achilles tendon with collagen deposition in group 3.

Figure 4. Moderate healing in Achilles tendon with collagen deposition.



It shows that BMSCs together with aquatic activity could increase the collagen deposition compared to other groups while the difference was statistically significant too.

DISCUSSION

Our findings suggested that transplantation of BMSCs along with aquatic activities could be effective in repair of Achilles tendon injuries. The histological factors in assessment of healing of Achilles tendon injuries showed that the number of fibroblasts in transplantation of BMSCs along with aquatic activities had a statistically significant higher healing property in comparison to other groups. In response to tendon damage, fibroblasts migrated to the inflamed wound area together with type III collagen which later was substituted by type I collagen (Lisa and Zena, 2002). It was shown that the mechanical pressures due to the physical activities affected the collagens homeostasis that increased in the injured tissues, such as tendons and ligaments (Heinemeier, 2007). Injection of BMSCs along

J HELLENIC VET MED SOC 2019, 70(1) ПЕКЕ 2019, 70(1)

Variables		Collagen			
Group	Fibroblasts	deposition	Cellularity	Tendon diameter	Total
1: Aquatic activity	51.25±10.8	1±0.0	2±0.0	1704.78±412.1	1759.03±408.6
2: BMSC transplant	48.75±7.9	2.5±0.75	1.2±0.46	1589.9±307.5	1642.49±308.5
3: BMSC transplant +Aquatic activity	65.25±2.9	2.5±0.53	1.6±0.51	2187.32±689.9	2256.7±690.39
4: Tendon injury without treatment	50.75±6.3	1±0.0	1.87±0.35	1485.97±307.08	1539.6±310.5
5: Control with no tendon injury	51.62±5.2	2.87±0.35	1.5±0.53	1582.33±128.37	1638.33±12.13

Table 2. Mean and standard deviation of effective factors related to the healing of Achilles tendon in studied groups.

with aquatic activities or injection of BMSCs alone was demonstrated to increase collagens deposition significantly in comparison to injury group. Physical activity was shown to increase the number of fibroblasts and as a result, the collagen deposition.

Tendons are considered as facilitators of force transfer and the arrangement of tendon fibers is important in passive forces and pressures absorbing, as tendon strength is dependent on orientation and length of collagen fibers and consistency of transplanted tendons. The collagen and transplanted tissue metabolism was shown to be influenced by the amount of physical activity. Physical activity was demonstrated to increase the differentiation and change of the transplanted tissue in the tendons denoting to the physiological adaptations and repair of the damages on the extracellular matrix based on physical activities (Kjaer, 2004). In our study, injection of cells alone lead to an increase in collagen deposition, which substituted various cellular sources for tendon repair. Similarly, the role of MSCs in tendon repair was shown before, due to their high proliferation capacity and differentiation properties (Harris et al., 2004).

We reported a significant increase in cellularity in

the group undergoing aquatic activity, compared to the control group that underwent just injury. Platelet rich plasma (PRP) was identically used to increase the performance and cellularity of the tendons and the amount of collagens and cellularity that were due to the increase in the metabolic activity and its positive impact on tissue repair (Lane et al., 2013). Our findings are also in accordance with the aforementioned study on the increase in cellularity.

We showed that the tendon diameter in the group undergoing aquatic activity together with cell transplantation revealed a significant increase. Similarly, it was also demonstrated that physical activity on treadmill for a week could increase the size of collagen fibers and tendon diameter in rats (Michna and Hartmann, 1989). In another study, an intense endurance exercise was found to increase in collagen deposition in Achilles tendon (Curwin et al., 1988). The physical exercise was noted to increase several growth factors in Achilles tendon which could stimulate the collagen synthesis and cell proliferation (Wang, 2006).

Rajabi et al. (2015) in their study on the healing effects of aquatic activities and allogenic injection of PRP on injuries of Achilles tendon based on the number of fibroblast, cellular density, collagen deposition, and tendon diameter, showed that aquatic activity together with PRP injection was the therapeutic measure of choice enhances healing in tendon injuries. Godwin et al. in their study on implantation of BMSCs in horse with overstrain injury of the superficial digital flexor tendon showed that cell implantation was safe and appeared to reduce the re-injury rate (Godwin et al., 2012). Lacitignola and colleagues (2008) in cell therapy for tendinitis in horses by BMSCs together with exercise program stated that re-injury rate reduced and good to excellent outcome in term of athletic success were noted.

He et al. (2015) by applying BMSCs revealed attenuated adhesions in the early time point following flexor tendon repair and healing in rabbit flexor tendon. Human adipose stem cells were also shown as another cell source as a potential approach for tendon tissue repair (Vuornos et al., 2016). Transplantation of tendon-derived stem cells pretreated with connective tissue growth factor and ascorbic acid as a different source of cells was found to promote better tendon repair in a patellar tendon window injury rat model (Lui et al., 2016). Umbilical cord blood-derived mesenchymal stem cells in a rabbit model with full-thickness rotator cuff tendon tear demonstrated the effectiveness of local injection of MSCs into the rotator cuff tendon to be a useful conservative treatment for full-thickness rotator cuff tendon tear repair (Park et al., 2015).

CONCLUSION

Identical to many studies that did not report pain for experimental animals, this limitation existed in our study too. We showed that injection of MSCs together with the aquatic activity could promote healing process that may be due to accumulation of monocytes, proliferation and differentiation of stem cells, migration of fibroblasts and angiogenesis. So based on our findings on the number of fibroblast, cellularity, collagen deposition, and tendon diameter, it was shown that aquatic activity together with MSC injection was a therapeutic of choice enhancing the healing of tendon injuries that can open a window in treatment of damages to tendons.

Disclosure of potential conflicts: All authors declare no financial or other conflicts of interest.

ACKNOWLEDGMENTS

The authors thank Office of Vice Chancellor for Research of Shiraz University of Medical Sciences for financial support Grant no. 1394).

CONFLICT OF INTEREST STATEMENT

No potential conflict of interest relevant to this article was reported.

REFERENCES

- Aliborzi G, Vahdati A, Hossini SE, Mehrabani D (2015) Evaluation of bone marrow-derived mesenchymal stem cells from Guinea pigs. Onl J Vet Res 19:450-459.
- Akinbo S, Aiyegbusi A, Duru F, Noronha C, Okanlawon A (2008) The efficacy of uktrasound therapy on the acute tendon injury. Internet J Orthop Surg 13:1-5.
- Chong AK, Ang AD, Goh JC, Hui JH, Lim AY, Lee EH, Lim BH (2007) Bone marrow-derived mesenchymal stem cells influence early tendon-healing in a rabbit Achilles tendon model. Bone Joint Surg Am 89:74-81.
- Curwin SL, Vailas AC, Wood J (1988) Immature tendon adaptation to strenuous exercise. J Appl Physiol 65:2297-2301.
- Faramarzi H, Mehrabani D, Fard M, Akhavan M, Zare S, BakhshalizadehSh, Manafi A, Kazemnejad S, Shirazi R (2016) The potential of menstrual blood-derived stem cells in differentiation to epidermal lineage: a preliminary report. World J Plast Surg 5:26-31.
- Gashmardi N, Hosseini SE, Mehrabani D, Edalatmanesh MA, Khodabandeh Z (2017) Impacts of bone marrow stem cells on caspase-3 levels after spinal cord injury in mice. Iran J Med Sci 42:593-598.
- Godwin EE, Young NJ, Dudhia J, Beamish IC, Smith RK (2012) Implantation of bone marrow-derived mesenchymal stem cells demonstrates improved outcome in horses with overstrain injury of the superficial digital flexor tendon. Equine Vet J 44:25-32.
- Harris MT, Butler DL, Boivin GP (2004) Mesenchymal stem cells used for rabbit tendon repair can form ectopic bone and express alkaline phosphatase activity in constructs. J Orthop Res 22:998-1003.
- Hart KJ, Shaw JM, Vajda E, Hegsted M, Miller SC (2001) Swim-trained rats have greater bone mass, density, strength, and dynamics. J Appl Physiol 91:1663-1668.
- He M, Gan AW, Lim AY, Goh JC, Hui JH, Chong AK (2015) Bone marrow derived mesenchymal stem cell augmentation of rabbit flexor tendon healing. Hand Surg 20:421-429.
- Heinemeier KM (2007) Adaptation of tendon and muscle connective tissue to mechanical loading. Ph.D. thesis.
- Hosseinkhani M, Mehrabani D, Karimfar MH, Bakhtiyari S, Manafi A, Shirazi R (2014) Tissue engineered scaffolds in regenerative medicine. World J Plast Surg 3:3-7.
- Kjaer M (2004) Role of extracellular matrix in adaptation of tendon and skeletal muscle to mechanical loading. Physiol Rev 84:649-698.
- Lacitignola L, Crovace A, Rossi G, Francioso E (2008) Cell therapy for tendinitis, experimental and clinical report. Vet Res Commun 32:S33-38.
- Lane JG, Healey RM, Chase DC, Amiel D (2013) Use of platelet-rich plasma to enhance tendon function and cellularity. Am J Orthop 42:209-214.
- Lisa M, Zena W (2002) Inflammation and cancer. Nature 420:860-867.
- McGoldrick R, Chattopadhyay A, Crowe C, Chiou G, Hui K, Farnebo S, Davis C, Le Grand A, Jacobs M, Pham H, Chang J (2017) The tissue engineered tendon bone interface: In Vitro and In Vivo synergistic effects of adipo-derived stem cells, platelet rich plasma and extracellular matrix hydrogel. Plast Reconstr Surg 140:1169-1184.
- McVeigh J, Kingsley S, Gray D, Loram LC (2010) Swimming enhances bone mass acquisition in growing female rats. J Sports Sci Med 9:612-619.
- Lui PP, Wong OT, Lee YW (2016) Transplantation of tendon-derived

stem cells pre-treated with connective tissue growth factor and ascorbic acid in vitro promoted better tendon repair in a patellar tendon window injury rat model. Cytotherapy 18:99-112.

- Mehrabani D, Hassanshahi MA, Tamadon A, Zare S, Keshavarz S, Rahmanifar F, Dianatpour M, Khodabandeh Z, Jahromi I, Tanideh N, Ramzi M, Aqababa H, Kuhi-Hoseinabadi O (2015a) Adipose tissue-derived mesenchymal stem cells repair germinal cells of seminiferous tubules of busulfan-induced azoospermic rats. J Hum Reprod Sci 8:103-110.
- Mehrabani D, Mahdiyar P, Torabi K, Robati R, Zare S, Dianatpour M, Tamadon A (2017) Growth kinetics and characterization of human dental pulp stem cells: Comparison between third molar and first premolar teeth. J Clin Exp Dent 9:e172-e177.
- Mehrabani D, Mehrabani G, Zare S, Manafi A (2013) Adipose-derived stem cells ADSC) and aesthetic surgery: a mini review. World J Plast Surg 2: 65-70.
- Mehrabani D, Rahmanifar F, Mellinizhad M, Tamadon A, Dianatpour M, Zare S, RazeghianJahromi I, Ghobadi F (2015b) Isolation, culture, characterization, and adipogenic differentiation of heifer endometrial mesenchymal stem cells. Comp Clin Pathol 24:1159-64.
- Michna H, Hartmann G (1989) Adaptation of tendon collagen to exercise. Int Orthop 13:161-165.
- Park GY, Kwon DR, Lee SC (2015) Regeneration of full-thickness rotator cuff tendon tear after ultrasound-guided injection with umbilical cord blood-derived mesenchymal stem cells in a rabbit model. Stem Cells Transl Med 4:1344-1351.
- Rahmanifar F, Tamadon A, Mehrabani D, Zare S, Abasi S, Keshavarz S, Dianatpour M, Khodabandeh Z, Jahromi IR, Koohi-Hoseinabadi O (2016) Histomorphometric evaluation of treatment of rat azoospermic seminiferous tubules by allotransplantation of bone marrow-derived mesenchymal stem cells. Iran J Basic Med Sci 19:653-661.
- Rajabi H, Sheikhani Shahin H, Norouzian M, Mehrabani D, Dehghani Nazhvani S (2015) The healing effects of aquatic activities and allogenic injection of platelet-rich plasma (PRP) on injuries of Achilles tendon in experimental rat. World J Plast Surg 4:66-73.
- RazeghianJahromi I, Mehrabani D, Mohammadi A, Dianatpour M, Tamadon A, Zare S, Ghahremani Seno M, Khodabandeh Z (2016) The effect of fetal rat brain extract on morphology of bone marrow-derived mesenchymal stem cells. Comp Clin Pathol 25: 343– 349.
- Rodeo SA, Delos D, Weber A (2010) What's new in orthopaedic research. J Bone Joint Surg Am 92:2491-2501.
- Romero A, Barrachina L, Ranera B, Remacha AR, Moreno B, de Blas I, Sanz A, Vázquez FJ, Vitoria A, Junquera C, Zaragoza P, Rodellar C (2017) Comparison of autologous bone marrow and adipose tissue derived mesenchymal stem cells, and platelet rich plasma, for treating surgically induced lesions of the equine superficial digital flexor tendon. Vet J 224:76-84.
- Shamosi A, Mehrabani D, Azami M, Ebrahimi-Barough S, Siavashi V, Ghanbari H, Sharifi E, Roozafzoon R, Ai J (2017) Differentiation of human endometrial stem cells into endothelial-like cells on gelatin/chitosan/bioglassnanofibrous scaffolds. Artif Cells Nanomed Biotechnol 45:163-173.
- ShaterzadehYazdi H, Mehrabani D, KhodakaramTafti A, Dianatpour M, Zare S, Tamaddon A, RazeghianJahromi I (2015) Osteogenic potential of subcutaneous adipose-derived stem cells in a rabbit model. Onl J Vet Res 19:436-445.

J HELLENIC VET MED SOC 2019, 70(1) ПЕКЕ 2019, 70(1) Vuornos K, Björninen M, Talvitie E, Paakinaho K, Kellomäki M, Huhtala H, Miettinen S, Seppänen-Kaijansinkko R, Haimi S (2016) Human adipose stem cells differentiated on braided polylactide scaffolds is a potential approach for tendon tissue engineering. Tissue Eng Part A 22:513-523.

Wang JHC (2006) Mechanobiology of tendon. J Biomech 39:1563-1582.