


Journal of the Hellenic Veterinary Medical Society

Vol 70, No 1 (2019)



Effects of Rapid and Ultra-Rapid Fluid Resuscitation Guided by Blood Lactate Clearance Rate in Diarrheic Calves

H. ERDOGAN, H. VOYVODA

doi: [10.12681/jhvms.20349](https://doi.org/10.12681/jhvms.20349)

Copyright © 2019, H. ERDOGAN, H. VOYVODA



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

To cite this article:

ERDOGAN, H., & VOYVODA, H. (2019). Effects of Rapid and Ultra-Rapid Fluid Resuscitation Guided by Blood Lactate Clearance Rate in Diarrheic Calves. *Journal of the Hellenic Veterinary Medical Society*, 70(1), 1413–1420. <https://doi.org/10.12681/jhvms.20349>

■ Effects of Rapid and Ultra-Rapid Fluid Resuscitation Guided by Blood Lactate Clearance Rate in Diarrheic Calves

H. Erdogan¹, H. Voyvoda¹

¹Department of Internal Medicine, Faculty of Veterinary Medicine, University of Adnan Menderes, Aydin, Turkey

ABSTRACT. The assessment of the results of intravenous fluid treatment in diarrheal calves is difficult under field conditions. To determine the effects of ultra-rapid (1 h) and rapid (3 h) intravenous infusions with 0.9% NaCl and 1.3% NaHCO₃ solutions at a dose of 60 ml/kg on lactate clearance (LC) in calves with diarrhea. Sixty calves, including a healthy control group (n=20) and a group of calves with diarrhea (n=40), were used. Diarrheic calves were divided into two groups (n=20) according to solution type, and each group was then divided into two equal subgroups according to the infusion rate (n=10). Clinical and laboratory inspections of diarrheic calves were performed pre- and post-infusion, and the healthy control group was examined once. LC rates were calculated in the 1 h and 3 h subgroups. Marked improvements in clinical findings related to dehydration were observed in all groups with neonatal diarrhea that were given intravenous infusions with 0.9% NaCl and 1.3% NaHCO₃ solutions. End of the infusion, the LC of the 0.9% NaCl and 1.3% NaHCO₃ solutions in the ultra-rapid groups was significantly increased by 36.4% and 31.8%, respectively. However, for rapid infusion of the same solutions, 13.6% and 31.8% increases were observed, but the differences were not significant. Under field conditions, the LC of L- and D-lactate varied with the infusion rate, and these variations were significant in ultra-rapid subgroups for both solutions. Further studies will be designed for fluid therapy in calves based on the calculation of LC.

Keywords: Calf diarrhea, blood lactate clearance, rapid infusion, sodium bicarbonate, sodium chloride.

Corresponding Author:

Hasan Erdogan, Department of Internal Medicine, Faculty of Veterinary Medicine,
Aydin Adnan Menderes University, Aydin, Turkey
Email address: hasan.erdogan@adu.edu.tr

Date of initial submission: 09-06-2018
Date of revised submission: 09-06-2018
Date of acceptance: 19-07-2018

INTRODUCTION

Major challenges in calf husbandry tend to occur in the neonatal period (0-28 day), and the primary problem during this period is calf diarrhea (Başoğlu et al., 2004; Sen and Constable, 2013). The high level of morbidity and mortality in the neonatal period is associated with complex etiological factors as well as the inability to perform effective treatments and the radical application of fluid therapy (Abeysekara et al., 2007). Metabolic acidosis and dehydration are other common problems that require specific treatments. Under field conditions, dehydration and metabolic acidosis of calves with diarrhea are estimated based on clinical signs due to the lack of laboratory equipment (Nakagawa et al., 2007). Metabolic acidosis occurs through the loss of bicarbonate through feces, which reduces the excretion of H ions from the kidneys and the accumulation of organic acids, such as lactate in the blood (Lorenz, 2009). Several studies have been published on lactate and D-lactatemia in calves with or without diarrhea (Omole et al., 2001; Ewaschuk et al., 2003), whereas the literature published on lactate in humans has focused on biomarker properties in sepsis and the assessment of trauma situations (Gomez and Kellum, 2015; Lee and An, 2016; Dekker et al., 2017).

The main goal of intravenous (IV) fluid therapy in diarrheic calves is to replace fluids, electrolytes, buffer and energy deficits. The route, amount, type and the rate of application of IV fluids for calves can change or are not suitable under field conditions (Kaske, 1994; Berchtold, 1999). Berchtold (1999) also stated that rapid infusion may cause cerebral edema, pseudoanemia, and hypoproteinemia. However, Kasari and Naylor (1985) as well as Kasari (1990) and Constable (2003) have recommended isotonic crystalloids for dehydrated calves at a maximum dose of 80 ml/kg/hr by an intravenous route. Another study found that an infusion rate of 80 ml/kg/hr may lead to hyponatremia and recommended that the dose be as low as 30 or 40 ml/kg/hr (Grove-White, 2007). Nevertheless, in children with neonatal diarrhea, Nager and Wang (2010) revealed the favorable effects of 1-h rapid infusions of isotonic crystalloid solutions at a dose of 60 ml/kg.

Lactate clearance (LC) is a simple and inexpensive clinical parameter based on the measurement of lactate levels between two serial times (Jones et al., 2010). LC indicates kinetic alterations in anaerobic metabolism, tissue hypoxia and septic conditions (Nguyen et al., 2010). In recent years, research has focused on this biomarker to discover the effects of fluid therapy in trauma and sepsis patients (James et al., 2011; Yu et al., 2013; Ghneim et al., 2013; Lyu et al., 2015). Interestingly, lactate clearance has not been systematically studied in fluid resuscitation in calves with diarrhea.

In this study, we aimed to determine the efficacy or inadequacy of ultra-rapid (1 h) and rapid (3 h) intravenous infusions with 0.9% NaCl and 1.3% NaHCO₃ solutions administered at a dose of 60 ml/kg on lactate clearance in clinically dehydrated neonatal calves with diarrhea.

MATERIALS AND METHODS

Animals and clinical examination

Sixty animals, including healthy (n=20) and diarrheic (n=40) Holstein calves with moderate dehydration from both sexes and 1-3 weeks of age, were enrolled in the study. The clinically healthy control group calves were selected from 4 different dairy cattle farms in Aydin province in the first three-week period of their lives and included in the study. Diarrheic calves in the first three weeks of age that had moderate dehydration were referred to the clinic of researcher and author Erdogan for examination and treatment. Healthy calves were animals that received no treatment or had no infection history, whereas diarrheic calves had a treatment history.

Moderate dehydration (6 - 8%) was determined according to predictors, including skin turgor, dry mucous membranes and eyeball recession of 2 mm (Sen and Constable, 2013). Along with these clinical findings, the diarrheic calves were divided into two major groups (n=20 in each group) for the isotonic 0.9% NaCl solution and 1.3% NaHCO₃ solution. The major groups were divided into two subgroups (n=10 in each group) for comparison of the effects of rapid (3 hr) and ultra-rapid (1 hr) infusions. Pre-treatment (Pre) and post-treatment (Post) clinical (e.g. eyeball recession, skin turgor, ability to stand,

suckling reflex, and mucosal membrane moistness) and laboratory data (Lactate, and D-lactate) were recorded.

Infusion protocol and blood samples

Calves in both groups were weighed with an electronic veterinary scale (Vaw 300 4P, Turkey) before treatment. The amounts of fluid for rehydration for both the 0.9% NaCl and 1.3% NaHCO₃ solution groups were calculated based on the dosage of 60 ml/kg b.w. Fluid resuscitation was performed for all diarrheic calves in the Vena jugularis intravenously (Braun: 16 G 2" 1.7×50 mm 196 ml/min Lot: 1G16258253, Germany). For each diarrheic calf, the calculated amount of total fluid was infused with an infusion pump (Biocare IP12, Shenzhen Biocare Bio-Medical Equipment Co. Ltd, China). By changing the infusion rate of the pump, the total of both fluids (0.9% NaCl and 1.3% NaHCO₃) was infused ultra-rapidly (1 hr) and rapidly (3 hr) to all subgroups. Diarrheic calves were randomly selected for the subgroups of both isotonic crystalloid solutions.

Blood samples were collected in lithium heparin (BD, USA) and a heparinized syringe (Genject Ca₂+LH-100 I.U., Turkey) for biochemical and blood gas analyses prior to treatment and post-treatment. Post-treatment blood samples were taken in all infusion groups at the end of the fifth minute after the infusion completed. Heparinized blood samples taken pre- and post-treatment were centrifuged, and the obtained plasma was frozen at -20°C in Eppendorf tubes until analysis. L-lactate concentrations were detected via a hand held portable blood gas analyzer (EPOC, UK) and D-lactate concentrations were determined with a spectrophotometric method and ELISA using commercial test kits (Biovision, USA).

Lactate clearance

Lactate and D-lactate clearance (percent) were calculated by the following equation described by Nguyen et al.,(2004):

Clearance= $\frac{(\text{Lactate pre-treatment}) - (\text{Lactate post-treatment})}{\text{Lactate pre-treatment}} \times 100$

A negative value indicates reduction or clearance of L-lactate or D-lactate, whereas a positive value

indicates elevation of lactate or D-lactate pre- and post-treatment intervention.

Statistical analyses

The average (X), standard deviation (SD) and minimum-maximum values (Xmin-Xmax) of the analysis parameters for L-lactate and D-lactate were calculated for each sample interval in the groups. The distribution of numerical data was evaluated using a Shapiro-Wilk test. One-way analysis of variance (ANOVA) was used for comparison of more than two groups in normally distributed parameters, and a Tukey test was used for post hoc comparisons. Nonparametric methods were used for the analysis of non-normally distributed data. A Kruskal-Wallis test with Bonferroni correction was applied for post hoc comparisons for the comparison of parameters in more than two groups. In calves with diarrhea, the T-test and Wilcoxon test were performed on normally and non-normally distributed parameters, respectively, for comparison of pre- and post-treatment values of dependent groups. P values less than 0.05 were accepted as statistically significant ($p < 0.05$). SPSS 15.0 software was used to analyze the statistical data.

RESULTS

Calves treated with isotonic NaCl solution at a dose of 60 ml/kg in the ultra-rapid (1 hr) and rapid (3 hr) groups showed marked improvement in skin turgor, mucosal membrane moistness, and eyeball recession, whereas there was no significant improvement in suckling reflexes, which were reduced post-treatment. Seven of ten calves that received ultra-rapid infusion in the 0.9% NaCl group urinated post-treatment; among those treated with rapid infusion, three out of ten calves urinated (30%). In the ultra-rapid infusion (URI) group and rapid infusion (RI) group of 0.9% NaCl, six and five calves were able to stand by themselves, respectively. However, the remaining 9 calves from both 0.9% NaCl solution groups were unable to stand by themselves.

The ultra-rapid infusion subgroup of calves that received the isotonic (1.3%) NaHCO₃ solution showed marked improvements in dehydration-related clinical findings, mental status and suckling reflex

post-treatment. In the rapid infusion subgroup of the 1.3% NaHCO₃ solution group, these clinical improvements manifested in the second hour of infusion, and the sucking reflex recovered more slowly than in the ultra-rapid infusion group. At the second hour of rapid infusion of 1.3% NaHCO₃ solution, all calves urinated; for the ultra-rapid infusion group calves using the same solution, 80% of calves urinated at the end of the infusion. Calves in the 1.3% NaHCO₃ solution ultra-rapid infusion subgroup were able to stand by themselves, and the rapid infusion group calves were also able to stand during the second hour of the infusion. Two calves in the 1.3% NaHCO₃ URI subgroup had unstable walking. In all infusion groups of calves were rehydrated and the presence and severity of dehydration was clinically reduced.

were 36.4% and 31.8%, respectively, and were statistically significant. There was no significant difference in D-lactate concentrations between the control group and the infusion groups of calves pre-treatment to post-treatment. In the URI and RI groups that received 0.9% NaCl solution, the D-lactate clearance increased, but the D-lactate clearance was reduced in the URI and RI groups that received the 1.3% NaHCO₃ solution (Table 2).

DISCUSSION

Research has shown that lactate is a product of anaerobic glycolysis, and the production and clearance of lactate is balanced in body fluids. The daily production of lactate occurs in skeletal muscle, the brain, cardiac muscle, intestines, the renal medulla and erythrocytes; however, it is produced

Table 1. Mean Body weights, amount of solutions and infusion rates in diarrheic calf groups (\pm s).

<i>Group</i>	<i>n</i>	<i>Body weight (kg)</i>	<i>Amount of solution (mL)</i>	<i>Infusion rate (mL/min)</i>
<i>0,9% NaCl Sol. URI</i>	10	29,0 \pm 2,5	1742,4 \pm 147,8	29 \pm 2,4
<i>0,9% NaCl Sol. RI</i>	10	30,0 \pm 2,4	1798,2 \pm 144,5	10 \pm 0,8
<i>1,3% NaHCO₃ Sol URI</i>	10	31,4 \pm 3,2	1886,4 \pm 193,2	31 \pm 3,2
<i>1,3% NaHCO₃ Sol RI</i>	10	30,2 \pm 1,8	1813,2 \pm 108,0	10 \pm 0,6

There were no significant alterations in the amount of either ultra-rapid or rapid infusions of 0.9% NaCl solution and 1.3% NaHCO₃ solution, as shown in Table 1.

L-lactate concentrations were statistically significantly higher for the pre-treatment 0.9% NaCl solution URI, 1.3% NaHCO₃ solution URI and 1.3% NaHCO₃ solution RI groups than for the control group calves. In the pre-treatment and post-treatment comparisons, ultra-rapid infusion and rapid infusion of both 1.3% NaHCO₃ solution and 0.9% NaCl solution resulted in elevations in lactate concentration; these elevations were significant for ultra-rapid infusion of the 1.3% NaHCO₃ and 0.9% NaCl solutions. Marked increases in LC were detected in the 0.9% NaCl solution URI, 1.3% NaHCO₃ solution URI and 1.3% NaHCO₃ solution RI groups. The 0.9% NaCl solution URI and 1.3% NaHCO₃ solution URI group lactate clearance rates

in almost all tissues and organs during disease (Sharkey and Wellman, 2015). The assessment of lactate concentrations has been used as a significant therapeutic and prognostic biomarker for human and animal health for many years (Nguyen et al., 2004; Karagiannis et al., 2006; Figueiredo et al., 2006; Allen and Holm, 2008; Zacher et al., 2010; Tennent-Brown et al., 2010). Lactate is commonly used to determine sepsis and its severity (Gomez and Kellum, 2015; Lee and An, 2016) as well as for the prognostic evaluation of patients with sepsis (Yu et al., 2013; Lyu et al., 2015). For example, in horses with acute abdomen, the influence of perioperative plasma lactate concentrations on the surgical intervention and recovery periods has been shown to be especially important (Donawick et al., 1975; Moore et al., 1976; Genn and Hertsch, 1982), and similar results have been observed in dogs with gastric dilatation and volvulus (De Papp, 1999). The

Table 2. L and D-lactate concentrations and clearances levels in calves.

Parameter	Groups								
	Control	0,9% NaCl Sol. URI		0,9% NaCl Sol. RI		1,3% NaHCO ₃ Sol URI		1,3% NaHCO ₃ Sol RI	
		Pre	Post	Pre	Post	Pre	Post	Pre	Post
<i>L. Lactate (mmol/L)</i>	0,9 ± 0,6	2,2 ± 1,7 ^{‡a}	3,0 ± 2,3 ^{†b}	1,4 ± 1,0	1,4 ± 1,3	4,4 ± 2,8 ^{‡a}	5,8 ± 4,2 ^{‡b}	2,2 ± 2,1*	2,9 ± 3,2*
<i>D- lactate (mmol/L)</i>	0,4 ± 0,1	0,41 ± 0,02	0,43 ± 0,13	0,39 ± 0,03	0,42 ± 0,14	0,39 ± 0,04	0,38 ± 0,05	0,41 ± 0,05	0,39 ± 0,06
<i>L. Laktat clearans</i>		36,4%↑		13,6%↑		31,8%↑		31,8%↑	
<i>D- lactate clearans</i>		4,9%		7,7%		- 2,6%↓		- 4,9%↓	

*, †, ‡: Statistically significant from control group p<0,05, p<0,01 and p<0,001, respectively.

↑: Increase (Reduce in Lactate Clearance) ; ↓; Decrease (Increases in Lactate Clearance); alterations in bold are statistically significant.

^{a,b}: Different letters are statistically significant between pre and post treatment.

importance of lactate has emerged in studies that were performed in association with lactic acidosis of the acute and subacute rumen (Ewaschuk et al., 2005) and in diseases, such as abomasum displacement and respiratory tract infections in cattle (Coghe et al., 2000; Figueiredo et al., 2006). Research evaluating variations in lactate levels (Kaske, 1994) and non-metabolized D-lactate levels (Lorenz and Gentile, 2014) in neonatal calves with diarrhea has appeared. It has been shown that lactate values can be used to determine the condition of strong ion acidosis (SID) in calves with diarrhea (Ewaschuk et al., 2003; Müller et al., 2012). However, no studies in the literature have evaluated lactate measurement or the effect of rehydration on lactate concentrations in calves with diarrhea. In this study, the influence of 0.9% NaCl and 1.3% NaHCO₃ infusion solutions with a constant dose of 60 ml/kg on the pre-and post-treatment L-lactate and D-lactate concentrations was determined in moderately dehydrated calves with diarrhea. This is the first study comparing the effects of infusion rates of two isotonic solutions on lactate concentration and lactate clearance.

Over the last decade, it has been demonstrated that the result of a single lactate measurement can be contradictory for the interpretation of tissue perfusion (Zacher et al., 2010). Recently, research has indicated that conditions of cellular hypoxia and shock in humans must be evaluated over lactate clearance by measuring lactate concentrations at certain intervals (Vincent et al., 1983; Nguyen et al., 2004; Kamolz et al., 2005). In a study assessing the infusion rates of isotonic crystalloid and colloidal solutions of lactate

clearance in patients with trauma, it was reported that the hydroxyethyl starch (HES) colloidal solution had significantly decreased lactate clearance compared with that of saline solution and resulted in less renal damage in patients with penetrating trauma; however, the relevant solutions did not cause any variation in the specified parameters in patients with blunt trauma (James et al., 2011).

In a study evaluating patients with sepsis who were admitted to an emergency clinic, different treatment groups underwent an increase in the central venous pressure (CVP) to 8 mmHg or greater. Overall, 4.5 L of isotonic crystalloid solution were administered on average for 6 h in a patient group, and the sepsis was treated. Because the study results were based on variations in sepsis levels and were not obtained by evaluating the patients who received infusions and different therapy methods, the efficacy of bolus fluid therapy on the lactate clearance could not be determined in the study (Jones et al., 2010). In another study, an average dose of 3,375 ml of crystalloid fluid altered lactate clearance at levels of 38.1% ± 34.6 and 12.0% ± 51.6 in surviving and non-surviving patients with sepsis, respectively (Nguyen et al., 2004). In our study, the ultra-rapid infusion (1 h) and the rapid infusion (3 h) of 0.9% NaCl solution at a dose of 60 ml/kg reduced the rates of lactate clearance to 36.4% and 13.6% in the calves, respectively. During ultra-rapid and rapid administration of 1.3% NaHCO₃ solution to eliminate the dehydration as well as metabolic acidosis in calves with diarrhea, the variations in lactate clearance levels reduced to 31.8% in both

subgroups; however, the variations in the lactate concentrations were statistically significant in the ultra-rapid infusion group (Table 2). The relevant variations in lactate clearance were similar to those in human studies. The statistically significant reduce that occurred during ultra-rapid infusion of both solutions was considered to be due to the non-metabolizing of lactate that occurred within an hour despite eliminating dehydration and microcirculation. Furthermore, lactate clearance is closely related with capillary perfusion independent of hemodynamic variables (De Backer et al., 2006). The results in this study suggest that reducing in LC might be related to insufficient perfusion restoration, intravascular volume, and ailment energy metabolism. The outcome of statistically non-significant reduces in the clearance during ultra-rapid infusion was correlated with measurements in the third hour following fluid administration. In humans research based on the LC the sequential measurements of lactate is based on the least 6 hours intervals (Nguyen et al., 2004). In this study the time intervals between two lactate measurements were 3 hours. It is thought that the measurement of lactate clearance in such a short period of time is a limiting factor in the study.

Acute diarrhea in calves mainly causes dehydration, metabolic acidosis, electrolyte imbalance, prerenal azotemia and negative energy balance, which are independent of its etiology (Kaske, 1994; Berchtold, 1999), and these variations occur at different levels according to the duration and severity of diarrhea (Smith, 2009). Metabolic acidosis and dehydration are among the most significant complications of diarrhea (Smith, 2009). Additionally, metabolic acidosis may vary by age (Naylor, 1989). Trefz et al. (2012) found that median concentrations of D-lactate were detected at 0.8 mmol/l and 3.9 mmol/l in neonatal calves with diarrhea (age <7 days) and calves older than 7 days, respectively, and the age-dependent role of D-lactate was revealed from the development of metabolic acidosis. In this study, the level of D-lactate was in the range of 0.39-0.41 mmol/l in neonatal calves with diarrhea, and the contribution of D-lactate remained limited in the development of metabolic acidosis.

Although there are studies that have found associations between the elimination of D-lactate

from the body and the clinical findings in calves (Lorenz, 2004), there are no studies regarding the association of D-lactate with renal clearance (Zello et al., 2008). In the study by Zello et al. (2008) the mean renal clearance was detected as 7.4 ± 3.4 ml/min when the blood concentration of D-lactate reached maximum levels in the calves receiving a D-lactic acid infusion. In the present study, the variations associated with infusion rates of the D-lactate concentration were observed in the blood, and the variations were measured at levels of 4.9% and 7.7% during the ultra-rapid and rapid infusions of 0.9% NaCl solution, respectively. On the other hand, variations were detected at levels of 2.6% and 4.9% during ultra-rapid and rapid infusion of 1.3% NaHCO₃ solutions, respectively. In the calves treated with both infusion solutions, the low detectable concentrations of D-lactate, the low clearance of D-lactate in the blood and the low variation rates might be associated with the type of metabolic acidosis in calves with acute primary metabolic acidosis.

Several studies on different species have been conducted in the fields of human and veterinary medicine regarding the infusion rates of isotonic crystalloid solutions (Nager and Wang, 2010); however, the data that were obtained could not enhance routine practices. Because rapid infusion has risks, including hyperhydration and pulmonary edema, it has been suggested that the infusion rate should not be high. In addition to these side effects, cerebral edema, anemia and hypoproteinemia might also develop (Abeysekera et al., 2012). The maximum infusion rate of isotonic solutions has been reported to be as high as 80 ml/kg/h in dehydrated calves (Kasari, 1990; Constable, 2003). Grove-White (2007) found this rate to be risky in their study, and they argued there should be a lower infusion rate due to the risk of hyperhydration; they suggested lowering the rates to 30 or 40 ml/kg/h for a less risky and suitable infusion. In a study performed by Roussel and Kasari (1990), the best infusion rate was reported as 50 ml/kg/h in which the infusion of fluid was completed within 2 to 3 h, and the risk of hyperhydration was therefore eliminated. In present study, both 0.9 NaCl% and 1.3% NaHCO₃ solutions were applied at 60 ml/kg/h for ultra-

rapid infusion and 20 ml/kg/h for rapid infusion in neonatal calves with moderate diarrhea. At the end of the infusion, the clinical findings for ultra-rapid and rapid infusions of isotonic NaCl solution in both calf groups with diarrhea were associated with an exacerbation of metabolic acidosis according to the relevant solution instead of the infusion rate.

CONCLUSION

The clearance of L-lactate and D-lactate varied with the infusion rate in calves with diarrhea, and these variations were statistically significant in both of the ultra-rapid subgroups. This study offers insight for

further research on fluid therapy in calves based on the calculation of lactate clearance.

ACKNOWLEDGMENT

This study was partially summarized from the PhD thesis of the corresponding author. The project was funded by Adnan Menderes University Research Projects Funding Unit under project number VTF-11009.

CONFLICT OF INTEREST

The authors have no conflict of interest to declare for this manuscript. ■

Fig 1. Lactate concentrations in infusion groups





Fig 2. D-Lactate concentrations in infusion groups



REFERENCES

- Abeysekara S, Naylor JM, Wassef AWA, Isak U, Zello GA (2007) D-Lactic acid-induced neurotoxicity in a calf model. *Am J Physiol* 293:558–65.
- Abeysekara S, Zello GA, Lohmann KL, Alcorn J, Hamilton DL, Naylor JM (2012) Infusion of sodium bicarbonate in experimentally induced metabolic acidosis does not provoke cerebrospinal fluid (CSF) acidosis in calves. *Can J Vet Res* 76: 16-22.
- Allen SE, Holm JL (2008) Lactate: Physiology and clinical utility. *J Vet Emerg Crit Care* 18:123-132.
- Baçoğlu A, Sen I, Sevinç M, Simsek A (2004) Serum concentrations of tumor necrosis factor-alpha in neonatal calves with presumed septicemia. *J Vet Intern Med* 18:238-241.
- Berchtold J (1999) Intravenous fluid therapy for calves. *Vet Clin North Am Food Anim Pract*, 15:505-531.
- Coghe J, Uystepuyt CH, Bureau F, Detilleux J, Art T, Lekeux P (2000) Validation and prognostic value of plasma lactate measurement in bovine respiratory disease. *Vet J* 160: 139-146.
- Constable PD (2003) Fluid and electrolyte therapy in ruminants. *Vet Clin North Am Food Anim Pract* 19:557–97.
- De Papp E, Drobatz KJ, Hughes D (1999) Plasma lactate concentration as a good predictor of gastric necrosis and survival among dogs with gastric dilatation-volvulus: 102 cases (1995–1998). *J Am Vet Med Assoc* 215:49–52.
- De Backer D, Creteur J, Dubois MJ, Sakr Y, Koch M, Verdant C, Vincent JL (2006) The effects of dobutamine on microcirculatory alterations in patients with septic shock are independent of its systemic effects. *Crit Care Med* 34:403-408.
- Dekker SE, de Vries HM, Lubbers WD, van de Ven PM, Toor EJ, Bloemers FW, Boer C (2017) Lactate clearance metrics are not superior to initial lactate in predicting mortality in trauma. *Eur J Trauma Emerg Surg* 43:1-11.
- Donawick WJ, Ramberg CF, Paul SR, Hiza MA (1975) The diagnostic and prognostic value of lactate determinations in horses with acute abdominal crisis. *J S Afr Vet Assoc* 4: 22-25.
- Ewaschuk JB, Naylor JM, Zello GA (2003) Anion gap correlates with serum D- and DL- lactate concentration in diarrheic neonatal calves. *J Vet Intern Med* 17: 940-942.
- Ewaschuk JB, Naylor JM, Zello GA (2005) D-lactate in human and ruminant metabolism. *J Nutr* 135: 1619-1625.
- Figueiredo MD, Nydam DV, Perkins GA, Mitchell HM, Divers TJ (2006) Prognostic value of plasma L-lactate concentration measured cow-side with a portable clinical analyzer in Holstein dairy cattle with abomasal disorders. *J Vet Intern Med* 20:1463-1470.
- Genn HJ, Hertsch B (1982). Die diagnostische und prognos-

- tische Bedeutung des Laktatwertes im Blut sowie in der Bauchhöhlenflüssigkeit bei der Kolik des Pferdes. *Dtsch Tierärztl Wochenschr* 89:295–299.
- Ghneim MH, Regner JL, Jupiter DC, Kang F, Bonner GL, Bready MS, Davis ML (2013) Goal directed fluid resuscitation decreases time for lactate clearance and facilitates early fascial closure in damage control surgery. *Am J Surg* 206: 995-1000.
- Gomez H, Kellum JA (2015) Lactate in sepsis. *Jama*, 313(2):194-195.
- Grove-White D (2007) Practical intravenous fluid therapy in the diarrhoeic calf. *In Pract* 29:404–8.
- James MFM, Michell WL, Joubert IA, Nicol AJ, Navsaria PH, Gillespie RS (2011) Resuscitation with hydroxyethyl starch improves renal function and lactate clearance in penetrating trauma in a randomized controlled study: the FIRST trial (Fluids in Resuscitation of Severe Trauma). *Br J Anaesth* 107: 693-702.
- Jones AE, Shapiro NI, Trzeciak S, Arnold RC, Claremont HA, Kline JA (2010) Lactate clearance vs central venous oxygen saturation as goals of early sepsis therapy. *JAMA* (303):739–746.
- Kamolz LP, Andel H, Schramm W, Meissl G, Herndon DN, Frey M (2005) Lactate: early predictor of morbidity and mortality in patients with severe burns. *Burns* 31:986–990.
- Karagiannis MH, Reniker AN, Kerl ME, Mann FA (2006) Lactate measurement as an indicator of perfusion. *Compend Contin Educ Vet* 28:287–300.
- Kasari TR (1990) Metabolic acidosis in diarrheic calves: alkalinizing agents in therapy. *Vet Clin North Am Food Anim Pract* 6:29-43.
- Kasari TR, Naylor JM (1985). Clinical evaluation of sodium bicarbonate, sodium L–lactate, and sodium acetate for the treatment of acidosis in diarrheic calves. *J Am Vet Med Assoc* 187:392-397.
- Kaske M (1994). Pathophysiologische Aspekte der neonatalen Kälberdiarrhoe. *Tierärztl Umsc* 49:336–348.
- Lee SM, An WS (2016). New clinical criteria for septic shock: serum lactate level as new emerging vital sign. *J Thorac Dis* 8:1388.
- Lorenz I (2004) Influence of D-lactate on metabolic acidosis and on prognosis in neonatal calves with diarrhoea. *J Vet Med A Physiol Pathol Clin Med* 51:425–428.
- Lorenz I (2009) An update on calf diarrhoea. Part I: Pathophysiology and treatment. *Ir Vet J* 62:58-61.
- Lorenz I, Gentile A (2014) D-lactic acidosis in neonatal ruminants. *Vet Clin Food Anim Pract* 30:317-331.
- Lyu X, Xu Q, Cai G, Yan J, Yan M (2015) Efficacies of fluid resuscitation as guided by lactate clearance rate and central venous oxygen saturation in patients with septic shock. *Zhonghua yi xue za zhi* 95:496-500.
- Moore JN, Owen RR, Lumsden JH (1976) Clinical evaluation of blood lactate levels in equine colic. *Equine Vet J* 8:49–54.
- Müller KR, Gentile A, Klee W, Constable PD (2012) Importance of the effective strong ion difference of an intravenous solution in the treatment of diarrheic calves with naturally acquired acidemia and strong ion (metabolic) acidosis. *Vet Intern Med* 26:674-683.
- Nager AL, Wang VJ (2010) Comparison of ultrarapid and rapid intravenous hydration in pediatric patients with dehydration. *Am J Emerg Med* 28:123-129.
- Nakagawa M, Suzuki K, Taguchi K (2007) Relationship between depression score and acid-base status in Japanese Black calves with diarrhea. *J Vet Med Sci* 69:549–552.
- Naylor JM (1989) A retrospective study of the relationship between clinical signs and severity of acidosis in diarrheic calves. *Can Vet J* 30:577–580.
- Nguyen HB, Rivers EP, Knoblich BP, Jacobsen G, Muzzin A, Ressler JA, Tomlanovich MC (2004) Early lactate clearance is associated with improved outcome in severe sepsis and septic shock. *Crit Care Med* 32:1637-1642.
- Nguyen HB, Loomba M, Yang JJ, Jacobsen G, Shah K, Otero RM, Rivers EP (2010) Early lactate clearance is associated with biomarkers of inflammation, coagulation, apoptosis, organ dysfunction and mortality in severe sepsis and septic shock. *J Inflamm* 7:6–17.
- Omole OO, Nappert G, Naylor JM, Zello GA (2001) Both L- and D-Lactate contribute to metabolic acidosis in diarrheic calves *J Nutr* 131:2128-2131
- Roussel AJ, Kasari TR (1990) Using fluid and electrolyte replacement therapy to help diarrheic calves. *Vet Met Sci* 85:303–304.
- Sen I, Constable PD (2013) General overview to treatment of strong ion (metabolic) acidosis in neonatal calves with diarrhea. *Eurasian J Vet Sci* 29:14-120
- Sharkey LC, Wellman ML (2015) Use of lactate in small animal clinical practice. *Clin Lab Med* 35:567-577.
- Smith GW (2009) Treatment of calf diarrhea: oral fluid therapy. *Vet Clin North Am Food Anim Pract* 25:55-72.
- Tennent-Brown BS, Wilkins PA, Lindborg S, Russell G, Boston RC (2010). Sequential plasma lactate concentrations as prognostic indicators in adult equine emergencies. *J Vet Intern Med* 24:198-205.
- Trefz F, Lorch M, Feist A, Sauter-Louis C, Lorenz I. (2012) Metabolic acidosis in neonatal calf diarrhea—clinical findings and theoretical assessment of a simple treatment protocol. *J Vet Intern Med* 26:162-170.
- Vincent JL, Dufaye P, Berre J, Leeman M, Degaute JP, Kahn RJ (1983) Serial lactate determinations during circulatory shock. *Crit Care Med* 11:449–451.
- Yu B, Tian HY, Hu ZJ, Zhao C, Liu LX, Zhang Y, Li J (2013) Comparison of the effect of fluid resuscitation as guided either by lactate clearance rate or by central venous oxygen saturation in patients with sepsis. *Zhonghua Wei Zhong Bing Ji Jiu Yi Xue* 25:578-583.
- Zacher LA, Berg J, Shaw SP, Kudej RK (2010) Association between outcome and changes in plasma lactate concentration during presurgical treatment in dogs with gastric dilatation- volvulus: 64 cases (2002 – 2008). *J Am Vet Med Assoc* 236:892-897.
- Zello GA, Janzen A, Abeysekara S, Hamilton D, Naylor JM, Lohmann KL (2008) Urinary excretion of both D-and L-lactate using a calf-infusion model. *The FASEB Journal* 22:1205.