

# Journal of the Hellenic Veterinary Medical Society

Vol 71, No 1 (2020)



## Effects of Safranal on Tissue Oxidative Stress in Sub-Chronic Thinner-Addicted Rats

M. DÜZ, A. F. FIDAN

doi: [10.12681/jhvms.22942](https://doi.org/10.12681/jhvms.22942)

Copyright © 2020, M. DÜZ, A. F. FIDAN



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

### To cite this article:

DÜZ, M., & FIDAN, A. F. (2020). Effects of Safranal on Tissue Oxidative Stress in Sub-Chronic Thinner-Addicted Rats. *Journal of the Hellenic Veterinary Medical Society*, 71(1), 1997-2004. <https://doi.org/10.12681/jhvms.22942>

## Effects of Safranal on Tissue Oxidative Stress in Sub-Chronic Thinner-Addicted Rats

M. Düz<sup>\*1</sup>, A.F. Fidan<sup>2</sup>

<sup>1</sup> Department of Chemistry, Faculty of Science and Literature, Afyon Kocatepe University, Afyonkarahisar, Turkey

<sup>2</sup> Department of Biochemistry, Faculty of Veterinary Medicine, Afyon Kocatepe University, Afyonkarahisar, Turkey

**ABSTRACT.** The present study was carried out to determine the effects of sub-chronic thinner addiction on the oxidant-antioxidant balance and oxidative stress on certain tissues and the possible protective effect of safranal against thinner toxication in rats. Adult male Wistar albino rats were randomly divided into four groups of 10 animals each as follows: control (C), safranal (S), thinner (T) and thinner+safranal (T+S). The control group received 1cc saline by gastric gavage. Safranal was administered to S and T+S groups by using gastric gavage at a dose of 100 mg/kg/day and volume of 0.1 mL/kg/day. Thinner inhalation was applied to T and T+S groups in a container with NaOH tablets twice a day. Levels of malondialdehyde (MDA), reduced glutathione (GSH), nitric oxide (NO<sub>x</sub>) metabolites, total antioxidant capacity (TAS) and total oxidant capacity (TOS) were determined in liver, lung, brain, kidney and testis tissues of the rats. In the T+S group, it was observed that the MDA levels significantly decreased in all tissues, except the kidney, in comparison to the thinner inhalation group (p = 0.000). When the NO<sub>x</sub> levels of the T+S group were compared with the levels of the T group, it was concluded that there existed a statistically significant decrease in the NO<sub>x</sub> levels in all tissues (p = 0.000).

In T+S group, it was observed that safranal either eliminated or mitigated oxidative stress that developed in tissues through decreasing MDA and TOS levels and increasing GSH and TAS levels and caused significant decreases in NO<sub>x</sub> levels in all tissues. As a result, it was determined that safranal, although not uniform for all tissue types, had a protective potential against the damaging effects of oxidative stress caused by sub-chronic thinner inhalation.

**KeyWords:** Thinner inhalation, oxidative stress, safranal, tissue

*Corresponding Author:*  
Mürüvvet Düz, Department of Chemistry, Faculty of Science and Literature, Afyon Kocatepe University, Afyonkarahisar, Turkey  
E-mail address: muruvvetduz@aku.edu.tr

*Date of initial submission: 26-03-2019*  
*Date of revised submission: 24-10-2019*  
*Date of acceptance: 30-11-2019*

## INTRODUCTION

Thinner is an organic solvent which is commonly used for industrial applications and currently has an increasing improper use as a pleasure inducing and narcotic substance. Thinner is a mixture of toluene, benzene, acetone, methanol, hexane and other substances (Nedzvetskii et al., 2012). It was indicated that the thinner used in Turkey was composed of 63% of toluene, 13% of acetone, 10% of isobutyl acetate, 7.5% of isobutane and 6.5% of butylene glycol and its harmful effects were mostly associated with toluene, where the remaining substances activated these harmful effects (Cobanoglu et al., 2007).

Reactive oxygen (ROS) and nitrogen (RNS) derivatives occur in the organism due to exposure to environmental and chemical agents and the interaction of such derivatives with macromolecules leads to oxidative stress (Mattia et al., 2003). Similar to various internal and external toxic agents, the toxic effects of thinner also emerge through free radicals (Mattia et al. 1991, Martínez-Alfaro et al., 2011).

Long-term use of volatile organic solvents has been shown to cause permanent damage to the lung and liver (Marjot and McLeod, 1989; Al-Alousi, 1989). Mattia et al. (1991; 1993a,b) reported in vivo and in vitro studies that ROS formation is accelerated and lipid peroxidation is increased in brain, liver, kidney and lung tissues with the effect of intraperitoneal administration of toluene. Al-Alousi (1989) emphasized that pulmonary edema is the cause of death due to volatile substances. When the results of many studies are considered, it has been observed that there are many damages as a result of using thinner in industrial area and for narcotic use. Studies show that MDA levels increase and GSH levels decrease due to degradation of lipid peroxidation due to oxidative stress of organic solvents (Ulakoglu et al. 1998b, Dillioglugil et al. 2005, El-Nabi Kamel and Shehata 2008).

In recent years, research in phytotherapy focused on the effects of the components in plant extracts on various diseases, rather than the effect of plant extracts. Safranal is one of the components of saffron extract. Saffron is an herb with antioxidant effects, included in medical literature and is acknowledged to have an increasing significance due to its therapeutic effects (Moghaddasi, 2010; Assimopoulou et al., 2015). The radical scavenging activity of safranal results with antidiabetic, antioxidant, anticancer and hypotensive properties, thus various studies concluded that safranal could be used in industries such as

food, pharmaceuticals and cosmetics (Assimopoulou et al., 2005, Imenshahidi et al., 2010, Hazman ve Bozkurt, 2015). Karafakioglu et al. (2017) reported that safranal (200 mg / kg) treatment may have a protective effect against cisplatin-induced nephrotoxicity and oxidative stress in rats. Alayunt et al. (2019) observed that particularly high-dose (100 mg/kg) administration of safranal to rats promoted the antioxidant system and be inflammatory against TNF- $\alpha$  and IL-6 cytokines on damage induced by CCl<sub>4</sub>.

The main aim of our study is to investigate to what extent thinner, which is a mixture of organic solvent, affects brain, lung, liver, kidney, testis via respiratory system and to investigate the effect of safranal on this condition. The results of the study may contribute to the development of new approaches both in the protection of worker health on an industrial scale and in the treatment of addicts using volatile substances.

## MATERIALS AND METHODS

### Chemicals

Safranal (Cat no:Sigma-W338907), Total oxidant (TOS) and antioxidant status (TAS) commercial kits (Rell Assay:RL0024, RL0017) and the other chemical used in the study were purchased from Sigma -Aldrich (Sigma Aldrich Chemical Co. St. Louis, Missouri, ABD).

### Animals

In the present study, 40 Wistar albino male rats (weighing ~250-300 g) were used. The animals were housed under standard conditions of temperature (23 $\pm$ 2°C), humidity, and dark-light cycle (lights on from 6:00 am to 6:00 pm). Throughout the study, the rats were maintained on standard rat feed and tap water *ad libitum*. They were fed twice daily at 0900 and 1900. Before starting the thinner experiments, rats were subjected to preexperiments in order to prevent possible toxication severity complications. All the animals were carefully monitored and maintained in accordance with the ethical recommendation of the University of Afyon Kocatepe Animal Ethics Committee. The permission was gained on 7 May 2015 with the number 49533702/46.

### Experimental procedure

A total of 40 rats were included in the study. They were divided into 4 groups as follows: Control Group (C) (n=10), Sub-chronic Thinner Inhalation Group (T) (n=10), Sub-chronic Thinner Inhalation + Sa-

franal Group (T+S) (n=10) and Safranal Group (S) (n=10). Control group (C); comprised 10 healthy rats. The rats in the control group were administered 1 cc of physiological saline daily by gastric gavage for 8 weeks. Rats were fed with standard rat feed for 8 weeks and their weights were recorded regularly during the experiments. Sub-chronic Thinner Inhalation Group (T); rats were placed in a specially designed cage size 100x60x40 cm with air ventilation that allowed inhalation of controlled amounts of thinner (5ml). This application was repeated twice a day for 8 weeks and was terminated when 50% of the rats lost their reflexes to stand (About 5-6 min) (Konuk et al., 2012). Sub-chronic Thinner Inhalation+Safranal Group (T+S); treated as T group and additionally a dose of 100 mg/kg of safranal dissolved in 1 cc of physiological saline solution was administered via gastric gavage for 8 weeks. Commercial safranal (Sigma-w338907) was used in the study. Safranal Group (S); a dose of 100 mg/kg of safranal dissolved in 1 cc of physiological saline solution was administered via gastric gavage for 8 weeks. The dose of safranal used in this study was based on previous studies (Hosseinzadeh et al., 2013; Hazman ve Bozkurt, 2015; Hazman and Ovalı, 2015).

### Sample preparation

At the end of the experimental period, the rats were anesthetized and sacrificed by cervical dislocation. Rats were anesthetized with 10 mg/kg-im xylazine and 50 mg/kg-im ketamine injection. All rats were sacrificed on the 56th day (8 weeks) of the experiment and tissue samples (brain, lung, liver, kidney, testis) were collected for examination of clinical biochemical analysis. Tissue samples (0.5 g) were placed in homogenizer and were added 5 mL of phosphate buffer solution (PBS) (pH: 7.4) and were homogenized by means of the homogenizer and the sonicator. The homogenates were centrifuged at 15000rpm and +4°C for 10 minutes, and the supernatants were stored at -80 °C in deep freeze pending analysis.

### Measurement of parameters

In the obtained supernatants, the Malondialdehyde (MDA) levels were determined according to Ohkawa et al. (1979), the reduced glutathione (GSH) analyzes were conducted based on Beutler (1963) and the amounts of Nitric oxide (NOx) were determined via the "Vanadium chloride (III) - Gries Reaction" method of Miranda et al. (2001). Total oxidant (TOS) and antioxidant status (TAS) measurements were

performed via Eliza Reader (Biotek, ELx800) using commercial kits based on the method developed by Erel (2004, 2005). The amount of toluene in the blood was measured and the level of inhalation in T and T + S groups were determined. The toluene levels in the rats' blood were determined using the GS-MS in Acıbadem Labmed Central Laboratory (Istanbul, Turkey).

### Statistical analysis

ANCOVA method was used for statistical analysis and the difference between the groups was tested using the Bonferroni test. The live weight averages of the groups at the zeroth and eighth week were chosen as the covariate. Data were analyzed using the SPSS® 24 package program. The results were expressed as "mean±standard error" (X±SE) and p <0.05 was considered as the significance value.

## RESULTS

### Blood toluene levels

Blood toluene levels of rats that inhaled sub-chronic thinner are presented in Table 1. No statistically significant difference was determined for inhaled toluene levels in T and T+S groups.

### MDA levels measured in tissue samples

The changes in MDA, which is an indicator of from lipid peroxidation, in the liver, lung, kidney, brain and testis tissue samples retrieved from the four experiment groups constituted in the present study, are presented in Table 2. Comparing the MDA levels of the T group with that of the control group levels it could be observed that a significant increase of MDA levels were determined in liver, lung, testis and kidney tissues (p=0.000).

### GSH levels measured in tissue samples

The GSH levels in groups are presented in Table 3. It was observed that GSH levels in liver, kidney brain, and testis tissues was a significant decreased between T group and that of the control groups (Respectively p=0.013; p=0.000; p= 0.015; p= 0.007)

### TAS and TOS levels measured in tissue samples

Changes of TAS and TOS levels in all tissues in groups are presented in Table 4 and Table 5, respectively. Comparing the TAS levels of the T group with that of the control group levels it was determined that TAS levels only significantly decreased in liver and kidney tissues (p=0.000). It was identified that liv-

er and brain TOS levels only significantly increased between the T group and that of the control group ( $p=0.000$ ;  $p=0.001$ ). It was observed that the liver and kidney tissues TAS levels significantly increased between T+S group and that of the T group ( $p=0.000$ ). It was concluded that liver tissue TOS levels significantly decreased between the T+S group and the T group ( $p=0.000$ ).

### NO<sub>x</sub> levels measured in tissue samples

The changes in the NO<sub>x</sub> levels obtained from all tissues of the experiment groups are presented in Table 6. It was observed that the NO<sub>x</sub> level significantly increased only in the brain tissue between T group and the control group ( $p=0.000$ ). When the NO<sub>x</sub> levels of the T+S group were compared with the levels of the T group, it was concluded that there existed a statistically significant decrease in the NO<sub>x</sub> levels in all tissues ( $p=0.000$ ).

**Table 1.** Blood toluene levels of rats in T and T+S groups

Groups	C $\bar{x}\pm SE$	S $\bar{x}\pm SE$	T $\bar{x}\pm SE$	T+S $\bar{x}\pm SE$	p
Toluene (ng/ml)	-	-	1,525±0,071	1,481±0,071	0,730

C: Control Group; T : Sub-chronic Thinner Inhalation Group; T+S: Sub-chronic Thinner Inhalation + Safranal Group; S: Safranal Group; X ± SE: mean ± standard error;  $p<0.05$

**Table 2.** MDA levels measured in tissue samples

MDA (nmol/g tissue)	C $\bar{x}\pm SE$	S $\bar{x}\pm SE$	T $\bar{x}\pm SE$	T+S $\bar{x}\pm SE$	p
LIVER	34,82±1,60 <sup>bc</sup>	36,71±1,24 <sup>ab</sup>	41,81±1,18 <sup>a</sup>	29,38±1,46 <sup>c</sup>	0,000
LUNG	19,95±0,94 <sup>b</sup>	19,83±0,73 <sup>b</sup>	24,27±0,69 <sup>a</sup>	18,32±0,86 <sup>b</sup>	0,000
KIDNEY	24,18±1,81 <sup>b</sup>	27,32±1,39 <sup>ab</sup>	32,83±1,33 <sup>a</sup>	27,82±1,65 <sup>ab</sup>	0,000
BRAIN	27,32±0,88 <sup>a</sup>	22,72±0,68 <sup>b</sup>	28,35±0,65 <sup>a</sup>	17,09±0,79 <sup>c</sup>	0,000
TESTIS	20,28±2,05 <sup>b</sup>	19,58±1,58 <sup>b</sup>	29,54±1,51 <sup>a</sup>	17,97±1,87 <sup>b</sup>	0,000

MDA: Malondialdehyde; C: Control Group; T : Sub-chronic Thinner Inhalation Group; T+S: Sub-chronic Thinner Inhalation + Safranal Group; S: Safranal Group; X ± SE: mean ± standard error; a,b,c: different letters on the same line indicate a significant difference  $p<0.05$

**Table 3.** GSH levels measured in tissue samples

GSH (nmol/g tissue)	C $\bar{x}\pm SE$	S $\bar{x}\pm SE$	T $\bar{x}\pm SE$	T+S $\bar{x}\pm SE$	p
LIVER	19,93±1,05 <sup>a</sup>	19,25±0,81 <sup>ab</sup>	16,47±0,78 <sup>b</sup>	18,69±0,96 <sup>ab</sup>	0,013
LUNG	16,46±0,70 <sup>ab</sup>	16,71±0,54 <sup>a</sup>	17,37±0,52 <sup>a</sup>	13,64±0,64 <sup>b</sup>	0,000
KIDNEY	15,99±0,98 <sup>ab</sup>	17,39±0,76 <sup>a</sup>	12,91±0,72 <sup>c</sup>	14,09±0,89 <sup>bc</sup>	0,000
BRAIN	14,70±0,75 <sup>a</sup>	14,36±0,58 <sup>a</sup>	12,41±0,55 <sup>b</sup>	13,14±0,69 <sup>ab</sup>	0,015
TESTIS	18,8±1,6 <sup>a</sup>	18,71±1,22 <sup>ab</sup>	13,76±1,16 <sup>b</sup>	18,60±1,44 <sup>ab</sup>	0,007

GSH: reduced glutathione; C: Control Group; T : Sub-chronic Thinner Inhalation Group; T+S: Sub-chronic Thinner Inhalation + Safranal Group; S: Safranal Group; X ± SE: mean ± standard error; a,b,c: different letters on the same line indicate a significant difference  $p<0.05$

**Table 4.** TAS levels measured in tissue samples

TAS (mmol Trolox Equiv./L)	C $\bar{x}\pm SE$	S $\bar{x}\pm SE$	T $\bar{x}\pm SE$	T+S $\bar{x}\pm SE$	p
LIVER	0,89±0,07 <sup>a</sup>	0,39±0,05 <sup>b</sup>	0,38±0,05 <sup>b</sup>	0,80±0,06 <sup>a</sup>	0,000
LUNG	0,28±0,09	0,29±0,07	0,32±0,07	0,49±0,09	0,213
KIDNEY	0,56±0,07 <sup>a</sup>	0,45±0,06 <sup>a</sup>	0,15±0,06 <sup>b</sup>	0,49±0,07 <sup>a</sup>	0,000
BRAIN	0,35±0,07	0,22±0,05	0,32±0,05	0,35±0,06	0,158
TESTIS	0,43±0,05	0,33±0,04	0,30±0,03	0,38±0,04	0,060

TAS: Total antioxidant status; C: Control Group; T : Sub-chronic Thinner Inhalation Group; T+S: Sub-chronic Thinner Inhalation + Safranal Group; S: Safranal Group; X ± SE: mean ± standard error; a,b,c: different letters on the same line indicate a significant difference  $p<0.05$

**Table 5.** TOS levels measured in tissue samples

TOS ( $\mu\text{mol H}_2\text{O}_2$ Equiv./L)	C $\bar{x}\pm\text{SE}$	S $\bar{x}\pm\text{SE}$	T $\bar{x}\pm\text{SE}$	T+S $\bar{x}\pm\text{SE}$	P
LIVER	18,24 $\pm$ 2,03 <sup>b</sup>	28,82 $\pm$ 1,56 <sup>a</sup>	35,32 $\pm$ 1,49 <sup>a</sup>	22,51 $\pm$ 1,85 <sup>b</sup>	0,000
LUNG	13,24 $\pm$ 1,89	16,17 $\pm$ 1,46	16,75 $\pm$ 1,39	13,26 $\pm$ 1,73	0,129
KIDNEY	12,64 $\pm$ 2,31	12,62 $\pm$ 1,78	16,10 $\pm$ 1,70	12,56 $\pm$ 2,11	0,361
BRAIN	7,46 $\pm$ 1,98 <sup>b</sup>	13,14 $\pm$ 1,53 <sup>ab</sup>	17,26 $\pm$ 1,46 <sup>a</sup>	15,44 $\pm$ 1,81 <sup>ab</sup>	0,001
TESTIS	11,74 $\pm$ 1,25	11,74 $\pm$ 0,97	13,77 $\pm$ 0,92	10,91 $\pm$ 1,14	0,193

TOS: Total oxidant status; C: Control Group; T : Sub-chronic Thinner Inhalation Group; T+S: Sub-chronic Thinner Inhalation + Safranal Group; S: Safranal Group;  $\bar{x} \pm \text{SE}$ : mean  $\pm$  standard error; a,b,c: different letters on the same line indicate a significant difference  $p < 0.05$

**Table 6.** NO<sub>x</sub> levels measured in tissue samples

NO <sub>x</sub> ( $\mu\text{mol/g tissue}$ )	C $\bar{x}\pm\text{SE}$	S $\bar{x}\pm\text{SE}$	T $\bar{x}\pm\text{SE}$	T+S $\bar{x}\pm\text{SE}$	p
LIVER	7,23 $\pm$ 1,314 <sup>ab</sup>	12,09 $\pm$ 1,02 <sup>a</sup>	9,32 $\pm$ 0,97 <sup>a</sup>	3,00 $\pm$ 1,19 <sup>b</sup>	0,000
LUNG	4,32 $\pm$ 0,89 <sup>ab</sup>	6,86 $\pm$ 0,69 <sup>a</sup>	6,54 $\pm$ 0,66 <sup>a</sup>	2,85 $\pm$ 0,82 <sup>b</sup>	0,000
KIDNEY	5,09 $\pm$ 0,90 <sup>ab</sup>	8,63 $\pm$ 0,69 <sup>a</sup>	6,99 $\pm$ 0,67 <sup>a</sup>	2,29 $\pm$ 0,82 <sup>b</sup>	0,000
BRAIN	3,32 $\pm$ 1,01 <sup>b</sup>	7,71 $\pm$ 0,78 <sup>a</sup>	6,98 $\pm$ 0,74 <sup>a</sup>	3,62 $\pm$ 0,92 <sup>b</sup>	0,000
TESTIS	5,64 $\pm$ 0,95 <sup>ab</sup>	5,87 $\pm$ 0,73 <sup>a</sup>	6,69 $\pm$ 0,69 <sup>a</sup>	2,81 $\pm$ 0,86 <sup>b</sup>	0,006

NO<sub>x</sub>: Nitric oxide; C: Control Group; T : Sub-chronic Thinner Inhalation Group; T+S: Sub-chronic Thinner Inhalation + Safranal Group; S: Safranal Group;  $\bar{x} \pm \text{SE}$ : mean  $\pm$  standard error; a,b,c: different letters on the same line indicate a significant difference  $p < 0.05$

## DISCUSSION

In the present study, it was observed that rats did not exhibit any escape reflexes during the application in the first two weeks of the inhalation, however, rats became more aggressive and exhibited escaping movements after the third week, their urination and excretion increased, and the uncontrolled rotational movements and standing reflexes deteriorated. Such observations obtained during the present study supported the previous findings reported in the literature suggesting that thinner acted as a neurotoxic agent and there existed evidence of intoxication in the central nervous system due to thinner inhalation (Bölükbaş, 2005). It was also observed that the rats in the T+S group exhibited calmer behaviors contrary to the aggressive behaviors detected in the thinner inhalation group. Such finding could be interpreted as the mitigating or suppressing effect of safranal in case of the intoxication due to thinner inhalation. However, such finding should be supported through research conducted at molecular level. The difference between the blood toluene levels of the rats in the T and T + S groups indicates that the groups inhaled the same level of thinner.

Hosseinzadeh et al. (2013) suggest that the oral administration of safranal, in subacute toxicity test did

not induce any toxic effects in many organs. Safranal at doses of 100 mg/kg by oral route for 8 weeks did not cause any death in rats. However, an increase in the movement of animals was observed in the first 5 minutes after the application. This might be related to the irritant nature of most essential oils.

Higher doses and longer durations of thinner inhalation increases the monooxygenases based on CYP-450 and therefore results in ROS (Backes, 1993).. It is acknowledged that the toxic effects of organic solvents on cellular damage occurs due to the initiation of lipid peroxidation by free oxygen radicals such as superoxide anion ferryl and hydroxyl ions or more generally by the ROS (Mattia et al., 1991, 1993; Ahmed-Choudhury et al., 1998). Konuk et al. (2012) reported that sub-chronic thinner inhalation in rats increased free radical production. Additionally, exposure to sub-chronic toluene was reported not only to increase ROS production measures, but also to significantly affect various antioxidants (Kodavanti et al. 2015).

Given that the membrane phospholipids are the main target of free radicals, concentrations of MDA, one of the end products of lipid peroxidation (LP), were reported to be directly related to the severity of damage (Aleksandrovskii et al., 1988). In the pres-

ent study, once the MDA levels of the thinner group were compared with the MDA levels of the control group, it was observed that the MDA levels significantly increased in liver, lung, testis and kidney tissues, however, no significant increase was detected for the brain tissues despite the relative increase. Such findings were found to be consistent with the previous studies (Ulakoglu et al., 1998a, 1998b; Karaözler et al., 2002; Halifeoğlu et al., 2000). It was reported that *in vivo* and *in vitro* exposure to thinner or toluene accelerated the ROS formation in the brain, liver, kidney and lungs of rats and increased lipid (Mattia et al., 1991;1993; Halifeoğlu et al., 2000; Baydas et al., 2005). In parallel with these results, it is possible to state that sub-chronic thinner inhalation increased lipid peroxidation in tissues in the present study. It was reported that saffron and its components exhibited hydroxyl and DPPH radical scavenging activity (Hosseinzadeh et al., 2009; Assimopoulou et al., 2013). The literature review yielded no studies that focused on investigating the effects of safranal on thinner toxicity. In studies conducted through the use of safranal as an antioxidant molecule, it was stated that the increased MDA levels were decreased due to the antioxidative properties of safranal (Hosseinzadeh and Sadeghnia 2005; Mehdizadeh et al., 2013; Sadeghnia et al., 2013; Samarghandian et al., 2014, 2015). In the present study, the tissue MDA levels in T+S groups were found to be close to the levels of the control group and such finding indicated that safranal had a protective effect against the damaging effects of oxidation reactions in the organs and had an antioxidant potential in preventing lipid peroxidation, however such potentials did not exhibit uniform intensities for all tissue types.

GSH protects cells from oxidative damage through a direct interaction with the free radicals of the sulfhydryl group or through acting as a cofactor (Valko et al., 2006). In the present study, once the GSH levels of the tissue samples of the five organs of the T group were compared to those of the control group, it was determined that the GSH levels in the liver, testis, brain and kidney tissues were statistically significantly lower. On the other hand, there existed a statistically insignificant increase in the lung tissues, contrary to the findings in literature. Similar studies that focused on the effects of thinner on GSH levels reported that GSH levels decreased due to thinner inhalation (Baydas et al., 2003,2005; Ulakoğlu et al.,1998a; El-Nabı Kamel and Shehata 2008; Dillioglugil et al., 2005; Ahmadizadeh et al., 2014). In conclusion, the

decreasing GSH levels in all tissues except the lung were considered to be due to the free oxygen radicals such as superoxide anions, ferryl and hydroxyl ions formed by the catabolism of thinner metabolites such as benzyl alcohol and benzaldehyde, which increased due to thinner inhalation. Hazman and Bozkurt (2015) and Samarghandian et al. (2014) reported that safranal increased the GSH levels in diabetic rats and suppressed the oxidative stress based on diabetes. In the present study, it was observed that the GSH levels decreased in the liver, testis, brain and kidney tissues of the T group were back to the control levels in the T+S group. Therefore, the authors considered that the excessive amounts of free radicals due to thinner inhalation were neutralized by safranal and the oxidative stress levels were decreased or safranal increased the biosynthesis GSH.

It was reported that the antioxidant/oxidant status in the body was more simply evaluated by TAS and TOS measurements and gave an insight about the sensitive balance between the *in vivo* oxidants and antioxidants as a result of synergistic interactions (Ghiselli et al., 2000; Erel, 2004; Erel, 2005; Kayar et al., 2015). However, once the antioxidant/oxidant balance was deteriorated in favor of oxidants and oxidative stress occurred, a significant negative correlation could be observed between the TAS and TOS levels (Erel, 2005). Considering the TAS and TOS levels for all tissues, TAS levels were determined to numerically increase in the T+S group when compared to the T group, and the TOS levels were determined to decrease. Once the data from the Safranal group were compared to the data of the control group, the expected TAS-TOS negative correlation could not be observed. On the other hand, similar studies that focused on the effects of thinner on TAS and TOS levels indicated that the decrease in total antioxidant capacity levels due to thinner inhalation was a result of the oxidative stress (Bayil et al., 2008; Konuk et al.,2012 Hazman and Bozkurt (2015) suggested that safranal increased the TAS levels and decreased the TOS levels especially in diabetic rats, hence, acted exactly similar to an antioxidant. The present study also states that safranal has a protective effect against the oxidative damage due to thinner inhalation.

NO is a free radical that is defined as a biological signal molecule and plays an important role in both physiological and pathophysiological processes (Velayutham. and Zweier, 2013). In the presence of stressors, it could be produced by the catalytic ac-

tion of inducible NO-synthase (iNOS) and could be at higher concentrations. NO could lead to damage to proteins, lipids and the DNA through direct or superoxide reaction and thus lead to the formation of a highly reactive peroxide anion (Rahal et al., 2014). In the present study, the increased NO<sub>x</sub> levels in the T groups in comparison to the control group indicated the presence of thinner metabolites and their activated release. While Konuk et al. (2012) determined that nitric oxide metabolites in rats were higher in T groups when compared to the control, Maniscalco et al. (2004) concluded that NO<sub>x</sub> levels increased in the blood of shoe and leather workers, breathing organic solvents such as toluene, xylene and methyl-ethyl ketone. Once the NO<sub>x</sub> levels of the S group were compared to those of the control group, a numeric increase was observed for all tissues. Farahmand et al. (2013) stated that safranal could be effective as a hormetin and in cases of mild oxidative damage that causes the activation of the antioxidative enzymes. Several studies that investigated the anti-inflammatory and antioxidant potentials of safranal reported a decrease in NO<sub>x</sub> level due to safranal administration (Samarhandian et al., 2014; Bukhari et al., 2015; Hazman and Bozkurt 2015). In the present study, decreased NO<sub>x</sub> levels in the T+S group indicated that safranal decreased the thinner-based stressors.

## CONCLUSIONS

In conclusion, it was determined that the oxidative stress in liver, lung, brain, testis and kidney tissues developed due to thinner inhalation was eliminated or mitigated due to safranal administration, either by de-

creasing the NO<sub>x</sub>, MDA and TOS levels or by increasing the GSH and TAS levels. Safranal was considered to have important antioxidant potentials in protecting the studied tissues against the harmful effects of oxidative stress. However, it was evident that safranal did not provide a uniform effect for every tissue. Although safranal appears to be a biological antioxidant, there exists scarce evidence of its effect as a direct chemical antioxidant. However, in the present study, the comparison between the data of the safranal group and the control group indicated close values for NO<sub>x</sub> and TOS levels of the T group and the expected TAS-TOS correlation was not achieved. Such results could indicate the significance of safranal dose in medical applications. Therefore, it was considered that safranal, effective as a hormetin, could also act indirectly as an antioxidant. Within the scope of these outcomes, it becomes essential to consider not to exceed the specific limit values of solvent inhalation in several industries. It is as well considered significant that applying treatment methods that include safranal for the rehabilitation of volatile substance addicts could contribute to the development of new treatment modulations.

## ACKNOWLEDGEMENTS

The present study was supported by the Scientific Research Projects Commission of the Afyon Kocatepe University, with the ID number 15.SAĞ.BİL.13.

## CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

## REFERENCES

- Alayunt Ömer N., Aksoy L, Karafakıoğlu Y S., Sevımlı S (2019) Assessment of Anti-inflammatory and Antioxidant Properties of Safranal on CCl<sub>4</sub>-Induced Oxidative Stress and Inflammation in Rats. *An. Acad. Bras. Ciênc* 91:(2)
- Al-Alousi LM (1989). Pathology of volatile substance abuse: a case report and a literature review. *Med Sci Law*, 29: 189-208.
- Ahmadzadeh M, Amirmoezy C, Pole T (2014) Effects of Toluene on Rat Kidney. *Jundishapur J Health Sci* 6(1): 281-287.
- Ahmed-Choudhury J, Orsler DJ, and Coleman R (1998) Hepatobiliary effects of tertiary-butylhydroperoxide (tBOOH) in isolated rat hepatocyte couplets. *Toxicology and Applied Pharm* 152: 270-275.
- Aleksandrovskii IA, Poiurovskii MV, Neznamov GG, Seredeniia EA, Krasova SB (1988) Lipid peroxidation in emotional stress and neurotic disorders. *Zh Nevro patol Psikhiatr Im S S Korsakova* 88(11): 95-101.
- Assimopoulou AN, Sinakos Z and Papageorgiou VP (2005) Radical scavenging activity of *Crocus sativus* L. extract and its bioactive constituents. *Phytother Res* 19: 997-1000.
- Backes WL, Sequeira DJ, Cawley GF, Eyer CS (1993) Relationship between hydrocarbon structure and induction of P450: Effects on protein levels and enzyme activities. *Xenobiotica* 23:1353-66.
- Baydas G, Reiter RJ, Nedzvetskii VS (2003) Melatonin protects the central nervous system of rats against toluene-containing thinner intoxication by reducing reactive gliosis. *Toxicol Letter* 137: 169-74.
- Baydas G, Özveren F, Tuzcu M, Yaşar A (2005) Effects of thinner exposure on the expression pattern of neural cell adhesion molecules, level of lipid peroxidation in the brain and cognitive function in rats. *European J. of Pharm* 512: 181-187.
- Bayıl S, Cıcek H, Cimenci GI, Hazar M (2008) Effects Of Organic Compounds On Antioxidant Enzymes In Textile Workers. *Archives of Ind. Hygiene and Tox* 59: 283-287.
- Beutler E, Dubon OB and Kelly BM (1963) Improved method for the determination of blood glutathione. *J Lab Clin Med* 61: 882-888.
- Bölükbaşı S(2005) Effects of Thinner Inhalation on Larynx and Nasal Mucosa in Rats. Thesis, Haydarpaşa Numune Training and Research Hospital 2nd ENT Clinic Istanbul.
- Bukhari SI, Pattnaik B, Rayees S, Kaul S and Dhar MK (2015) Safranal of *Crocus sativus* L. Inhibits Inducible Nitric Oxide Synthase and Attenuates Asthma in a Mouse Model of Asthma. *Phytotherapy Res* 29: 617-627.



- Cobanoglu B, Ibrahim H, Ozercan M, Ozercan R, & Yalcın O (2007) The Effect of Inhaling Thinner and/or Cigarette Smoke on Rat Kidneys. *Inh Toxicology* 19 (3): 303-308.
- Dillioglugil MO, Ilgazli A, Maral H, Sengul C, Ozdemir G, Erinc C (2005) Protective effects of N-acetylcysteine on the peroxidative changes of rat lungs exposed to inhalation of thinners. *Respirology* 10: 615-19.
- El-Nabi K, Shehata M (2008) Effect of toluene exposure on the antioxidant status and apoptotic pathway in organs of the rat. *British J. of Biomedical Sci* 65(2): 75-79.
- Erel Ö (2004) A novel automated method to measure total antioxidant response against potent free radical reactions. *Clin Biochem* 37(2): 112-119.
- Erel Ö (2005) A new automated colorimetric method for measuring total oxidant status. *Clin Biochem* 38(12): 1103-1111.
- Farahmand S K, Samini F, Samini M, Samarghandian S (2013) Safranal ameliorates antioxidant enzymes and suppresses lipid peroxidation and nitric oxide formation in aged male rat liver. *Biogerontology* 14: 63-71.
- Ghiselli A, Serafini M, Natela F Scaccini C (2000) Total antioxidant capacity as a tool to assess redox status: critical view and experimental data. *Free Radical Biology & Med* 29: 1106-1114.
- Halifeoglu I, Canatan H, Ustundag B, Ilhan N, and Inanc F (2000) Effect of thinner inhalation on lipid peroxidation and some antioxidant enzymes of people working with paint thinner. *Cell Biochem Func* 18: 263-267.
- Hazman Ö and Bozkurt FM (2015) Anti-inflammatory and antioxidative activities of safranal in the reduction of renal dysfunction and damage that occur in diabetic nephropathy. *Inflammation* 38:4.
- Hazman Ö and Ovalı S (2015) Investigation of the anti-inflammatory effects of safranal on high-fat diet and multiple low-dose streptozotocin induced type 2 diabetes rat model. *Inflammation* 38, 1012-1019.
- Hosseinzadeh H, Shakib SS, Sameni AK, Taghiabadi E (2013) Acute and Subacute Toxicity of Safranal, a Constituent of Saffron, in Mice and Rats. *Iran J Pharm Res Winter* 12:93
- Hosseinzadeh H, Sadeghnia HR (2005) Safranal, a constituent of *Crocus sativus* (saffron), attenuated cerebral ischemia induced oxidative damage in rat hippocampus. *J. of Pharmacy & Pharmaceutical Sci* 8: 394-9.
- Hosseinzadeh H, Shamsaie F, Soghra M (2009) Antioxidant activity of aqueous and ethanolic extracts of *Crocus sativus* L. stigma and its bioactive constituent, crocin and safranal. *Pharmacognosy Magazine* J 5: 419-424.
- Imenshahidi M, Hosseinzadeh H, Javadvpour Y (2010) Hypotensive effect of aqueous saffron extract (*Crocus sativus* L.) and its constituents, safranal and crocin, in normotensive and hypertensive rats. *Phytotherapy Res* 24: 990-994.
- Karafakioğlu YS, Bozkurt MF, Hazman Ö, Fidan A. F (2017) Efficacy of safranal to cisplatin-induced nephrotoxicity. *Biochem J* 474 (7): 1195-1203. doi: <https://doi.org/10.1042/BCJ20160971>
- Karaözler AA, Mehmet N, Batcioglu K (2002) Effects of long-term solvent exposure on blood cytokine levels and antioxidant enzyme activities in house painters. *J of Tox and Env Health* 13 (65): 1237-46.
- Kayar A, Dokuzeylül B, Kandemir, FM, Kirbas A, Bayrakal A, Or ME (2015) Total oxidant and antioxidant capacities, nitric oxide and malondialdehyde levels in cats seropositive for the feline coronavirus. *Vet Medicina* 60(5): 274-281.
- Kodavanti PRS, Royland JE, Moore-Smith DA, Besas J, Richards JE, Beasley TE, Evansky P, Bushnell P (2015) Acute and subchronic toxicity of inhaled toluene in male Long-Evans rats: Oxidative stress markers in brain. *J. NeuroToxicology* 51 10-19.
- Konuk M, Çiğerci H, Şahin T, Fidan FA, Korcan E (2012) Effects of  $\alpha$ -lipoic acid on DNA damage, protein oxidation, lipid peroxidation, and some biochemical parameters in sub-chronic thinner-addicted rats. *Turkish J of Biology* 36: 702-710c.
- Manscalco M, Grieco L, Galdi A, Lundberg JO, Sofia M (2004) Increase in exhaled nitric oxide in shoe and leather workers at the end of the workshift 54 (6): 404-7
- Martinez-Alfaro M, Alcaraz-Contreras Y, Cárabez-Trejo A, & Leo-Amarador GE (2011) Oxidative stress effects of thinner inhalation *Ind.J. of Occup. and Env Med* 15(3): 87-92.
- Marjot R and McLeod AA (1989). Chronic non-neurological toxicity from volatile substance abuse. *Human Toxicol*, 8: 301-306.
- Mattia CJ, Adams JD, Bondy SC (1993a) Free radical induction in the brain and liver by products of toluene catabolism. *Biochem Pharm* 46: 103-110.
- Mattia CJ, Ali SF, Bondy SC (1993b) Toluene induced oxidative stress in several brain regions and other organs. *Mol Chem Neuropathology* 18, 313-28.
- Mattia CJ, LeBel CP, and Bondy SC (1991) Effect of toluene and its metabolites on cerebral oxygen radical formation. *Biochem Pharmacol* 42(25): 879-82.
- Mehdizadeh R, Parizadeh MR, Khooei AR, Mehri S, Hosseinzadeh H (2013) Cardioprotective Effect of Saffron Extract and Safranal in Isoproterenol-Induced Myocardial Infarction in Wistar Rats. *Basic Medical Sci* 16:56
- Miranda KM, Espey MG, Wink DA (2001) A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide - Biology and Chem* 5 (1): 62-71.
- Moghaddasi MS (2010) Saffron chemicals and medicine usage. *Journal of Medicined Plants Res* 4: (6) 427-430.
- Nedzvetskii V, Kirichenko S, Baydas G, Nerush O (2012) Effects of Melatonin on Memory and Learning Deficits Induced by Exposure to thinner. *Neurophysiology* 44: 1.
- Ohkawa H, Ohishi N and Yagi K (1979) Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem* 95: 351-358.
- Rahal A, Kumar A, Singh V, Yadav, B, Tiwari R, Chakraborty S, Dhama K (2014) Oxidative Stress, Prooxidants, and Antioxidants: The Interplay. *Bio Med Research International* 1:19.
- Sadeghnia HR, Kamkar M, Assadpour E, Boroushaki MT, Ghorbani A (2013) Protective effect of safranal, a constituent of *Crocus sativus*, on quinolinic acid-induced oxidative damage in rat hippocampus. *Iranian J. of Basic Medical Sci* 16: 73-82.
- Samarghandian S, Afshari R, Sadati A (2014) Oxidative Stress Indices for Assessing the Preventing Effects of Safranal on Respiratory Distress in Diabetic Rats. *Hindawi Publishing Corporation Scientific World* J 6
- Samarghandian S, Azimi-Nezhad M & Samini F (2015) Preventive effect of safranal against oxidative damage in aged male rat brain. *Exp Animals* 64(1): 65-71.
- Ulakoğlu EZ, Saygı A, Gümüstas MK, Öztekin İ, Kökoğlu E (1998b) Alterations in superoxide dismutase activities, lipid peroxidation and glutathione levels in thinner inhaled rat lung: relationship between histopathological properties. *Pharm Res* 38: (3) 209-214.
- Ulakoğlu EZ, Saygı A, Gümüstas MK, Zor E, Akaya A, Kökoğlu E (1998a) The effect of chronic thinner inhalation on lipid peroxidation in rat lung and liver. *Cerrahpasa J of Med* 29: 75-8.
- Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M (2006) Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chemico-Biological Interactions* 160: 1-40.
- Velayutham M and Zweier JL (2013) Nitric Oxide Signaling in Biology. *Messenger* 2:1.