

Journal of the Hellenic Veterinary Medical Society

Vol 73, No 2 (2022)



Effects of L-carnitine and cryodevices on the vitrification and developmental competence of invitro fertilized buffalo oocytes

Mohammed A Elmetwally, Amira M Mostagir, Fuller W Bazer, Abdelmonem Montaser, Magdy Badr, Wael Eldomany, Ashraf Eldesoky, Mona M Elghareeb, Daniela Jaimes, Yaser Y Lenis, Samy M Zaabel

doi: [10.12681/jhvms.24581](https://doi.org/10.12681/jhvms.24581)

Copyright © 2022, Mohammed A Elmetwally



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

To cite this article:

Elmetwally, M. A., Mostagir, A. M., Bazer, F. W., Montaser, A., Badr, M., Eldomany, W., Eldesoky, A., Elghareeb, M. M., Jaimes, D., Lenis, Y. Y., & Zaabel, S. M. (2022). Effects of L-carnitine and cryodevices on the vitrification and developmental competence of invitro fertilized buffalo oocytes. *Journal of the Hellenic Veterinary Medical Society*, 73(2), 3961-3970. <https://doi.org/10.12681/jhvms.24581>

Effects of L-carnitine and different cryodevices on the vitrification and developmental competence of *invitro* fertilized buffalo oocytes

Mohammed A Elmetwally^{1,2}, Amira M Mostagir^{1,2}, Fuller W Bazer³,
Abdelmonem Montaser^{1,2}, Magdy Badr⁴, Wael Eldomany^{1,2}, Ashraf Eldesoky^{1,2},
Mona M Elghareeb⁵, Daniela Jaimes⁶, Yaser Y Lenis⁷, Samy M Zaabel^{1,2}

¹Department of Theriogenology, Veterinary Medicine Faculty, Mansoura University, Mansoura 36615, Egypt;

²Center for Reproductive Biotechnology, Veterinary Medicine Faculty, Mansoura University, Mansoura 36615, Egypt;

³Department of Animal Science, Texas A&M University, College Station, TX, 77843, USA 77843, USA

⁴Artificial Insemination and Embryo Transfer Department, Animal Reproduction Research Institute, Al Haram, Giza, Egypt

⁵Department of Physiology, Veterinary Medicine Faculty, Mansoura University, Mansoura 36615, Egypt

⁶Faculty of Agricultural Sciences, University of Applied and Environmental Sciences U.D.C.A, Bogota, Colombia

⁷Research group OHVRI, School of Veterinary Medicine, Faculty of Agrarian Science, Universidad de Antioquia, Calle 70, No. 52-21, Medellín, Colombia

ABSTRACT: The cryopreservation of immature oocytes from buffalo provides a continuous non-seasonal source of female gametes. The current study compared different cryodevices and L-carnitine (LC) supplementation to the vitrification medium on the viability and developmental competence of buffalo oocytes. The immature oocytes were vitrified in TCM119 medium supplemented with 0.0, 0.3, 0.6 or 1.2 mg/ml of LC in either a straw (SD), open pulled-straw (OPS), or solid surface device (SSD). Vitrification in a straw resulted in higher rates of recovery (52.7 %) and viability (85.5 %) of oocytes in medium containing 0.6 mg/ml LC ($P<0.05$). Use of OPS resulted in greater recovery (50.2 %) and viability rates (79.7%) in medium containing 0.3 mg/ml LC ($P<0.05$). There were greater recovery (54.1%) and viability (83.3%) rates for medium containing 0.6 mg/ml LC and SSD ($P<0.05$). Abnormalities of the cytoplasm (all cryodevices) and zona pellucida (SD) were higher when medium contained 1.2 mg/ml LC ($P<0.05$). Furthermore, the maturation rates of oocytes were the greatest when using OPS and 0.3 mg LC (46.9 %, $P<0.05$). The rates of fertilization (60.5%), cleavage (27.9 %) and development of fertilized oocytes to blastocysts (16.3 %) were the greatest when using OPS and vitrification medium with 0.3 mg LC ($P<0.05$). In conclusion, the proper LC concentration differs at different cryodevices. The rates of recovery and viability were the greatest for medium with 0.6 mg/ml LC in either SD or SSD. However, when using OPS the best results were recorded when the medium contained 0.3 mg/ml LC.

Keywords: L Carnitine, Buffalo Oocytes, Straw, Open Pulled-Straw, Solid Surface Device

Corresponding Author:

Mohammed A. Elmetwally, DMV, PhD, Department of Theriogenology, Faculty of Veterinary Medicine, Mansoura University, Mansoura, 35516, Egypt
E-mail address: : mmetwally@mans.edu.eg

Date of initial submission: 25-08-2020
Date of acceptance: 01-07-2021

INTRODUCTION

Buffalos have a prominent role in the economy in many developing countries including Egypt. They have many advantages in comparison to cattle such as resistance to many diseases, the ability to utilize low quality roughages, and the production of high-quality milk (Choudhary et al., 2016).

Cryopreservation is used to enhance the handling of gametes from buffalo for both research and commercial purposes (Attanasio et al., 2010). Moreover, cryopreservation of oocytes collected from slaughtered animals of high genetic value offers a chance to overcome the loss of valuable germplasm and to enhance genetic improvement of this species (Wani et al., 2004; Moawad et al., 2019).

There are two main techniques to cryopreserve gametes and embryos: controlled slow freezing (Gautam et al., 2008) and vitrification (Yamada et al., 2007; Busardò et al., 2016). The controlled slow freezing method for cryopreservation of oocytes and embryos frequently has disadvantages due to osmotic shock and intracellular ice crystallization which may severely damage the cells (Ledda et al., 2007).

Vitrification is a valuable method for cryopreservation as cells are exposed to high concentrations of cryoprotectants and ultra-rapid cooling rates prevent formation of icecrystals (Bus et al., 2019; Chang et al., 2019; Gutnisky et al., 2019). Furthermore, vitrification is a simple and fast technique of cryopreservation that does not require expensive equipment (Scholz and Navas 2014; Zolini et al., 2019; Gutnisky et al., 2019).

The high lipid content in buffalo oocytes makes them sensitive to oxidative damage (Liebermann 2002; Asgari et al., 2012; Schiwe et al., 2017). Moreover, vitrification of oocytes decreases the antioxidant glutathione while increases amounts of hydrogen peroxide (H_2O_2) (Kelly et al., 2006; Schiwe et al., 2017). Therefore, the increase in the amount of intracellular lipids in oocytes reduces their rate of survival and embryo quality after cryopreservation (Romek et al., 2009). As physical changes to lipids leads to cryoinjury, large intracellular lipid droplets resolve into small lipid droplets that alter the morphology of oocytes during cryopreservation (Romek et al., 2009).

The mechanical removal of lipid droplets (Nagashima et al., 1995; Bartolac et al., 2018) or the decreasing of their abundance by using chemical agents

(Men et al., 2006) are promising approaches to improve the tolerance of oocytes to cryopreservation. L-carnitine (LC) enhances lipid metabolism in cells of animals and is very important in the transport of fatty acids from the cytosol to the mitochondria to undergo beta-oxidation (Men et al., 2006). Supplementation of embryo culture medium with L-carnitine reduces lipid content in bovine embryos and increases tolerance of bovine oocytes to cryopreservation (Chankitisakul et al., 2013). L-carnitine also has antioxidant activity that protects cells from DNA damage (Jiang et al., 2012; Lahneche et al., 2019). The double role of LC as an antioxidant and an essential component of lipid metabolism makes it a promising agent for improving oocyte cryopreservation efficacy and subsequent embryonic development (El-Shalofy et al., 2017; Moawad et al., 2019).

The present study evaluated the effects of different concentrations of L-carnitine (LC) in vitrification medium and different cryo-devices on the efficacy of oocyte vitrification and warming by evaluating effects on rates of recovery, viability, morphological abnormalities, and developmental competence.

MATERIAL AND METHODS

The present study was carried out in the Theriogenology Department, Faculty of Veterinary Medicine, Mansoura University, Egypt, in collaboration with the Department of Artificial Insemination and Embryo Transfer, Animal Reproduction Research Institute (ARRI), Al-Haram, Giza between January and December 2017. The chemicals used in this study were purchased from Sigma Chemical Co (St. Louis, MO, USA). All experimental procedures were approved by the Ethics Committee of the Faculty of Veterinary Medicine, Mansoura University, 35516 Mansoura, Egypt.

Recovery and classification of immature oocytes

Ovaries from apparently normal reproductive organs were collected from buffalo heifers and adults of unknown breeding history within 30 min following slaughter and evisceration in the EL-sharkawy abattoir as described previously (Mostagir et al., 2019). Briefly, the ovaries were kept in a thermos flask containing warm normal saline during transport to the laboratory within 2-3 h after slaughter. In the laboratory, ovaries were washed in fresh, sterile physiological saline to further remove any contaminants and were dried with sterile paper towels. Follicles (2 to 8 mm) were aspirated by the use of an 18-gauge needle attached to a

10 ml syringe. The oocytes were pooled in a 15 ml falcon tube and allowed 10-15 min to settle to the bottom of the tube. Then, about 5ml of the medium into which the oocytes settled was aspirated and placed into a sterile 100 mm polystyrene petri dish. The cumulus-oocyte complexes (COCs) were recovered and selected by using a stereomicroscope (Nikon, Tokyo, Japan). The collected COCs were transferred into another dish containing fresh pre-warmed washing medium. The oocytes were then classified into one of three grades based on their morphological appearance as previously described by Sadeesh et al., (2016).

Grade a oocytes had more than 4 layers of cumulus cells and homogenous cytoplasm. Grade b oocytes had complete or partial (2-4) layers of cumulus cells and homogenous cytoplasm. Grade c oocytes were denuded with non-homogeneous cytoplasm. Non-spherical, vacuolated, or fragmented oocytes were classified as degenerated. Good quality COCs were those having homogenous cytoplasm and dense compact cumulus cells. Grades a and b oocytes were used in this study. The effects of LC supplementation in vitrification medium and different cryodevices on developmental competence of the immature buffalo oocytes were checked in three replicates (Figure 1).




Figure 1: recovery and classification of immature buffalo oocytes

Vitrification of immature oocytes

Equilibration and vitrification of selected oocytes were done as described by El-Shalofy et al., (2017). In brief, equilibration and vitrification solutions were prepared using TCM 119 medium supplemented with 10% fetal calf serum (FCS) as the basic medium. Five COCs were placed in 50 μ l drops of basic medium (TCM119) for 1 min and then transferred to an equilibrated solution [100 μ l drops of TCM 119 containing

10% EG (ethylene glycol) and 10% dimethylsulfoxide (DMSO)] for 10 min. Then equilibrated COCs were transferred to the vitrification solution (100 μ l drops of TCM 119 containing 20%EG and 20% DMSO), as well as L-carnitine (LC) at 0.0, 0.3, 0.6 or 1.2 mg/ml for 1 min (Phongnimitr et al., 2013). The cryo-devices used for the vitrification process were: i) Straws (SD): five to ten oocytes were immediately loaded in 0.25-ml mini-straws in the middle column of the vitrifying solution separated by air bubbles and sealed with polyvinyl alcohol powder. The straws were pre-cooled by keeping them in liquid nitrogen vapor at a height of about 5 cm from the surface of the liquid nitrogen level (LN₂) to avoid cracking of straws for 1 min, following which they were dipped vertically in LN₂. ii) Open pulled straws (OPS): Oocytes were immediately loaded in open pulled straw (OPS) as described previously (Vajta et al., 1998a; Attanasio et al., 2010) with slight modifications. Briefly, 0.25 ml semen straws were modified by removing the cotton plug and heating them over a hot plate to approximately 100°C for 10-15 seconds and then stretching them manually until the inner diameter was approximately one-half the original diameter. The straws were cooled in the air for 10 sec before being cut in the middle using a razor plate. Subsequently, oocytes of each group were loaded in OPSs by capillary action and pre-cooled by keeping them in liquid nitrogen vapor at a height of about 5 cm from the surface of the LN₂ for 1 min, followed by dipping vertically into the LN₂ tank. iii) Solid surface device (SSD) for cryopreservation was as described by Gupta et al., (2007). In brief, oocytes from a group of 5 COCs with vitrification solution were aspirated into a glass pipette and the solution containing the COCs were put into a dry surface of a hollow metal cube pre-cooled by immersion in LN₂. The pellets of pre-cooled oocytes were placed into cryotubes and immersed into liquid nitrogen.

Warming

Straw: after a period of storage, the vitrified oocytes in straws were thawed in a water bath at 37°C for 1 min before the content of each straw was expelled into a tissue culture dish containing 1, 0.5, 0.25 M sucrose solutions, respectively and held for 3 min in each concentration of sucrose for stepwise rehydration of oocytes. Finally, the COCs were incubated in basic medium (TCM119) for 3min at room temperature.

Open pulled straw: the loaded ends were sub-

merged into 1ml sucrose for 3 min at 37°C; then moved to decreasing concentrations of sucrose solution as described for straws and finally to the TCM 119 for 3min at room temperature.

Solid surface device: the vitrified droplets were warmed by putting them into 1ml of sucrose for 3min at 37°C and then in decreasing concentrations of sucrose as noted previously.

Evaluation of vitrified/warmed immature oocytes quality

Oocyte recovery rate and morphological assessment

The recovery rate was defined as the number of oocytes counted after the end of rehydration compared to the total number of oocytes vitrified (Moawad et al., 2019). Vitrified /warmed oocytes were examined under a stereomicroscope for evaluation of their morphological characteristics as reported by Sharma et al., (2010). The oocytes with a spherical and symmetrical shape and no signs of lyses/degeneration were considered normal, whereas oocytes with ruptured zona pellucida, fragmented cytoplasm, or degenerative signs were classified as abnormal and discarded (Figure 2)

Evaluation of oocytes' viability

The trypan blue exclusion test was used for the evaluation of oocytes' viability. Trypan blue stain is a useful and quick method to assess the initial quality and viability of oocytes, because it provides an assessment of cell membrane integrity, as those cells with damaged or non-intact cell membranes permit the passage of the trypan blue toward the nucleus of the oocyte (Fig. 2). Trypan blue solutions (0.05%) were prepared by dissolving trypan blue in phosphate buffer saline (PBS; pH = 7.0). Oocyte staining was performed at room temperature for 2 min (Mostagir et al., 2019; Moawad et al., 2019). Oocytes (n= 806) were categorized based on the degree of dye exclusion. Unstained oocytes were classified as live and fully stained oocytes were considered dead (Moawad et al., 2019).

In vitro maturation and fertilization

Oocyte maturation: The vitrified-warmed good quality COCs (homogenous oocyte cytoplasm and dense compact cumulus cells, n= 173) were rinsed three times in maturation medium and 10- 15 oocytes were transferred to a 100 μ l drop of maturation medium under mineral oil and placed in an incubator for 20-22 h at 38.5°C and an atmosphere of 5% CO₂ in the air with maximum humidity.

Assessment of maturation rate: Maturation rates were determined under a stereomicroscope by assessing the expansion of cumulus cells. Oocytes with complete or moderate cumulus expansion and oocytes with slight or no cumulus cell expansion, but with the first polar body extruded into the perivitelline space, were considered mature (Fig. 2, Mostagir et al., 2019).

Semen preparation and oocyte fertilization: First, the fertilization dish was prepared by adding 50 μ l drops of modified Tyrode's medium (TALP) into a sterile disposable petri dish and then covered with mineral oil and incubated at 38.5°C with an atmosphere of 5% CO₂ in air with maximum humidity. Mature oocytes were subjected to repeated gentle pipetting to remove surrounding cumulus cell to allow easy penetration of the sperm and washed three times in TALP medium before adding to the fertilization drops (10 oocytes/drop).

Spermatozoa were capacitated in vitro in the TALP medium. Three straws of frozen semen with known concentrations of sperm (2×10^6 sperm/ml) were




Figure 2: Experimental design of the present study (first panel, different treatments). Evaluation of vitrified/warmed immature oocytes quality (second panel, left); oocytes morphology (second panel, middle), cytoplasmic and zona pellucida abnormalities (second panel, right); viability of oocytes (third panel, left); the maturation and fertilization rates (third panel, middle), and cleavage and blastocyst (third panel, right)

thawed for 1 min in a 37°C water bath. Immediately after thawing, the most motile spermatozoa were separated in sperm-TALP medium containing 6 mg/ml BSA for 1 h by the swim-up technique (Mostagir et al., 2019). Then, the uppermost layer of the medium containing the most motile spermatozoa were collected using a plastic Pasteur pipette and transferred into small test tubes containing 3 ml of SP-TALP medium. The freezing medium was washed out by centrifugation at 2,000 rounds per minute (rpm) for 5 min to form sperm pellets. The supernatant was discarded and 1 ml of TALP added and mixed with the pellet before being centrifuged again and the supernatant was discarded again to completely remove the freezing medium. Next, 1 ml TALP medium containing 10 mg/ml heparin (for in vitro capacitation of sperm) was added, mixed with the sperm pellet and incubated at 38.5°C in an atmosphere of 5 % of CO₂ in air with maximum humidity for 10 min before being evaluated for motility. If motility was acceptable (60-80%), 20 µl of semen was co-incubated with each fertilization drop for 18 h.

***In vitro* culture and embryo development**

The fertilization and embryo culture was carried out as described previously (Chung et al., 2007; Räty et al., 2011). The culture dish was prepared by adding 50 µl drops of culture medium (synthetic oviduct fluid - SOF) into a sterile disposable petri dish, covered with mineral oil and incubated at 38.5°C in an atmosphere of 5% CO₂ in air with maximum humidity. The oocytes were freed from loosely bound spermatozoa and any remaining attached cumulus cells by gentle pipetting. The culture medium was replaced every 48 h with fresh medium until day 7 post-incubation to prevent accumulation of ammonium resulting from amino acid degradation and removal of oocytes that had not cleaved. During the culture period, the gentle shaking of the culture dish was done to allow a uniform environment among fertilized oocytes. The percentage of cleaved oocytes was recorded 48 h post-insemination and those that developed to the blastocyst stage were recorded at 5 and 7-days post-insemination, respectively.

Assessment of fertilization and embryonic development

The assessment of fertilization and embryonic development in the present study was carried out as described previously (Chung et al., 2007; Räty et al., 2011). The presumptive zygotes (n= 79) were denud-

ed from surrounding cumulus cells and mounted on slides with coverslips at 18 h post-insemination. Subsequently, they fixed in acetic acid/ethanol (1:3) solution for at least 24 h. The presumptive zygotes were stained with 1% orcein dissolved in 45% acetic acid solution and examined for evidence of fertilization based on the presence of decondensed sperm heads or male pronuclei with their accompanying sperm tails in the cytoplasm. Oocytes with two pronuclei and a clear second polar body were considered to have been fertilized normally. The percentage of cleaved oocytes was assessed at 48 h post-insemination (Fig. 2; Chung et al., 2007).

Statistical analysis

The normality of quantitative parameters was assessed using normal probability plots and the Kolmogorov-Smirnov test generated with the UNIVARIATE procedure of SAS (Elmetwally, 2012; Gohar et al., 2018; Elmetwally et al., 2018). The experimental data are expressed as means ± SEM (standard error of the mean). To determine the effect of cryodevices as well as the different concentrations of L-carnitine on the vitrification and developmental competence of *in vitro* fertilized buffalo oocytes, a one-way analysis of variance (ANOVA) was used followed by Duncan's multiple comparison test. The rates of recovery, viability, abnormalities of cytoplasm and abnormalities of zona pellucida are expressed as percentages. Statistical analyses were done using SAS® (version 9.2, SAS Institute, Cary, NC, USA). Differences were considered to be significant at $P \leq 0.05$.

RESULTS

The recovery rate, the viability rate and certain abnormalities rates (zona pellucida and cytoplasmic) of buffalo oocytes vitrified by using different cryodevices and different L-carnitine treatments

The recovery rate, the rate of normal and abnormal (zona pellucida and cytoplasmic abnormalities) buffalo oocytes after vitrification in straws (SD), open pulled straws (OPS) and solid surface devices (SSD) with different concentrations of L-carnitine (0.0, 0.3, 0.6 and 1.2 mg/ml) are presented in Table 1. The highest recovery rate was recorded when 0.6 mg LC was used at straw and solid surface device groups and when 0.3 mg/ml LC was used at OPS group.

The greatest viability rates of oocytes were observed when using 0.6 mg/ml LC at SD and SSD groups and 0.3 mg LC at OPS group. Viability rates

Table 1: Recovery and Viability of vitrified buffalo oocytes, as well as abnormalities in their cytoplasm and zona pellucida following different treatments with L carnitine and various cryodevices

LC concentration (mg/ml)	SD (straw)	OPS (open pulled straw)	SSD (solid surface device)
oocytes recovery rate % (n/total n)			
controls	35.2 (61/173) ^{b, β}	31.4 (52/166) ^{b, β}	40.8 (53/130) ^{b, α}
0.3	43.7 (86/197) ^{ab, β}	50.2 (74/147) ^{a, α}	48.6 (82/169) ^{ba, α}
0.6	52.7 (95/181) ^{a, α}	39.7 (51/128) ^{b, β}	54.1 (99/183) ^{a, α}
1.2	31.6 (48/152) ^{b, α}	26.8 (42/155) ^{bc, β}	32.1 (63/196) ^{c, α}
Morphologically normal oocytes % (n/total n)			
controls	71.1 (43/61) ^{b, α}	57.6 (30/52) ^{b, β}	62.3 (33/53) ^{c, β}
0.3	70.1 (72/86) ^{b, β}	79.7 (59/74) ^{a, α}	77.8 (64/82) ^{b, α}
0.6	85.5 (81/95) ^{a, α}	61.1 (31/51) ^{b, β}	83.3 (84/99) ^{a, α}
1.2	70.8 (34/48) ^{b, α}	41.1 (17/42) ^{c, β}	66.9 (42/63) ^{c, α}
Zona Pellucida abnormalities % (n/total n)			
Controls	27.6 (48/173) ^{ba, α}	25.8 (43/166) ^{a, α}	23.9 (31/130) ^{a, β}
0.3	24 (47/197) ^{b, α}	19.1 (28/147) ^{b, β}	20.7 (35/169) ^{b, α}
0.6	21.5 (39/181) ^{bc}	18.8 (24/128) ^b	18 (33/183) ^b
1.2	29.8 (45/152) ^{a, α}	25.2 (39/155) ^{a, α}	22.4 (44/196) ^{a, β}
Cytoplasmic abnormalities % (n/total n)			
controls	34 (59/173) ^a	30.7 (51/166) ^c	36.2 (47/130) ^b
0.3	32.2 (64/197) ^{ab}	30.7 (45/147) ^c	30.8 (52/169) ^c
0.6	25.8 (47/181) ^{b, β}	41.3 (53/128) ^{b, α}	27.9 (51/183) ^{c, β}
1.2	38.6 (59/152) ^{a, γ}	84.2 (75/155) ^{a, α}	45.4 (89/196) ^{a, β}

^{a,b,c} significant differences in the same column, ^{a,β,γ} significant differences in the same row

were also high when 0.3 mg/ml was used at SSD groups. The viability rates were similar when using 0.3 and 1.2 mg/ml L-carnitine when compared with control for SD group. The percentage of zona pellucida abnormalities for oocytes after vitrification and warming at all cryo-device groups was higher in 1.2 mg/ml LC concentration and in controls. The highest percentage of cytoplasmic abnormalities was recorded when 1.2 mg/ml LC was used at all cryo-devices groups. The cytoplasmic abnormalities were the highest (84.2%) at OPS when 1.2 mg/ml LC was used (Table 1). Furthermore, the rates of cytoplasmic abnormalities were also high in controls at SD and SSD.

Effects of LC on maturation, fertilization, cleavage, and blastocyst rates of vitrified warmed buffalo oocytes using different cryodevices

The maturation rate of oocytes was greater when using 0.3 mg/ml LC at OPS group (46.9%, OPS, $P<0.05$) than control at the same cryo-device group (30.4%). However, there was no other significant effect of LC concentration or cryo-device on maturation rate. It is important to notice that other groups had also increased maturation rates (0.6 mg/ml LC at SD group: 37.1% and 0.6 mg/ml LC at SSD group: 41.9%). The greatest fertilization rates ($P<0.05$) was observed when 0.3 mg/ml LC and OPS was used (60.5%)

compared to controls of the same group (38.2%), to 0.6 mg/ml LC at SD group (31.7%) and at SSD group (48.7%). The cleavage rate was the greatest ($P<0.05$) for both 0.3 mg/ml LC at OPS group (27.9%) and 0.6 mg/ml LC at the SSD group (30.8%) compared to controls (14.4%; $P<0.05$) and 0.6 mg/ml at SSD group (17.1%).

The rate of blastocyst development was the greatest in 0.3 mg/ml LC at OPS group (16.3%) compared to control group (3.7%; $P<0.05$) and 0.6 mg/ml in SD group (7.3%). Good blastocyst rate was also recorded when 0.6 mg/ml LC was used at SSD group (12.82%) compared to controls.

DISCUSSION

Cryopreservation of certain mammalian embryos is now a common procedure; however, there are substantial differences of the efficiency of this method depending on the stage, species, and origin of cells produced either in vivo or in vitro. Factors associated with variations in those differences include the number of intracellular lipid droplets and the different microtubular structures, chilling injury, and volume/surface ratio, all of which play a crucial role in penetration of the cryoprotectant (Vajta et al., 1998b).

The recovery rate of buffalo oocytes after vitri-

fication and warming by using straws and the solid surface device was increased in medium with 0.6 mg/ml LC compared with the absence of LC and 0.3 or 1.2 mg/ml LC; whereas the recovery rate of oocytes when using OPS was greater in 0.3 mg/ml LC group compared with control, 0.6 and 1.2 mg/ml LC. The significant effect of LC could be due to the amelioration of detrimental effects of vitrification on mitochondrial function, which causes ATP loss and reduces the competence of vitrified buffalo oocytes (Zhao et al., 2011b; Zhao et al., 2011a). LC supplementation improves ATP production in mouse cells (Zare et al., 2017; Zolini et al., 2019). The metabolism of lipids in rats and bovine cells is promoted by L-carnitine which is considered a beta-oxidation factor via fatty acids transport from the cytosol to the mitochondria (Kerner and Hoppel 2000; Chankitisakul et al., 2013). Cryoprotection devices play a crucial role in the success of *in vitro* fertilization procedures. Additionally, the extent of ice crystals contributes to the success of cryobiology processes. In the present study, we found a significant increase in viability rate of oocytes when using 0.6 mg/ml (SD, SSD) or 0.3 mg/ml (OPS) LC compared with control and 1.2 mg/ml LC concentrations. As concern OPS, lower concentration of LC was enough because of the minimum surface area in OPS, which reduces the volume of vitrification solution and allows direct contact between oocytes and liquid nitrogen compared to those parameters when using SD (Vajta et al., 1998b; Attanasio et al., 2010), because the surface area of straw and solid surface device are large; thus, this theoretically needs more L-carnitine concentration to have a potential antioxidant effect. Smaller drop size and a rapid cooling rate improve vitrification success (Arav and Zeron 1997; Arav and Natan 2019). Previous studies on the viability rate of oocytes in buffalo (Sharma et al., 2010; Choudhary et al., 2016), pigs (Cuello et al., 2010), and mice (Zhou et al., 2007) indicated better results when OPS was used.

Oocytes cryopreservation-induced cryodamage causes morphological and functional damage to oocytes and results in poor development. Oocyte cryopreservation is frequently associated with alterations of the zona pellucida and cytoplasm of oocytes after freezing and thawing (Huang et al., 2008; Son et al., 2019). In the present study, we found a higher rate of zona pellucida and cytoplasmic abnormalities to be associated with the highest concentration of LC (1.2 mg) and all cryodevices used. The concentration of 0.6 mg/ml L carnitine showed the lowest rate of zona

pellucida abnormalities when using SD, OPS or SSD; furthermore the concentrations of 0.3 and 0.6 mg/ml when using OPS were also related to low rate of zona pellucida abnormalities, probably because the diameter of OPS is smaller than SD and SSD. Accordingly, the L-carnitine concentrations could inversely related to the surface area. At the same time, the higher zona pellucida abnormalities when using 1.2 mg/ml L-carnitine, may be due to reduced lipid density that is unfavourable for the development competence of oocytes. Consistent with this result, Spricigo et al., (2017) reported that lower concentrations of LC were associated with a significant improvement of vitrification of immature bovine oocytes. Similarly, excessive concentrations of LC may reduce lipid density that is unfavorable to *in vitro* development of bovine embryos (Sutton-McDowall et al., 2012). Therefore, in the current study, we investigated various concentrations of LC to determine the effects on zona pellucida abnormalities and found that 1.2 mg/ml LC was detrimental when compared to 0.3 and 0.6 mg/ml L-carnitine.

In the present study, the use of low doses of L-carnitine improved recovery and survival rates of oocytes and minimized abnormalities in zona pellucida and cytoplasm compared to the absence of LC. However, there is a different threshold that LC affects positively oocyte quality according to the cryo-device used. L-carnitine stimulates lipid metabolism in animal cells because of its role in the transport of fatty acids from the cytosol to the mitochondria for beta-oxidation (Kerner and Hoppel, 2000). Consequently, L-carnitine improves ATP production in animal cells (Vanella et al., 2000). Similar results were obtained after adding LC to the maturation medium before vitrification of *in vitro*-matured calf oocytes (Spricigo et al., 2017).

The success of oocytes vitrification is determined by the developmental competence including *invitro* maturation and fertilization rates, as well as cleavage and blastocysts formation rates. Results of the present study demonstrated that the maturation rate of oocytes was significantly increased when 0.3 mg/ml LC and OPS were used. This may be attributed to the damage of certain cytoplasmic components when no LC added in vitrification medium and the maturation rate and developmental competence of vitrified oocytes was decreased compared to low doses of LC. Ruppert-Lingham et al., (2003) reported that vitrification of germinal vesicle oocytes causes damage to the

surrounding cumulus cells, which consequently disrupts their communication with enclosed oocytes that is very important for the successful completion of *in vitro* maturation and subsequent embryonic development (Mostagir et al., 2019). Thus, supplementation of the basic vitrification medium with LC may avoid that damage (El-Shalofy et al., 2017). The oxidative activity of mitochondria is decreased mildly by vitrification, but increased by adding LC to vitrification medium in mice (Moawad et al., 2019). Furthermore, it is important to notice that according to our results, there is different threshold of LC for different cryo-devices.

There are several reports about lower maturation rates of immature oocytes following cryopreservation by using basic media alone in human (Khalili et al., 2017; Hatirnaz et al., 2018; Son et al., 2019), mouse (Yeo et al., 2008), cow (Isachenko et al., 2001) and ewe (Peng et al., 2013) oocytes. In the present study, the use of OPS to vitrify immature buffalo oocytes improved the maturation rate after warming, perhaps because the tip of the OPS has a small diameter and thin wall to enhance cryopreservation efficiency. Oocytes held in OPS with a very small volume of vitri-

fication solution undergo a faster rate of cooling and warming (a theoretical rate of 20, 000°C/min) than those in conventional straws (2,500°C/min) (Vajta et al., 1998b). Moreover, oocytes in vitrification medium (1-2 µl) in OPS are warmed directly during dilution, quickly expelled (within 1 sec) and immediately diluted. This reduces exposure to inappropriate temperatures. OPS decreases chilling injury, and toxic and osmotic damage of oocytes (Vajta et al., 1998b).

CONCLUSION

In conclusion, it seems that different LC concentrations are positively affect oocyte quality after vitrification at different cryodevices. In the present study, the vitrification of buffalo oocytes could be improved by using OPS and 0.3 mg LC in combination with EG and DMSO as cryoprotectants. Further research is needed to understand the mechanism by which different LC concentrations act at different cryopreservation procedures.

CONFLICT OF INTEREST

None to declare.

REFERENCES

Arav A, Natan Y (2019) The Near Future of Vitrification of Oocytes and Embryos: Looking into Past Experience and Planning into the Future. *Transfus Med Hemother* 46:182-187

Arav A, Zeron Y (1997) Vitrification of bovine oocytes using modified minimum drop size technique (MDS) is effected by the composition and the concentration of the vitrification solution and by the cooling conditions. *Theriogenology* 47 (1):341

Asgari V, Hosseini SM, Ostadhosseini S, Hajian M, Azhdari ZT, Mosaie M, Nasr-Esfahani MH (2012) Specific activation requirements of in vitro-matured sheep oocytes following vitrification-warming. *Mol Reprod Dev* 79:434-444

Attanasio L, Boccia L, Vajta G, Kuwayama M, Campanile G, Zicarelli L, Neglia G, Gasparini B (2010) Cryotop vitrification of buffalo (*Bubalus bubalis*) in vitro matured oocytes: effects of cryoprotectant concentrations and warming procedures. *Reprod Domest Anim* 45:997-1002

Bartolac LK, Lowe JL, Koustas G, Grupen CG, Sjöblom C (2018) Effect of different penetrating and non-penetrating cryoprotectants and media temperature on the cryosurvival of vitrified invitro produced porcine blastocysts. *Anim Sci J* 89:1230-1239

Berkane N, Liere P, Oudinet JP, Hertig A, Lefèvre G, Pluchino N, Schumacher M, Chabbert-Buffet N (2017) From pregnancy to preeclampsia: A key role for estrogens. *Endocr Rev* 38:123-144

Bus A, Langbeen A, Martin B, Leroy JLMR, Bols PEJ (2019) Is the pre-antral ovarian follicle the 'holy grail' for female fertility preservation? *Anim Reprod Sci* 207:119-130

Busardò FP, Vergallo GM, Turillazzi E, Bolino G, Vullo A, Frati P (2016) Accidental Thawing of Embryos, Cryopreserved for Transfer. Two Italian cases, Milan and Rome. *Curr Pharm Biotechnol* 17:321-325

Chang H, Chen H, Zhang L, Wang Y, Xie X, Zhang Y, Quan F (2019) Effect of oocyte vitrification on DNA damage in metaphase II oocytes and the resulting preimplantation embryos. *Mol Reprod Dev* 86:1603-1614

Chankitisakul V, Somfai T, Inaba Y, Techakumphu M, Nagai T (2013) Supplementation of maturation medium with L-carnitine improves cryo-tolerance of bovine in vitro matured oocytes. *Theriogenology* 79:590-598

Choudhary KK, Kavya KM, Jerome A, Sharma RK (2016) Advances in reproductive biotechnologies. *Vet World* 9:388-395

Chung J-T, Tosca L, Huang TH, Xu L, Niwa K, Chian RC (2007) Effect of polyvinylpyrrolidone on bovine oocyte maturation in vitro and subsequent fertilization and embryonic development. *Reprod Biomed Online* 15:198-207

Cuello C, Sanchez-Osorio J, Almiñana C, Gil MA, Parrilla I, Roca J, Vazquez JM, Martinez EA, Rodriguez-Martinez H (2010) Superfine open pulled straws vitrification of porcine blastocysts does not require pretreatment with cytochalasin B and/or centrifugation. *Reprod Fertil Dev* 22:808-817

El-Shalofy AS, Moawad AR, Darwish GM, Ismail ST, Badawy ABA, Badr MR (2017) Effect of different vitrification solutions and cryo-devices on viability and subsequent development of buffalo oocytes vitrified at the germinal vesicle (GV) stage. *Cryobiology* 74:86-92

Elmetwally MA (2012) Clinical applicability of non-invasive Doppler ultrasonography in small ruminants throughout pregnancy. Doctoral dissertation, Hanover University of Veterinary Medicine. <https://d-nb.info/1024417344/34>

Elmetwally MA, Lenis Y, Tang W, Wu G, Bazer FW (2018) Effects of catecholamines on secretion of interferon tau and expression of genes for synthesis of polyamines and apoptosis by ovine trophectoderm. *Biol Reprod* 99:611-628

Gautam SK, Verma V, Palta P, Chauhan MS, Manik RS (2008) Effect of type of cryoprotectant on morphology and developmental competence of in vitro-matured buffalo (*Bubalus bubalis*) oocytes subjected to slow freezing or vitrification. *Reprod Fertil Dev* 20:490-496

Gohar MA, Elmetwally MA, Montaser A, Zaabel SM (2018) Effect of Oxytetracycline Treatment on Postpartum Reproductive Performance in Dairy Buffalo-Cows with Retained Placenta in Egypt. *JVHC* 1:45-53

Gupta MK, Uhm SJ, Lee HT (2007) Cryopreservation of immature and in vitro matured porcine oocytes by solid surface vitrification. *Theriogenology* 67:238-248

Gutnisky C, Morado S, Gadze T, Donato A, Alvarez G, Dalvit G, Cetica PI (2019) Morphological, biochemical and functional studies to evaluate bovine oocyte vitrification. *Theriogenology* 143:18-26

Haturnaz S, Ata B, Haturnaz ES, Dahan MH, Tannus S, Tan J, Tan SL (2018) Oocyte in vitro maturation: A systematic review. *Turk J Obstet Gynecol* 15:112-125

Huang JYJ, Chen HY, Park JY, Tan SL, Chian RC (2008) Comparison of spindle and chromosome configuration in in vitro- and in vivo-matured mouse oocytes after vitrification. *Fertil Steril* 90:1424-1432

Isachenko V, Isachenko E, Michelmann HW, Alabart JL, Vazquez I, Bezugly N, Nawroth F (2001) Lipolysis and ultrastructural changes of intracellular lipid vesicles after cooling of bovine and porcine GV-oocytes. *Anat Histol Embryol* 30:333-338

Jain JK, Paulson RJ (2006) Oocyte cryopreservation. *Fertil Steril* 86:1037-1046

Jiang Y, Han W, Shen T, Wang M-H (2012) Antioxidant Activity and Protection from DNA Damage by Water Extract from Pine (*Pinus densiflora*) Bark. *Prev Nutr Food Sci* 17:116-121

Kelly J, Kleemann D, Kuwayama M, Walker S (2006) effect of cysteamine on survival of bovine and ovine oocytes vitrified using the minimum volume cooling (mvc) cryotop method. *Reprod Fertil Dev* 18:158

Kerner J, Hoppel C (2000) Fatty acid import into mitochondria. *Biochim Biophys Acta* 1486:1-17

Khalili MA, Shahedi A, Ashourzadeh S, Nottola SA, Macchiarelli G, Palmerini MG (2017) Vitrification of human immature oocytes before and after in vitro maturation: a review. *J Assist Reprod Genet* 34:1413-1426

Lahneche AM, Boucheham R, Ozen T, Altun M, Boubekri N, Demirtas I, Bicha S, Bentamene A, Benayache F, Benayache S, Zama D (2019) In vitro antioxidant, DNA-damaged protection and antiproliferative activities of ethyl acetate and n-butanol extracts of *Centaurea sphaerocephala*L. *An Acad Bras Cienc* 91:e20180462

Ledda S, Bogliolo L, Succu S, Ariu F, Bebbere D, Leoni GG, Naitana S (2007) Oocyte cryopreservation: oocyte assessment and strategies for improving survival. *Reprod Fertil Dev* 19:13-23

Liebermann J (2002) Potential Importance of Vitrification in Reproductive Medicine. *Biology of reproduction* 67:1671-1680

Men H, Agca Y, Riley LK, Critser JK (2006) Improved survival of vitrified porcine embryos after partial delipidation through chemically stimulated lipolysis and inhibition of apoptosis. *Theriogenology* 66:2008-2016

Moawad M, Hussein HA, Abd El-Ghani M, Darwish G, Badr M (2019) Effects of cryoprotectants and cryoprotectant combinations on viability and maturation rates of *Camelus dromedarius* oocytes vitrified at germinal vesicle stage. *Reprod Domest Anim* 54:108-117

Mostagir A, Elmetwally M, Montaser A, Zaabel S (2019) Effects of L Carnitine and Cryoprotectants on Viability Rate of Immature Buffalo Oocytes in vitro After Vitrification. *AJVS* 62:45

Nagashima H, Kashiwazaki N, Ashman RJ, Grupen CG, Nottle MB (1995) Cryopreservation of porcine embryos. *Nature* 374:416

Peng J, Li Q, Wigglesworth K, Rangarajan A, Kattamuri C, Peterson RT, Eppig JJ, Thompson TB, Matzuk MM (2013) Growth differentiation factor 9:bone morphogenetic protein 15 heterodimers are potent regulators of ovarian functions. *Proc Natl Acad Sci USA* 110:E776-85

Phongnimitr T, Liang Y, Srirattana K, Panyawai K, Sripunya N, Treetampinich C, Parmpai R (2013) Effect of L-carnitine on maturation, cryo-tolerance and embryo developmental competence of bovine oocytes. *Anim Sci J* 84:719-725

Räty M, Ketoja E, Pitkänen T, Ahola V, Kananen K, Peippo J (2011) In vitro maturation supplements affect developmental competence of bovine cumulus-oocyte complexes and embryo quality after vitrification. *Cryobiology* 63:245-255

Romek M, Gajda B, Krzysztofowicz E, Smorag Z (2009) Lipid content of

non-cultured and cultured pig embryo. *Reprod Domest Anim* 44:24-32

Ruppert-Lingham CJ, Paynter SJ, Godfrey J, Fuller BJ, Shaw RW (2003) Developmental potential of murine germinal vesicle stage cumulus-oocyte complexes following exposure to dimethylsulphoxide or cryopreservation: loss of membrane integrity of cumulus cells after thawing. *Hum Reprod* 18:392-398

Sadeesh EM, Sikka P, Balhara AK, Balhara S (2016) Developmental competence and expression profile of genes in buffalo (*Bubalus bubalis*) oocytes and embryos collected under different environmental stress. *Cytotechnology* 68:2271-2285

Schiwe MC, Zozula S, Nugent N, Waggoner K, Borba J, Gamboa L, Whitney JB (2017) Modified microsecure vitrification: A safe, simple and highly effective cryopreservation procedure for human blastocysts. *J Vis Exp* 121: 54871

Scholz EC, Navas CG (2014) The Mainining of Cryopreservation for in-vitro Fertilization Patients. Recent advances in cryopreservation, Hideaki Yamashiro, IntechOpen, DOI: 10.5772/58746. Available from: <https://www.intechopen.com/books/recent-advances-in-cryopreservation/the-mainining-of-cryopreservation-for-in-vitro-fertilization-patients>

Sharma GT, Dubey PK, Chandra V (2010) Morphological changes, DNA damage and developmental competence of in vitro matured, vitrified-thawed buffalo (*Bubalus bubalis*) oocytes: A comparative study of two cryoprotectants and two cryodevices. *Cryobiology* 60:315-321

Son W-Y, Henderson S, Cohen Y, Dahan M, Buckett W (2019) Immature oocyte for fertility preservation. *Front Endocrinol (Lausanne)* 10:464

Spricigo JF, Morató R, Arcarons N, Yeste M, Dode MA, López-Bejar M, Mogas T (2017) Assessment of the effect of adding L-carnitine and/or resveratrol to maturation medium before vitrification on in vitro-matured calf oocytes. *Theriogenology* 89:47-57

Sutton-McDowall ML, Feil D, Robker RL, Thompson JG, Dunning KR (2012) Utilization of endogenous fatty acid stores for energy production in bovine preimplantation embryos. *Theriogenology* 77:1632-1641

Vajta G, Holm P, Kuwayama M, Booth PJ, Jacobsen H, Greve T, Callesen H (1998a) Open pulled straw (OPS) vitrification: A new way to reduce cryoinjuries of bovine ova and embryos. *Mol Reprod Dev* 51:53-58

Vajta G, Holm P, Kuwayama M, Booth PJ, Jacobsen H, Greve T, Callesen H (1998b) Open pulled straw (OPS) vitrification: A new way to reduce cryoinjuries of bovine ova and embryos. *Mol Reprod Dev* 51:53-58

Vanella A, Russo A, Acquaviva R, Campisi A, Di Giacomo C, Sorrenti V, Barcellona ML (2000) L-propionyl-carnitine as superoxide scavenger, antioxidant, and DNA cleavage protector. *Cell Biol Toxicol* 16:99-104

Wani NA, Misra AK, Maurya SN (2004) Maturation rates of vitrified-thawed immature buffalo (*Bubalus bubalis*) oocytes: effect of different types of cryoprotectants. *Anim Reprod Sci* 84:327-335

Yamada C, Caetano HVA, Simões R, Nicacio AC, Feitosa WB, Assumpção ME, Visintin JA (2007) Immature bovine oocyte cryopreservation: comparison of different associations with ethylene glycol, glycerol and dimethylsulfoxide. *Anim Reprod Sci* 99:384-388

Yeo CX, Gilchrist RB, Thompson JG, Lane M (2008) Exogenous growth differentiation factor 9 in oocyte maturation media enhances subsequent embryo development and fetal viability in mice. *Hum Reprod* 23:67-73

Zare Z, Abouhamzeh B, Masteri Farahani R, Salehi M, Mohammadi M (2017) Supplementation of L-carnitine during in vitro maturation of mouse oocytes affects expression of genes involved in oocyte and embryo competence: An experimental study. *Int J Reprod Biomed (Yazd)* 15:779-786

Zhao X-M, Du W-H, Wang D, Hao HS, Liu Y, Qin T, Zhu HB (2011a) Recovery of mitochondrial function and endogenous antioxidant systems in vitrified bovine oocytes during extended in vitro culture. *Mol Reprod Dev* 78:942-950

Zhao X-M, Du W-H, Wang D, Hao HS, Liu Y, Qin T, Zhu HB (2011b) Effect of cyclosporine pretreatment on mitochondrial function in vitrified bovine mature oocytes. *Fertil Steril* 95:2786-2788

Zhou C, Zhou G-B, Zhu SE, Hou YP, Jin F, Zhao XM, Hong QH (2007) Open-pulled straw (OPS) vitrification of mouse hatched blastocysts. *Anim Biotechnol* 18:45-54

Zolini AM, Carrascal-Triana E, Ruiz de King A, Hansen PJ, Alves Torres CA, Block J (2019) Effect of addition of l-carnitine to media for oocyte maturation and embryo culture on development and cryotolerance of bovine embryos produced invitro. *Theriogenology* 133:135-143