



## Journal of the Hellenic Veterinary Medical Society

Vol 71, No 3 (2020)



## To cite this article:

KOLYGAS, M., LAMPOU, E., DOUKAS, D., TONTIS, D., PAPPAS, I., GOURZIOTI, E., BAKOPOULOS, V., KAKAVA, E., & ATHANASSOPOULOU, F. (2020). Squamous Cell Carcinoma (SCC) in Brown meagre (Sciaena umbra Linnaeus, 1758), a new candidate species for aquaculture in Mediterranean. *Journal of the Hellenic Veterinary Medical Society*, *71*(3), 2399–2406. https://doi.org/10.12681/jhvms.25104

# Squamous Cell Carcinoma (SCC) in Brown meagre (*Sciaena umbra* Linnaeus, 1758), a new candidate species for aquaculture in Mediterranean

M.N. Kolygas<sup>1</sup>, E. Lampou<sup>1</sup>, D. Doukas<sup>2</sup>, D. Tontis<sup>2</sup>, I. S. Pappas<sup>3</sup>, E. Gourzioti<sup>1</sup>, V. Bakopoulos<sup>4</sup>, E. Kakava<sup>1</sup>, F. Athanassopoulou<sup>1</sup>

<sup>1</sup> Laboratory of Pathology of Aquatic Animals, Ichthyology & Aquaculture, Faculty of Veterinary Medicine, University of Thessaly, Greece

<sup>2</sup>Laboratory of Pathology, Faculty of Veterinary Medicine, University of Thessaly, Greece

<sup>3</sup>Laboratory of Veterinary Pharmacology and Toxicology, University of Thessaly, Greece

<sup>4</sup>Department of Marine Sciences, School of Environment, University of The Aegean, Lesvos, Greece

**ABSTRACT:** A case of epithelial tissue tumors on the body of two adult Brown meagres (*Sciaena umbra* Linnaeus, 1758) is reported. Brown Meagre population of 20.000 was imported to sea cages on September of 2003 at 3g, to check the potential use of this species as a new candidate in Mediterranean aquaculture. The population had been kept for 8 years to serve as a genetic pool for further zootechnical investigation. The main pathology observed after 8 years in the sea cages were mainly ocular lesions (cataract) possibly due to natural aging and/or long culture period under high natural light intensity (UV radiation) and/or net injuries at a percentage of 1,32% of total cultured population. Apart from that, on two fish, reddish- white tumorous growths were evident on skin, macroscopically. Histologically, the tumorous growths were diagnosed as Squamous cell carcinoma. Numerous rounded pearls and nests of epidermal proliferated cells were evident, some of which had a non-cellular center. There were also foci of necrosis and irregular deposits of keratin within the tumor tissue. Special stains (Ayoub-Shklar, Haematoxylin-Eosin & Schiff's Periodic Acid) were applied to the tissue for comparison. To our knowledge this is the first presented report of SCC in this species.

Keywords: Squamous cell carcinoma, Ayoub-Shklar Stain, keratin, Sciaena umbra, Brown meagre

Corresponding Author:

Markos N. Kolygas, Laboratory of Pathology of Aquatic Animals, Ichthyology & Aquaculture, Faculty of Veterinary Medicine, University of Thessaly, Greece E-mail address: kolygas@vet.uth.gr

Date of initial submission: 29-09-2019 Date of revised submission: 17-05-2020 Date of acceptance: 11-06-2020

#### INTRODUCTION

he epidermis in teleost fish is a non-cornified strat-L ified squamous epithelium that varies in thickness (from 3 up to 25 cells) (Genten et al., 2009) and is capable of producing intermediate filaments (tonofilaments) of keratinous origin (Garcia et al., 2005) which represent the most abundant proteins (nearly 80% of total protein content) in stratified epithelia of vertebrates (Rao et al., 2015).Intermediate filaments (IFs) are cytoskeletal proteins generating complex networks that support the cell's shape and conjunctions to other cells (maculae adherents or desmosomes) (Schaffeld& Markl on Omary&Coulomb 2004). In cases of abnormal manifestations on fish epidermis, IFs and especially cytokeratins can be a major diagnostic tool on malignancies' differentiation, characterization and origin.

Teleost fish, like other vertebrates can develop benign, premalignant and/or malignant cellular deformities. The etiological agents or suspected factors of such manifestations are contributed to environmental (pollution, carcinogenic chemicals, radiation, toxins), hereditary (genetic predisposition), enzymatic, immunological, traumatic, viral& other constituents(Coffee et al. 2013,Pinkney *et al.* 2004, Landsberg 2002, Pinkney *et al.* 2001, Morrison *et al.* 1996, Harshbarger *et al.* 1993, Ahne *et al.* 1990, Sano *et al.* 1985, Harhsbarger 1972), while main predisposing factors are considered to be age, sex and immunological competence.

In fish, protruding skin tumors are more common in freshwater species (Ribelin & Migaki 1975) while malignant tumors are rarer than papillomata (Poochirian et al. 2011). Squamous cell carcinoma a malignant neoplasm, has been reported in many fish species (Table 1) in a variety of loci.

#### **MATERIALS & METHODS**

#### **Case history**

A cultured population of brown meagres reared in sea cages, had been reported to present bilaterally clouded-opaque eye lenses (Fig.1d) in a percentage of 1,32% of total cultured population, which was subsequently affecting the wholesale price. This population of 20.000 brown meagres was transferred to sea cages on September of 2003 at 3g, to check the potential use of this species as a new candidate in Mediterranean aquaculture. The population had been kept for 8 years to serve as a genetic pool for further zootechnical investigation. After 8 years, when the G2 generation was ready to substitute the G1 broodstock, the whole population of almost 20.000 brown meagres was harvested. The main pathology observed after 8 years in the sea cages were mainly ocular lesions(cataract) possibly due to natural aging and/or long culture period under high natural light intensity and/or net injuries. Apart from that, tumors were evident. During the scheduled harvesting, fish sampling took place during the packaging process in order to investigate the cause of the problem.In the course of fish sampling, two (2)brown meagres had distinctive solitary protruding lesions (one on each fish) similar in texture (Fig.1). More specifically, the first fish (fish No1) had a protruding formation of solid texture on the proximal ventral left site of juncture between os angulo-articulare and os dentale (Fig1b, arrow). Circumferential skin was pale with no further obvious anomalies (ulcers or trauma). The second fish (fish No2) had a large hemispherical, ulcerative mass ( $\approx$ 5,5x3,1x2,9cm) that was posteriorly covering the biggest area of the left pectoral fin and anteriorly was extended up to the distal end of the left operculum. The dorsal site of the mass extended near the fish's left lateral line. The lesion was strongly attached to the subjacent tissue, with no indication of fluctuated content, but with multifocal necrosis (Fig1c, arrows) and ulcerative processes (Fig1c, white asterisk).

#### Microbiology

Prior to parasitological examinations, tryptone soy agar (TSA) and thiosulphate citrate bile salt agar (TCBS) media, inoculated from kidney and spleen samples, while *Flexibacter Maritimus Medium* (FMM) and Marine Agar (MA) media inoculated from skin and gill samples. All petri dishes were incubated at 21<sup>o</sup>C (sea water temperature). Bacterial Identification was assessed biochemically using API 20E®, BioMérieux®.

#### Parasitology

Skin examination was initially conducted macroscopically. Microscopic examination of wet mount scrapings was performed only from suspicious skin loci. Parasitological examination was conducted in gills, skin, intestine and bile. Specially, bilateral gill epithelium scrapings from the first and second gill arches, was examined under light microscope for parasites. Concerning to intestine's parasitological examinations, wet mount scrapings from pars anterior, medial and posterior *canalis intestinalis* was performed. Bile was excised and handled similarly to the intestine.

Species			
Common name	Scientific name	Habitat	Reference
Bream	Abrami ssp.	Freshwater	Hanjavanit & Mulcahy 2004
Yellowbullhead	Ameiurus natalis	Freshwater	Hanjavanit & Mulcahy 2004
Brownbullhead	Ameiurus nebulosus	Freshwater	PInkney et al 2011,
Blackbullhead	Ameiuru smelas	Freshwater	R.T.L.A. (Sterling VA) Hanjavanit & Mulcahy 2004
American eel	Anguilla rostrata	Catadromous	Hanjavanit & Mulcahy 2004
Oscar	Astronotus ocellatus	Freshwater	Rahmati-Holasoo & Hobbenaghi 2010
Barbel	Barbus barbus	Freshwater	Keysselitz 1908, Clunet 1910 (on Ribelin & Migaki 1975)
Gulf menhaden	Brevoortia patronus	Marine	Fournie et al. 1987
Goldfish	Carrassius auratus	Freshwater	Hanjavanit & Mulcahy 2004
Whitesucker	Catostomus commersoni	Freshwater	Hanjavanit & Mulcahy 2004
-	Chondrostoma soetta	freshwater	Mazzarelli, 1910 (on Ribelin & Migaki 1975)
Mirrorcarp	Cyprinus carpio	Freshwater	Manera & Biavati 1994, Wildgoose 1992
Malawicichlid	Cyrtocara moorii	Freshwater	Hanjavanit & Mulcahy 2004
Saithe	Gadus virens	Marine	Williams 1929 (on Ribelin & Migaki 1975)
Gudgeon	Gobio gobio	Freshwater	Mawdesley-Thomas & Bucke 1967
Black Tetra	Gymnocorymbus ternetzi	Freshwater	Sharifpour et al. 2014
Hybrid sunfish	Lepomi ssp.	Freshwater	Fitzgerald et al. 1991
Ide	Leuciscus idus	Freshwater	Mawdesley-Thomas 1972b (on Ribelin & Migaki 1975)
Whiting	Merlangius merlangus	Marine	Johnstone,1923-25,1924,1924-25 (on Ribelin & Migaki 1975)
Grey mullet	Mugil cephalus	Marine-Freshwater	Poochirian et al. 2011,
Chinooksalmon	Onchorhynchus tshawytscha	Anadromous	Hanjavanit & Mulcahy 2004
Medaka	Oryzias latipes	Amphidromous	Hanjavanit & Mulcahy 2004
European smelt	Osmerus eperlanus	Marine-Freshwater	Breslauer, 1915-16 (on Ribelin & Migaki 1975)
Atlantic rainbow smelt	Osmerus mordax	Anadromous	Hanjavanit & Mulcahy 2004, Herman 1988, Morrison & MacDonald 1995
Pudgy cuskeel	Parabassogigas crassus (Invalid) Spectrunculus grandis (valid)	Marine	Mawdesley- Thomas 1971a (on Ribelin & Migaki 1975)
Congo tetra	Phenacogrammus interruptus	Freshwater	Hanjavanit & Mulcahy 2004
Roach	Rutilus rutilus	Freshwater	Hanjavanit & Mulcahy 2004
Salmon	Salmo salar	Anadromous	Roberts 1972
Rudd	Scardinius erythrophthalmus	potamodromous	Hanjavanit et al. 1990, Hanjavanit & Mulcahy 2004
Brown meagre	Sciaena umbra	Marine	Kolygas et al (Current article)
Sea stickleback	Spinachia spinachia	Marine	Murray, 1908 (on Ribelin & Migaki 1975)
Alaska pollock	Theragra chalcogramma	Marine	Takahashi, 1929 (on Ribelin & Migaki 1975)

## Table 1. Cases of Squamous cell Carcinoma in fish



Figure 1: a) spherical, ulcerated tumor at the level of the lateral line in *Sciaena umbra*, b) small protruding formation on the left mandible, c) tumorous growth, note the necrotic loci (arrows) and the ulcerative site (white asterisk), d) cloudy-opaque eye lenses was the initial reason of examination.

#### Histopathology

Both lesions were excised, and tissue blocks were fixed in 10% formalin and Davidson's fixative. Tissue process was according to routine protocols. All tissue sections (5µm) were stained with Haematoxylin - Eosin (H&E), Periodic-Acid Schiff's(PAS) and Ayoub-Shklar (A-S)(Bancroft & Gamble 2008, Rao et al. 2005). In the A-S staining procedure no phosphotungstic acid(PTA) was used, the reason for this change in A-S staining protocol is explained in discussion's chapter. Sections were deparaffinized and hydrated to distilled water. Acid fuchsin5% added for 3min and slides were transferred directly to 0,5% aniline blue-1% orange G solution for 45min. Dehydration was performed on two 5min changes on 95% alcohol followed by one 3min, 100% EtOH change. After dehydration, two 3min changes on Xylene followed and eventually slides were mounted. Moreover, after necropsy, tissues from main internal organs such as liver, spleen and kidney were also selected for histopathology. Blood smears were impossible to be conducted given the fact that fish sampling took place during manufacture packaging process.

#### RESULTS

#### Microbiology

After a 72h incubation period of FMM and MA inoculated media from skin samples of both fish, colonies of distinct bacteria resembling to *Tenacibaculum maritimum* (Pazos et al., 1996) were observed and confirmed by the biochemical profile of the bacterium (Kolygas et al., 2012, Gourzioti et al., 2016). After a 48h incubation period of TSA and TCBS inoculated media from spleen and kidney, growth of *Listonella* (*Vibrio*) anguillarum was observed on fish No2samples.

#### Parasitology

Parasitological examination of the gills indicated a mild infection from monogenean parasites of the genus *Lamellodiscus* (5,25±2,12 parasites/gill arch). Wet skin and bile mounts showed no presence of parasites. In contrast, wet mounts of intestine from all three parts on both fish revealed scolices (mean intensity 0,43 parasites/ optical field).

Microscopical studies from both fish lesions showed similar histopathological pattern of well differentiated squamous cell carcinomas (Fig.2).More specifically, multiple aggregates of tumor cells nests (Fig2.c), with pleomorphic characteristics (Fig.2a) were abundant. Certain areas of well differentiated squamous epithelial cells (resembling more or less to normal fish epidermis) forming projections into the chorium, exhibited a pattern with basosquamous cells on the nest's periphery (Fig.2a, arrow), and squamous keratinogenous cells at the center (Fig.2b, asterisk). The intermediate space between the nests, contained layers of connective tissue infiltrated by inflammatory cells (lymphocytes). Tissue from liver, spleen and kidney did not indicate any histopathological findings of neoplasm metastasis.



Figure 2: a) pleomorphic epithelial cells (arrow) inside a nest (H&E stain), b) wide areas of necrotic tissue (arrows) surrounding the well differentiated clusters of tumor cells (tumor cell nests). Note the squamous keratinogenous cells at the center of these nests (asterisk), (H&E stain), c) tumor's diathesis on creating digitiform tumors strands, (H&E stain), d) tumor's diathesis on creating satellite growths. The mature keratinogenous nest (center) gave early, small proximal satellite foci (asterisk) tumors (H&E stain).

#### **Comparison between stains**

Standard H&E stain on low magnifications (40x-100x) displayed a clear differentiation between the connective tissue and the tumorous growths, mainly because of the more compact distribution of epithelial cells and the hyperchromatic diathesis, of the latter. In higher magnification powers ( $\geq$ 400x) early development of tumorous cell nests (Fig.2d, asterisk), was difficult to diagnose. The transitional states of cell nests and keratin accumulations on the more matured ones, was problematic to distinguish. Some cells nests had higher stain accumulations than other, and that can be mistakenly attributed to respective keratin accumulation, but in fact this correlation is precarious as necrotic content within the cell nests

can give the same results (Fig.3, H&E, 100x &400x). Staining with PAS, gave similar results to H&E on lower magnification. More specifically, discernment between the peri-tumorous connective tissue and the tumorous cells nests was fair in low magnifications (40x, 100x). Higher magnification powers ( $\geq$ 400x) gave the same disadvantages as H&E stain. Ayoub-Shklar stain displayed better diagnostic potentials in comparison to afore mentioned stains, as it was able not only to highlight early developmental tumorous cell nests and early satellite growths (Fig.3, A-S, 100x) but also to distinguish their transitional level with escalating stain intensities on keratin accumulation (Fig.3, A-S, 400x).



Figure 3. Pictomicrograph comparison of SCC stained with H&E, Schiff's Periodic Acid (PAS), and Ayoub-Shklar (A-S), respectively at different magnification powers.

#### DISCUSSION

Both cases of Brown meagres lesions, exhibit tumorous growth diagnosed as Squamous Cell Carcinomas. To our knowledge this is the first occurrence of malignancy of this type on Brown meagres. The causative agent of this malignant manifestation in these two fish is unknown, and only speculations on the possible etiology can be made. Many authors have indicated several possible agents that can give rise to such cellular abnormal proliferations (Roberts, 2001, Hanson et al, 2011). The microbiological and parasitological findings are considered of minor importance and at some level expected given the health status of both fish. Regarding the staining methods, Ayoub-Shklar stain exhibited higher specificity than H&E and PAS in squamous cell carcinomas. Considering that the tumors were well differentiated, the risk of misinterpreted diagnosis with H&E and PAS stain was low, but in poor differentiated tumorous growths, an accurate diagnosis is highly compromised under the use of unspecific keratin stains. Keratin as a marker can provide not only concluding diagnosis in SCC but also key differential histological evidence between mesenchymal and epithelial malignancies (Ramulu et al. 2013). Immunohistochemical methods are more sensitive diagnostic tools for keratin demonstration (Karantza 2011) but are expensive and time consuming (Rao et

al. 2014).Moreover, specific stains can demonstrate histologically the degree of keratinization (Ramulu et al, 2013). Diagnostic misinterpretations between a) mature versus early SCC satellite keratin nests and b) keratin accumulations versus necrotic content within the nests, in histological sections are likely to occur when basic H&E protocols are used, in contrast to Ayoub-Sklar which is a quick, cost effective stain that can provide adequate specificity on keratin exposition in SCC cases and set definitive diagnosis on SCC in fish.

Another crucial parameter that always must be taken into consideration when cases of SCC are suspected is whether to apply or not phosphotungstic acid in Ayoub-Shklar staining protocol. In fact, the presence of PTA can dramatically change the final stain outcome of A-S on keratin demonstration of SCC cases. During a comparative staining trial on keratin demonstration it is mentioned that in some SCC cases, A-S stain failed to uniformly stain malignant manifestations of keratin pearl demonstrations (Ramulu et al 2013). Moreover, during a similar comparative staining trial of keratin demonstration, the conclusion was that H&E's staining specificity and intensity is highest when compared to special stains such as Ayoub-Shklar (Rao et al 2014), which is in contrast to the conclusions of the present study. The reason

for these dissenting results could be that the presence of PTA during A-S stain on afore mentioned studies, possibly reduces stain's overall specificity, as far as keratin demonstration is concerned. The presence of PTA possibly transitions the staining outcome closer to Masson's Trichrome stain, a stain that is used to distinguish collagen fibers versus smooth muscles and not keratinogenous cells, keratin and prekeratin. In fact, the role of PTA in histology is to bind to prestained specific endocellular and ectocellular molecules in order to selectively decolorize them (Everett & Miller 1974). In trichrome stains like Masson & Mallory, PTA is used to mordant anionic dyes to connective tissue fibers. In the case of A-S stain protocol, PTA might have the same effect on some keratin nests of certain maturity states, reducing the overall specificity. To either accept or reject this hypothesis, further studies on different staining protocols need to be made.

Conclusively, the above A-S stain protocol can be adopted in in-house labs as a quick and cost-effective diagnostic procedure especially on broodstock facilities where early intervention is of major importance on this high-value livestock.

#### **CONFLICT OF INTEREST**

None declared by the authors.

#### REFERENCES

- Ahne, W., Anders, M., Halder, M., Yoshimizu, M. (1990). Isolation of picornavirus-like particles from the European smelt, Osmeruseperlanus (L.). Journal of Fish Diseases 13:167–168.
- Bancroft, J.D. & Gamble, M. (2008) Theory and Practice of Histological Techniques. 6th Edition, Churchill Livingstone, Elsevier, China.
- Bragulla, H. H., &Homberger, D. G. (2009). Structure and functions of keratin proteins in simple, stratified, keratinized and cornified epithelia. *Journal of Anatomy*, 214(4), 516-559. doi:10.1111/j.1469-7580.2009.01066.x
- Coffee, L. L., Casey, J. W., & Bowser, P. R. (2013). Pathology of tumors in fish associated with retroviruses: A review. *Veterinary Pathology*, 50(3), 390-403. doi:10.1177/0300985813480529
- Everett, M. M., & Miller, W. A. (1974). The role of phosphotungstic and phosphomolybdic acids in connective tissue staining I. histochemical studies. *The Histochemical Journal*, 6(1), 25-34. doi:10.1007/ BF01011535
- Fitzgerald, S.D., Carlton, W.W., Sandusky, G. (1991). Metastatic squamous cell carcinoma in a hybrid sunfish. *Journal of Fish Diseases*. 14:481–488.
- Fournie, J.W., Vogelbein, W.K., Overstreet, R.M. (1987). Squamous cell carcinoma in the gulf menhaden, *Brevoortiapatronus Goode. Journal of Fish Diseases* 10:133–136. doi:10.1111/j.1365-2761.1987. tb00728.x
- García, D. M., Bauer, H., Dietz, T., Schubert, T., Markl, J., &Schaffeld, M. (2005). Identification of keratins and analysis of their expression in carp and goldfish: Comparison with the zebrafish and trout keratin catalog. *Cell and Tissue Research*, 322(2), 245-256. doi:10.1007/

s00441-005-0031-1

- Genten, F., Terwinghe, E., Danguy, A. (2009) Atlas of fish Histology, Science Publishers
- Gourzioti, E., Kolygas, M.N., Athanassopoulou, F., Babili, V. (2016). Tenacibaculosis in aqauaculture farmed marine fish. *Journal of Hellenic Veterinary Medical Society*, 67(1):21-31
- Quaedvlieg, P. J. F., Creytens, D. H. K. V., Epping, G. G., Peutz-Kootstra, C. J., Nieman, F. H. M., Thissen, M. R. T. M., &Krekels, G. A. (2006). Histopathological characteristics of metastasizing squamous cell carcinoma of the skin and lips. *Histopathology*, 49(3), 256-264. doi:10.1111/j.1365-2559.2006.02472.x
- Hanjavanit, C., & Mulcahy, M. F. (2004). Squamous cell carcinoma in ruddscardinius erythrophthalmus: Histopathology, ultrastructure, and transmission. Diseases of Aquatic Organisms, 61(3), 215-226.
- Hanjavanit, C., Mulcahy, M.F., Doyle, C.T. (1990). Squamous-cell carcinoma in ruddScardinius erythrophthalmus. Dis Aquat Org 8:155–156
- Hanson, L., Dishon, A., & Kotler, M. (2011). Herpesviruses that infect fish. Viruses, 3(11), 2160-2191. doi:10.3390/v3112160
- Harshbarger, J.C. (1972) Work of the Registry of Tumours in Lower Animals with emphasis on fish neoplasms. Symposia of the Zoological Society of London 30:285–303.
- Harshbarger, J.C., Spero, P.M., Wolcott, N.M. (1993). Neoplasms in wild fish from the marine ecosystem emphasizing environmental interaction. In: Couch J.A., Fournie J.W. (eds) Pathobiology of marine and estuarine organisms. CRC Press, Boca Raton, FL.
- Herman, R.L. (1988). Squamous cell carcinoma in rainbow smelt, Osmerusmordax. Journal of Diseases of Aquatic Organisms 5:71–73

- Karantza, V. (2011). Keratins in health and cancer: More than mere epithelial cell markers. *Oncogene*, 30(2), 127-138. doi:10.1038/ onc.2010.456
- Kolygas, M. N., Gourzioti, E., Vatsos, I. N., &Athanassopoulou, F. (2012). Identification of tenacibaculummaritimum strains from marine farmed fish in greece. *Veterinary Record*, 170(24), 623. doi:10.1136/ vr.100778
- Landsberg, J.H. (2002). The effects of harmful algal blooms on aquatic organisms. *Reviews in Fisheries Science* 10(2):113–390.
- Manera, M., &Biavati, S. (1994). An immuno-histochemical technique used to demonstrate the transition form of a squamous cell carcinoma in a mirror carp, *cyprinuscarpio* L. *Journal of Fish Diseases*, 17(1), 93-96. doi:10.1111/j.1365-2761.1994.tb00349.x
- Matsui, T., &Amagai, M. (2015). Dissecting the formation, structure and barrier function of the stratum corneum. *International Immunology*, 27(6), 269-280. doi:10.1093/intimm/dxv013
- Mawdesley-Thomas, L. E., &Bucke, D. (1967). Squamous cell carcinoma in a gudgeon (gobiogobio, L.). *PathologiaVeterinaria*, 4(5), 484-489..
- Morrison, C.M., Leggiadro, C.T., Martell, D.J. (1996). Visualisation of viruses in tumors of rainbow smelt, Osmerusmordax. Diseases of aquatic organisms 26:19–23.
- Morrison, C. M., & Macdonald, C. A. (1995). Epidermal tumors on rainbow smelt and atlantic salmon from nova scotia. *Journal of Aquatic Animal Health*, 7(3), 241-250.
- Omary B. M., & Coulombe P. (2004). Intermediate filament Cytosceleton. Academic Press. Volume 78, 1st Edition
- Pazos, F., Santos, Y., Macias, A.R., Nuñez, S., Toranzo, A.E. (1996). Evaluation of media for the successful culture of *Flexibactermaritimus*. *Journal of Fish Diseases* 19: 193–197.
- Pinkney, A. E., Harshbarger, J. C., Karouna-Renier, N. K., Jenko, K., Balk, L., Skarphéinsdóttir, H., Rutter, M. A. (2011). Tumor prevalence and biomarkers of genotoxicity in brown bullhead (ameiurus nebulosus) in chesapeake bay tributaries. *Science of the Total Environment, 410-411*, 248-257. doi:10.1016/j.scitotenv.2011.09.035
- Pinkney, A.E., Harshbarger. J.C., May, E.B., Reichert, W.L. (2004). Tumor prevalence and biomarkers of exposure and response in brown bullhead (*Ameiurusnebulosus*) from the Anacostia river Washington, DC and Tuckahoe River, Maryland, USA. *Environmental Toxicology* and Chemistry 23(3):638–647.
- Pinkney, A.E., Harshbarge, J.C., May, E.B., Malancon, M.J. (2001). Tumor prevalence and biomarkers of exposure in brown bullheads (*Am-eiurusnebulosus*) from the Tidal Potomac River Watershed. *Environmental Toxicology and Chemistry* 20:1196–1205.

Poochirian, J. K., Maruthamuthu, N., Sahoo, P. K., & Govindarajan, T.

(2011). Squamous cell carcinoma in grey mullet, mugil cephalus L. (perciformes: Mugilidae). *Comparative Clinical Pathology*, 20(6), 673-675. doi:10.1007/s00580-011-1215-6

- Rahmati-Holasoo, H., Hobbenaghi, R., Tukmechi, A., &Morvaridi, A. (2010). The case report on squamous cell carcinoma in oscar (astronotusocellatus). *Comparative Clinical Pathology*, 19(4), 421-424. doi:10.1007/s00580-010-0963-z
- Ramulu, S., Kale, A. D., Hallikerimath, S., &Kotrashetti, V. (2013). Comparing modified papanicolaou stain with ayoub-shklar and haematoxylin-eosin stain for demonstration of keratin in paraffin embedded tissue sections. *Journal of Oral and Maxillofacial Pathology*, 17(1), 23-30. doi:10.4103/0973-029X.110698
- Rao, R.S., Patil, S., Majumdar, B., Oswal, R.G., (2015). Comparison of special stains for keratin with routinehematoxylin and eosin stain. *Journal of International Oral Health*. 7(3):1-5.
- Ribelin, W. E., & Migaki, G. & Registry of Comparative Pathology (1975). The Pathology of fishes: proceedings of a symposium. University of Wisconsin Press, Madison
- Roberts, R.J. (1972). Oral carcinomata in a salmon (Salmosalar L.). Veterinary Record, 91:199.
- Roberts, R.J. (2001). Neoplasia of teleosts. In: Roberts RJ (ed) Fish Pathology, 3rd edn. WB Saunders, Edinburg, 151–168.
- Rogers, M. A., Winter, H., Langbein, L., Wollschläger, A., Praetzel-Wunder, S., Jave-Suarez, L. F., & Schweizer, J. (2007). Characterization of human KAP24.1, a cuticular hair keratin-associated protein with unusual amino-acid composition and repeat structure. *Journal* of Investigative Dermatology, 127(5), 1197-1204. doi:10.1038/sj. jid.5700702
- Sano, T., Fukuda, H., Furukawa, M. (1985). *Herpesviruscyprini*: biologic and oncogenic properties. *Fish Pathology*20:381–388.
- Schaffeld, M., Markl, J.(2004). Fish Keratins, research article on bookIntermediate filament cytosceleton, Academic Press, Volume 78, Pages 627-671, ISSN 0091-679X, ISBN 9780125641739
- Sharifpour, M. F., Hobbenaghi, R., Alyali, P., &Morvaridi, A. (2014). Squamous cell carcinoma in a black tetra, gymnocorymbusternetzi (boulenger). *Journal of Fish Diseases*, 37(11), 985-987. doi:10.1111/ jfd.12191
- Verzijl, D., Riedl, T., Parren, P. W. H. I., & Gerritsen, A. F. (2017). A novel label-free cell-based assay technology using biolayer interferometry. *Biosensors and Bioelectronics*, 87, 388-395. doi:10.1016/j. bios.2016.08.095
- Wildgoose, W. H. (1992). Papilloma and squamous cell carcinoma in koi carp (cyprinuscarpio). VeterinaryRecord, 130(8), 153-1