



Journal of the Hellenic Veterinary Medical Society

Vol 73, No 1 (2022)



To cite this article:

Gholami-Ahangaran, M., Dehkordi, M. K., Ostadpoor, M., Dastgerdi, A. A., Mahdi, O. A. ., Jawad, M. A. ., & Mustafa, Y. F. . (2022). The synergistic effect of probiotic and phytobiotic for improving growth performance and biological indices in broiler chickens. *Journal of the Hellenic Veterinary Medical Society*, *73*(1), 3681–3688. https://doi.org/10.12681/jhvms.25492

Research article Ερευνητικό άρθρο

The synergistic effect of probiotic and phytobiotic for improving growth performance and biological indices in broiler chickens

M. Gholami-Ahangarn^{1*}, M. Karimi-Dehkordi¹, M. Ostadpoor², A. Ahmadi-Dastgerdi³, O. A. Mahdi⁴, M. A. Jawad⁵, Y. F. Mustafa⁶

¹ Department of Clinical Sciences, Faculty of Veterinary Medicine, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran

² Graduated from Veterinary Medicine Faculty, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran

³ Department of Food Science and Technology, Ardestan Branch, Islamic Azad University, Ardestan, Iran

⁴ Anesthesia Techniques Department, Al-Mustaqbal University College, Babylon, Iraq

⁵ Al-Nisour University College, Baghdad, Iraq

⁶ Department of Pharmaceutical Chemistry, College of Pharmacy, University of Mosul, Mosul-41001, Iraq

ABSTRACT: The use of phytobiotics and probiotics show positive effect on the performance in poultry production and can produce healthy protein sources for humans. In this study, the effects of both commercial probiotic and phytobiotic were evaluated on a chicken experimental model. A total of 300 chicks were divided into 4 groups that fed the basal diet, diet containing probiotic (Protexin®), phytobiotic (garlic extract and Tichoke®), and probiotic plus phytobiotic all over the growing period. The growth indices were measured weekly, analyzed at the 21 and 42 days of age. At 42 days of age, blood samples were collected from all chickens. The concentration of liver enzymes, lipid profiles and antioxidant status were measured in blood samples. Results showed that the weight gain was significantly higher and FCR significantly lower in chickens which received probiotic alone or probiotic plus phytobiotic complex in comparison with chickens which received phytobiotic alone or control chickens (p < 0.05). Furthermore, the addition of phytobiotic plus probiotic showed a significantly increase of blood GPx and TAS level in comparison to chickens that received probiotic or phytobiotic, alone. The level of TG, CHL and LDL were lower and the HDL value was higher in chickens which received phytobiotic or probiotic plus phytobiotic in comparison to chickens fed with probiotic or control chickens (p < 0.05). The level of AST, ALT and ALP showed a significant decrease in chickens fed with probiotic plus phytobiotic. In conclusion, the continuous administration of the studied probiotic and phytobiotic in broiler diets can induce a synergistic effect on growth improvement, increasing of antioxidant capacity, reducing of serum lipids, and improvement of liver function in broiler chickens.

Keywords: Antioxidant, Chicken, Growth Performance, Phytobiotic, Probiotic.

E-mail address: mgholami6@gmail.com

Date of initial submission: 06-12-2020 Date of revised submission: 30-03-2021 Date of acceptance: 17-04-2021

Corresponding Author:

Majid Gholami-Ahangaran, Department of Poultry Diseases, Faculty of Veterinary Medicine, Shahrekord Branch, Islamic Azad University, P.O. Box:166, Shahrekord, Iran

INTRODUCTION

Antibiotics are commonly used in the poultry industry to prevent, and treat infectious diseases (Ben *et al.*, 2019). Some antibiotics promote growth of chickens; reduce mortality by reducing the level or activity of gastrointestinal pathogens, ultimately reducing the absorption of bacterial toxins, improving digestion, and absorption of nutrients (Gholami-Ahangaran *et al.*, 2019; Barton, Hart 2001). In addition, some antibiotics, which play a key role in boosting the immune system, are thought to stimulate the immune response (Gholami-Ahangaran *et al.*, 2020).

Probiotics have long been used as a live beneficial microbe in livestock. In poultry diets, it has positive effects improving the microbial balance of the gastrointestinal tract, (Sekhin et al., 2010). According to the WHO (2001), probiotics are living microorganisms that can cause positive effects in the host. Probiotics may contain one or more strains of bacteria or yeast (Sánchezet al., 2017). The bacteria species commonly used as probiotics include Bacillus, Bifido bacterium, Lactobacillus, Enterococcus, Pediococcus, Leuconostoc, Escherichia coli and Streptococcus, whereas the yeast species, mainly include Saccharomyces cerevisiae and other Saccharomyces species (Pandey et al., 2015; Fijan, 2014, Villena, Kitazawa, 2017). Probiotics benefits include the reduction of the dominance of pathogenic bacteria, balance of microbial populations, and stimulation of immune responses (Sekhon et al., 2010).

Another alternative to the use of antibiotics can be herbs, and their derivatives (Ghoalmi-Ahangaran *et al.*, 2020). The use of medicinal plants in poultry production provides beneficial effects for poultry due to the presence of phenolic, terpenoid and alkaloid compounds (Gheisar, Kim 2018). Phytobiotics include a wide range of plant-derived products which are added to poultry diets to enhance growth performance, improve meat quality, showing immunostimulatory, antimicrobial, anti-inflammatory, and antioxidant properties (Windisch *et al.*, 2008; Kumar *et al.*, 2014). The chemical composition and expected effects of the used phytobiotics depends on the botanical family, part of the plant used (flowers, leaves, seeds, etc.), geographical origin, and harvest season (Bakkali *et al.*, 2008).

Phytobiotics can increase growth by improving taste, increasing the secretion of digestive enzymes (Czech *et al.*, 2009), and modulation of enteric pathogenic bacteria (Ghoalmi-Ahangaran *et al.*, 2020). The aim of our study was to evaluate the effectiveness of a

commercial standard probiotic (named Protexin[®]) and a phytobiotic (an artichoke extract named Tichoke[®] plus garlic extract) on several health indices in broiler chickens.

MATERIAL ANDMETHODS

Study design

The study was undertaken with approval from the Islamic Azad University, Shahrekord Branch ethics committee for care and use of animal for research (Ethical No.: 95-711).

In this study, 300 one-day old broiler chicks (Ross 308) were randomly divided into 4 groups with 5 replicates, so that 15 chickens were allocated in each replicate until 42 days of age. All groups of chickens received freely feed, which was balanced according to Ross 308 production manual (Ross 308 Broiler Nutrition Specification, 2019), ad libitum sanitized drinking water, reared under same growing condition comprising continuous lighting program, mechanical ventilation, at least 50% air humidity, and comfortable temperature (Starting from 32 °C and gradually decreased to 21 °C until end of growing period). All chickens received Newcastle disease (ND) vaccines at 7 (Hiprviar B1, B1 strain), 18 (Nobilis ND Clone 30, Clone 30 strain), 35 (Cevac New L, LaSota strain) days of age.

Chickens of the group one (G1) received a commercial probiotic (Protexin[®], Probiotics International Ltd., UK) according to the manufacturer's recommendation. Chickens of the group2 (G2) received phytobiotic (complexing of prepared garlic extract 100 mg/L, and artichoke extract; Tichoke[®], Goldaru, Iran, 100mg/L). Chickens of the group 3 (G3) received the mentioned probiotic and mentioned phytobiotic by using both mentioned doses. Finally, chickens of the group4 (G4) did not receive any additives in the basal diet and were considered as negative control. The weight gain (WG), feed intake (FI), feed conversion rate (FCR) were measured weekly, and calculated at the 21 and 42 days of age.

At the 42 days of age, chickens of the four groups were weighed, and non-heparinized and heparinized blood samples were taken from wing vein. The serum samples obtained from the non-heparinized were utilized for measuring of humoral antibody against ND vaccine by using the hemagglutination inhibition test according to Allan and Gough (1976). For separation of plasma, the heparinized blood samples were centrifuged at 3, 000 \times g for 15 min at 4°C. All samples were stored at -80°C until analysis was carried out.

The concentration of plasma total protein (TP), triglyceride (TG), cholesterol (CHL), high-density lipoprotein (HDL), low-density lipoprotein (LDL), aspartate transferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP) levels were determined by commercial kits (Pars-Azmoon Co., Tehran, Iran) in a spectrophotometer (Technicon RA1000, H83014 model, Technicon Industrial Systems, Tarrytown, NY), according to the instructions of the manufacturer. Furthermore, the plasma total antioxidant status (TAS) was determined using Randox total antioxidant capacity test kit (Randox Laboratories Ltd., Crumlin, UK) as described by Miller et al. (1993). The blood superoxide dismutase (SOD) activity was measured by Ransod spectrophotometric kit (Ransod, Randox Laboratories Ltd., Crumlin, UK), according to the method previously described by Woolliams et al. (1983). The blood Glutathione peroxidase (GPx) activity was assessed by Ransel spectrophotometric kit (Ransel, Randox Laboratories Ltd., Crumlin, UK) as described by Paglia and Valentine (1967).

Statistical analysis

All data were analyzed with the One-way ANOVA method, using SPSS (version 22) statistical package (SPSS Inc., Chicago, IL, US). Significant differences among the treatments were recognized at P < 0.05 using Tukey test.

RESULTS

Growth performance

At the growth indices in 21 and 42 days of age there was no statistical differences between the chickens of the different groups during the growing period, while the WG and FCR showed statistically significant differences between both ages.

At the 21 days of age, the weight gain was significantly higher in chickens which received the probiotic, or the combination of the probiotic plus phytobiotic in comparison with chickens which only received the phytobiotic or control chickens (P < 0.05).

The comparison of WGs in all treated groups at 42 days of age showed the highest WG and it was obtained in chickens which received both probiotic and phytobiotic that it did not show significant differences with chickens fed with probiotic only. The FCR data in 21 and42 days of age representsthe same pattern as the lowest FCR was noted in chickens which received both probiotic and phytobiotic, and showing no statistically significant differences with those chickens which received probiotic only (Table 1).

Biochemical parameters

The TP in all treated chickens was higher than the TP in the control group. In chickens which received both probiotic and phytobiotic, the TG, CHL, LDL were lower, and the HDL was higher than those parameters observed in chickens only fed with probiotic or the control chickens (p<0.05) (Table 2).

The comparison of the ALT, AST and ALP between the different treated groups showed that chickens that received diet supplemented with phytobioticor probiotic did not have any significant differences with those parameters of the control chickens or chickens which received both probioticand phytobiotic. The minimal value of the ALT, AST and ALP was noted in chickens which received both probiotic and phytobiotic that was significantly lower than the ones of the control group (P<0.05) (Table 2).

The concentration of SOD in all treated groups did not have any significant difference with SOD in control chickens. The GPx and TAS concentration was significantly higher in chickens fed with probiotic alone, phytobiotic alone and probiotic plus phytobiotic. Indeed, the addition of both phytobiotic and probiotic can significantly increase the activity of GPx and TAS in comparison with chickens that received probiotic or phytobiotic alone (P<0.05) (Table 2).

Finally, The IgG antibody titer against ND vaccine in control group was significantly lower than other groups (P<0.05). There was no significant difference in ND titers between chickens which received probiotic alone, phytobiotic alone and probiotic plus phytobiotic (Table 2).

The growth parameters in oroner emercins red with provide and phytoprote at the 21 and 12 augs of age									
Index/ Groups	Feed intake (gr)		Weight gain (gr)		FCR				
	21 days of age	42 days of age	21 days of age	42 days of age	21 days of age	42 days of age			
Probiotic	858±20ª	3620±150 ^a	729±9ª	2234±81ª	1.16±0.04 ^b	1.59 ± 0.03^{b}			
Phytobiotic	863±31ª	3510±183ª	701±15 ^b	2001±83 ^b	1.22 ± 0.03^{a}	1.73±0.02ª			
Probiotic plus	873±30ª	3610±186ª	743±19ª	2284±90ª	$1.16{\pm}0.04^{b}$	$1.58{\pm}0.03^{b}$			
Phytobiotic									
Control	882±18ª	3588±202ª	710±18 ^b	2055±121 ^b	1.24±0.02ª	$1.70{\pm}0.03^{a}$			

Table 1. The growth parameters in broiler chickens fed with probiotic and phytobiotic at the 21 and 42 days of age

• The different superscript in each column represents significant differences between treatment group (P < 0.05).

Table 2. The biochemical parameters in broiler chickens fed with probiotic, phytobiotic and probiotic plus phytobiotic at the 42 days of age

Index/Group	Probiotic	Phytobiotic	Probiotic plus Phytobiotic	Control
TP (gr/dl)	$4.14{\pm}0.50^{a}$	3.61 ± 0.42^{ab}	4.11 ± 0.45^{a}	3.11±0.22 ^b
TG (mg/dl)	94±11 ^b	72±15°	59±19°	125±14 ^a
CHL (mg/dl)	153±26ª	130±20 ^b	133±22 ^b	169±41ª
HDL (mg/dl)	69±15 ^b	75 ± 23^{ab}	$82{\pm}17^{a}$	65±23 ^b
LDL (mg/dl)	63±19ª	44±14 ^b	43±19 ^b	78±20ª
ALT (U/L)	$4.50{\pm}0.7^{ab}$	$4.30{\pm}0.6^{\text{ab}}$	$3.90{\pm}0.7^{\rm b}$	$5.00{\pm}0.8^{a}$
AST (U/L)	152±26 ^{ab}	135±43 ^{ab}	130±46 ^b	166±32ª
ALP (U/L)	2.90±0.23ª	2.77 ± 0.34^{a}	2.53±0.23 ^b	2.95±0.25ª
SOD (U/mg Hb)	1145±90 ^a	1168±103ª	1175±127ª	1106±90ª
GPx (U/mg Hb)	188±26 ^b	139±27 ^b	200±20ª	103±24°
TAS (mmol/L)	$0.74{\pm}0.06^{b}$	$0.77 {\pm} 0.06^{b}$	$0.88{\pm}0.08^{a}$	$0.49{\pm}0.08^{\circ}$
HI titer (NDV)	$4.04{\pm}1.00^{a}$	4.22±1.01ª	4.69±1.54ª	2.30±1.32 ^b

• The different superscript in each line represents significant differences between treatment group (P < 0.05).

DISCUSSION

In our study, the continuous administration of probiotics (Protexin[®]) or probiotic (Protexin[®]) plus phytobiotic (garlic extract and Tichoke[®]) in poultry diets increases growth indices by improving FCR and WG. The consumption of this phytobiotic has no effect on growth indices when was administrated alone in broiler chickens in our experimental model.

The current literature related to the effect of probiotics on growth indices showed very diverse results of using probiotic in poultry diets from not affecting on growth indices to improvement in all growth indices of treated birds (Gunal et al., 2006; Shargh et al., 2012). Gunal et al. (2006) and Shargh et al. (2012) reported that the use of probiotic has no effect on growth indices, while Khosravi et al. (2010) showed that it has no effect on feed intake and final weight but increases food efficiency. In another report, Murry et al. (2006) stated that Lactobacillus-based probiotics increased food efficiency and reduced FI. Also, Awad et al. (2009) noted the improvement in all growth indices following dietary supplementation with a Lactobacillus-based probiotic in chickens. The wide variation in the results of these studies on probiotics seems to be affected by several factors such as growing conditions, diet ingredients, types of probiotics, gastrointestinal pH, presence of stressors, administrated dose, and period of probiotic administration (Wang *et al.*, 2017).

Addition of phytobiotic complex in broiler diets enhanced growth performance at 21 and 42 days of age. The beneficial effect of phytobiotic complex may be partially related to garlic. In a study, Ziarlarimi et al. (2011) stated that garlic powder included many organosulfur compounds such as allicin, alliin, ajoene, diallylsulfide, dithiin, and S-allylcysteine, so that garlic as a natural plant-derived feed additive may be successfully used to improve broiler growth performance. The obtained results are in agreement with the findings of Khan et al. (2012) who observed that broiler treated with garlic powder had significantly higher live weight gain than control group. The same trend was also found by Suriya et al. (2012) who used 0.5% garlic powder in broiler diet. The better effect of garlic as natural feed additives might be due to increased enzymes activity of pancreas, which offer a better environment for digestion and absorption of nutrients (Ismail et al., 2021).

Various studies have been performed to elucidate

the effects of the artichoke plant in poultry (Nadia et al., 2007; Fallah et al., 2013; Khoramshahi et al., 2015; Mirderikvandi et al., 2016). The role of this plant was previously studied in liver protection of Japanese quails (Nateghi et al., 2013; Khoramshahi et al., 2015), improving the performance index of laying hens (Nadia et al., 2007; Yildiz et al., 2006), and lowering cholesterol (Fallah et al., 2013; Abdo et al., 2007) in chickens. Several studies evaluated the effect of dried-leaf-powder or extract of artichoke on FI, WG and FCR in broilers, but showing a wide range on the effect of artichoke on growth indices. These results varied from no effect on growth indices (Tajodini et al., 2015 Mirderikvandi et al., 2016) to decrease of the FCR at the end of the growing period (Rouzmehr et al., 2014), increase of the body weight (Lertpatarakomo et al., 2015), increase of the WG and FI (Boroumandnia et al., 2014), increase body weight, FI and decrease of the FCR (Shokri et al., 2018), and decrease of final weight (Abdo et al., 2007). However, the results of our study are consistent with previously reported works showing no effect on growth indices after administration of artichoke (Mirderikvandi et al., 2016; Tajodini et al., 2015). Several studies have been performed in order to elucidate the effects of probiotics on the antioxidant system in birds (Cross et al. 2002; Erdogan et al. 2010). Amaretti et al. (2013) showed that the use of probiotics increases antioxidant capacity, decreases oxygen radicals and reduces oxidative stress. In fact, probiotics produce butyric acid, hydrogen, which may play a stimulating role in the production of antioxidants and free radical scavenging (Zheng et al., 2019). In our study, the use of probiotics had no effect on the SOD but increased the GPx, and TAS, significantly. Cross *et al.* (2002) and Erdogan et al. (2010) showed that probiotics had no effect on GPx level. In addition, Aluwong et al. (2013) showed that the use of yeast probiotics significantly increased GPx activity without affecting SOD in broilers. In another study, Bai et al. (2016) reported a Bacillus subtilis based probiotic increased antioxidants in broiler breast muscle. This is associated with increased mRNA expression of antioxidant genes, decreased oxidative damage in the pectoral muscle. Later, the same authors (Bai et al., 2017) also demonstrated that the higher expression of SOD and GPx genes in mitochondria of the hepatic cells was directly related to feeding with probiotics in chickens.

In this study, we demonstrated that the use of artichoke extract alone could increase GPx and the combination of the probiotic and phytobiotic (garlic extract and artichoke extract) also have synergistic activity in increasing GPx and TAS, which even showed significant differences with the groups that received probioticor phytobiotic alone. There is scarcity of clinical research studies on the antioxidant properties of artichoke in poultry because most of the published studies have been performed in laboratory animals (Jimene-Escrig et al., 2003). In rats, Jimene-Escrig et al. (2003) studied the antioxidant properties of artichoke and highlighted the in vitro antioxidant properties of this plant detecting increased blood GPx and had no effect on catalase and SOD, which is consistent with the findings of our study. In broilers, Mirderikvandi et al. (2016) stated that adding 500 mg/L of artichoke extract in drinking water for 2 weeks had no effect on MDA and reduced GPx. However, we found the increase in antioxidant activity after the use of the artichoke extract is related to the level of scavenging activity of AE. The antioxidant property of artichoke extract could be related to RSA (40% of AE content) and phenolic contents (5% of AE content, in the form of chlorogenic acid). The role of chlorogenic acid as a potent antioxidant has been previously demonstrated under in vivo and in vitro conditions (Sato et al., 2011).

In the chickens received phytobiotic complex, the levels of TAS and GPx were significantly increased while no effect on SOD. Our results agreed those obtained by Nasiroleslami *et al.* (2018) who showed that broiler fed diet supplemented with guanidinoace-tic acid significantly improved GPx activity in broiler liver and decreased serum MDA. Furthermore, Alagawany *et al.* (2007) demonstrated that dietary supplementation of garlic in diet had positive impact on SOD and TAS activities.

Several properties of probiotics in chickens have been previously studied (Makled *et al.*, 2019, Yan *et al.*, 2019; Ashayerizadeh *et al.*, 2011). The positive effects of probiotics in chickens are the improvement on growth indices, specific and non-specific immune responses, gastrointestinal health (Makled *et al.*, 2019), laying performance, bone strength (Yan *et al.*, 2019), but there is little information about the effect of probiotics in improving liver health in chickens. In our study, we found that the continuous consumption of the studied commercial probiotic can improve liver enzymatic activity, reduce total cholesterol and increase total protein. Several studies on the effect of probiotics on serum lipids in poultry were previously published (Ashayerizadeh *et al.* 2011; Abdulrahim *et*

al., 1996; Mohan et al., 1996). Ashayerizadeh et al. (2011) demonstrated that adding probiotic to broiler diet decrease serum cholesterol level in these animals. In addition, dietary supplementation with probiotic containing Saccharomyces cerevisiae was demostrated to decrease cholesterol in both egg yolk (Abdulrahim et al., 1996) and chicken serum (Mohan et al., 1996). Amer and Khan (2012) showed that the supplementation with probiotic containing Lactobacillus acidophilus, B. subtilis, S. cerevisiae and Aspergillus orvzae decreases cholesterol in chicken serum after 6 weeks of initiate the addition of probiotics. Further to the ability of probiotic in elimination of lipids, these microorganisms can adsorb, detoxify the microbial toxin in the gastrointestinal tract and prevent the intestinal absorption. Detoxification of poisons in GIT inhibits the effect of toxins on hepatocytes (Markowiaket al., 2019). In our study, the increasing of TP may be related to the influence of probiotic on secretory function of the gastrointestinal tract that leads to increase digestion, adsorption, and subsequently can elevate total protein in plasma. It seems that the positive effects of probiotics on physiological function lead to an improvement in growth indices in chickens.

There are few studies previously reported on the effect of probiotics on liver enzyme profiles (Bityutsky *et al.* 2019; Gholami-Ahangaran *et al.*, 2016). Bityutsky *et al.* (2019) reported that that the use of probiotics in quail could reduce the levels of liver enzymes of ALT and AST. The damage to the liver cell membrane causes these enzymes to be released into the bloodstream (Gholami-Ahangaran *et al.*, 2016). Therefore, the lack of increase in liver enzymes indicates no liver damage following probiotic supplementation.

In this study, AE was able to have effect on serum lipid profile, reducing CHL, TG, LDL and increasing HDL. Regarding the effect of artichoke on lipid metabolism, several studies in humans, mammals and poultry agreed that artichoke can positively affect lipid metabolism. Rouzmehr *et al.* (2014) reported that, although the addition of 200 g per ton of dried artichokes to the diet can increase WG and FI, reduced abdominal fat and blood CHL. Also, Abdo *et* al. (2007) stated that consumption of 6% dried artichoke leaves in the diet causes weight loss. However, AE seems to reduce plasma cholesterol levels by increasing bile secretion and decreasing cholesterol biosynthesis (Edward et al., 2015). In addition, there is evidence that the active ingredients in artichokes have the ability to inhibit HMG-CoA reductase (Gebhardt, 1998). In our study, the decrease of hepatic enzymes including ALT, AST and ALP, and decreased plasma lipid profiles may support this hypothesis because if lipids accumulated in hepatocytes, hepatic complications would manifest as elevated hepatic enzymes in plasma. Certainly, this improve of liver health is in accordance to the increase of antioxidant capacity and it can be due to the increased ability of antioxidants to protect liver cells against toxins and oxidants.

Findings of our study showed that supplementing diets with garlic and artichoke extract decreased blood TG and LDL, and increased TP, and HDL. These results are partially in agreement with Rahimi et al. (2011) who found in broilers that blood TG, and LDL concentrations were decreased, while HDL increased when fed enriched-diet with 0.1% garlic powder. At this regard, Issa and Omar (2012) showed that addition of garlic powder at 0.2% and 0.4% reduced significantly concentrations of TG, CHL, LDL in broilers, and increasing the level of HDL compared to control. The potential effect of garlic products may be due to depressing the lipogenic and cholesterogenic activity of liver enzymes such as fatty acid synthase, glucose-6-phosphatase dehydrogenase, malic enzyme, and 3-hydroxy-3-methylglutaryl-CoA reductase, consequently, the mechanism of hypocholesterol and hypolipid syntheses (Mahmoud et al., 2010).

In conclusion, our study demonstrated that the continuous administration of the studied commercial probiotic along with artichoke and garlic extract in broiler chicken diets improved growth indices, increased antioxidant capacity, reduced serum lipids and improved liver function in the experimentally studied broiler chickens.

CONFLICT OF INTEREST

None declared.

REFERENCES

- Abdo MA, Radwan NL, Selim NA (2007) The effect of artichoke leaves meal on the utilization of dietary energy for broiler chickens. Int J Poult Sci 12: 973-982.
- Abdulrahim SM, Haddad MSY, Hashlamoun EAR, Robinson RK (1996) The influence of Lactobacillus acidophilus, Bacitracin on layer performance of chickens, cholesterol content of plasma, egg yolk. BrJ Poult Sci 37: 341-346.
- Alagawany M, Ashour E, Reda FM (2016) Effect of dietary supplementation of garlic (*Allium sativum*) and turmeric (*Curcuma longa*) on growth performance, carcass traits, blood profile and oxidative status in growing rabbits. Ann Anim Sci 16:489-505.
- Allan WH, Gough RE (1976) A comparison between the haemagglutination inhibition, complement fixation tests for Newcastle disease. Res Vet Sci 20: 101-103.
- Aluwong T, Kawu M, Raji M, Dzenda T, Govwang F, Sinkalu V, Ayo J (2013) Effect of Yeast Probiotic on Growth, Antioxidant Enzyme Activities, Malondialdehyde Concentration of Broiler Chickens. Antioxidants 2: 326-339.
- Amaretti A, Di Nunzio M, Pompei A, Raimondi S, Rossi M, Bordoni A (2013) Antioxidant properties of potentially probiotic bacteria, in vitro, in vivo activities. ApplMicrobiol Biotechnol 97: 809-817.
- Amer MY, Khan SH (2012) A comparison between the effects of a probiotic, an antibiotic on the performance of Desi chickens. Vet World 5: 160.
- Ashayerizadeh A, Dabiri N, Mirzadeh KH, Ghorbani MR (2011) Effect of dietary supplementation of probiotic, prebiotic on growth indices, serum biochemical parameters of broiler chickens. J Cell Anim Biol 5: 152-156.
- Awad WA, Ghareeb K, Abdel-Raheem SM, Böhm J (2009) Effects of dietary inclusion of probiotic, synbiotic on growth performance, organ weights, intestinal histomorphology of broiler chickens. Poult Sci 88: 49-56.
- Bai WK, Zhang FJ, He TJ, Su PW, Ying XZ, Zhang LL, Wang T (2016) Dietary Probiotic Bacillus subtilis Strain Increases Antioxidant Capacity, Oxidative Stability of Chicken Breast Meat during Storage. PLoS ONE 11: e0167339.
- Bai K, Huang Q, Zhang J, He J, Zhang L, Wang T (2017) Supplemental effects of probiotic *Bacillus subtilis* on growth performance, antioxidant capacity, meat quality of broiler chickens. Poult Sci96: 74-82.
- Bakkali F, Averbeck S, Averbeck D, Idaomar M (2008) Biological effects of essential oils-a review. Food Chem. Toxicol 46: 446-475.
- Barton MD, Hart WS 2001. Public health risks: antibiotic resistance-review. Asian-Australs J AnimSci 14: 414-422.
- Ben Y, Fu C, Hu M, Liu L, Wong MH, Zheng C (2019) Human health risk assessment of antibiotic resistance associated with antibiotic residues in the environment: a review. Environ Res 169: 483-493.
- Bityutsky V, Tsekhmistrenko S, Tsekhmistrenko O, Melnychenko O, Kharchyshyn V (2019) Effects of Different Dietary Selenium Sources Including Probiotics Mixture on Growth Performance, Feed Utilization, Serum Biochemical Profile of Quails. In: Modern Development Paths of Agricultural Production. Springer, Cham:pp 623-632.
- Boroumandnia Z, Mohammadzadeh S, Azizi M, Khaldari M (2014) The effect of adding artichoke extract to drinking water on performance, carcass characteristics, blood lipoprotein concentration in broiler chickens. Iranian JAppl Anim Sci 3: 71-79.
- Cross ML (2002) Microbes versus microbes: immune signals generated by probiotic lactobacilli, their role in protection against microbial pathogens. FEMS Immunol Med Microbiol 34: 245-253.
- Czech A, Kowalczuk E, Grela ER (2009) The effect of an herbal extract used in pig fattening on the animal performance, blood components. Ann Univ Mariae Curie Sklodowska 27: 25-33.
- Edward SE, da Costa Rocha I, Williamson EM, Heinrich M (2015) Phytopharmacy: An evidence-based guide to herbal medicinal products. UK:John Wiley, Sons.218 pp.
- Erdoğan Z, Erdoğan S, Aslantaş Ö, Çelik S (2010) Effects of dietary supplementation of synbiotics, phytobiotics on performance, caecal coliform population, some oxidant/antioxidant parameters of broilers. J

Anim Physiol Anim Nut 94: e40-e48.

- Fallah R, Kiani A, Azarfar A (2013) Effect of artichoke leaves meal, mentha extract (Mentha piperita) on immune cells, blood biochemical parameters of broilers. Global Vet 10: 99-102.
- FAO/WHO (2001) Report of a joint FAO/WHO expert consultation on evaluation of health, nutritional properties of probiotics in food including powder milk with live lactic acid bacteria.Cordoba: FAO/ WHO Publishing.23.
- Gebhardt R (1998) Inhibition of cholesterol biosynthesis in primary cultured rat hepatocytes by artichoke (*Cynara scolymus* L.) extracts. J Pharmacol Exp Therap 286: 1122-1128.
- Gheisar MM, Kim IH (2018) Phytobiotics in poultry, swine nutrition a review. Italian J AnimSci 17: 92-99
- Gholami-Ahangaran M, Rangsaz N, Azizi S (2016) Evaluation of turmeric (*Curcuma longa*) effect on biochemical, pathological parameters of liver, kidney in chicken aflatoxicosis. Pharm Biol 5: 780-787.
- Gholami-Ahangaran M, Peimani N, Dastgerdi AA (2019) The effect of thyme (*Thymus daenensis*) supplement on growth, hygienic parameters of broilers meat. Iraqi J Vet Sci 33: 87-92.
- Gholami-Ahangaran M, Ahmadi-Dastgerdi A, Karimi-Dehkordi M (2020) Thymol, carvacrol, as antibiotic alternative in green healthy poultry production. Plant Biotechnol Persa 2: 22-25.
- Gilliland SE, Nelson CR, Maxwell C (1985) Assimilation of cholesterol by Lactobacillus acidophilus. ApplEnviron Microbiol 49: 377-381.
- Gunal M, Yayli G, Kaya O, Karahan N, Sulak O (2006) The effects of antibiotic growth promoter, probiotic or organic acid supplementation on performance, intestinal microflora, tissue of broilers. Int J PoultSci 5: 149-155.
- Issa KJ, Omar JMA (2012) Effect of garlic powder on performance and lipid profile of broilers. Open J Anim Sci2:62-8.
- Ismail IE, AlagawanyM, Taha AE, Puvača N, Laudadio V, Tufarelli, V (2021) Effect of dietary supplementation of garlic powder and phenyl acetic acid on productive performance, blood haematology, immunity and antioxidant status of broiler chickens. AnimBiosci 34 (3): 363-370.
- Jimene-Escrig A, Dragsted LO, Daneshvar B, Saura F, Saura-Calixto F (2003) In vitro antioxidant activities of edible artichoke (*Cynara* scolymus L.), effect on biomarkers of antioxidant in rats. J AgricFood Chem 51: 5540-5545.
- Khan RU, Nikousefat Z, Tufarelli V, Naz S, Javdani M, Laudadio V (2012) Garlic (*Allium sativum*) supplementation in poultry diets: effect on production and physiology. Worlds Poult Sci J 68:417-424.
- Khoramshahi M, Samadi F, Ganji F, Zerehdaran S, Samadi S (2015) The effect of artichoke leaf powder on performance, internal organs weight, some blood parameters of Japanese quail poisoned with carbon tetrachloride. Anim SciRes 25: 57-67.
- Khosravi A, Boldaji F, Dastar B, Hasani S (2010) Immune response, performance of broiler chicks fed protexin, propionic acid. Int JPoult Sci 9: 188-191.
- Kumar M, Kumar V, Roy D, Kushwaha R, Vaiswani S (2014) Application of herbal feed additives in animal nutrition-a review. Int J Lives Res 4: 1-8.
- Lertpatarakomo R, Jaipeng P, Sirimongkolworaku S, Paditporn K, Mitchaothai J (2015). Effect of artichoke (*Cynara scolymus* L.) extract supplementation in drinking water on growth performance of broiler chickens, In: 20th European Symposium on Poultry Nutrition, Prague, Czech Republic: pp220-222.
- Mahmoud KZ, Gharaibeh SM, Zakaria HA, Qatramiz AM (2010) Garlic (*Allium sativum*) supplementation: influence on egg production, quality, and yolk cholesterol level in layer hens. Asian-Australas J Anim Sci 23:1503-1509.
- Makled MN, Abouelezz KFM, Gad-Elkareem AEG, Sayed AM (2019) Comparative influence of dietary probiotic, yoghurt, sodium butyrate on growth performance, intestinal microbiota, blood hematology, immune response of meat-type chickens. Trop AnimHealth Prod 51: 2333-2342.

J HELLENIC VET MED SOC 2022, 73(1) ПЕКЕ 2022, 73(1)

Markowiak, P, Śliżewska K, Nowak A, Chlebicz A, Żbikowski A,

Pawłowski K, Szeleszczuk P (2019) Probiotic microorganisms detoxify ochratoxin A in both a chicken liver cell line, chickens. J Sci Food Agricult 99: 4309-4318.

- Miller JK, Brzezinska-Slebodzinska E, Madsen FC (1993) Oxidative stress, antioxidants, animal function. J Dairy Sci 76: 2812-2823.
- Mirderikvandi M, Kiani A, Khaldari M, Alirezaei M (2016) Effects of artichoke (*Cynara scolymus* L.) extract on antioxidant status in chicken thigh meat. Iranian J Vet Med 10: 73-81.
- Mohan B, Kadrivel R, Natarajan A, Bhaskaran M (1996) Effect of probiotic supplementation on growth, itrogen utilization, serum cholesterol in broilers. Br Poult Sci 37: 395-401.
- Murry AC, Hinton Jr A, Buhr RJ (2006) Effect of botanical probiotic containing lactobacilli on growth performance, populations of bacteria in the ceca, cloaca, carcass rinse of broiler chickens. Int J Poult Sci 5: 344-350.
- Nadia LR, Zeinab M, Abdo A, Hassan RA (2007) Effect of feeding artichoke leaves meal on productive, reproductive performance of Mandarah hens. Int J Poult Sci 6: 826-834.
- Nasiroleslami M, Torki M, Saki AA, Abdolmohammadi AR (2018) Effects of dietary guanidinoacetic acid and betaine supplementation on performance, blood biochemical parameters and antioxidant status of broilers subjected to cold stress. J Appl Anim Res46:1016-1022.
- Nateghi R, Samadi F, Ganji F, Zerehdaran S (2013) Hepatoprotective effects of Cynara scolymus L. extract on CCl4 induced liver injury in broiler chickens. Int J Agricult Sci Res 3: 678-688.
- Paglia DE, Valentine WN (1967) Studies on the quantitative, qualitative characterization of erythrocyte glutathione peroxidase.J Lab Clin Med 70: 158-169.
- Pandey KR, Naik SR, Vakil BV (2015) Probiotics, prebiotics, synbiotics-a review. J Food Sci Technol 52: 7577-7587.
- Rahimi S, Teymouri Zadeh Z, Karimi Torshizi MA, Omidbaigi R, Rokni H (2011) Effect of the three herbal extracts on growth performance, immune system, blood factors and intestinal selected bacterial population in broiler chickens. J Agric Sci Technol13:527-539.
- Rai S, Wahile A, Mukherjee K, Saha BP, Mukherjee PK (2006) Antioxidant activity of *Nelumbo nucifera* (sacred lotus) seeds. J Ethnopharm 104: 322-327.
- Rouzmehr F, Mohit A, KoshSekeh M, Hassanzade M (2014) The effect of the additive containing artichoke extract (APC) on growth, blood cholesterol level, carcass characteristics, immune system of broiler chickens.Iranian JVet Clin Pathol 8 (1): 357-366.
- Sato Y, Itagaki S, Kurokawa T, Ogura J, Kobayashi M, Hirano T, Iseki K

(2011) In vitro, in vivo antioxidant properties of chlorogenic acid, caffeic acid. Int JPharm 403: 136-138.

- Sekhon BS Jairath S (2010) Prebiotics probiotics, synbiotics: an overview. JPharm Edu Res 1: 13.
- Shargh MS, Dastar B, Zerehdaran B, Khomeiri M, Moradi A (2012) Effects of using plant extracts, a probiotic on performance, intestinal morphology, , microflora population in broilers. J ApplPoult Res 21: 201-208.
- Shokri A, Akbari Gharaei M, Varmaghani S, Taherpour K, Khatibjoo A, Soltani M (2018) Effect of *Cynara scolymus* powder on performance, ascites syndrome in broiler chickens. J Anim Prod 20: 55-67.
- Suriya R, Zulkifli I, Alimon AR (2012) The effect of dietary inclusion of herbs as growth promoter in broiler chickens. J Anim Vet Adv11:346-350.
- Tajodini M, Samadi F, Hasani S, Hashemi SR, Samadi S (2015) Influence of Artichoke (*Cynara scolymus*) Leaf Powder on Growth Performance, Carcass Traits, Blood Parameters in Broiler Chickens. Iranian J ApplAnim Sci 5: 141-146.
- Wang Y, Wu Y, Wang Y, Xu H, Mei X, Yu D, Wang Y, Li W (2017) Antioxidant properties of probiotic bacteria. Nutrients 9: 521.
- Windisch W, Schedle K, Plitzner C, Kroismayr A (2008) Use of phytogenic products as feed additives for swine, poultry. J Anim Sci 86: 140-148.
- Woolliams JA, Wiener G, Anderson PH, Mcmurray CH (1983) Variation in the activities of glutathioneperoxidase, superoxide-dismutase, in the concentration of copper in the blood in various breed crosses of sheep. Resp J Vet Sci 34: 253-256.
- Yan FF, Murugesan GR, Cheng HW (2019) Effects of probiotic supplementation on performance traits, bone mineralization, cecal microbial composition, cytokines, corticosterone in laying hens. Animal; an Int JAnimBiosci 13: 33.
- Yildiz G, Sacakli P, Gungor T (2006) The effect of dietary Jerusalem artichoke (*Helianthus tuberosus* L.) on performance, egg quality characteristics, egg cholesterol content in laying hens. Czech J Anim Sci 51: 349-354.
- Zheng W, Dawei Z, Wenjun Y, Changhai Q, Jiang Y (2019) Probiotic Cocktails Alleviate Heat-induced Intestinal Barrier Dysfunction, Oxidative Stress in Broiler Chickens. Anim Husb Feed Sci 11: 107-114.
- Ziarlarimi A, Irani M, Gharahveysi S, Rahmani Z (2011) Investigation of antibacterial effects of garlic (*Allium sativum*), mint (*Menthe spp.*) and onion (*Allium cepa*) herbal extracts on *Escherichia coli* isolated from broiler chickens. Afr J Biotechnol 10:10320-10322.