

Journal of the Hellenic Veterinary Medical Society

Vol 73, No 1 (2022)



Relationships between glucose-6-phosphate dehydrogenase, glutathione peroxidase, reduced nicotinamide adenine dinucleotide phosphate, total protein, malondialdehyde, total glutathione and vitamin C parameters in goat milk cells

Filiz Kazak, Yasemin Karafakioglu, Nuri Başpınar, Pınar Peker Akalın

doi: [10.12681/jhvms.25512](https://doi.org/10.12681/jhvms.25512)

Copyright © 2022, Filiz Kazak, Yasemin Karafakioglu, Nuri Başpınar, Pınar Peker Akalın



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

To cite this article:

Kazak, F., Karafakioglu, Y., Başpınar, N., & Peker Akalın, P. (2022). Relationships between glucose-6-phosphate dehydrogenase, glutathione peroxidase, reduced nicotinamide adenine dinucleotide phosphate, total protein, malondialdehyde, total glutathione and vitamin C parameters in goat milk cells. *Journal of the Hellenic Veterinary Medical Society*, 73(1), 3699–3706. <https://doi.org/10.12681/jhvms.25512>

Relationships between glucose-6-phosphate dehydrogenase, glutathione peroxidase, reduced nicotinamide adenine dinucleotide phosphate, total protein, malondialdehyde, total glutathione and vitamin C parameters in goat milkcells

F. Kazak¹, Y. Karafakioğlu*², N. Başpınar³, P. Coşkun¹

¹ Department of Biochemistry, Faculty of Veterinary Medicine, Mustafa Kemal University, Hatay, Turkey

*² Department of Science Education, Faculty of Education, Uşak University, Uşak, Turkey

³ Department of Biochemistry, Faculty of Veterinary Medicine, Selçuk University, Konya, Turkey

ABSTRACT: In this study, to reveal the antioxidant potential of goat milk cells, the activities of glucose-6-phosphate dehydrogenase (G6PD), glutathione peroxidase (GPx) and the levels of reduced nicotinamide adenine dinucleotide phosphate (NADPH), total glutathione (tGSH), malondialdehyde (MDA), vitamin C (Vit C) and total protein (TP) in goat milk cells were determined and correlations between these parameters were evaluated. Milk samples were collected from 19 clinically healthy goats from a private goat farm. Briefly, milk cells were collected from milk by centrifugation and then they were sonicated. Supernatant G6PD, GPx activities and NADPH, tGSH, MDA, Vit C and total protein levels were determined by spectrophotometric methods. As regards correlations: milk cell MDA levels were positively correlated with milk cell tGSH ($r=0.725$, $p<0.01$), milk cell Vit C ($r=0.622$, $P<0.01$) and milk cell NADPH ($r=0.763$, $P<0.01$) levels. There was a positive correlation between milk cell GPx activity and milk cell NADPH levels ($r=0.659$, $P<0.01$). Milk cell tGSH levels were positively correlated with milk cell Vit C ($r=0.615$, $P<0.05$) and milk cell NADPH ($r=0.846$, $P<0.01$) levels. Milk cell NADPH levels was positively correlated with milk cell Vit C levels ($r=0.791$, $P<0.01$). As a conclusion, the antioxidant potential of goat milk cells were evaluated and discusses.

Keywords: Goat milk cell, GPx, G6PD, NADPH, tGSH

Corresponding Author:

Y. Karafakioğlu, Department of Science Education, Faculty of Education, Uşak University, Uşak, Ankara-İzmir Road 8, Km 1, Eylül Campus, 64200, Turkey
E-mail address: yasemin.sunucu@usak.edu.tr

Date of initial submission: 09-12-2020
Date of revised submission: 01-04-2021
Date of acceptance: 05-05-2021

INTRODUCTION

Though goat milk meets about 2% of the world's total annual milk supply, its contribution to human nutrition and economic well-being is extremely important in many parts of the world, particularly in the Mediterranean countries and the Middle East (Park 1994a, Park 1994b, Park and Haenlein 2007). Aleppo (Damascus) (Barıtcı and Adıgüzel, 2017) and Kilis goats (Keskin and Tüney, 2015) are two breeds in the Mediterranean and Southeastern Anatolia regions. Aleppo goat is a native breed of Syria and other Near East countries. It has been improved for over 40 years through genetic selection for milk and meat production and it has been given a high priority by the Food and Agriculture Organization (Mavrogenis et al., 2006). Kilis goat possesses crucial advantages in terms of high milk yield and reproductive traits with good toleration for warm environments and drought under extensive or semiintensive management conditions in arid and hot climates (Keskin et al., 2017). There is also an increasing demand for goat milk and its products in developed countries because people demand healthy food and also people who are allergic or have intolerance to cow milk turn towards goat milk and its products (Park, 1990; Park, 1994a; Park, 1994b; Chandan et al., 1992; Tziboula-Clarke, 2003). Although the general composition of goat milk is similar to cow milk, it is differentiated from cow milk with some features; goat milk contains on average 3.8% fat, 3.5% protein, 4.1% lactose and 0.8% ash; that is, more fat, protein and ash and less lactose than cow milk (Haenlein and Caccese, 1984). As goat milk contains a higher percentage of small-diameter fat globules than cow milk, and because of the qualitative and quantitative differences in milk proteins, especially in α -1 casein, it is easier to digest goat milk. Goat milk is rich in short and medium-chain mono and polyunsaturated fatty acids (caproic, caprylic, capric) (Djordjevic et al., 2019).

Healthy goat's milk has a higher number of somatic cells ($270-2000 \times 10^3$ cells/ml, Souza et al 2012) than cow's milk does ($<200 \pm 10^3$ cells/ml, Dang et al., 2008). The content of the somatic cells also vary between goat and cow's milk. While 40% to 87% of somatic cells in goat's milk are formed by neutrophils from PMNs (Souza et al., 2012; Shah et al., 2017), the rate of neutrophils in cow milk cells varies between 5% and 20% (Sarıkaya, 2006; Souza et al., 2012) and macrophages are the predominant cell type in healthy cow milk (Osstenson 1993).

Goats are reported to be more resistant to infectious conditions such as mastitis, than other species, presumably as they have a higher number of neutro-

phils (Tian et al., 2005; Souza et al., 2012). Besides, antioxidants such as reduced nicotinamide adenine dinucleotide phosphate (NADPH), reduced glutathione (rGSH), glucose-6-dehydrogenase (G6PD), glutathione peroxidase (GPx) and vitamin C (Vit C), have important roles in a cell's defence mechanism (Ralat et al., 2006; Stincone et al., 2015; Carole et al., 2007; Ighodaro and Akinloye, 2018). In a previous study (Akalin et al., 2019), we revealed the antioxidant potential of Holstein-Friesian cow milk cells by determining the levels of NADPH, GSH, GPx and G6PD and the relation of these parameters with mastitis, for the first time. Since it is thought that the antioxidant potential of the cells will change as the somatic cell count and somatic cell component change, we thought that revealing the antioxidant potential of goat milk cells will offer an insight to understand the goat milk cell defence mechanism.

Therefore, in order to investigate the antioxidant potential of goat milk cells, in the current study, it was aimed to determine G6PD and GPx activities and NADPH, tGSH, MDA, Vit C and total protein levels in the cells of milks obtained from clinically healthy goats' udders and to investigate the relationships between the related parameters.

MATERIALS AND METHODS

Milk samples were collected from 19 healthy goats (10 Aleppo, 9 Kilis goats) aged between 1-4 and were fed in the same care and nutritional conditions in a private goat farm in April, in Hatay region ($36^{\circ} 11' 56''$ North, $36^{\circ} 9' 38''$ East). Milk was obtained while the owner was milking the goats returned from the pasture in the morning. Milk was taken from the right lobe of clinically healthy udders. During milking, the first 2-3 squeezes of milk were thrown away after the teat was wiped with 70% alcohol cotton, and milk samples were tested by California Mastitis Test (CMT) and CMT negative samples were included in the study. Collected milk samples were brought to the biochemistry laboratory in the cold chain and stored at -20°C until the treatment. Milk samples (14 ml each) were thawed and centrifuged at $600 \times g$ for 10 min at 4°C . After centrifugation, the supernatant was eliminated by removing the upper layer of fat with a cotton pad; the remaining cell pellet was washed twice with cold phosphate-buffered saline (PBS) and centrifuged at $600 \times g$ for 10 min at $+4^{\circ}\text{C}$. Finally, the supernatant was removed, and the remaining pellet was completed to 2 ml with PBS and sonicated (Bandelin Sonopuls HD 2070, Germany) (Akalin et al., 2016) for 5 repeti-

tions of 10 sec each, with a 30 sec cooling period (on ice) between each repetition. By this process, milk cells in 14 ml of milk was concentrated in 2 ml PBS. After sonication, the homogenates were centrifuged at 13,000×g for 15 min at +4°C. Milk cell supernatant was used for the further analysis.

Determination of G6PD activity

Milk cell G6PD activity was determined by using the method developed by Beutler (1971) and calculated by the spectrophotometric measurement of the absorbance difference in optical density caused by the conversion of NADP⁺ to NADPH at 340 nm (UV 2100 UV-VIS Recording Spectrophotometer Shimadzu, Japan). Results are presented as IU/g protein.

Determination of GPx activity

Determination of GPx activity in milk cells was done according to the method described by Beutler (1975). According to this method, GPx catalyzes the conversion of rGSH to oxidizedglutathione (GSSG) in the presence of hydrogen peroxide (H₂O₂). Oxidized glutathione, formed by GPx in an environment where H₂O₂ is present, is converted back to GSH with the help of glutathionereductase and NADPH. The activity was calculated by the spectrophotometric measurement of the absorbance difference in optical density caused by the conversion of NADPH to NADP⁺ at 340 nm. Results are presented as IU/mg protein.

Determination of NADPH Levels

NADPH levels in cell supernatants were determined spectrophotometrically using a commercial kit (Sigma MAK038). The reaction principle is based on the spectrophotometric (with ELISA reader, BioTek Instruments, µQuant, U.S.A) analysis of the reduction of formazan dye by NADPH at 565 nm, which is synthesized enzymatically in the pentose phosphate pathway. Results are presented as nmol/mg protein.

Determination of tGSH Levels

Total glutathione (GSSG+rGSH) levels in cell supernatants and milk serum were calculated using a commercial kit (Sigma CS0260). It is a kinetic method based on the principle of the reduction of 5,5'-dithiobis (2-nitrobenzoic) acid to trinitrobenzoate (TNB) by glutathione. When oxidized glutathione is regenerated by glutathione reductase and NADPH, TNB absorbance at 412 nm can be measured by spectrophotometry. Results are presented as nmol/mg protein.

Determination of MDA Levels

Tissue MDA levels were determined spectrophotometrically according to the method proposed by Ohkawa et al. (1979). Principle: Tissue MDA determination; it is based on spectrophotometric measurement at 532 nm of the pink complex formed by MDA with TBA, which is the secondary product of lipid peroxidation, as a result of incubation of tissue homogenate in a boiling water bath for one hour under aerobic conditions and at pH:3.5. Results are presented as nmol/mg protein.

Determination of Vit C Levels

Milk vitamin C levels were calculated according to the manual spectrophotometric method of Haag (1985). Principle: Ascorbic acid (Vitamin C) is converted to dehydro ascorbic acid with mild oxidizing agents, dehydro ascorbic acid slowly converts to diketogulonic acid in mild acid solutions. Dehydro-ascorbic acid and diketogulonic acid react with 2,4-dinitrophenylhydrazine (DNPH) to form bis 2,4-dinitrophenylhydrazone. The results are given as mg/milk cells of 1 ml milk.

Determination of Total Protein Levels

Total protein levels in milk cell supernatants were determined by the Bradford (1976) method (Coomassie Brilliant Blue G, Sigma 27815-100 G). Protein concentration was determined spectrophotometrically by determining the absorbance at 595 nm. Bovine serum albumin (Merck 112018) was used as a standard. The results are given as mg/milk cells of 1 ml milk.

Statistics

The values obtained were evaluated by SPSS 22.0 program and descriptives (Mean ± Standard Error) were evaluated. Pearson correlation was performed for correlation analysis and P<0.05 indicated as statistically significant.

RESULTS

As shown in Table 1, goat milk cell G6PD and GPx activities were determined to be 1.81±0.48IU/g protein and 0.38±0.02 IU/mg protein, respectively. NADPH levels were 1.18±0.23 nmol/mg protein, tGSH levels were 101.73±25.50 nmol/mg protein, MDA levels were 1.43±0.35 nmol/mg protein, Vit C levels were 0.45±0.09 µg/milk cell of 1 ml milk and TP levels were 0.09±0.001 mg/milk cell of 1 ml milk.

Table 1. Some Biochemical Parameters in Goat Milk Cells

	Mean	SE	n
G6PD (IU/g protein)	1.81	0.48	19
GPx (IU/mg protein)	0.38	0.02	19
NADPH (nmol/mg protein)	1.18	0.23	17
tGSH (nmol/mg protein)	101.73	25.50	14
MDA (nmol/mg protein)	1.43	0.35	19
Vit C (µg/ milk cell of 1 ml milk)	0.45	0.09	19
TP (mg/ milk cell of 1 ml milk)	0.09	0.001	19

G6PD:Glucose-6-phosphate dehydrogenase, GPx: Glutathione peroxidase, NADPH:Reduced nicotinamideadenin dinucleotide, tGSH: Total Glutathione, VitC: Vitamin C, TP: Total Protein

Table 2. Correlations between the parameters (Spearman's correlation test)

	TP mg/ml	GPx U/ml	tGSH nmol/ml	Vit C µg/ml	NADPH pmol/ml	G6PD U/ml
MDA nmol/ml	0,454	0,408	0,725**	0,622**	0,763**	0,226
TP mg/ml		0,392	-0,150	0,063	0,377	-0,228
GPx U/ml			0,412	0,426	0,659**	-0,344
tGSH nmol/ml				0,615*	0,846**	0,266
Vit C µg/ml					0,791**	0,065
NADPH pmol/ml						-0,032

G6PD: Glucose-6-phosphate dehydrogenase, GPx: Glutathione peroxidase, NADPH: Reduced nicotinamideadenin dinucleotide, tGSH: Total Glutathione, VitC: Vitamin C, TP: Total Protein *P<0.05,**P<0.01.

As shown in Table 2, milk cell MDA levels were positively correlated with milk cell tGSH ($r=0,725$, $p<0,01$), milk cell Vit C ($r=0.622$, $P<0.01$) and milk cell NADPH ($r=0.763$, $P<0.01$) levels. There was a positive correlation between milk cell GPx activity and milk cell NADPH levels ($r=0.659$, $P<0.01$). Milk cell tGSH levels were positively correlated with milk cell Vit C ($r=0.615$, $P<0.05$) and milk cell NADPH ($r=0.846$, $P<0.01$) levels. Milk cell NADPH levels were positively correlated with milk cell Vit C levels ($r=0.791$, $P<0.01$).

DISCUSSION

Milk cells are composed of epithelial cells inter-fused to the milk from the breast tissue and leukocytes (neutrophils, lymphocytes and macrophages) that pass from the blood to the mammary gland and then into the milk. The content and the number of somatic cells in healthy goat milk differ from that in cow milk (Sarıkaya, 2006; Dang et al., 2008; Souza et al., 2012; Shah et al., 2017). Studies determining the protein levels of milk cells are quite limited. In the study conducted on cow milk somatic cells (Akalin et al., 2019), the total protein levels were determined as 0.374mg/1 ml cell supernatant in the cell pellet obtained by taking 50 ml of cow milk and concen-

trating it into 2 ml PBS. When the total protein levels in the present study were calculated with dilution coefficients (14 ml milk concentrated with 2 ml PBS: 0.09 mg/1 ml cell supernatant), they were found to be slightly lower than the levels in the cow milk (approximately 0.321 mg/1 ml cell supernatant). No study investigating total protein levels of goat milk somatic cells has been found. The major component of goat milk somatic cell, unlike cow's milk is PMN. While 40% to 87% of somatic cells in goat milk (Souza et al., 2012; Shah et al., 2017) are formed by neutrophils from PMNs, the rate of neutrophils in cow milk cells varies between 5% and 20% (Sarıkaya, 2006; Souza et al., 2012). Moreover, 15% to 41% of somatic cells in healthy udder consist of macrophages whereas 9% to 20% of them consist of lymphocytes and 1% to 6% of them consist of epithelial cells (Paape et al., 2001). Determination of low total protein levels in goat milk cell compared to cow milk may be related to the difference in milk cell components.

The main goal of the pentose phosphate pathway is to produce NADPH, which has reducing power, and ribose-5-phosphate, which is the building block of DNA and RNA. NADPH is produced in the pentose phosphate pathway by glucose-6-phosphoglucanate dehydrogenase (6PGD) and G6PD, from

NADP⁺ (oxidized NADPH) (Reuter et al., 1990; Stincone et al., 2015). While NADPH levels in cow milk cell were 4.24 nmol/mg protein (Akalin et al., 2019) and 5.99 nmol/mg protein (Akin et al. 2019), in the current study, the levels in goat milk (1.18 nmol / mg protein) were found to be lower than in the cow milk. As reported by Akin et al. (2019) and Akalin et al. (2019), a significant positive correlation was determined between cow milk cell NADPH levels and G6PD activity. However, in the current study no correlation was observed between these parameters in goat's milk. It can be speculated that NADPH is not controlled only by G6PD but also by 6PGD in the pentose phosphate pathway in the goat milk cell. It is suggested to evaluate 6PDG levels in goat milk cells to reveal the NADPH synthesis.

While positive correlations of milk leukocyte NADP⁺reductase activity (Ritter et al., 1977) and milk somatic cell G6PD activity (Akalin et al., 2019) with milk cell total protein levels in cows were reported, no correlation was found between total protein levels and G6PD activity in the current study. Glutathione protects cells against free radicals, reactive oxygen species, endogenous and exogenous toxic compounds. During the reduction, oxidized glutathione (GSSG) is formed and this molecule is reduced by GR enzyme which uses NADPH. While GPx provides detoxification of H₂O₂ in the cells, it allows conversion of rGSH to GSSG (Ighodaro and Akinloye, 2018). In the current study, tGSH levels (101.73±25.5 nmol/mg protein) were found to be slightly lower than the levels reported for cow milk cells (122.88±13.08 and 142.16±37.06 nmol/mg protein) (Akalin et al., 2019; Akin et al., 2019, respectively). Also, goat milk cell NADPH levels were positively correlated with milk cell tGSH levels and GPx activity. While Akin et al. (2019) reported no significant correlation of NADPH with rGSH and GPx activity in cowmilk cells, Akalin et al. (2019) reported only a weak positive correlation between NADPH levels and GPx activity, and they thought that GSH levels and GPx activity in cow milk cells were also related to other mechanisms, and these parameters were not directly dependent on NADPH and G6PD. In goat milk cell, it can be concluded that NADPH is related to tGSH and GPx, unlike cow milk cell. While in their study, Akalin et al. (2019) determined GPx activity as 4.44±0.36 IU/mg protein in cow milk cells, GPx activity was measured as 0.38±0.02 IU/mg protein in the current study conducted with goat milk cells. Just as in the cow milk cells (Akalin et al., 2019; Akin et al., 2019), no correlation was re-

ported between total protein and GPx activity in the goat milk cells.

Vitamin C (ascorbic acid, ascorbate) is a water-soluble vitamin that causes the reduction of compounds such as molecular oxygen, nitrate, cytochrome a and c and is capable of reacting with free radicals in the aqueous environment. It reacts with superoxide and hydroxyl radicals and forms the first antioxidant defence against oxidant agents (Carole et al., 2007). In the current study, a significant positive correlation was determined between Vit C levels and tGSH levels. To our knowledge, no study was reported regarding correlation of milk cell Vit C and GSH levels. In a study conducted on lymphocytes, which were isolated from human blood, a significant and high level of positive correlation was found between tGSH and ascorbate (Lenton et al., 2000). Reduced glutathione is used to reduce oxidized Vit C (dehydroascorbic acid) to reduced form (ascorbic acid, ascorbate) (Winkler et al., 1994; May et al., 1997). The direct correlation between Vit C and GSH suggests that these two antioxidant agents may work synergistically with each other in the cell (Lenton et al., 2000). In the current study, a significant and high level of correlation was determined between Vit C and NADPH levels. There was no literature on the correlation between Vit C and NADPH levels in milk cells. On the other hand, *ex vivo* in human neutrophils, it has been reported that the activity of NADPH-oxidase enzyme, which uses NADPH as a substrate and enables the formation of superoxide radicals from molecular oxygen (Lassegue et al., 2012), is inhibited by the lipophilic derivative of ascorbic acid, thereby reducing the formation of superoxide radicals (Schmid et al., 1994). In the current study, the existence of a positive correlation between Vit C and NADPH in the absence of a correlation between Vit C and G6PD activity suggests that this might be due to NADPH-oxidase. It has been reported that thioredoxin reductase enzyme, which enables the conversion of dehydroascorbic acid to ascorbat in rat liver, uses NADPH as a reducing agent (May et al., 1997). Also, the positive correlation between Vit C and NADPH may be different in healthy and inflammatory environment. Malondialdehyde is the final peroxidation product of fatty acids with multiple double bonds found in cell and organelle membranes. Increasing peroxidation of lipids by free radicals in membranes causes an increase in MDA level. Malondialdehyde and other lipid peroxides can react with DNA or proteins and disrupt their structure (Gawel et al., 2004). The level MDA is served as a re-

liable biomarker of lipid peroxidation (LPO) and usually served as a marker of LPO (Acaroz et al., 2018). No study on MDA levels in milk cells was found in the literature review. Goat milk cell MDA levels were determined as 1.43 ± 0.35 nmol/mg protein. In addition, positive correlations of goat milk cell MDA levels with reducing agents such as tGSH, Vit C and NADPH were revealed. As MDA levels increase in goat milk cells, the increase seen in antioxidant molecules may indicate that these molecules play a role in the defence mechanism against oxidation and against the formation of MDA.

CONCLUSION

In conclusion, positive correlations of milk cell MDA levels with milk cell tGSH, Vit C and NADPH levels, positive correlations of milk cell NADPH lev-

els with milk cell tGSH, GPx and Vit C levels, positive correlation between milk cell Vit C and milk cell tGSH levels suggest that goat milk cells have an efficient antioxidant mechanism in healthy conditions which is some different from cow milk cells.

ACKNOWLEDGEMENTS

Pınar Peker Akalın and Filiz Kazak contributed to the study by conducting the study, experimental design, project design and data analysis. Nuri Başpınar, and Yasemin Karafakıoğlu have contributed to experimental animal practices, laboratory studies and data analysis. No potential conflict of interest was reported by the authors.

CONFLICT OF INTEREST

None declared by the authors.

REFERENCES

- Acaröz U, İnce S, Arslan-Acaröz D, Gürler Z, Küçük Kurt I, Demirel HH, Arslan HO, Varol N, Zhu K (2018) The ameliorative effects of boron against acrylamide-induced oxidative stress, inflammatory response, and metabolic changes in rats. *Food and Chemical Toxicology*. 118:745-752. doi: 10.1016/j.fct.2018.06.029.
- Akalın PP, Başpınar N, Çoyan K, Bucak MN, Güngör Ş, Öztürk C (2016) Effects of ultrasonication on damaged spermatozoa and mitochondrial activity rate. *Turkish Journal of Veterinary and Animal Sciences*. 40:195-199. doi:10.3906/vet-1507-18.
- Akalın PP, Ergün Y, Başpınar N, Doğruer G, Küçükgül A, Cantekin Z, İşgör M, Sarıbay M, Koldaş E., Baştan A, Salar S, Pehlivanlar S (2019) Glucose 6 phosphate dehydrogenase glutathione peroxidase total glutathione and reduced nicotinamide dinucleotide phosphate in milk cells of subclinical mastitic cows. *Polish Journal of Veterinary Sciences* 22 (2) :271-278. doi:10.24425/pjvs.2019.129216.
- Akın Z, Akalın PP, Kazak F, Ergün Y, Başpınar, N (2019) Glukoz-6-fosfat dehidrogenaz, glutatyon peroksidaz, nikotinamid adenin dinükleotid fosfat ve glutatyon ile süt kalite parametreleri arasındaki ilişkiler. *Atatürk University Journal of Veterinary Sciences* 14 (2) :193-200. doi:10.17094/ataunivbd.555594.
- Barıtcı İ, Adıgüzel C (2017) Aleppo (Damascus) Goat Breeding. *Dicle University Journal of the Institute of Natural and Applied Science*. 6:39-42.
- Beutler E (1971) Red cell metabolism, In: *A Manual of Biochemical Methods*, Academic, Press, London. pp 68-70.
- Beutler E (1975) Red cell metabolism, In *A Manual of Biochemical Methods*, 2nd ed., Grune and Stratton Press, New York.
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*. 72:248-254. doi:10.1006/abio.1976.9999.
- Carole L, Linster CL, Van Schaftingen E (2007) Vitamin C: Biosynthesis, recycling and degradation in mammals. *FEBS Journal*. 274 (1) :1-22. doi:10.1111/j.1742-4658.2006.05607.x
- Chandan RC, Attaie R, Shahani KM (1992) Nutritional aspects of goat milk and its products, In: *Proceedings of the Vth International Conference on Goats*, New Delhi, India. vol.II, Part II, pp. 399.
- Dang AK, Kapila S, Singh C, J.P. Sehgal JP (2008). Milk differential cell counts and compositional changes in cows during different physiological stages. *Milchwissenschaft*. 63 (3) :239-242.
- Djordjevic J, Ledina T, Baltic MZ, Trbovic D, Babic M, Bulajic S (2019) Fatty acid profile of milk. *IOP Conf. Series: Earth and Environmental Science* 333:012057. doi:10.1088/1755-1315/333/1/012057.
- Gawel S, Wardas M, Niedworok E, Wardas P (2004) Malondialdehyde (MDA) as a lipid peroxidation marker. *Wiadomosci Lekarskie*. 57 (9-10) :453-5.
- Haag W (1985) Zur methodik und praktischen Bedeutung der Vitamin C - Bestimmung bei im Rind in Vergangenheit und Gegenwart. *Inaugural Dissertation*. Justus Liebig Universität, Giessen.
- Haenlein GFW, Caccete R (1984) Goat Milk Versus Cow Milk, In: Haenlein GFW, Ace DL. (editors) *Extension Goat Handbook*, USDA Publications, Washington, DC, E-1, p.1.
- Ighodaro OM, Akinloye OA (2018) First line defence antioxidants-superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX) : Their fundamental role in the entire antioxidant defence grid. *Alexandria Journal of Medicine*. 54:287-293. doi: 10.1016/j.ajme.2017.09.001.
- Keskin M, Tüney D (2015) Relationship between body condition score and reproductive characteristics in Kilis goat. *Journal of Agricultural Faculty of Mustafa Kemal University*. 20 (2) :60-65.
- Keskin M, Gül S, Biçer O, Daşkıran İ (2017) Some reproductive, lactation, and kid growth characteristics of Kilis goats under semiintensive conditions. *Turkish Journal of Veterinary and Animal Sciences*. 41:248-254.
- Lassegue B, Martin AS, Griendling KK (2012) Biochemistry, physiology and pathophysiology of NADPH oxidases in the cardiovascular system. *Circulation Research*. 110: (10) 1364-1390. doi:10.1161/CIRCRESAHA.111.243972
- Lenton KJ, Theriault H, Cantin AM, Fülöp T, Payette H, Fülöp T, Payette H, Wagner JR (2000) Direct correlation of glutathione and ascorbate and their dependence on age and season in human lymphocytes. *The American Journal of Clinical Nutrition*. 71:1194-200. doi:10.1093/ajcn/71.5.1194.
- Mavrogenis A, Antoniadis N, Hooper R (2006) The Damascus (Shami) goat of Cyprus. *Animal Genetic Resources Information*. (38) : 57-65. doi:10.1017/S1014233900002054.
- May JM, Mendiratta S, Hill KE, Burk RF (1997) Reduction of dehydroascorbate to ascorbate by the seleno enzyme thioredoxin reductase. *The Journal of Biological Chemistry*. 272: (36) 22607-22610. doi:10.1074/jbc.272.36.22607.
- Ohkawa H, Ohishi N, Yagi K (1979) Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analytical Biochemistry*. 95:351-358. doi:10.1016/0003-2697(79)90738-3.
- Osstenson K (1993) Variations during lactation in total and differential leukocyte counts, N-acetyl-d-glucosaminidase, antitrypsin and serum albumin in foremilk and residual milk from non-infected quarters in the bovine. *Acta Veterinaria Scandinavica*. 34:83-93.
- Paape MJ, Poutrel B, Contreras A, Marco JC, Capuco AV (2001) Milk somatic cells and lactation in small ruminants. *Journal of Dairy Science*. 84:237-244.
- Park YW (1990) Nutrient profiles of commercial goat milk cheeses manufactured in the United States. *Journal of Dairy Science*. 73:3059.
- Park YW (1994a) Nutrient and mineral composition of commercial US goat milk yogurts. *Small Ruminant Research*. 13:63-70.
- Park YW (1994b) Hypo-allergenic and therapeutic significance of goat milk. *Small Ruminant Research*. 14: (2) 151-159. doi:10.1016/0921-4488(94)90105-8.
- Park YW, Haenlein GFW (2007) Goat milk, its products and nutrition, In: Hui YH (editor) *Handbook of Food Products Manufacturing*, John Wiley & Sons, Inc., New York, p:447-486.
- Ralat LA, Manevich Y, Fisher AB, Colman RF (2006) Direct evidence for the formation of a complex between l-cysteine peroxiredoxin and glutathione S-transferase pi with activity changes in both enzymes. *Biochemistry*. 45:360-372. doi:10.1021/bi0520737.
- Reuter R, Naumann M, Metz P, Kopperschläger G (1990) Purification and characterization of glucose-6-phosphate dehydrogenase from *Pseudomonas W6*. *Biomedica Biochimica Acta*. 49: (7) 539-546.
- Ritter C, Conti A, Morse GE (1977) Hexose shunt dehydrogenase activity in leukocytes isolated from bovine milk. *Journal of Dairy Science*. 60:1987-1990. doi:10.3168/jds.S0022-0302(77)84133-7.
- Sarıkaya H (2006) Somatic cell populations in milk: Importance in mammary gland physiology and behaviour during technological processing, *Technical University of Munich, Science Center for Nutrition, Land Use and Environment, Department of Physiology, PhD Thesis*, pp:27-28.
- Schmid E, Figala V, Ullrich V (1994) Inhibition of NADPH-oxidase activity in human polymorphonuclear neutrophils by lipophilic ascorbic acid derivatives. *Molecular Pharmacology*. 45: (5) 815-825.
- Shah A, Darzi MM, Kamil SA, Mir MS, Maqbool R, Ali R, Kashani B, Wani H, Bashir A, Aijaz A, Dar AA, Qureshi S (2017) Somatic cell alteration in healthy and mastitic milk of sheep and goats. *Journal of*

- Entomology and Zoology Studies. 5 (6) : 27-33.
- Souza FN, Blagitz MG, Penna CF, Della Libera AM, Heinemann MB, Cerqueira MM (2012) Somatic cell count in small ruminants: Friend or foe?. *Small Ruminant Research*. 107:65-75.
- Stincone A, Prigione A, Cramer T, Wamelink MMC, Campbell K, Cheung E, Olin-Sandoval V, Grüning NM, Krüger A, Alam MT, Keller MA, Breitenbach M, Brindle KM, Rabinowitz JD, Ralser M (2015) The return of metabolism: biochemistry and physiology of the pentose phosphate pathway. *Biological Reviews of the Cambridge Philosophical Society*. 90 (3) :927-963. doi: 10.1111%2Fbrv.12140.
- Tian SZ, Chang CJ, Chiang CC, Peh HC, Huang MC, Lee JW, Zhao X (2005) Comparison of morphology, viability and function between blood and milk neutrophils from peak lactating goats. *Canadian Journal of Veterinary Research*. 69:39-45.
- Tziboula-Clarke A (2003) Goat milk, In: Roguiski H, Fuquay J, Fox P. (editors). *Encyclopedia of Dairy Sciences*, Academic Press. pp.1270-1279.
- Winkler BS, Orselli SM, Rex TS (1994) The redox couple between glutathione and ascorbic acid: a chemical and physiological perspective. *Free Radical Biology and Medicine*. 17:333-49. doi:10.1016/0891-5849 (94) 90019-1.