

Journal of the Hellenic Veterinary Medical Society

Vol 72, No 2 (2021)



Current research and application of stem cells in the dog and cat

A.A. ANATOLITOU, K.I. SIDERI, N.N. PRASSINOS

doi: [10.12681/jhvms.27512](https://doi.org/10.12681/jhvms.27512)

Copyright © 2021, A.A. ANATOLITOU, K.I. SIDERI, N.N. PRASSINOS



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

To cite this article:

ANATOLITOU, A., SIDERI, K., & PRASSINOS, N. (2021). Current research and application of stem cells in the dog and cat. *Journal of the Hellenic Veterinary Medical Society*, 72(2), 2791–2802. <https://doi.org/10.12681/jhvms.27512>

Current research and application of stem cells in the dog and cat

A.A. Anatolitou¹, K.I. Sideri², N.N. Prassinos³

¹ *Kastella Veterinary Centre, Small Animal Clinic, Piraeus, Greece*

² *Faculty of Veterinary Medicine, School of Health Sciences, University of Thessaly, Karditsa, Greece*

³ *School of Veterinary Medicine, Faculty of Health Sciences, Aristotle University, Thessaloniki, Greece*

SUMMARY: Stem cells (SCs) are multipotent cells with differentiation and proliferation capacities in many cell lineages. The majority of literature applications are about bone marrow derived stem cells (BMSCs) and adipose derived stem cells (ADSCs). Most clinical trials have been done for the treatment of musculoskeletal and neurological problems in canine patients. Hematopoietic SCs (HSCs), synovium (SDSCs) and cartilage (CSPCs) - derived SCs, umbilical cord blood-derived SCs (UCSCs), muscle, dental, cardiac and hepatic SCs have been used with promising results. Despite the overall progress crucial questions about SCs remain unanswered. It is still unclear if the regeneration mechanism of SCs owed to their differentiation into specific progenitor cells or due to their immunomodulatory and anti-inflammatory secretions. Also, there are questions about the best origin of stem cells, whether they should be delivered in situ or systemically, if they should be embedded into scaffolds or not and which is the suitable transplantation stem cells number. Many of the published studies have limitation regarding to sample size, blind randomization, control groups and homogeneity of population. In addition, the long-term efficacy and safety of MSCs need further evaluation. This review is an update in usage of SCs, mainly focused on BMSCs and ADSCs in small animals and its purpose is to present the late developments in this field. Also, the advantages, the disadvantages and the limitations of the current literature review are discussed.

Keywords: ADSCs, cats, dogs, MSCs, stem cells, transplantation.

Corresponding Author:
Anthi Anatolitou, Myrtidiotissis 8, Kastella, 18533 Piraeus, Greece
E-mail address: anthianatol@yahoo.gr

Date of initial submission: 10-01-2020
Date of revised submission: 10-01-2020
Date of acceptance: 20-12-2020

INTRODUCTION

Stem cells (SCs) are multipotent cells with differentiation and proliferation capacities in many cell lineages. They produce factors with immunomodulatory and angiogenesis properties (Takemitsu et al., 2012). They divided into embryonic SCs (ESCs) and adult mesenchymal SCs (MSCs). The latter can be harvested by mesodermal, endodermal or ectodermal tissues (de Bakker et al., 2013). Depending on their origin they have advantages and limitations. For example, bone marrow - derived mesenchymal SCs (BMSCs) have the higher differentiation capacity in musculoskeletal cells, but their culture is time consuming (Fortier and Travis, 2011). Their osteogenic potential is better than adipose tissue - derived SCs (ADSCs) (Park et al., 2012), while the latter exhibit higher proliferation rate (Webb et al., 2011). Also, allogeneic versus autologous transplantation is still debatable (Zhang et al., 2015). In vitro studies proved that allogeneic transplantation is safe (Ryan et al., 2005), due to a protective mechanism against T and B cells that forms a safe micro-environment (Ryan et al., 2005). Moreover, xenogeneic SCs transplantation was also presented. Human SCs promoted healing in dogs with bone defects (Cruz et al., 2015, Zang et al., 2016), dermatological problems (Ferrer et al., 2015) and neurological diseases (Lee et al., 2009, Chung et al., 2013).

SCs usage in tissue engineering is a progressive technique that continues to expand increasingly in the veterinary world. Many scientific papers have been published over the last 10 years (Arinze et al., 2003, Yan et al. 2007, Cui et al. 2007, Wang et al. 2014, Nantavisai et al., 2019, Voga et al. 2020).

ADVANCES IN STEM CELLS

Adipose tissue - derived stem cells (ADSCs)

Dogs

Musculoskeletal applications

In 2007, Cui et al. (2007) showed that autologous ADSCs enhanced bone regeneration in critical sized bone defects in canine models. Same conclusions retrieved from trials on mandibular (Haghighat et al., 2011) and long bone defects (Bigham-Sadegh et al., 2012). In 2013, Lui et al. repeated the same model with allogeneic ADSCs and presented similar results without adverse reactions (Liu et al., 2013).

The application of autologous ADSCs in coxofemoral (Black et al., 2007) and humeroradial joints with osteoarthritis (Black et al., 2008, Guercio et al., 2012) resulted in clinical improvement. Another study reinforced these results with kinetic force assessments (Vilar et al., 2014). Equally effective was the application of allogeneic ADSCs (Harman et al., 2016). Moreover, the combination of autologous ADSCs with platelet-rich plasma (PRP) (Vilar et al., 2013, Cuervo et al., 2016, Yun et al., 2016) reduced pain in cases of osteoarthritis, but the studies lacked of control groups.

In 2014, an innovative administration of allogeneic ADSCs for hip dysplasia, in acupuncture points, showed clinical improvement (Marx et al., 2014). Also, promising was the administration of allogeneic ADSCs in elbow dysplasia (Kriston-Pal et al., 2017). Nevertheless, these studies did not present arthroscopic findings. Recently, another research demonstrated that the local injection of ADSCs was superior compared to intravenous for degenerative arthritis (Shah et al., 2018). Unfortunately, the outcomes were based only on clinical evaluation.

ADSCs combined with PRP accelerated the reconstruction of partial ruptured anterior cruciate ligament (Canapp et al., 2016a). In contrast, the use of ADSCs during tibial tuberosity advancement procedure did not diminish the healing time (Dos Santos et al., 2018). The results about ADSCs advantages were controversial, but both studies had heterogeneous populations and different rehabilitation protocols. Furthermore, preliminary results proposed the combination of ADSCs with PRP for refractory supraspinatus tendinopathy (Canapp et al. 2016b), semitendinosus myopathy (Gibson et al., 2017) and tracheal cartilage defects (Hashemibeni et al., 2012). Lastly, the combination of ADSCs with growth factor showed promising results in a canine tendon injury model (Sheh et al., 2018).

Neurological applications

ADSCs were used for spinal cord injury (SCI) (Roszek et al., 2016). Canine models investigated ADSCs capacities in situ (Ryu et al., 2009). Interestingly, ADSCs promoted functional improvement after SCI (Park et al., 2012) and accelerated disk regeneration after disk degeneration disease (DDD) (Ganey et al., 2009). However, the results were based on experimental conditions. In clinical field, two studies described the benefits of ADSCs transplantation in dogs with thoracolumbar DDD (Kim et al., 2016)

and peripheral (facial) nerve trauma (Ghoreishian et al., 2013). Both confirmed clinical improvement, but lacked of histopathological confirmation.

Dentistry and ophthalmology applications

ADSCs were used successfully in periodontal trauma (Tobita et al., 2013), atonic oral ulcers (Alamoudi et al., 2014) and as an osseointegration factor for dental implants (Bressan et al., 2015). To the authors' knowledge, no clinical studies are currently available. After the proof of ADSCs safety inperiocular (Wood et al., 2012) and intra-lacrimal (Park et al., 2013) injections, interest was concentrated in ADSCs usage against keratoconjunctivitis sicca (KCS) (Villatoro et al., 2015). Peri- and intra-lacrimal ADSCs transplantation led to KCS remission (Bittencourt et al., 2016). However, possible adverse reactions were not studied.

Other applications

The in vitro differentiation of ADSCs into cells similar to hepatocytes (Banas et al. 2007) triggered further studies. In one preclinical study, ADSCs decreased hepatic parameters after liver trauma (Teshima et al., 2017), and increased survival time in one dog with hepatocutaneous syndrome (Nam et al., 2017). Also, ADSCs were employed for dogs with inflammatory bowel disease (Perez-Merino et al., 2015). The conclusions are suggestive, as these studies lacked of long term efficacy and safety data. Despite the fact that ADSCs improve the cardiac function after myocardial infarction, clinical trials failed to prove their benefits in dilated cardiomyopathy (Pogue et al., 2013). A current comparison of ADSCs and BMSCs intrarenal injection in a canine renal injury model revealed better level of protection for ADCSs (Osman et al. 2020). Lastly, ADSCs application was beneficial for one dog with an atonic ulcer (Han et al., 2015) and another with pemphigus (Zubin et al., 2014), but was not helpful for allergies (Hall et al., 2010).

Cats

One pilot study showed improvement in renal function in cats with chronic kidney failure after intrarenal administration of autologous ADSCs (Quimby et al., 2011). However, the intravenous administration of allogeneic ADSCs was associated with adverse effects (Quimby et al., 2013). Further research in order to attempt to minimize these reactions failed (Quimby et al., 2016). The intravenous injection of ADSCs in

cats with inflammatory bowel disease led to (Webb and Webb, 2015) signs remission. Also, the intraperitoneal injection of ADSCs was shown to be safe and possibly effective for various feline diseases (Parys et al., 2016). ADSCs were employed successfully in autoimmune diseases like gingivostomatitis (Arzi et al., 2016), asthma (Trzil et al., 2015) and eosinophilic keratitis (Villatoro et al., 2018). Nevertheless, larger sample sizes are needed to obtain more reliable results.

Bone marrow-derived mesenchymal stem cells (BMSCs)

Dogs

Musculoskeletal applications

The implantation of autologous BMSCs embedded into various scaffolds, encouraged bone formation in alveolar (Kim et al., 2009), mandible (Hu et al., 2014), orbital (Yang et al., 2014) and long bone defects (Ozdal-Kurt et al., 2014). Also, the combination of cell sheet and scaffolds resulted in healing in mandible injuries (Shan and Hu, 2017). All the macroscopic results were co-evaluated with imaging and histological findings. Despite the fact that these were control studies and their results were significant different between groups, their sample sizes were limited. Few experimental works studied the importance of scaffolds and the migration of systemically administered BMSCs. The intra-osseous injection of allogeneic BMSCs into femurs promoted bone regeneration in mandible defects (Liu et al., 2014). Similar were the results of intra-arterial infusion of autologous BMSCs in dogs with femoral head necrosis (Jin et al., 2016). Both studies concluded that the technique was not only promising for bone density problems, but it was also safe. Moreover, genetic modification of BMSCs enhanced specific capacities. The induced expression of vascular endothelial growth factor 165 (VEGF₁₆₅) (Hang et al., 2012) or bone morphogenetic protein -2 (BMP-2) (Peng and Wang, 2017) stimulated the quantity and quality of new formed bone in femoral head osteonecrosis. These suggestions resulted from control-based studies with big sample sizes. Additionally, smaller studies concluded that microRNA-31 and dentin matrix protein-1 (DMP-1) manipulated BMSCs enhanced bone formation with superior features compared to simple BMSCs (Deng et al., 2014, Liu et al., 2016). BMSCs have also been

used for cartilage repair (Bornes et al., 2014). In 2005, Wayne et al. proposed that the injection of BMSCs in canine joints with articular defects promoted superior quality of new formed cartilage. Concurrently, the outcomes from intralesional injection of bone marrow into traumatized meniscus, implied healing properties of BMSCs (Abdel-Hamid et al., 2005). Equivalent findings concluded from the intravenous and intrarticular injection of BMSCs, in dogs with partial cruciate ligament rupture, despite the limited sample and volume of synovial fluid available for tests (Muir et al., 2016). Following studies focused on PRP combination with BMSCs, because it improved their proliferation rate (Chen et al., 2014) and chondrogenetic capacity (Kazemi et al., 2017). A large retrospective study suggested that their injection in dogs with supraspinatus tendinopathy resulted in clinical and ultrasonography improvement (McDougall et al., 2018). However, there are no immunohistochemical assessments of these findings. Lastly, BMSCs were used successfully in tendon repair not only in vitro (Ozasa et al., 2014) but also in vivo (Case et al. 2013) and esophagus reconstruction (Tan et al., 2013). Nevertheless, the data were preliminary.

Neurological applications

BMSCs were also helpful for neurological disorders (Kamishina et al., 2006). In 2008, Hiyama et al. (2008) investigated the intralesional autologous BMSCs injection in a canine model of DDD. The histological and imaging analysis showed alteration of the micro-environment and inflammation's inhibition (Hiyama et al., 2008). However, this was an experimental model. Therefore, scientists investigated the safety of the procedure in clinical cases of DDD. Their results were controversial. One study presented improvement of nociception and proprioception after autologous BMSCs transplantation (Besalti et al., 2015). Another study concluded that the procedure may be safe, but not useful (Steffen et al., 2017). Both reports failed to make a statement due to limited samples and heterogenous populations. Furthermore, clinical results highlighted BMSCs efficacy after decompression surgery, in SCI (Nishida et al., 2012, Besaltiet al., 2016) even in chronic cases (Nishida et al., 2011). The improvement was similar after autologous or allogeneic BMSCs implantation (Jung et al., 2009, Sarmiento et al., 2014). Concurrently investigation about the most suitable time for BMSCs application (Penha et al., 2014) and the number of implanted cells (Serigano et al., 2010) was made. These

parameters affected the outcomes, but no protocols were proposed. Recently, Wu et al. (2018) created a canine model with complete transection of spinal cord and presented that BMSCs could be beneficial even for devastating cases. Moreover, autologous BMSCs combined with scaffolds were used successfully in sciatic (Ding et al., 2010) and ulnar nerve defects (Kaizawa et al., 2016). Degeneration was accelerated functionally and histologically. Lastly, BMSCs were proposed for cases of autoimmune meningoencephalomyelitis. Although the diagnosis was made by clinical examination, there was noted signs remission (Zeira et al., 2015).

Dentistry and ophthalmology applications

One report studied autologous BMSCs implantation into alveolar clefts combined with scaffolds, PRP and bone grafts (Yuanzhneng et al., 2015). The combination of BMSCs with PRP showed the best results. The findings were equivalent to studies that support PRP's proliferation properties in BMSCs (Chen et al., 2014, Kazemi et al., 2017, McDougall et al. 2018). Another study proposed autologous, alveolar BMSCs as a different origin of BMSCs, due to their access and osteogenic potentials (Wang et al., 2018). Nevertheless, the conclusions were preliminary. Additionally, large oral (Aly et al., 2014), vocal fold (Kanemaru et al., 2003) and laryngeal ulcers (Irvani et al., 2017) were favoured by the intra-lesional administration of autologous BMSCs. These results were supported by histological findings, but more research is needed. In veterinary ophthalmology experimental models of corneal ulcer (Tognoli et al., 2008) and retinal degeneration (Tracy et al., 2016), showed that autologous BMSCs were safe and anti-inflammatory.

Other applications

Canine models of cirrhosis (Matsuda et al., 2017), renal trauma (Lim et al., 2016) and chronic enteritis (Xu et al., 2016) were used to investigate other effects. Autologous BMSCs improved function in liver failure, inhibited fibrosis in renal injuries and diminished clinical signs in post radiation enteritis (Xu et al., 2016). Also, the usage of allogeneic BMSCs promoted healing in cutaneous wounds without adverse effects (Kim et al., 2013). Furthermore, Memon et al. (2005) presented that the intra-lesional injection of BMSCs accelerated regeneration in ischemic myocardium. Also, their intracoronary injection reduced infarctions' areas, confirmed by histology and immunohistochemistry analysis (Hao et al., 2015). In the same

model, the adjunct of basic fibroblast growth factor augmented the engraftment and the differentiation of BMSCs (Wang et al., 2015). Undoubtedly, the previous studies had preliminary value and more research should be done in dosage, time points and administration routes.

Cats

Feline BMSCs morphology and isolation are similar to human and rodents, (Martin et al., 2002, Zhang et al., 2011, Munoz et al., 2012) with higher neurogenic properties (Zhang et al., 2011). Also, c-kit⁺ feline BMSCs differentiated into cardiac myocytes with cardiac action potentials (Kubo et al., 2009). However, feline BMSCs had culture limitations referred to the donor health status (Quimby et al., 2011). In vivo applications are few. A randomized clinical trial failed to demonstrate significant effect of BMSCs in acute kidney injury. Also, it left unanswered questions about the ideal routes, doses and frequency of BMSCs administration (Rosselli et al., 2016). Additionally, a case report of a cat with lumbar fracture that gained clinical rehabilitation after intraoperative BMSCs transplantation reinforced their beneficial effects in SCI (Penha et al., 2012).

Different tissue derived mesenchymal stem cells

Synovium (SDSCs) and cartilage (CSPCs) - derived stem cells

SDSCs showed osteogenic capacity similar to BMSCs and rapid proliferation like ADSCs (Bearden et al., 2017). As referred to chondrogenic differentiation, SDSCs were found to be superior to ADSCs and BMSCs (Sasaki et al., 2018). In fact, the intra-articular injection of autologous SDSCs combined with hyaluronic acid in a canine model for cartilage repair, resulted in better macroscopic and histological scores compared to the control group (Miki et al., 2015). However, the studies lacked of statistical analysis. Until now, only autologous CSPCs harvested from ear cartilage showed capacity to regenerate both elastic and hyaline chondral tissue (Mizuno et al., 2014).

Muscle stem cells

Satellite cells become activated after trauma, as muscle precursor cells (MPCs). Satellite cells have high myogenic capacities, but they are no good candidates for transplantation, because of harvesting, survival and migration problems. However, their good

expanding and long term survival in vivo (Eberli et al., 2012) promoted their usage for functional regulation of urinary sphincter (Eberli et al., 2012) and re-innervated thyroarytenoid muscle (Paniello et al., 2018) in canine models. Histological analysis of both studies revealed the formation of new innervated muscle fibres. Muscle-derived stem cells (MDSCs) can be differentiated not only in myogenic, but also in other lineages, such as osteogenic, chondrogenic and tenocyte-like cells. In fact, MDSCs were superior compared to BMSCs for tendon repair (Ozasa et al., 2014). Nevertheless, these findings were concluded in vitro. The majority of studies investigated MDSCs efficacy in dystrophic models. In 2011, a study proposed the intra-arterial administration of allogeneic MDSCs in dystrophic dogs to promote muscle regeneration (Rouger et al., 2011). Despite the limitation of small sample size, clinical and histopathological outcomes were favourable. Additionally, MDSCs were studied in molecular lever (Robriquet et al., 2015, Lardenois et al., 2016). Furthermore, a trial to make their usage easier showed that transient immunosuppression of the hosts was sufficient (Lorant et al., 2018). Undoubtedly, more research is needed about the immunology features of MSDSs.

Cardiac derived stem cells (CSCs)

Despite the fact that the majority of cardiac cells have not potentials to further differentiation, there is a population of CSCs with capacities to form myocytes and promote angiogenesis. In contrast to MSDSs, which could differentiate into muscle cells when injected into the myocardium (Yoon et al., 1995), CSCs could enhance both cardiac regeneration and function by producing functional cardiac cells (Welt et al., 2013, Taghavi et al., 2015). Autologous CSCs helped cardiac remodelling in infraction canine model (Welt et al., 2013) and improved cardiac function in feline cardiomyopathy model (Taghavi et al., 2015). However, in both studies the delivery method was invasive and new methods should be tried in order to make CSCs usage a possible clinical approach.

Hepatic stem cells

Hepatic progenitor cells (HPCs) differentiated into hepatocytes and cholangiocytes (Kruitwagen et al., 2014). The activation, reaction and identification of HPCs were evaluated in normal dogs (Ijzer et al., 2010) and cats (Ijzer et al., 2009). Also, HPCs were assessed in dogs with various liver diseases, such as acute and chronic hepatitis, cooper toxicosis, cancer

or biliary problems (Ijzer et al., 2010, Kruitwagen et al. 2019) and in cats with feline cholangitis (Otte et al., 2018) and lipidosis (Valtolina et al., 2018). Despite the fact that HPCs could be an alternative to liver transplant, which is not feasible in veterinary medicine, no current trials of HPCs usage have been reported.

Dental pulp (DSCs), skin and hair follicle (HFSCs) stem cells

DSCs have high proliferation rate, easy accessibility and differentiation potentials into many cell lineage (Ashiry et al., 2018). Autologous DSCs were used successfully in canine models for pulp regeneration (Nakashima and Iohara, 2014, Chen et al., 2015, Ashiry et al., 2018) and periodontal healing (Khor-sand et al., 2013). Also, allogeneic DSCs application was efficacious for periodontitis management (Iohara et al., 2018). Moreover, allogeneic DSCs were used in degenerative heart disease (Petchdee and Sompee-wong, 2016) and xenogeneic (human) DSCs in SCI in dogs (Feitosa et al., 2017). Despite the promising results, both studies lacked histological and imaging assessments. Also, periodontal stem cells (PDLSCs) were isolated as another tool for periodontitis therapy, but the studies were in vitro (Khoshhal et al., 2017). To the authors' knowledge skin and hair follicle stem cells were identified and characterized in dogs (Brachelente et al., 2013), but they have not been used in vivo.

Umbilical cord blood-derived stem cells (UCSCs)

UCSCs exhibit less immunogenicity, higher plasticity and production of cells per volume compared to MSCs (Jin et al., 2008). UCSCs were used mainly in SCI models (Lim et al., 2007, Park et al. 2011, Ryu et al. 2012) in order to promote clinical improvement and regulate inflammation (Park et al., 2011). However, histopathology and imaging evidence is needed (Lim et al., 2007, Ryu et al., 2012).

Hematopoietic stem cells (HSCs)

HSCs originate from bone marrow, umbilical cord and peripheral blood. HSCs form lymphoid and myeloid progenitor cells (Gomes et al., 2017). In 1962 and 1974, bone marrow transplantation in dogs implied its benefits (Thomas et al., 1962, Epstein et al., 1967, Dale and Graw, 1974). Later, studies used autologous hematopoietic stem cells transplantation (HSCT), against auto-immune diseases (Stolfi et al., 2016). Also, allogeneic HSCT was employed in the

modification of host-versus-graft reaction (Zorn et al., 2011, Mathes et al., 2014, Vrecenak et al., 2014, Rosinski et al., 2015). Interestingly, allogeneic HSCT improved hosts' tolerance towards skin allografts, by T-cells regulation (Mathes et al., 2014). The first clinical case of allogeneic HSCT was completed in a dog with lymphoma (Lupu et al., 2006). Then, two studies tried autologous HSCT for lymphoma (Escobar et al., 2012, Willcox et al., 2012) and acute leukemia (Suter et al., 2015). All agreed that HSCT increased the duration of signs remission. However, according to Schaefer et al. (2016) allogeneic HSCT had problems related to HSCs ability to reach bone marrow without being trapped in other tissues. Lange et al. (2017) tried to identify the best route of administration by comparing the intravenous and intraosseous HSCT. However, the study failed to prove superiority of one method over the other. Few are known about feline HSCs, which have similar proliferative capacities to canine (Abkowitz et al., 1993, Abkowitz et al., 1995). Feline HSCs were used after genetical manipulation for inhibition of coronavirus replication and development of feline infectious peritonitis (Anis et al., 2017).

CONCLUSIONS

In conclusion, literature review cannot still answer questions regarding the best origin of SC, the most suitable route of administration, whether they should be embedded into scaffolds or not as well as which is the ideal number of transplanted cells. Also, it remains unanswered if the therapeutic potentials of SCs exist due to their differentiation into progenitor cells or due to their immunomodulatory secretions. Many experimental models are used for preliminary data, while the clinical use of SCs is in early stages. Most of these studies focused on the usage of BMSCs and ADSCs, because of their easy collection and culture, their sufficient differentiation capacity in musculoskeletal cells and high proliferation rate. However published data have limitations because of their small sample size, heterogeneity of population, lack of blind randomization and control groups. Therefore, a future research plan should involve methods and techniques capable to collect answers for all these questions.

CONFLICT OF INTEREST

None declared.

REFERENCES

- Abdel-Hamid M, Hussein MR, Ahmad AF, Elgezawi EM (2005) Enhancement of the repair of meniscal wounds in the red-white zone (middle third) by the injection of bone marrow cells in canine animal model. *Int J Exp Pathol* 86(2):117-123.
- Abkowitz JL, Linenberger ML, Persik M, Newton MA, Guttorp P (1993) Behavior of feline hematopoietic stem cells years after busulfan exposure. *Blood* 82(7):2096-2103.
- Abkowitz JL, Persik MT, Shelton GH, Ott RL, Kiklevich JV, Catlin SN, Guttorp P (1995) Behavior of hematopoietic stem cells in a large animal. *Proc Natl Acad Sci U S A* 92(6):2031-2035.
- Alamoudi NM, El Ashiry EA, Farsi NM, El Derwi DA, Atta HM (2014) Treatment of oral ulcers in dogs using adipose tissue-derived mesenchymal stem cells. *J Clin Pediatr Dent* 38(3):215-222.
- Aly LA, Menoufy H, HS, Ragae A, Sabry D (2014) Efficiency of systemic versus intralesional bone marrow-derived stem cells in regeneration of oral mucosa after induction of formocresol induced ulcers in dogs. *Dent Res J* 11(2): 212-221.
- Anis EA, Dhar M, Legendre AM, Wilkes RP (2017) Transduction of hematopoietic stem cells to stimulate RNA interference against feline infectious peritonitis. *J Feline Med Surg* 19(6):680-686.
- Arinze TL, Peter SJ, Archambault MP, van den Bos C, Gordon S, Kraus K, Kadiyala S (2003) Allogeneic mesenchymal stem cells regenerative bone in a critical-sized canine segmental defect. *J Bone Joint Surg Am* 85 (10):1927-1935.
- Arzi B, Mills-Ko E, Verstraete FJ, Kol A, Walker NJ, Badgley MR, Fazel N, Murphy WJ, Vapniarsky N, Borjesson DL (2016) Therapeutic Efficacy of Fresh, Autologous Mesenchymal Stem Cells for Severe Refractory Gingivostomatitis in Cats. *Stem Cells Transl Med* 5(1):75-86.
- Ashiry EA El, Alamoudi NM, Ashiry MK El, Bastawy HA, Derwi DA El, Atta HM (2018) Tissue Engineering of Necrotic Dental Pulp of Immature Teeth with Apical Periodontitis in Dogs: Radiographic and Histological Evaluation. *J Clin Pediatr Dent* 42(5):373-382.
- Banas A, Teratani T, Yamamoto Y, Tokuhara M, Takeshita F, Quinn G, Okochi H, Ochiya T (2007) Adipose tissue-derived mesenchymal stem cells as a source of human hepatocytes. *Hepatology* 46(1):219-228.
- Bearden RN, Huggins SS, Cummings KJ, Smith R, Gregory CA, Saunders WB (2017) In-vitro characterization of canine multipotent stromal cells isolated from synovium, bone marrow, and adipose tissue: a donor-matched comparative study. *Stem Cell Res Ther* 8(1):218.
- Besalti O, Aktas Z, Can P, Akpinar E, Elcin AE, Elcin YM (2016) The use of autologous neurogenically-induced bone marrow-derived mesenchymal stem cells for the treatment of paraplegic dogs without nociception due to spinal trauma. *J Vet Med Sci* 78(9):1465-1473.
- Besalti O, Can P, Akpinar E, Aktas Z, Elcin AE, Elcin YM (2015) Intra-spinal Transplantation of Autologous Neurogenically-Induced Bone Marrow-Derived Mesenchymal Stem Cells in the Treatment of Paraplegic Dogs without Deep Pain Perception Secondary to Intervertebral Disk Disease. *Turk Neurosurg* 25(4):625-632.
- Bigham-Sadegh A, Mirshokraei P, Karimi I, Oryan A, Aparviz A, Shafiei-Sarvestani Z (2012) Effects of adipose tissue stem cell concurrent with greater omentum on experimental long-bone healing in dog. *Connect Tissue Res* 53(4):334-342.
- Bittencourt MKW, Barros MA, Martins JFP, Vasconcellos JPC, Morais BP, Pompeia C, Bittencourt MD, Evangelho KS, Kerkis I, Wenceslau VC (2016) Allogeneic Mesenchymal Stem Cell Transplantation in Dogs With Keratoconjunctivitis Sicca. *Cell Med* 8(3): 63-77.
- Black LL, Gaynor J, Adams C, Dhupa S, Sams AE, Taylor R, Harman S, Gingerich DA, Harman R (2008) Effect of intraarticular injection of autologous adipose-derived mesenchymal stem and regenerative cells on clinical signs of chronic osteoarthritis of the elbow joint in dogs. *Vet Ther* 9(3):192-200.
- Black LL, Gaynor J, Gahring D, Adams C, Aron D, Harman S, Gingerich DA, Harman R (2007) Effect of adipose-derived mesenchymal stem and regenerative cells on lameness in dogs with chronic osteoarthritis of the coxofemoral joints: a randomized, double-blinded, multicenter, controlled trial. *Vet Ther* 8(4): 272-284.
- Bornes TD, Adesida AB, Jomha NM (2014) Mesenchymal stem cells in the treatment of traumatic articular cartilage defects: a comprehensive review. *Arthritis Res Ther* 16(5):432.
- Brachelente C, Porcellato I, Sforza M, Lepri E, Mechelli L, Bongiovanni L (2013) The contribution of stem cells to epidermal and hair follicle tumours in the dog. *Vet Dermatol* 24: 188-194.
- Bressan E, Botticelli D, Sivoletta S, Bengazi F, Guazzo R, Sbricoli L, Ricci S, Letizia Ferroni L, Gardin C, Velez JU, Zavan B (2015) Adipose-Derived Stem Cells as a Tool for Dental Implant Osseointegration: an Experimental Study in the Dog. *Int J Mol Cell Med* 4(4): 197-208.
- Canapp SO JR, Canapp DA, Ibrahim V, Carr BJ, Cox C, Barrett JG (2016) The Use of Adipose-Derived Progenitor Cells and Platelet-Rich Plasma Combination for the Treatment of Supraspinatus Tendinopathy in 55 Dogs: A Retrospective Study. *Front Vet Sci* 3:61b.
- Canapp SO JR, Leasure CS, Cox C, Ibrahim V, Carr BJ (2016) Partial Cranial Cruciate Ligament Tears Treated with Stem Cell and Platelet-Rich Plasma Combination Therapy in 36 Dogs: A Retrospective Study. *Sherman O. Front Vet Sci* 3: 112 a.
- Case JB, Palmer R, Valdes-Martinez A, Egger EL, Haussler KK (2013) Gastrocnemius tendon strain in a dog treated with autologous mesenchymal stem cells and a custom orthosis. *Vet Surg* 42(4):355-360.
- Chen KS, Chien KC, Huang YS, Chang CLT, Chang SC (2014) Proliferative effect of platelet rich fibrin on canine bone marrow derived stromal cells. *Taiw Vet J* 40(3):151-161.
- Chen YJ, Zhao YH, Zhao YJ, Liu NX, Lv X, Li Q, Chen FM, Zhang M (2015) Potential dental pulp revascularization and odonto-/osteogenic capacity of a novel transplant combined with dental pulp stem cells and platelet-rich fibrin. *Cell Tissue Res* 361(2):439-455.
- Chung WH, Park SA, Lee JH, Chung DJ, Yang WJ, Kang EH, Choi CB, Chang HS, Kim DH, Hwang SH, Han H, Kim HY (2013) Percutaneous transplantation of human umbilical cord-derived mesenchymal stem cells in a dog suspected to have fibrocartilaginous embolic myelopathy. *J Vet Sci* 14(4):495-497.
- Cruz AC, Caon T, Menin A, Granato R, Boabaid F, Simões CM (2015) Adipose-derived stem cells incorporated into platelet-rich plasma improved bone regeneration and maturation in vivo. *Dent Traumatol* 31(1):42-48.
- Cuervo B, Rubio M, Sopena J, Dominguez JM, Vilar J, Morales M, Cugat R, Carrillo JM (2014) Hip osteoarthritis in dogs: a randomized study using mesenchymal stem cells from adipose tissue and plasma rich in growth factors. *Int J Mol Sci* 15(8):13437-13460.
- Cui L, Liu B, Liu G, Zhang W, Cen L, Sun J, Yin S, Liu W, Cao Y (2007). Repair of cranial bone defects with adipose derived stem cells and coral scaffold in a canine model. *Biomaterials* 28(36): 5477-5486.
- Dale DC, Graw JR RG (1974) Transplantation of Allogeneic Bone Marrow in Canine Cyclic Neutropenia. *Science* 183(4120): 83-84.
- de Bakker E, Van Ryssen B, De Schauwer C, Meyer E (2013) Canine mesenchymal stem cells: state of the art, perspectives as therapy for dogs and as a model for man. *Vet Q* 33(4):225-233.
- Deng Y, Zhou H, Gu P, Fan X (2014) Repair of canine medial orbital bone defects with miR-31-modified bone marrow mesenchymal stem cells. *Invest Ophthalmol Vis Sci* 55(9):6016-6023.
- Ding F, Wu J, Yang Y, Hu W, Zhu Q, Tang X, Liu J, Gu X (2010) Use of tissue-engineered nerve grafts consisting of a chitosan/poly(lactic-co-glycolic acid)-based scaffold included with bone marrow mesenchymal cells for bridging 50-mm dog sciatic nerve gaps. *Tissue Eng Part A* 16(12):3779-3790.
- Dos Santos RC, da Rocha Filgueiras R, Furtado Malard P, Rodrigues da Cunha Barreto-Vianna A, Nogueira K, da Silva Leite C, Maurício

- Mendes de Lima E (2018) Mesenchymal stem cells in osteotomy repair after tibial tuberosity advancement in dogs with cranial cruciate ligament injury. *J Exp Orthop* 5(1):17
- Eberli D, Aboushwareb T, Soker S, Yoo JJ, Atala A (2012) Muscle precursor cells for the restoration of irreversibly damaged sphincter function. *Cell Transplant* 21(9):2089-2098.
- Epstein RB, Bryant J, Thomas ED (1967) Cytogenetic demonstration of permanent tolerance in adult outbred dogs. *Transpl* 5 (2):267.
- Escobar C, Grindem C, Neel JA, Suter SE(2012) Hematologic changes after total body irradiation and autologous transplantation of hematopoietic peripheral blood progenitor cells in dogs with lymphoma. *Vet Pathol* 49(2):341-343.
- Feitosa MLT, Sarmiento CAP, Bocabello RZ, Beltrão-Braga PCB, Pignatari GC, Giglio RF, Miglino MA, Orlandin JR, Ambrósio CE (2017) Transplantation of human immature dental pulp stem cell in dogs with chronic spinal cord injury. *Acta Cir Bras* 32(7):540-549.
- Ferrer L, Kimbrel EA, Lam A, Falk EB, Zewe C, Juopperi T, Lanza R, Hoffman A (2016) Treatment of perianal fistulas with human embryonic stem cell-derived mesenchymal stem cells: a canine model of human fistulizing Crohn's disease. *Regen Med* 11(1):33-43.
- Fortier LA, Travis AJ (2011) Stem cells in veterinary medicine. *Stem Cell Res Ther* 2 (3):9-14.
- Ganey T, Hutton WC, Moseley T, Hedrick M, Meisel HJ (2009) Intervertebral disc repair using adipose tissue-derived stem and regenerative cells: experiments in a canine model. *Spine* 34(21):2297-2304.
- Ghoreishian M, Rezaei M, Beni BH, Javanmard SH, Attar BM, Zalzali H (2013) Facial nerve repair with Gore-Tex tube and adipose-derived stem cells: an animal study in dogs. *J Oral MaxillofacSurg* 71(3):577-587.
- Gibson MA, Brown SG, Brown NO (2017) Semitendinosus myopathy and treatment with adipose-derived stem cells in working German shepherd police dogs. *Can Vet J* 58(3):241-246.
- Gomes IS, de Oliveira VC , Pinheiro AO, Roballo KCS, de Araujo GSM, Veronezi JC, Martins DS, Ambrósio CE (2017) Bone marrow stem cell applied in the canine veterinary clinics. *Pesq Vet Bras* 37(10): 1139-1145.
- Guercio A, Di Marco P, Casella S, Cannella V, Russotto L, Purpari G, Di Bella S, Piccione G (2012) Production of canine mesenchymal stem cells from adipose tissue and their application in dogs with chronic osteoarthritis of the humeroradial joints. *Cell Biol Int* 36(2):189-194.
- Haghighat A, Akhavan A, Hashemi-Beni B, Deihimi P, Yadegari A, Heidari F (2011) Adipose derived stem cells for treatment of mandibular bone defects: An autologous study in dogs. *Dent Res J* 8(Suppl 1):51-57.
- Hall MN, Rosenkrantz WS, Hong JH, Griffin CE, Mendelsohn CM (2010) Evaluation of the potential use of adipose-derived mesenchymal stromal cells in the treatment of canine atopic dermatitis: a pilot study. *Vet Ther* 2010 11(2):1-14.
- Han SM , Kim HT, Kim KW, Jeon KO, Seo KW, Choi EW, Youn HY (2015) CTLA4 overexpressing adipose tissue-derived mesenchymal stem cell therapy in a dog with steroid-refractory pemphigus foliaceus *BMC Vet Res* 11:49
- Hang D, Wang Q, Guo C, Chen Z, Yan Z (2012) Treatment of osteonecrosis of the femoral head with VEGF165 transgenic bone marrow mesenchymal stem cells in mongrel dogs. *Cells Tissues Organs* 195(6):495-506.
- Hao L, Hao J, Fang W, Han C, Zhang K, Wang X (2015) Dual isotope simultaneous imaging to evaluate the effects of intracoronary bone marrow-derived mesenchymal stem cells on perfusion and metabolism in canines with acute myocardial infarction. *Biomed Rep* 3(4):447-452.
- Harman R, Carlson K, Gaynor J, Gustafson S, Dhupa S, Clement K, Hoelzler, McCarthy MT, Schwartz P, Adams C (2016) A Prospective, Randomized, Masked, and Placebo-Controlled Efficacy Study of Intra-articular Allogeneic Adipose Stem Cells for the Treatment of Osteoarthritis in Dogs *Front Vet Sci* 3: 81.
- Hashemibeni B, Goharian V, Esfandiari E, Sadeghi F, Fasihi F, Alipur R, Valiani A, Ghorbani M, Emami ZM, Shabani F, Goharian M (2012) An animal model study for repair of tracheal defects with autologous stem cells and differentiated chondrocytes from adipose-derived stem cells. *J PediatrSurg* 47(11):1997-2003.
- Hiyama A, Mochida J, Iwashina T, Omi H, Watanabe T, Serigano K, Tamura F, Sakai D (2008) Transplantation of mesenchymal stem cells in a canine disc degeneration model. *J Orthop Res*:26(5):589-600.
- Hu YC, Liu X, Shen JJ, He JC, Chen QE (2014) [Experimental study of canine bone marrow mesenchymal stem cells combined with calcium phosphate cement for repair of mandibular bone defects in Beagle dogs]. [Article in Chinese] *Shanghai Kou Qiang Yi Xue* 23(4):402-408.
- Ijzer J, Kisjes JR, Penning LC, Rothuizen J, van den Ingh TS (2009) The progenitor cell compartment in the feline liver: an (immuno)histochemical investigation. *Vet Pathol* 46(4):614-621.
- Ijzer J, Schotanus BA, Vander Borgh S, Roskams TA, Kisjes R, Penning LC, Rothuizen J, van den Ingh TS (2010) Characterisation of the hepatic progenitor cell compartment in normal liver and in hepatitis: an immunohistochemical comparison between dog and man. *Vet J* 184(3):308-314.
- Iohara K, Utsunomiya S, Kohara S, Nakashima M (2018) Allogeneic transplantation of mobilized dental pulp stem cells with the mismatched dog leukocyte antigen type is safe and efficacious for total pulp regeneration. *Stem Cell Res Ther* 9(1):116.
- Iravani K, Sobhanmanesh A, Ashraf MJ, Hashemi SB, Mehrabani D, Zare S (2017) The Healing Effect of Conditioned Media and Bone Marrow-Derived Stem Cells in Laryngotracheal Stenosis: A Comparison in Experimental Dog Model. *World J PlastSurg* 6(2):190-197.
- Jin GZ, Yin XJ, Yu XF, Cho SJ, Choi EG, Lee YS, Jeon JT, Yee ST, Kong IK (2008) Generation of neuronal-like cells from umbilical cord blood-derived mesenchymal stem cells of a RFP-transgenic cloned cat *J Vet Med Sci* 70(7):723-726.
- JinH, XuT, Chen Q, Wu C, Wang P, Mao Q, Zhang S, Shen J, Tong P (2016) The Fate and Distribution of Autologous Bone Marrow Mesenchymal Stem Cells with Intra-Arterial Infusion in Osteonecrosis of the Femoral Head in Dogs. *Stem Cells International* 2016:1-10.
- Jung DI, Ha J, Kang BT, Kim JW, Quan FS, Lee JH, Woo EJ, Park HM (2009) A comparison of autologous and allogeneic bone marrow-derived mesenchymal stem cell transplantation in canine spinal cord injury. *J Neurol Sci* 285(1-2):67-77.
- Kaizawa Y, Kakinoki R, Ikeguchi R, Ohta S, Noguchi T, Oda H, Matsuda S (2016) Bridging a 30 mm defect in the canine ulnar nerve using vessel-containing conduits with implantation of bone marrow stromal cells. *Microsurgery* 36(4):316-324.
- Kamishina H, Deng J, Oji T, Cheeseman JA, Clemmons RM (2006) Expression of neural markers on bone marrow-derived canine mesenchymal stem cells. *Am J Vet Res* 67(11):1921-1928.
- Kanemaru S, Nakamura T, Omori K, Kojima H, Magruffov A, Hiratsuka Y, Hirano S, Ito J, Shimizu Y (2003) Regeneration of the vocal fold using autologous mesenchymal stem cells. *AnnOtolRhinolLaryngol* 112(11):915-920.
- Kazemi D, Shams Asenjan K, Dehdilani N, Parsa H (2017) Canine articular cartilage regeneration using mesenchymal stem cells seeded on platelet rich fibrin: Macroscopic and histological assessments. *Bone Joint Res* 6(2):98-107.
- Khorsand A, Eslaminejad MB, Arabsofghar M, Paknejad M, Ghaedi B, Rokn AR, Moslemi N, Nazarian H, Jahangir S (2013) Autologous dental pulp stem cells in regeneration of defect created in canine periodontal tissue. *J Oral Implantol* 39(4):433-443.
- Khoshhal M, Amiri I, Gholami L (2017) Comparison of in vitro properties of periodontal ligament stem cells derived from permanent and deciduous teeth. *J Dent Res Dent Clin Dent Prospects* 11(3):140-148.
- Kim JW, Lee JH, Lyoo YS, Jung DI, Park HM (2013) The effects of topical mesenchymal stem cell transplantation in canine experimental cutaneous wounds. *Vet Dermatol* 24(2):242-253.
- Kim SH, Kim KH, Seo BM, Koo KT, Kim TI, Seol YJ, Ku Y, Rhyu IC, Chung CP, Lee YM (2009) Alveolar bone regeneration by transplantation of periodontal ligament stem cells and bone marrow stem cells in a canine peri-implant defect model: a pilot study. *J Periodontol*

- 80(11):1815-1823.
- Kim Y, Lee SH, Kim WH, Kweon OK (2016) Transplantation of adipose derived mesenchymal stem cells for acute thoracolumbar disc disease with no deep pain perception in dogs. *J Vet Sci* 17(1): 123-126.
- Kriston-Pál É, Czibula Á, Gyuris Z, Balka G, Seregi A, Sükösd F, Süth M, Kiss-Tóth E, Haracska L, Uher F, Monostori É (2017) Characterization and therapeutic application of canine adipose mesenchymal stem cells to treat elbow osteoarthritis. *Can J Vet Res* 81(1):73-78.
- Kruitwagen HS, Bart Spee B, Schotanus BA (2014) Hepatic progenitor cells in canine and feline medicine: potential for regenerative strategies *BMC Vet Res* 10: 137.
- Kruitwagen HS, Fieten H, Penning LC (2019) Towards bioengineered liver stem cell transplantation studies in a preclinical dog model for inherited copper toxicosis *Bioeng* 2019;6(4):88.
- Kubo H, Berretta RM, Jaleel N, Angert D, Houser SR (2009) c-Kit⁺ bone marrow stem cells differentiate into functional cardiac myocytes. *Clin Transl Sci* 2(1):26-32.
- Lange S, Steder A, Killian D, Knuebel G, Sekora A, Vogel H, Lindner I, Dunkelmann S, Prall F, Murua H, Mathias E, Junghans FC (2017) Biology of Blood and Marrow Transplantation Biology Engraftment Efficiency after Intra-Bone Marrow versus Intravenous Transplantation of Bone Marrow Cells in a Canine Nonmyeloablative Dog Leukocyte Antigen-Identical Transplantation Model. *Biology of Blood and Marrow Transplantation* 23 (2): 247-254.
- Lardenois A, Jagot S, Lagarrigue M, Guével B, Ledevin M, Larcher T, Dubreil L, Pineau C, Rouger K, Guével L (2016) Quantitative proteome profiling of dystrophic dog skeletal muscle reveals a stabilized muscular architecture and protection against oxidative stress after systemic delivery of MuStem cells. *Proteomics* 16(14):2028-2042.
- Lee JH, Chang HS, Kang EH, Chung DJ, Choi CB, Lee JH, Hwang SH, Han H, Kim HY (2009) Percutaneous transplantation of human umbilical cord blood-derived multipotent stem cells in a canine model of spinal cord injury. *J Neurosurg Spine* 11(6):749-757.
- Lim CY, Han JI, Kim SG, Lee CM, Park HM (2016) Evaluation of autologous bone marrow-derived mesenchymal stem cells on renal regeneration after experimentally induced acute kidney injury in dogs. *Am J Vet Res* 77(2):208-217.
- Lim JH, Byeon YE, Ryu HH, Jeong YH, Lee YW, Kim WH, Kang KS, Kweon OK (2007) Transplantation of canine umbilical cord blood-derived mesenchymal stem cells in experimentally induced spinal cord injured dogs. *J Vet Sci* 8(3):275-282.
- Liu G, Zhang Y, Liu B, Sun J, Li W, Cui L (2013) Bone regeneration in a canine cranial model using allogeneic adipose derived stem cells and coral scaffold. *Biomaterials* 34(11):2655-2664.
- Liu X, Li Q, Wang F, Wang Z (2016) Maxillary sinus floor augmentation and dental implant placement using dentin matrix protein-1 gene-modified bone marrow stromal cells mixed with deproteinized bovine bone: A comparative study in beagles. *Arch Oral Biol* 64:102-108.
- Liu X, Liao X, Luo E, Chen W, Bao C, Xu HH (2014) Mesenchymal stem cells systemically injected into femoral marrow of dogs home to mandibular defects to enhance new bone formation. *Tissue Eng Part A* 20(3-4):883-892.
- Liu Z, Yin X, Ye Q, He W, Ge M, Zhou X, Hu J, Zou S (2016) Periodontal regeneration with stem cells-seeded collagen-hydroxyapatite scaffold. *J Biomater Appl* 31(1):121-131.
- Lorant J, Larcher T, Jaulin N, Hedan B, Lardenois A, Leroux I, Dubreil L, Ledevin M, Goubin H, Moullec S, Deschamps JY, Thorin C, André C, Adjali O, Rouger K (2018) Vascular Delivery of Allogeneic MuStem Cells in Dystrophic Dogs Requires Only Short-Term Immunosuppression to Avoid Host Immunity and Generate Clinical/Tissue Benefits. *Cell Transplant* 27(7):1096-1110.
- Lupu M, Sullivan EW, Westfall TE, Little MT, Weigler BJ, Moore PF, Stroup PA, Zellmer E, Kuhr C, Storb R (2006) Use of multigeneration-family molecular dog leukocyte antigen typing to select a hematopoietic cell transplant donor for a dog with T-cell lymphoma. *J Am Vet Med Assoc* 228(5):728-732.
- Martin DR, Cox NR, Hathcock TL, Niemeyer GP, Baker HJ (2002) Isolation and characterization of multipotential mesenchymal stem cells from feline bone marrow. *Exp Hematol* 30(8):879-886.
- Marx C, Silveira MD, Selbach I, da Silva AS, Gomes de Macedo Braga LM, Camassola M, Nardi NB (2014) Acupoint Injection of Autologous Stromal Vascular Fraction and Allogeneic Adipose-Derived Stem Cells to Treat Hip Dysplasia in Dogs. *Stem Cells Intern* 2014:391274.
- Mathes DW, Chang J, Hwang B, Graves SS, Storer BE, Butts-Miwongtum T, Sale GE, Storb R (2014) Simultaneous transplantation of hematopoietic stem cells and a vascularized composite allograft leads to tolerance. *Transplantation* 98(2):131-138.
- Matsuda T, Takami T, Sasaki R, Nishimura T, Aibe Y, Paredes BD, Quintanilha LF, Matsumoto T, Ishikawa T, Yamamoto N, Tani K, Terai S, Taura Y, Sakaida I (2017) A canine liver fibrosis model to develop a therapy for liver cirrhosis using cultured bone marrow-derived cells. *Hepato Comm* 1(7):691-703.
- McDougall RA, Canapp SO, Canapp DA (2018) Ultrasonographic Findings in 41 Dogs Treated with Bone Marrow Aspirate Concentrate and Platelet-Rich Plasma for a Supraspinatus Tendinopathy: A Retrospective Study. *Front Vet Sci* 5,98.
- Memon IA, Sawa Y, Miyagawa S, Taketani S, Matsuda H (2005) Combined autologous cellular cardiomyoplasty with skeletal myoblasts and bone marrow cells in canine hearts for ischemic cardiomyopathy. *J Thorac Cardiovasc Surg* 130(3):646-653.
- Miki S, Takao M, Miyamoto W, Matsushita T, Kawano H (2015) Intra-Articular Injection of Synovium-Derived Mesenchymal Stem Cells with Hyaluronic Acid Can Repair Articular Cartilage Defects in a Canine Model. *J Cell Res Ther* 5:314.
- Mizuno M, Kobayashi S, Takebe T, Kan H, Yabuki Y, Matsuzaki T, Yoshikawa HY, Nakabayashi S, Ik LJ, Maegawa J, Taniguchi H (2014) Brief report: reconstruction of joint hyaline cartilage by autologous progenitor cells derived from ear elastic cartilage. *Stem Cells* 32(3):816-821.
- Muir P, Hans EC, Racette M, Volstad N, Sample SJ, Heaton C, Holzman G, Schaefer SL, Bloom DD, Bleedorn JA, Hao Z, Amene E, Suresh M, Hematti P (2016) Autologous Bone Marrow-Derived Mesenchymal Stem Cells Modulate Molecular Markers of Inflammation in Dogs with Cruciate Ligament Rupture. *PLoS One* 11(8):1-21.
- Munoz JL, Greco SG, Sherman LS, Bhatt S, Bhatt RS, Shrensel JA, Guan YZ, Xie G, Ye JH, Rameshwar P, Siegel A (2012) Feline bone marrow-derived mesenchymal stromal cells (MSCs) show similar phenotype and functions with regards to neuronal differentiation as human MSCs. *Differentiation* 84(2): 214-222.
- Nakashima M, Iohara K (2014) Mobilized dental pulp stem cells for pulp regeneration: initiation of clinical trial. *J Endod* 40(4 Suppl):26-32.
- Nantavisai S, Egusa H, Thanaphum O, Sawangmake C (2019) Mesenchymal stem cell-based bone tissue engineering for veterinary practice. *Heliyon* 5(11):e 02808
- Nam A, Han SM, Go DM, Kim DY, Seo KW, Youn HY (2017) Long-Term Management with Adipose Tissue-Derived Mesenchymal Stem Cells and Conventional Treatment in a Dog with Hepatocutaneous Syndrome. *J Vet Intern Med* 31(5):1514-1519.
- Nishida H, Nakayama M, Tanaka H, Kitamura M, Hatoya S, Sugiura K, Suzuki Y, Ide C, Inaba T (2011) Evaluation of transplantation of autologous bone marrow stromal cells into the cerebrospinal fluid for treatment of chronic spinal cord injury in dogs. *Am J Vet Res* 72(8):1118-1123.
- Nishida H, Nakayama M, Tanaka H, Kitamura M, Hatoya S, Sugiura K, Harada Y, Suzuki Y, Ide C, Inaba T (2012) Safety of autologous bone marrow stromal cell transplantation in dogs with acute spinal cord injury. *Vet Surg* 41(4):437-442.
- Osman Y, Hamed SH, Barakat NM, Khater S, Gabr M, Mosbah A, Gaballah MA, Shaaban A (2020) Prophylaxis against renal ischemia-reperfusion injury in canine model: Stem cell approach. *Indian J Urol* 36(1):44-49.
- Otte CM, Valtolina C, Vreman S, Hubers S, van Wolferen ME, Favier RP, Rothuizen J, Penning LC (2018) Immunohistochemical evaluation of the activation of hepatic progenitor cells and their niche in feline lymphocytic cholangitis. *J Feline Med Surg* 20(1):30-37.
- Ozasa Y, Gingery A, Thoreson AR, An KN, Zhao C, Amadio PC (2014) A

- comparative study of the effects of growth and differentiation factor 5 on muscle-derived stem cells and bone marrow stromal cells in an in vitro tendon healing model *J Hand Surg Am* 39(9):1706-1713.
- Özdamar Kurt F, Tuğlu I, Vatanserver HS, Tong S, Delilöğlu-Gürhan SI (2015) The effect of autologous bone marrow stromal cells differentiated on scaffolds for canine tibial bone reconstruction. *Biotech Histochem* 90(7):516-528.
- Paniello RC, Brookes S, Bhatt NK, Bijangi-Vishehsaraei K, Zhang H, Halum S (2018) Improved adductor function after canine recurrent laryngeal nerve injury and repair using muscle progenitor cells. *Laryngoscope* 128(7):241-246.
- Park SA, Reilly CM, Wood JA, Chung DJ, Carrade DD, Deremer SL, Seraphin RL, Clark KC, Zwingenberger AL, Borjesson DL, Hayashi K, Russell P, Murphy CJ (2013) Safety and immunomodulatory effects of allogeneic canine adipose-derived mesenchymal stromal cells transplanted into the region of the lacrimal gland, the gland of the third eyelid and the knee joint. *Cytotherapy* 15(12):1498-1510.
- Park SS, Byeon YE, Ryu HH, Kang BJ, Kim Y, Kim WH, Kang KS, Han HJ, Kweon OK (2011) Comparison of canine umbilical cord blood-derived mesenchymal stem cell transplantation times: involvement of astrogliosis, inflammation, intracellular actin cytoskeleton pathways, and neurotrophin-3. *Cell Transplan* 20(11-12):1867-1880.
- Park SS, Lee YJ, Lee SH, Lee D, Choi K, Kim WH, Kweon OK, Han HJ (2012) Functional recovery after spinal cord injury in dogs treated with a combination of Matrigel and neural-induced adipose-derived mesenchymal Stem cells. *Cytotherapy* 14(5):584-597.
- Parys M, Nelson N, Koehl K, Miller R, Kaneene JB, Kruger JM, Yuzbasiyan-Gurkan V (2016) Safety of Intraperitoneal Injection of Adipose Tissue-Derived Autologous Mesenchymal Stem Cells in Cats. *J Vet Intern Med* 30(1):157-163.
- Peng WX, Wang L (2017) Adenovirus-Mediated Expression of BMP-2 and BFGF in Bone Marrow Mesenchymal Stem Cells Combined with Demineralized Bone Matrix For Repair of Femoral Head Osteonecrosis in Beagle Dogs. *Cell PhysiolBiochem* 43(4):1648-1662.
- Penha EM, Aguiar PHP, Barrouin-Melo SM, de Lima RS, da Silveira ACC, Oteleto ARS, Pinheiro CMB, dos-Santos RR, Soares MBP (2012) Clinical Neurofunctional Rehabilitation of a Cat with Spinal Cord Injury after Hemilaminectomy and Autologous Stem Cell Transplantation. *Int J Stem Cells* 5(2): 146-150.
- Penha EM, MeiraCS, Guimaraes ET, Mendonça MVP, Gravelly FA, Pinheiro CMB, Pinheiro TMB, Barrouin-Melo SM, dos-Santos RR, Soares MBP (2014) Use of Autologous Mesenchymal Stem Cells Derived from Bone Marrow for the Treatment of Naturally Injured Spinal Cord in Dogs *Stem Cells International* 2014, 437521.
- Pérez-Merino EM, Usón-Casaús JM, Zaragoza-Bayle C, Duque-Carrasco J, Mariñas-Pardo L, Hermida-Prieto M, Barrera-Chacón R, Gualtieri M (2015) Safety and efficacy of allogeneic adipose tissue-derived mesenchymal stem cells for treatment of dogs with inflammatory bowel disease: Clinical and laboratory outcomes. *Vet J* 206(3):385-390.
- Petchdee S, Sompeewong S (2016) Intravenous administration of puppy deciduous teeth stem cells in degenerative valve disease *Vet World* 2016 9(12): 1429-1434.
- Pogue B, Estrada AH, Sosa-Samper I, Maisenbacher HW, Lamb KE, Mincey BD, Erger KE, Conlon TJ (2013) Stem-cell therapy for dilated cardiomyopathy: a pilot study evaluating retrograde coronary venous delivery. *J Small Anim Pract* 54(7):361-366.
- Quimby JM, Webb TL, Gibbons DS, Dow SW (2011) Evaluation of intrarenal mesenchymal stem cell injection for treatment of chronic kidney disease in cats: a pilot study. *J Feline Med Surg* 13(6):418-426.
- Quimby JM, Webb TL, Gibbons DS, Dow SW (2011) Evaluation of intrarenal mesenchymal stem cell injection for treatment of chronic kidney disease in cats: a pilot study. *J Feline Med Surg* 13(6):418-426.
- Quimby JM, Webb TL, Habenicht LM, Dow SW (2013) Safety and efficacy of intravenous infusion of allogeneic cryopreserved mesenchymal stem cells for treatment of chronic kidney disease in cats: results of three sequential pilot studies. *Stem Cell Res Ther* 4(2): 48.
- Quimby JM, Webb TL, Randall E, Marolf A, Valdes-Martinez A, Dow SW (2016) Assessment of intravenous adipose-derived allogeneic mesenchymal stem cells for the treatment of feline chronic kidney disease: a randomized, placebo-controlled clinical trial in eight cats. *J Feline Med Surg* 18(2):165-171.
- Robriquet F, Lardenois A, Babarit C, Larcher T, Dubreil L, Leroux I, Zuber C, Ledevin M, Deschamps JY, Fromes Y, Cherel Y, Guevel L, Rouger K (2015) Differential Gene Expression Profiling of Dys-trophic Dog Muscle after MuStem Cell Transplantation. *PLoS One* 10(5): 1-24.
- Rosinski SL, Stone B, Graves SS, Fuller DH, De Rosa SC, Spies GA, Mize GJ, Fuller JT, Storb R (2015) Development of a Minor Histocompatibility Antigen Vaccine Regimen in the Canine Model of Hematopoietic Cell Transplantation. *Transplantation* 99(10):2083-2094.
- Rosselli DD, Mumaw JL, Dickerson V, Brown CA, Brown SA, Schmiedt CW (2016) Efficacy of allogeneic mesenchymal stem cell administration in a model of acute ischemic kidney injury in cats. *Res Vet Sci* 108:18-24.
- Roszek K, Makowska N, Czarnecka J, Porowińska D, Dąbrowski M, Danielewska J, Nowak W (2017) Canine Adipose-Derived Stem Cells: Purinergic Characterization and Neurogenic Potential for Therapeutic Applications. *J Cell Biochem* 118(1):58-65.
- Rouger K, Larcher T, Dubreil L, Deschamps JY, Guiner C, Jouvion G, Delorme B, Lieubeau B, Carlus M, Fornasari B, Theret M, Orlando P, Ledevin M, Zuber C, Leroux I, Deleau S, Guigand L, Testault I, Rumeur E, Fiszman M, Chérel Y (2011) Systemic Delivery of Allogeneic Muscle Stem Cells Induces Long-Term Muscle Repair and Clinical Efficacy in Duchenne Muscular Dystrophy Dogs. *Am J Pathol* 179(5): 2501-2518.
- Ryan JM, Barry FP, Murphy JM, Mahon BP (2005) Mesenchymal stem cells avoid allogeneic rejection. *J Inflamm* 2:1-11.
- Ryu HH, Kang BJ, Park SS, Kim Y, Sung GJ, Woo HM, Kim WH, Kweon OK (2012) Comparison of mesenchymal stem cells derived from fat, bone marrow, Wharton's jelly, and umbilical cord blood for treating spinal cord injuries in dogs. *J Vet Med Sci* 74(12):1617-1630.
- Ryu HH, Lim JH, Byeon YE, Park JR, Seo MS, Lee YW, Kim WH, Kang KS, Kweon OK (2009) Functional recovery and neural differentiation after transplantation of allogeneic adipose-derived stem cells in a canine model of acute spinal cord injury. *J Vet Sci* 10(4):273-284.
- Sarmento CA, Rodrigues MN, Bocabello RZ, Mess AM, Miglino MA (2014) Pilot study: bone marrow stem cells as a treatment for dogs with chronic spinal cord injury. *Regen Med Res* 2(1):9.
- Sasaki A, Mizuno M, Ozeki N, Katano H, Otabe K, Tsuji K, Koga H, Mochizuki M, Sekiya I (2018) Canine mesenchymal stem cells from synovium have a higher chondrogenic potential than those from infrapatellar fat pad, adipose tissue, and bone marrow. *PLoS One* 13(8):e0202922.
- Schaefer S, Werner J, Lange S, Machka C, Knuebel G, Sekora A, Rooff C, Winkler T, Murua EH, Nolte I, Junghanss C (2016) Graft-versus-Host Disease in a Dog After Reduced-intensity Hematopoietic Stem Cell Transplantation from a DLA-identical Littermate. *In Vivo* 30(4):427-432.
- Serigano K, Sakai D, Hiyama A, Tamura F, Tanaka M, Mochida J (2010) Effect of cell number on mesenchymal stem cell transplantation in a canine disc degeneration model. *J Orthop Res* 28(10):1267-1275.
- Shah K, Drury T, Roic I, Hansen P, Malin M, Boyd R, Sumer H, Ferguson R (2018) Outcome of Allogeneic Adult Stem Cell Therapy in Dogs Suffering from Osteoarthritis and Other Joint Defects. *Stem Cells Int* 2018:1-7.
- Shan X, Hu D (2017) Bone engineering by cell sheet technology to repair mandibular defects. *Exp Ther Med* 14(5): 5007-5011.
- Shen H, Jayaram R, Yoneda S, Linderman SW, Sakiyama-Elbert SE, Xia Y, Gelberman RH, Thomopoulos S (2018) The effect of adipose-derived stem cell sheets and CTGF on early flexor tendon healing in a canine model. *Sci Rep* 23;8(1):11078.
- Steffen F, Smolders LA, Roentgen AM, Bertolo A, Stoyanov J (2017) Bone Marrow-Derived Mesenchymal Stem Cells as Autologous Therapy in Dogs with Naturally Occurring Intervertebral Disc Disease: Feasibility, Safety, and Preliminary Results. *Tissue Eng Part C*

- Methods 23(11):643-651.
- Stolfi JL, Pai CC, Murphy WJ (2016) Preclinical modeling of hematopoietic stem cell transplantation - advantages and limitations. *FEBS J* 283(9):1595-1606.
- Suter SE, Hamilton MJ, Sullivan EW, Venkataraman GM (2015) Allogeneic hematopoietic cell transplantation in a dog with acute large granular lymphocytic leukemia. *J Am Vet Med Assoc* 246(9):994-997.
- Taghavi S, Sharp TE, Duran JM, Makarewich CA, Berretta RM, Starosta T, Kubo H, Barbe M, Houser SR (2015) Autologous c-Kit⁺ Mesenchymal Stem Cell Injections Provide Superior Therapeutic Benefit as Compared to c-Kit⁺ Cardiac-Derived Stem Cells in a Feline Model of Isoproterenol-Induced Cardiomyopathy. *Clin Transl Sci* 8(5):425-431.
- Takemitsu H1, Zhao D, Yamamoto I, Harada Y, Michishita M, Arai T (2012) Comparison of bone marrow and adipose tissue-derived canine mesenchymal stem cells. *BMC Vet Res* 8 (1):150-159.
- Tan B, Wei RQ, Tan MY, Luo JC, Deng L, Chen XH, Hou JL, Li XQ, Yang ZM, Xie HQ (2013) Tissue engineered esophagus by mesenchymal stem cell seeding for esophageal repair in a canine model. *J Surg Res* 182(1):40-48.
- Teshima T, Matsumoto H, Michishita M, Matsuoka A, Shiba M, Nagashima T, Koyama H (2017) Allogenic Adipose Tissue-Derived Mesenchymal Stem Cells Ameliorate Acute Hepatic Injury in Dogs. *Stem Cells Int* 2017:3892514.
- Thomas ED, Collins JA, Herman EC JR, Ferrebee JW (1962) Marrow Transplants in Lethally Irradiated Dogs Given Methotrexate. *Blood* 19:217-228
- Tobita M, Uysal CA, Guo X, Hyakusoku H, Mizuno H (2013) Periodontal tissue regeneration by combined implantation of adipose tissue-derived stem cells and platelet-rich plasma in a canine model. *Cytotherapy* 15(12):1517-1526.
- Tognoli GK, Olsson DC, Martins DB, Júnior EBS, Salbego FZ, de Oliveira GK, Braga FVA, Raiser AG, Dezengrini R, da Cruz FSF, de Castro V MB, Rosa MC, Carregaro AB, Pippi N L (2008) Transplante autólogo de células mononucleares da medula óssea em úlcera de córnea experimental em cães Bone marrow mononuclear cells autotransplant in experimental corneal ulcer in dogs. *Cienc Rural* 39 (1): 148-155.
- Tracy CJ, Sanders DN, Bryan JN, Jensen CA, Castaner LJ, Kirk MD, Katz ML (2016) Intravitreal Implantation of Genetically Modified Autologous Bone Marrow-Derived Stem Cells for Treating Retinal Disorders. *Adv Exp Med Biol* 854:571-577.
- Trzil JE, Masseur I, Webb TL, Chang CH, Dodam JR, Liu H, Quimby JM, Dow SW, Reiner CR (2016) Intravenous adipose-derived mesenchymal stem cell therapy for the treatment of feline asthma: a pilot study. *J Feline Med Surg* 18(12):981-990.
- Valtolina C, Robben JH, Favier RP, Rothuizen J, Grinwis GC, Schotanus BA, Penning LC (2018). Immunohistochemical characterisation of the hepatic stem cell niche in feline hepatic lipidosis: a preliminary morphological study. *J Feline Med Surg* "in press".
- Vilar JM, Batista M, Morales M, Santana A, Cuervo B, Rubio M, Cugat R, Sopena J, Carrillo JM (2014) Assessment of the effect of intraarticular injection of autologous adipose-derived mesenchymal stem cells in osteoarthritic dogs using a double blinded force platform analysis. *BMC Vet Res* 10:143.
- Vilar JM, Morales M, Santana A, Spinella G, Rubio M, Cuervo B, Cugat R, Carrillo JM (2013) Controlled, blinded force platform analysis of the effect of intraarticular injection of autologous adipose-derived mesenchymal stem cells associated to PRGF-Endoret in osteoarthritic dogs. *BMC Vet Res* 9:131.
- Villatoro AJ, Claros S, Fernández V, Alcholado C, Fariñas F, Moreno A, Becerra J, Andrades JA (2018) Safety and efficacy of the mesenchymal stem cell in feline eosinophilic keratitis treatment. *BMC Vet Res* 14(1):116.
- Villatoro AJ, Fernández V, Claros S, Rico-Llanos GA, Becerra J, Andrades JA (2015) Use of Adipose-Derived Mesenchymal Stem Cells in Keratoconjunctivitis Sicca in a Canine Model. *BioMed Res Int* 2015:527926.
- Voga M, Adamic N, Vengust M, Majdic G (2020) Stem cells in Veterinary Medicine - Current state and treatment options. *Frontiers in Veterinary Science* 7:278.
- Vrecenak JD, Pearson EG, Santore MT, Todorow CA, Li H, Radu A, Bhatti T, Peranteau WH, Johnson MP, Flake AW (2014) Stable long-term mixed chimerism achieved in a canine model of allogeneic in utero hematopoietic cell transplantation. *Blood* 124(12):1987-1995.
- Wang Y, Bi X, Zhou H, Deng Y, Sun J, Xiao C, Fan X (2014) Repair of orbital bone defects in canines using grafts of enriched autologous bone marrow stromal cells. *J Transl Med* 12(1):123.
- Wang F, Zhou Y, Zhou J, Xu M, Zheng W, Huang W, Zhou W, Shen Y, Zhao K, Wu Y, Zou D (2018) Comparison of Intraoral Bone Regeneration with Iliac and Alveolar BMSCs. *J Dent Res* 97(11):1229-1235.
- Wang X, Zhen L, Miao H, Sun Q, Yang Y, Que B, Lopes Lao EP, Wu X, Ren H, Shi S, Lau WB, Ma X, Ma C, Nie S (2015) Concomitant Retrograde Coronary Venous Infusion of Basic Fibroblast Growth Factor Enhances Engraftment and Differentiation of Bone Marrow Mesenchymal Stem Cells for Cardiac Repair after Myocardial Infarction. *Theranostics* 5(9):995-1006.
- Wayne JS, McDowell CL, Shields KJ, Tuan RS (2005) In vivo response of poly(lactide acid-alginate) scaffolds and bone marrow-derived cells for cartilage tissue engineering. *Tissue Eng* 11(5-6):953-963.
- Webb TL, Webb CB (2015) Stem cell therapy in cats with chronic enteropathy: a proof-of-concept study. *J Feline Med Surg* 17(10):901-908.
- Welt FG, Gallegos R, Connell J, Kajstura J, D'Amario D, Kwong RY, Coelho-Filho O, Shah R, Mitchell R, Leri A, Foley L, Anversa P, Pfeffer MA (2013) Effect of cardiac stem cells on left-ventricular remodeling in a canine model of chronic myocardial infarction. *Circ Heart Fail* 6(1):99-106.
- Willcox JL, Pruitt A, Suter SE (2012) Autologous peripheral blood hematopoietic cell transplantation in dogs with B-cell lymphoma. *J Vet Intern Med* 26(5):1155-1563.
- Wood JA, Chung DJ, Park SA, Zwingenberger AL, Reilly CM, Ly I, Walker NJ, Vernau W, Hayashi K, Wisner ER, Cannon MS, Kass PH, Cherry SR, Borjesson DL, Russell P, Murphy CJ (2012) Periocular and intra-articular injection of canine adipose-derived mesenchymal stem cells: an in vivo imaging and migration study. *J Ocul Pharmacol Ther* 28(3):307-317.
- Wu GH, Shi HJ, Che MT, Huang MY, Wei QS, Feng B, Ma YH, Wang LJ, Jiang B, Wang YQ, Han I, Ling EA, Zeng X, Zeng YS (2018) Recovery of paralyzed limb motor function in canine with complete spinal cord injury following implantation of MSC-derived neural network tissue. *Biomaterials* 181:15-34.
- Xu W, Chen J, Liu X, Li H, Qi X, Guo X (2016) Autologous bone marrow stromal cell transplantation as a treatment for acute radiation enteritis induced by a moderate dose of radiation in dogs. *Transl Res* 171:38-51.
- Yang Y, Bi X, Zhou H, Deng Y, Sun J, Xiao C, Gu P, Fan X (2014) Repair of orbital bone defects in canines using grafts of enriched autologous bone marrow stromal cells. *J Transl Med* 12(1):123.
- Yoon PD, Kao RL, Magovern GJ (1995) Myocardial regeneration. Transplanting satellite cells into damaged myocardium. *Tex Heart Inst J* 22(2):119-125.
- Yuan J, Cui L, Zhang WJ, Liu W, Cao Y (2007) Repair of canine mandibular bone defects with bone marrow stromal cells and porous beta-tricalcium phosphate. *Biomaterials* 28(6):1005-1013.
- Yuanzheng C, Yan G, Ting L, Yanjie F, Peng W, Nan B (2015) Enhancement of the repair of dog alveolar cleft by an autologous iliac bone, bone marrow-derived mesenchymal stem cell, and platelet-rich fibrin mixture. *Plast Reconstr Surg* 135(5):1405-1412.
- Yun S, Ku SK, Kwon YS (2016) Adipose-derived mesenchymal stem cells and platelet-rich plasma synergistically ameliorate the surgical-induced osteoarthritis in Beagle dogs. *J Orthop Surg Res* 11:9.
- Zang S, Jin L, Kang S, Hu X, Wang M, Wang J, Chen B, Peng B, Wang Q (2016) Periodontal Wound Healing by Transplantation of Jaw Bone Marrow-Derived Mesenchymal Stem Cells in Chitosan/Anorganic Bovine Bone Carrier Into One-Wall Infrabony Defects in Beagles. *J*

- Periodontol 87(8):971-981.
- Zeira O, Asiag N, Aralla M, Ghezzi E, Pettinari L, Martinelli L, Zahirpour D, Dumas MP, Lupi D, Scaccia S, Konar M, Cantile C (2015) Adult autologous mesenchymal stem cells for the treatment of suspected non-infectious inflammatory diseases of the canine central nervous system: safety, feasibility and preliminary clinical findings. *J Neuroinflammation* 12: 181.
- Zhang J, Huang X, Wang H, Liu X, Zhang T, Wang Y, Hu D (2015) The challenges and promises of allogeneic mesenchymal stem cells for use as a cell-based therapy. *Stem Cell Res Ther* 6: 234.
- Zhang Z, Maiman DJ, Kurpad SN, Crowe MJ, Alexanian AR (2011) Feline bone marrow-derived mesenchymal stem cells express several pluripotent and neural markers and easily turn into neural-like cells by manipulation with chromatin modifying agents and neural inducing factors. *Cell Reprogram* 13(5):385-390.
- Zorn J, Schwamberger S, Panzer W, Adler H, Kolb HJ (2011) Transplantation of CD6-depleted peripheral blood stem cells after DLA-haploidentical bone marrow transplantation contributes to engraftment and tolerance in a preclinical model of stem cell transplantation. *Vet Immunol Immunopathol* 144(1-2):27-35.
- Zubin E, Conti V, Leonardi F, Zanichelli S, Ramoni R, Grolli S (2015) Regenerative therapy for the management of a large skin wound in a dog. *Clin Case Rep* 3(7): 598-603.