



Journal of the Hellenic Veterinary Medical Society

Vol 73, No 2 (2022)



To cite this article:

Bayraktar, M., Hussein, M., & Hassan, H. (2022). Determination genetic variation and phylogenetic analysis in Echinococcus granulosus isolated from Iraqi sheep. *Journal of the Hellenic Veterinary Medical Society*, *73*(2), 4245–4252. https://doi.org/10.12681/jhvms.28066



Determination genetic variation and phylogenetic analysis in *Echinococcus* granulosus isolated from Iraqi sheep

M. Bayraktar^{1*}, M. Hussein², H. Hassan²

^{1*}Department of Animal Science, College of Agriculture, University of Çukurova, Adana, Turkey

²Department of Biyology, College of Science, University of Kirkuk, Kirkuk, Iraq

ABSTRACT : *Echinococcus granulosus* causes cystic echinococcosis in intermediate hosts, such as sheep, cattle, goats, camels, and horses. Cystic echinococcosis is an essential parasitic disease that affects humans and animals and constitutes a significant public health and economic problem worldwide. This study aimed to determine the molecular characterization and phylogenetic analysis of *Echinococcus granulosus* in sheep's livers and lungs. The genetic variation was performed by directly sequencing the mitochondrial DNA (mtDNA) genes coding of the NADH dehydrogenase subunit 1 (ND1). The evidence from molecular studies and DNA sequencing revealed that the isolates of *Echinococcus granulosus* in Kirkuk city are related to the sheep strain G1 genotype. The alignment analysis showed that there is a concordance between isolates with each other by 99%. When comparing the results with Genbank, it indicated the occurrence of genetic variations between nucleotides represented by eleven codons, seven of which represented transversion mutations, and four were transition mutations.

Keywords: Echinococcus granulosus, Cystic Echinococcosis, Sheep, ND1 gene, Sequencing

Corresponding Author: Mervan Bayraktar E-mail address: mbayraktar@cu.edu.tr

Date of initial submission: 25-09-2021 Date of acceptance: 24-01-2022

INTRODUCTION

Chinococcus granulosus (E. granulosus) is one L of the smallest parasitic tapeworms that dwell as adult worms in the small intestine of the definitive hosts (dogs, wolves, hyenas, leopards, and lions), and as hydatid cysts in the intermediate hosts (sheep, goats, cattle, camels, buffalos, horses, donkeys, pigs, rabbits and humans) (Abdulla et al., 2020; Hammad et al., 2018; Hasan et al., 2016). E.granulosus causes a zoonotic parasitic disease called cystic echinococcosis (CE). CE also called echinococcosis, hydatidosis and unilocular hydatidosis. CE is a severe epidemic disease that affects humans and causes significant economic losses to livestock. This disease spreads almost all over the world. Still, it is more common in rural areas, especially in grazing areas. The reason is due to the presence of the intermediate host (cattle and sheep) and the definitive host (dogs), which helps to complete the life cycle of this parasite (Alvi et al., 2020; Kadhimand & Al-Mayali 2020; Mahdi et al., 2020). CE grows and develops slowly, as the disease is characterized by the absence of symptoms in the early stages of infection. Symptoms depend on the affected organ, the liver being the most affected organ, with a rate of about 60-70%, followed by the lungs 20-22%, spleen, heart, muscles, eyes, thyroid, kidneys, brain, and bones (Saadi 2020; Sharma et al., 2013). Some studies have recorded more than 50 cases per 100.000 people per year in endemic areas. The disease also kills about one million people annually worldwide and causes a loss of about three

billion US\$ (Fallahizadeh et al., 2019; Alsaady and Al-Quzweeni 2019). Based on nucleotide sequences analysis of two mitochondrial regions, cytochrome c oxidase subunit 1 (CO1) and NADH dehydrogenase subunit 1 (ND1) and one ribosomal region, internal transcribed spencer 1 (ITS1), ten distinct genotypes or strain (G1-G10) have been described for E.granulosus. This includes sheep strain G1, Tasmanian sheep strain G2, buffalo strain G3, horse strain G4, cattle strain G5, camel strain G6, pig strain G7, cervid strain G8, the human polish strain G9 and the fennoscandian cervid strain G10 (Lavikainen et al., 2003; Piccoli et al., 2013; Yan et al., 2018). Iraq is an endemic country for CE due to the spread of stray dogs infected with parasites. Although there are methods available to control the disease, it still remains a significant health problem. This investigation aimed to characterize the genetic variation and phylogenetic analysis of E.granulosus in terms of ND1 gene isolated from sheep liver.

MATERIALS AND METHODS

Sample collection

Eight samples of hydatid cysts isolated from sheep livers and lungs were used in this study. The samples were obtained from the Kirkuk slaughterhouse from 10/29/2019 to 2/16/2020 under the supervision of veterinarians (Figure 2). The ages of the animals ranged from 7 months to 1 year. The samples were transported by icebox to preserve the vitality of the protoscoleces to the laboratory (Figure 1).



Figure 1: Hydatid cysts obtained from sheep's livers and lungs



Figure 2: The geographical area from which the samples were collected

Isolation of protoscoleces

Protoscoleces were isolated according to the Smyth (1985) method. Hydatid cysts were cleaned with 70% ethyl alcohol and washed with phosphate buffer saline (PBS), and then hydatid fluid was with-drawn from the cyst. Germinal Layer was extracted and washed with PBS. The hydatid fluid was with-drawn and centrifuged at 3000 rpm for 15 minutes. The filtrate was removed, the precipitate was taken, and then washed with PBS. The samples were stored at -20 °C.

DNA extraction from protoscoleces

DNA extraction was done for each sample using Genomic DNA Mini Kit (Wizard, Korea) according to the manufacturer's instructions. The purity and concentration of extracted DNA samples were analyzed by Nanodrop.

PCR assay

The PCR was performed in a total of 25µl, which contained 12.5µl Go Tag Green Master Mix (Promega, USA), 1µl of 10 pmol/µl of each primer, 2µl of the genomic DNA, and 8.5µl nuclease-free water. PCR conditions were as follows: initial denaturation at 95 °C for 5 min; 30 cycles of denaturation at 95 °C for 30 sec, annealing at 58 °C for 30 sec and elongation at 72 °C for 30 sec and a final extension at 72 °C for 5 min. To amplified the ND1 gene (800 bp) was used forward (5'-GTTTTTGGGTTAGTCTCTGG-3') and reverse (5'-ATCATAACGAACACGTGG-3') primers (Sánchez *et al.*, 2012). PCR products were separated at 2% agarose gel in 80 V for 60 min. The gel was stained by ethidium bromide and results were checked under ultraviolet light.

Sequences and phylogenetic analysis

The PCR products of the ND1 gene for eight samples were sequenced. The DNA sequencing was performed by Macrogen Corporation (Korea) using the 3730 XL DNA Analyzers (Applied Biosystems, Foster City, California, USA). The reference sequences of the ND1 gene of *E.granulosus* were obtained from GenBank (https://www.ncbi.nlm.nih.gov/). Reference and isolate sequences were used in multiple alignments, and phylogenetic distances were calculated using BioEdit software. Phylogenetic trees were constructed using the neighbor-joining methods in MEGA version 6.0. All sequences were uploaded to GenBank and assigned GenBank number MW980133, MW980134, MW980135, MW980136, MW980137, MW980138, MW980139 and MW980140.

RESULTS

800 bp of PCR product was amplified of the ND1 gene (Figure 3). All eight samples were genotype G1 strain (100%). Alignment analysis demonstrates the similarity of 99% (for 6 samples) and 100% (for 2 samples). When comparing the results with Genbank, it shows the occurrence of mutations between nucleotides represented by eleven codons, 7 of which are transversion mutations and 4 transition mutations. In sample A, four mutations were found, three of which were transversion mutations and one transition mutation. Mutations were; $(T \setminus G \rightarrow TGT \setminus GGT)$ at site [572], $(T\setminus G \rightarrow TTG\setminus TGG)$ at site [574], $(G\setminus A \rightarrow AGA\setminus AAA)$ at site [617] and $(T/G \rightarrow AGT \setminus AGG)$ at site [651]. The predicted effect was missense with anidentical ratio of 99% (Table 1). No mutation was found in sample B; the identical percentage was 100%. Sample C showed three transversion mutations in the site [607] (G\C \rightarrow -GAG\CAG), (G\T \rightarrow AGT\ATT) in site [650] and (T\ $A \rightarrow TTT \setminus ATT$) in site [673]. The mutation was an effective missense, and the identical rate between them was 99%. The percentage of congruence was 100% in sample D and no mutation was found. In sample E, one transversion mutation $G \to CAG \to CAG \to CAT$) was identified in the site [150]. The mutation was missense, with anidentical percentage of 99%. About samples (F, G, H), transition mutation $(T \land C \rightarrow TAT \land$

TAC) was detected at sites [285]. The mutation was silent and the identical was 99%. The phylogenetic tree results showed identical between samples (G, H, F) as 100%, also, the identical between samples (D, B) were 100%. The identical was between samples (A, C, E) as %99, whereas the identical between samples (D, B) and samples (H, F, G) were 99% (Figure 4, Figure 6).Based on a comparison between the sequences of ND1 gene for Iraqi E.granulosus isolates with other strains from different countries were 99% match with the common sheep strain (G1). It appeared that the highest percentage of identical (99%) was with China, Nigeria, Algeria, Greece, Tunisia, India, Mexico, Kazakhstan, Iran, Mongolia, Argentina, Brazil, Chile, Turkey, and Pakistan. While the countries samples were identical to each other by 100% as well (Figure 5).







Figure 4: Neighbor-joining tree showing the identical ratio between the species of isolates







Figure 6: Multiple sequences alignment of ND1 gene

Table 1: Genetic variations of E.granulosus according to ND1 gene							
Sample Number	Type of substitution	Location	Nucleotie	Nucleotide change	Predicted effect	Sequence ID with compare	Identities
А	Transversion	572	T\G	TGT\GGT	Missense	ID: <u>MN269986.1</u>	99%
	Transversion	574	T\G	TTG\TGG	Missense		
	Transition	617	G\A	AGA\AAA	Missense		
	Transversion	651	T\G	AGT\AGG	Missense		
В						ID: <u>MN269986.1</u>	100%
С	Transversion	607	G\C	GAG\CAG	Missense		
	Transversion	650	$G \setminus T$	AGT\ATT	Missense	ID: <u>MN269986.1</u>	99%
	Transversion	673	T\A	TTT\ATT	Missense		
D						ID: <u>MN269986.1</u>	100%
Е	Transversion	150	G\T	CAG\CAT	Missense	ID: <u>MN269986.1</u>	99%
F	Transition	285	T\C	TAT\TAC	Silent	ID: <u>MN269986.1</u>	99%
G	Transition	285	Т\С	TAT\TAC	Silent	ID: <u>MN269986.1</u>	99%
Н	Transition	285	T\C	TAT\TAC	Silent	ID: <u>MN269986.1</u>	99%

J HELLENIC VET MED SOC 2022, 73(2) ПЕКЕ 2022, 73(2)

DISCUSSION

Although cystic hydatidosis gives rise to significant and critical public health and economic problems in Africa, Asia, and the Middle East, current efforts to control the disease are inadequate (Wen et al., 2019). Because of the wide variety of cystic larval stages. Moreover, the control of hydatidosis is usually hampered due to ignorance of the true prevalence of the disease and the invalidity they cause (Saadi 2020). In addition, accurate diagnosis of hydatid cyst disease is often difficult because some of the esoteric diseases resemble those found in patients with hydatidosis (Kern et al., 2017). Therefore, there has been a great need for a more reliable laboratory test to diagnose hydatidosis in this area. The PCR amplification and DNA sequencing of specific regions of the ND1 gene have been shown to provide reliable alternatives to more traditional methods for the specific identification of Echinococcus strains from different intermediate hosts (Nakao et al., 2013). Based on the results above, it can be suggested that G1 or sheep genotype is the most important variant of *E.granulosus*. This finding should be considered during the planning and implementation of the *E.granulosus* control program region. The results demonstrated in this investigation agree with Iraqi isolates. Hamoo and Abdulraheem (2019) showed that the local isolate is G1 genotype (sheep strain), and the phylogenetic results were compatible with North Africa and the Middle East isolates as 95% and 96%.Fadhil & A'aiz (2016) reported three strains of the *E.granulosus* parasite, sheep strain (G1) 40%, buffalo strain (G3) 48% and camel strain (G6) 12%. Whereas, sheep were G1 (80%) and G3 (20%). Mustafa et al., (2019), AL-Asadi et al., (2021), Muhaidi et al., (2017), Muhaidi et al., (2018) identified E.granulosus in terms of ND1 gene as G1 genotype sheep strain (100%). The Iraqi isolates are consistent with the isolates of other countries, such as, Turkey (Barazesh et al., 2019), Iran (Barazesh et al. 2019;

Hajialilo et al., 2012; Pezeshki et al., 2013; Sharbatkhori et al., 2016), Sudan (Ahmed et al., 2018), Egypt (Amer et al., 2015), Uzbekistan (Kim et al., 2020), Peru (Sánchez et al. 2012; Sánchez et al., 2010) and China (Ohiolei et al., 2019). This may be due to the sensitivity of sheep to strain G1 E.granulosus and the short life span of parasite depending on sheep slaughtering age compared to other animals such as sheep, goats and cattle (Larrieu et al., 2019). In addition, an essential factor that plays a key role in the spread of disease is the fertility rate in the protoscoleces in livestock farms. It was observed that the fertility rate in the protoscoleces observed in the sheep is more significant than that observed in the cystic sacs where cattle and goats live (Mokhtaria et al., 2013). The phylogenetic tree that has been constructed in this study to confirm the relationship between Iraqi E.granulosus strain G1 and the other global isolates can be predicted. Therefore, Echinococcus that infect humans in Kirkuk, Iraq is a G1 sheep strain. The ND1 gene could be considered a powerful tool for accurate molecular identification and phylogenetic differentiation of this parasite.

CONCLUSIONS

E.granulosus isolates taken from sheep slaughtered in Kirkuk were identified as *E.granulosus* strain G1. These results contribute to the understanding of the epidemiology of *E.granulosus* in Iraq. The epidemiology of *E. granulosus* must be clarified in order to implement successful control programs and to help improve public health. For this reason, it is necessary to collect more information on this topic by genotyping the parasite and larval forms detected in different intermediate and definitive host populations in different geographical regions.

CONFLICT OF INTEREST

The authors declared no conflict of interest.

REFERENCES

- Abdulla R G, Mageed S N, Obed C E, Jumaa J A (2020) Molecular characterization of fertile hydatid cysts from the liver of the sheep and cows and associated environmental influence factors. Iraqi Journal of Veterinary Sciences 34:321-27.
- Hammad S J, Cavallero S, Milardi G L, Gabrielli S, Al-Nasiri F S (2018) Molecular genotyping of Echinococcus granulosus in the North of Iraq. Veterinary parasitology 15:82-87.
- Hasan H F, Fadhil M H, Fadhil Z H (2016) Molecular characterization of Echinococcus granulosus isolated from human and domestic animals in Kirkuk, Iraq. Animal Research International 13:2544-47.
- Alvi M A, Ohiolei J A, Saqib M, Li L, Tayyab M H, Alvi A A, Wu Y-T, Fu B-Q, Yan H-B, Jia W-Z (2020) Echinococcus granulosus (sensu stricto) (G1, G3) and E. ortleppi (G5) in Pakistan: phylogeny, genetic diversity and population structural analysis based on mitochondrial DNA. Parasites & vectors 13:1-10.
- Kadhimand H A A, Al-Mayali H M H (2020) Molecular characterizations of Echinococcus granulosus isolated from human and sheep in Euphrates region of Iraq. International Journal of Pharmaceutical Research 1:2296-303.
- Mahdi Z M S, Al-Hamairy A K, Al-Rubaiey H M (2020) Genotyping of Echinococcusgranulosus Isolates from Human, Sheep and Cattles Hydatid Cysts in Some Central Euphrates Provinces, Iraq. Medico Legal Update 20:570-75.
- Saadi A. 2020. Contribution to the evaluation of the control program of cystic echinococcosis in Morocco. University of Liège, Liege, Belgium
- Sharma M, Sehgal R, Fomda B A, Malhotra A, Malla N (2013) Molecular characterization of Echinococcus granulosus cysts in north Indian patients: identification of G1, G3, G5 and G6 genotypes. PLoS neglected tropical diseases 7:1-6.
- Fallahizadeh S, Arjmand R, Jelowdar A, Rafiei A, Kazemi F (2019) Determination of Echinococcus granulosus genotypes in livestock slaughtered in Shush County, Southwest Iran using PCR-RFLP. Helminthologia 56:196-201.
- Alsaady H A M, Al-Quzweeni H A N (2019) Molecular study of Echinococcus Granulosus in Misan Province, south of Iraq. Indian Journal of Public Health 10:1047.
- Lavikainen A, Lehtinen M, Meri T, Hirvelä-Koski V, Meri S (2003) Molecular genetic characterization of the Fennoscandian cervid strain, a new genotypic group (G10) of Echinococcus granulosus. Parasitology 127:207-15.
- Piccoli L, Bazzocchi C, Brunetti E, Mihailescu P, Bandi C, Mastalier B, Cordos I, Beuran M, Popa L, Meroni V (2013) Molecular characterization of Echinococcus granulosus in south-eastern Romania: evidence of G1-G3 and G6-G10 complexes in humans. Clinical Microbiology and Infection 19:578-82.
- Yan B, Liu X, Wu J, Zhao S, Yuan W, Wang B, Wureli H, Tu C, Chen C (2018) Genetic diversity of Echinococcus granulosus genotype G1 in Xinjiang, Northwest of China. The Korean journal of parasitology 56:391-96.

Smyth J (1985) In vitro culture of Echinococcus spp. Proceedings of the

13 th Int. Congr. Hydatidology. Madrid 84-9.

- Sánchez E, Cáceres O, Náquira C, Miranda E, Samudio F, Fernandes O (2012) Echinococcus granulosus genotypes circulating in alpacas (Lama pacos) and pigs (Sus scrofa) from an endemic region in Peru. Memórias do Instituto Oswaldo Cruz 107:275-78.
- Wen H, Vuitton L, Tuxun T, Li J, Vuitton D A, Zhang W, Mcmanus D P (2019) Echinococcosis: advances in the 21st century. Clinical microbiology reviews 32:1-39.
- Kern P, Da Silva A M, Akhan O, Müllhaupt B, Vizcaychipi K, Budke C, Vuitton D (2017) The echinococcoses: diagnosis, clinical management and burden of disease. Advances in parasitology 96:259-369.
- Nakao M, Lavikainen A, Iwaki T, Haukisalmi V, Konyaev S, Oku Y, Okamoto M, Ito A (2013) Molecular phylogeny of the genus Taenia (Cestoda: Taeniidae): proposals for the resurrection of Hydatigera Lamarck, 1816 and the creation of a new genus Versteria. International Journal for Parasitology 43:427-37.
- Hamoo R, Abdulraheem S (2019) Nad1 gene analysis of Echinococcus granulosus from sheep in Aqrah city, Iraq. Iraqi Journal of Veterinary Sciences 33:341-45.
- Fadhil S, A'aiz N (2016) Genotyping of cystic echinococcosis isolates from clinical samples of human and domestic animals. Iraqi Journal of Veterinary Sciences 30:33-39.
- Mustafa J Y, Al-Azizz S A, Lazim A R (2019) Molecular study of hydatid cysts in human, sheep and donkeys at basra city, southern iraq. Biochemical and Cellular Archives 19:4155-61.
- Al-Asadi S A M, Hansh W J, Awad A-H H (2021) Employing NADH Dehydrogenase Subunit 1 in the Determination of Echinococcus granulosus Strain in Sheep, Cattle and Human in Thi-Qar Province, Iraq. Baghdad Science Journal 18:0238-38.
- Muhaidi M, Maysam A, Mohammed D (2017) Determination the causative strain for hydatid cyst in Iraqi cattle by using ND1 gene. Iraqi Journal of Veterinary Medicine 41:11-16.
- Muhaidi M, Ahmed M, Aziz L (2018) Identification of Echinococcus Granulosus Genotype in Iraqi's Sheep by Using Nd1 Gene. Journal of Global Pharma Technology 10:261-66.
- Barazesh A, Sarkari B, Sarısu G, Hami M, Mikaeili F, Aydın A, Ekici A, Ebrahimi S (2019) Comparative genotyping of Echinococcus granulosus infecting livestock in Turkey and Iran. Turkish Journal of Parasitology 43:123-29.
- Hajialilo E, Harandi M F, Sharbatkhori M, Mirhendi H, Rostami S (2012) Genetic characterization of Echinococcus granulosus in camels, cattle and sheep from the south-east of Iran indicates the presence of the G3 genotype. Journal of helminthology 86:263-70.
- Pezeshki A, Akhlaghi L, Sharbatkhori M, Razmjou E, Oormazdi H, Mohebali M, Meamar A (2013) Genotyping of Echinococcus granulosus from domestic animals and humans from Ardabil Province, northwest Iran. Journal of helminthology 87:387-91.
- Sharbatkhori M, Tanzifi A, Rostami S, Rostami M, Harandi M F (2016) Echinococcus granulosus sensu lato genotypes in domestic livestock and humans in Golestan Province, Iran. Revista do Instituto de Medicina Tropical de São Paulo 58:1-7.

- Ahmed M E, Salim B, Grobusch M P, Aradaib I E (2018) First molecular characterization of Echinococcus granulosus (sensu stricto) genotype 1 among cattle in Sudan. BMC veterinary research 14:1-5.
- Amer S, Helal I B, Kamau E, Feng Y, Xiao L (2015) Molecular characterization of Echinococcus granulosus sensu lato from farm animals in Egypt. PLoS One 10:1-12.
- Kim H-J, Yong T-S, Shin M H, Lee K-J, Park G-M, Suvonkulov U, Kovalenko D, Yu H S (2020) Phylogenetic Characteristics of Echinococcus granulosus sensu lato in Uzbekistan. The Korean journal of parasitology 58:205-10.
- Sánchez E, Cáceres O, Náquira C, Garcia D, Patiño G, Silvia H, Volotão A C, Fernandes O (2010) Molecular characterization of Echinococcus granulosusfrom Peru by sequencing of the mitochondrial cyto-

chrome C oxidase subunit 1 gene. Memórias do Instituto Oswaldo Cruz 105:806-10.

- Ohiolei J A, Xia C-Y, Li L, Liu J-Z, Tang W-Q, Wu Y-T, Zhu G-Q, Shi B, Fu B-Q, Yin H (2019) Genetic variation of Echinococcus spp. in yaks and sheep in the Tibet Autonomous Region of China based on mitochondrial DNA. Parasites & vectors 12:1-10.
- Larrieu E, Gavidia C M, Lightowlers M W (2019) Control of cystic echinococcosis: background and prospects. Zoonoses and public health 66:889-99.
- Mokhtaria K, Fatima B-K, Abboud B, Ammar S S M (2013) Cystic echinococcosis in small ruminants in Tiaret (Algeria). Global Veterinaria 11:753-58.