



# Journal of the Hellenic Veterinary Medical Society

Vol 73, No 4 (2022)



# To cite this article:

Olgun, O., Gül, E., & Yildiz, A. (2023). Influence of Organic Iron Enriched Diets on Performance, Egg Quality, Blood Haematological and Biochemical Constituents in Quails. *Journal of the Hellenic Veterinary Medical Society*, *73*(4), 4899–4904. https://doi.org/10.12681/jhvms.28232

# Influence of Organic Iron Enriched Diets on Performance, Egg Quality, Blood Haematological and Biochemical Constituents in Quails

# O. Olgun<sup>\*</sup>, E.T. Gül<sup>®</sup>, A.Ö. Yıldız<sup>®</sup>

Department of Animal Science, Faculty of Agriculture, University of Selcuk, Selcuklu, Konya, Turkey

**ABSTRACT**: The current research was carried out to investigate the effect of diets enriched with different levels of organic iron on performance, egg quality, blood haematological parameters, and serum biochemical constituents in laying Japanese quails. In the study, 120 female quails at 20 weeks of age were allocated to 5 treatment groups with 6 replicates of 4 quails ( $249\pm11g$ ) in each. Experimental diets were formed by adding 0, 50, 100, 150, and 200 mg/kg of iron (iron-glycine) to the diet containing 96 mg/kg of iron. Performance parameters of quails were not affected by the supplementation of organic iron to diet (P>0.05). The supplementation of organic iron affected as quadratic only the eggshell breaking strength among the egg external quality parameters, and it was maximum at 150 mg/kg organic iron addition (P<0.01). Egg yolk a\* value, which is one of the egg internal quality parameters, was linearly affected by organic iron addition, but other parameters were not affected (P<0.01). The administration of organic iron to the diet linearly increased the serum iron level(P<0.01), the serum phosphorus level was affected quadratically(P<0.05) and reached a maximum at 150 mg/kg. Among the blood haematology parameters, white blood cell and monocyte counts linearly increased with the addition of organic iron to the diet (P<0.01), but other parameters were not affected. According to the results obtained from current research, it can be said that enrichment of laying quail diets with organic iron up to 150 mg/kg level improved eggshell breaking strength without affecting performance, however, it negatively affected blood parameters by increasing white blood cell and monocyte without changing blood red cell parameters.

Keywords: Egg quality, haematology, organic iron, performance, quail, serum.

Corresponding Author: Corresponding Author: Osman Olgun. Department of Animal Science, Faculty of Agriculture, University of Selcuk, Selcuklu, 42130 Konya, Turkey. E-mail address: oolgun@selcuk.edu.tr

Date of initial submission: 13-10-2021 Date of acceptance: 18-11-2021

## **INTRODUCTION**

ron is an essential element of vital importance in Lall animals. The reason for that is of iron is found in the haemoglobin structure in the organism, it carries out many functions such as oxygen and electron transport (Leeson and Summers, 2001), enzymatic activities (Chen et al., 2008), acts as prooxidant (Rajpathaket al., 2009), energy production (Mackenzie et al., 2008), and collagen synthesis (McDowell, 2003).

Since vegetable raw materials are insufficient in terms of iron due to factors such as soil, climate and phytate content of the plant (Yu et al., 2000; Gupta et al., 2008), iron deficiency occurs in poultry fed with these materials (Gupta et al., 2008). In these mentioned feed sources, both the iron level and its bio-availability are low, as well as high calcium, phytic acid, and iron levels in the diet affects its availability (Vieira, 2008; Gibson et al., 2010; Xie et al., 2019a,b).

Accordingly, this element is supplemented in the diet with premix in order to ensure that the iron needs of the poultry are exactly covered. Commonly, iron sulphate, which is a cost-efficient and low bioavailability source (form), is generally used in premixes (Ma et al., 2014; Xie et al., 2019a).For that purpose, addition of amino acid chelates (such as iron glycine) to diet instead of inorganic forms or enrichment diets in terms of iron become importance for the sustainability of the performance. Xieet al., (2019a) stated that the lower level (20 mg/kg) of organic source (iron glycine) can be used instead of iron sulphate for the permanence of egg quality in laying hens. Results reported by Sarlak et al. (2021) were similar. In another study (Paik et al., 2009), it was stated that there was an improvement in egg weight and Haugh unit in laying hens fed with diets enriched with organic iron source (iron proteinate). Compared to other essential trace

elements, the number of studies on iron in poultry is less. Therefore, the aim of this research was to examine the effects of practical diets enriched with organic iron on performance, internal and external qualities of egg, blood haematology and serum biochemical constituents in layer quails.

## **MATERIALS AND METHODS**

#### **Ethical Approval**

Criteria specified by the National Institute of Health Guide for the Care and Use of Laboratory Animals were followed during the study period.

#### **Animals and Feed Materials**

In the present experiment, 120 female Japanese quails at the age of 20 weeks were fed ad-libitum for 70 days with 5 treatment diets that included 0, 50, 100, 150 or 200 mg/kg organic iron (iron-glycine) to the basal diet based on corn-soybean meal (Table 1). The study was conducted in 5 experimental groups consisting of 6 replicates, each containing 4 female quails. A 16-hour lighting program was applied to quails.

### Method

#### **Determination of performance parameters**

At the beginning of the experiment, the quails were placed in the cages by group weighing and at the end of the experiment, the groups were weighed again, to determine changes in body weight. Experimental feeds were given to the treatment subgroups by weighing and feed intake (FI) were assessed as g/day/quail. Eggs collected at the same time of each day were recorded and egg production (EP) was calculated as %. Egg weight (EW), on the other hand, was determined as g by weighing all eggs collected in the last three days of the experiment. From these data, also, egg mass (EM)

fable 1. Basal diet and its calculated nutrient contents								
Ingredients	g/kg	Nutrient contents	g/kg					
Corn	544.0	Metabolizable energy, kcal ME/kg	2900					
Soybean meal (46%)	343.7	Crude protein	200					
Soybean oil	36.7	Calcium	25.0					
Limestone	56.0	Available phosphorus	3.5					
Dicalcium phosphate	11.5	Lysine	10.9					
Salt	3.5	Methionine	4.5					
Premix <sup>1</sup>	2.5	Cystine	3.7					
DL methionine	2.1	Methionine+cystine	8.2					
Total	1000.0	Iron	0.96					

<sup>1</sup>Premix is supplied that per kg of diet; Manganese: 80 mg, Iron: 60 mg, Copper: 5 mg, Iodine: 1 mg, Selenium: 0.15 mg, Vitamin A: 8.800 IU, Vitamin D<sub>3</sub>: 2.200 IU, Vitamin E: 11 mg, Nicotine acid: 44 mg, Cal-D-Pan: 8.8 mg, Riboflavin: 4.4 mg, Thiamine: 2.5 mg, Vitamin B<sub>12</sub>: 6.6 mg, Folic acid: 1 mg, Biotin: 0.11 mg, Choline: 220 mg.

> J HELLENIC VET MED SOC 2022, 73 (4) ПЕКЕ 2022, 73 (4)

was calculated as g/day/quail with the formula (*EP x EW*) / 100 and feed conversion ratio (FCR) was determined as g feed/g egg with FI / EM formula.

#### Determination of egg quality parameters

During the experiment, broken, cracked and damaged eggs were recorded and calculated as % of the number of eggs. Egg internal and external quality parameters were determined from all eggs collected in the last three days of the experiment. Eggshell breaking strength was assessed by applying supported-systematic pressure to the blunt of the eggs (Egg Force Reader, Orka Food Technology, Israel).Immediately after the determination of the eggshell breaking strength, the eggs were broken on a clean, glass surface, and after the residues in the eggshell were cleaned, the shells were dried at room temperature for three days and weighed, and relative weights were calculated as a ratio (%) of the egg weight. Eggshell thickness was calculated by averaging the measurements obtained from three sections (equator, blunt, and pointed parts) of the eggshell using a micrometre (Mitutoyo, 0.01 mm, Japan). Eggs, which external quality characteristics were determined, were broken on a surface and their albumen and yolk heights were measured with a height gauge and their length and width were measured with the calliper. The parameters calculated from these data and the formulas used are follows: albumen index with the formula *albumen* height/ ((albumen width + albumen length) / 2)  $\times$ 100, yolk index with the (yolk height / yolk diameter)  $\times$  100, and finally Haugh unit with the 100 x log (albumen height +  $7.57-1.7 \times EW^{0.37}$ ). In addition, L\*, a\* and b\* a values of yolk were measured with a colorimeter (MinoltaChromaMeter CR 400 (MinoltaCo., Osaka, Japan) (Romero et al., 2002).

#### **Blood haematological analysis**

At the end of the experiment, blood were collected from the wing veins of 6 quails randomly selected from each treatment group (one quail from each subgroup) into heparinized tubes for blood haematological analysis. Blood haematological analyses were performed in an auto-analyser in a commercial laboratory according to Campbell (1988).

#### Serum biochemical analysis

At the end of the experiment (10<sup>th</sup>week), 3 ml of blood was taken from one random quail (30 in total) of similar body weight from each subgroup to determine serum parameters. Blood were centrifuged at 4000 rpm for 10 minutes and serum were extracted. The serums were stored at -20°C until analysis, and the concentrations of glucose, triglyceride, cholesterol, aspartate aminotransferase (AST), alanine aminotransferase (ALT), total protein, albumin, globulin, urea, creatinine, calcium, phosphorus and iron in the serum were determined (Choi et al., 2013) in an auto-analyser device using commercial kits in a commercial laboratory (DDS® SpectrophotometricKits, Diesis DiagnosticSystemsCo., Istanbul, Turkey).

#### **Statistical analysis**

Data were analysed in the SPSS 18.0 software package (SPSS Inc., Chicago, IL, USA) with model of one-way ANOVA, using the group mean as an experimental unit. A probability value of P<0.05 was considered statistically significant. Orthogonal polynomial contrasts were used to evaluate the significance of linear and quadratic models to determine the response of the dependent variable to an increasing organic iron level.

#### **RESULTS AND DISCUSSION**

#### **Performance Parameters**

Effects of organic iron supplemented to commercial quail diets at different levels (0, 50, 100, 150 or 200 mg/kg) on performance were demonstrated in Table 2. Administration of organic iron to laying quail diets did not significantly affect body weight change,

T.1.1. A T.A	· · · · · · · · · · · · · · · · · · ·	C	· · · · · · · · · · · · · · · · · · ·	
Table 7. Infilience of diefa	ry sunniementation of	t organic iron on the	nertormance i	harameters in laver dualis
i ubic 2. innuciec of ulcu	i j supprementation of	i of Same if on on the	per tor munee p	fur uniceers in injer quans

· · · · ·		0					<b>U</b>	
		Organic iron (mg/kg)					P-value	of contrast
Parameters	0	50	100	150	200	SEM*	Linear	Quadratic
Body weight change, g	-3.17	-11.92	-2.00	1.17	-5.52	2.056	0.188	0.051
Egg production, %	90.43	92.00	89.13	89.91	92.02	0.739	0.841	0.464
Egg weight, g	12.30	12.27	12.02	12.38	11.46	0.135	0.097	0.289
Egg mass, g/d/quail	11.12	12.29	10.71	11.12	10.54	0.151	0.226	0.697
Feed intake, g/d/quail	29.45	30.67	28.13	29.34	29.02	0.268	0.205	0.742
Feed conversion ratio, g feed/g	2.65	2.74	2.63	2.64	2.77	0.038	0.627	0.543
egg								

\*Standard error means

EP, EW, EM, FI and FCR (P>0.05). These results are agreement with the reports of Buckiuniene et al. (2016), Bai et al. (2018), Gou et al. (2020) who stated that the addition of iron to the diet did not affect performance in laying birds, but not Paik et al. (2009), Xie et al. (2019a), Sarlak et al. (2021) who clarified that it affect positively performance. Iron is a mineral involved in the protein and energy metabolism, and transport of oxygen in animals (Leeson and Summers, 2001; Mackenzie et al., 2008). With these properties, it is a mineral that can affect the productivity of poultry. However, in the current study, the reason why enriched the diet with organic iron did not affect performance could be because the iron level in the basal diet was sufficient for the performance of quails.

### Egg quality parameters

Effects of organic iron supplementation to commercial quail diets on the egg external quality were given in Table 3, and the effects on egg internal quality were shown in Table 4.

Damaged egg rate, relative eggshell weight, and eggshell thickness were not affected by the treatments(P>0.05). However, addition of iron-glycine to laying quail diets quadratically affected the eggshell breaking strength (P<0.01). Eggshell breaking strength improved with the supplementation of organic iron to the diet and reached maximum at 150 mg/kg, but it was minimum at the level of 200 mg/ kg addition. In a related study, Gou et al. (2020) explained that addition of iron to the diets linearly and quadratically affected eggshell breaking strength, and it was maximum at 58 mg/kg and it minimum at 100 mg/kg level in broiler breeders. Sarlak et al. (2021), on the other hand, stated that the addition of inorganic iron(iron-sulphate) at the level of 30 and 60 mg/ kg and organic iron (iron-glycine) at the level of 30, 60, and 120 mg/kg increased the eggshell breaking strength in laying hens. However, Paik et al. (2009), Seo et al. (2010), Taschetto et al. (2017), Xie et al. (2019b) noted that the supplementation of organic iron at the level of 0-200 mg/kg to the diet of laying hens did not affect the eggshell breaking strength.

Among the egg internal quality parameters, albumen index, yolk index, Haugh unit, and L\* and b\* values of yolk did not considerably affected by the administration of organic iron to the diet(P>0.05). Yolk a\* value was linearly reduced by treatments(P<0.01) and it was minimum at 200 mg/kg iron-glycine addition to the diet. Since iron is included in the structure of haemoglobin and myoglobin, which colouring red to meat, it is known that added iron to the diet increases a\* value (redness) of meat (Lin et al., 2020). In addition, it has been reported that the supplementation of iron to the diet increased (Sarlak et al., 2021) or did not affect (Paik et al., 2009; Seo et al., 2010; Gou et al., 2020; Xie et al., 2019b) the colour of egg yolk(Roche scale). However, in the literature, there aren't data examining the effect of dietary iron level (additional) on a\* value, which is a measure of greenery and redness of egg yolk. Therefore, it is obvious that the linear decrease in the egg yolk a\* value(increasing greenery) by the risen iron level in the diet should be investigated in more detail.

Table 3. Influence of dietary supplementation of organic iron on the egg external quality in layer quails									
	Organic iron (mg/kg)						<i>P</i> -value of contrast		
Parameters	0	50	100	150	200	SEM*	Linear	Quadratic	
Damaged egg rate, %	0.28	0.35	0.82	0.14	0.00	0.155	0.503	0.242	
Eggshell breaking strength, kg	1.39	1.49	1.55	1.61	1.36	0.034	0.796	0.006	
Relative eggshell weight, %	8.07	8.19	7.99	8.16	8.33	0.073	0.359	0.466	
Eggshell thickness, µm	217	218	214	221	221	1.9	0.466	0.534	

\*Standard error means

	Organic iron (mg/kg)						P-value	of contrast
Parameters	0	50	100	150	200	SEM*	Linear	Quadratic
Albumen index	3.38	3.16	3.63	3.02	2.97	0.093	0.128	0.333
Yolk index	50.87	53.24	53.31	53.10	52.80	0.503	0.311	0.201
Haugh unit	95.84	94.94	97.91	93.26	93.11	0.642	0.100	0.229
L* value of yolk	51.96	53.51	53.93	53.31	53.98	0.290	0.061	0.237
a* value of yolk	0.20	-0.28	-0.53	-0.20	-0.65	0.092	0.008	0.347
b* value of yolk	24.76	27.23	24.29	25.40	25.46	0.401	0.878	0.810

\*Standard error means

#### **Blood haematology**

Effects of organic iron supplemented to commercial quail diets at different levels (0, 50, 100, 150 or 200 mg/kg) on blood haematological parameters were shown in Table 5.Among the blood haematological parameters, white blood cell and monocyte counts were linearly affected by the administration of organic iron to the diet(P<0.01), this effect was not observed in other parameters(P>0.05). White blood cell concentration showed a linear increase with the supplementation of iron-glycine to the diet and it was maximum at the level of 150 mg/kg. Supplementation of organic iron did linearly affect the monocyte level of laying quails, and while it was minimum with added organic iron at 50 mg/kg, it reached maximum at 200 mg/ kg. Seo et al. (2010) indicated that the administration of 100 mg/kg iron proteinate to laving hen diets did not affect serum white blood cell and monocyte levels. There is evidence that iron inhibits intracellular monocyte accumulation (Li and Frei, 2006; Scott et al., 2021). Therefore, enrichment of the diet with iron could diminish the number of intracellular monocytes in quails and caused an increase in monocytes in the blood. In parallel, it is seen that the increase in monocyte, which is one of the white blood cells, also increases the counts of white blood cell.

#### Serum biochemical constituents

Supplementation of organic iron to the laying quail diets did not statistically affect serum biochemical constituents except for phosphorus and iron levels. Serum phosphorus level was quadratically affected by the treatments (P<0.05), and it was maximum

in the group fed with diet added at lowest level (50 mg/kg) of iron-glycine, while it was minimum in the group added at highest level (200 mg/kg) of iron-glycine. This result was not in harmony with the reports demonstrated by Xieet al. (2019a).

Serum iron level of laying quails linearly increased the addition of organic source of iron to the diet and it reached a maximum at 200 mg/kg level. Similarly, Xie et al. (2019a) notified that the serum iron level augmented with the administration of organic source of iron (iron-glycine) at the level of 60 and 80 mg/kg to the laying hen diets. In addition, Bai et al. (2018) stated that high iron supplementation provided an increase in serum iron level. Similar results were reported by Guoet al. (2020) and Sarlak et al. (2021).

It was expected that the serum iron level would increase with the increasing iron level in the diet. Surprisingly, however, despite an antagonistic relationship between iron and phosphorus (Ashmead and Zunino, 1993), serum phosphorus levels increased with the addition of organic iron to the diet at 50, 100, and 150 mg/ kg. This increase in serum phosphorus level can be due to insufficient iron level for the antagonist between the two elements. As a matter of fact, when Table 6 is examined, it is seen that the serum phosphorus level tends to decrease with the addition of 200 mg/kg iron in the diet. It can also be predicted that this decrease in serum phosphorus level will be exacerbated by the addition of higher levels of iron to the diet.

#### CONCLUSIONS

In this study, the performance parameters of lay-

Table 5. Influence of dietary supplementation of organic iron on the blood haematological parameters in layer quails										
		Organ	ic iron (n	P-value of	<i>P</i> -value of contrast					
Parameters	0	50	100	150	200	SEM*	Linear	Quadradic		
White Blood Cell, 10 <sup>3</sup> /µL	14.26	17.42	18.10	21.43	19.25	0.671	0.001	0.093		
Neutrophil, 10 <sup>3</sup> /µL	0.632	0.753	0.656	0.989	0.803	0.0535	0.125	0.673		
Lymphocyte, 10 <sup>3</sup> /µL	13.73	15.78	13.46	17.14	15.65	1.608	0.355	0.870		
Monocyte, $10^{3}/\mu L$	0.058	0.043	0.122	0.100	0.147	0.0121	0.004	0.799		
Eosinophil, 10 <sup>3</sup> /µL	0.195	0.178	0.294	0.236	0.215	0.0287	0.648	0.472		
Basophil, 10 <sup>3</sup> /µL	1.300	0.528	1.048	0.488	0.835	0.2280	0.569	0.566		
Erythrocyte, 10 <sup>6</sup> /µL	2.97	3.07	2.94	2.67	3.10	0.063	0.777	0.321		
Haemoglobin, g/dL	16.52	17.73	16.96	17.68	17.09	0.472	0.473	0.914		
Haematocrit, %	43.05	45.10	44.14	42.25	47.18	0.599	0.173	0.301		
Mean corpuscular volume, µm <sup>3</sup>	146	147	150	150	147	1.2	0.525	0.223		
Mean corpuscular haemoglobin, pg	55.90	58.12	57.70	55.76	57.20	0.871	0.970	0.692		
Red cell distribution width, %	10.90	10.63	11.58	11.22	10.98	0.163	0.522	0.372		
Platelet, $10^{3}/\mu L$	29.94	43.55	38.22	41.56	28.52	3.470	0.848	0.144		
Haemoglobin/ Lymphocyte	1.22	1.16	1.50	1.07	1.18	0.147	0.754	0.517		

\*Standard error means

Table 6. Innuence of dietary supplementation of organic from on the serum parameters in fayer quans									
	Organic iron (mg/kg)						P-value	of contrast	
Parameters	0	50	100	150	200	SEM*	Linear	Quadratic	
Glucose, mg/dL	294	287	255	280	295	6.2	0.933	0.057	
Triglyceride, mg/dL	1197	1290	1318	1164	1231	33.1	0.813	0.414	
Cholesterol, mg/dL	183	159	172	145	164	4.9	0.118	0.247	
AST, U/L	259	251	238	240	281	11.6	0.714	0.218	
ALT, U/L	1.50	1.33	1.67	2.50	2.00	0.251	0.245	0.939	
Total protein, g/dL	4.10	3.77	4.30	4.18	4.20	0.098	0.385	0.952	
Albumin, g/dL	1.45	1.40	1.58	1.50	1.47	0.035	0.602	0.442	
Globulin, g/dL	2.65	2.37	2.72	2.68	2.73	0.070	0.340	0.634	
Urea, mg/dL	11.33	10.17	11.33	9.67	12.50	0.472	0.585	0.199	
Creatinine, mg/dL	0.308	0.275	0.297	0.680	0.292	0.0805	0.522	0.612	
Calcium, mg/dL	20.20	17.78	22.02	20.23	19.12	0.590	0.945	0.150	
Phosphorus, mg/dL	7.60	8.70	8.30	8.53	7.07	0.256	0.487	0.039	
Iron, μg/dL	389	495	509	541	610	18.7	< 0.001	0.632	

\*Standard error means, AST: aspartate aminotransferase, ALT: alanine aminotransferase.

ing quails were not affected by the addition of organic iron to the diet. Eggshell resistance and white blood cell counts increased with the addition of organic iron to the diet at the level of 150 mg/kg. In addition, serum iron concentration linearly increased with rising dietary organic iron supplementation. These results

showed that laying quails can be fed with diets enriched with organic iron at the level of 150 mg/kg.

#### **CONFLICT OF INTEREST**

The authors declare no conflict of interest.

# REFERENCES

- Ashmead HD, Zunino H (1993) Factors which affect the intestinal absorption of minerals. The roles of amino acid chelates in animal nutrition. Noyes Publications, New Jersey, pp: 21-46. Bai S, Zhang K, Ding X, Wang J, Zeng Q, Peng H, Xuan Y, Su Z, Bai J
- (2018) High dietary iron differentially influences the iron distribution in the livers and the spleens of laying hens after salmonella typhimurium infection. Biol Trace Elem Res 185: 497-508.
- Buckiuniene V, Grashorn MA, Gruzauskas R, Kliseviciute V, Raceviciute-Stupeliene A, Svirmickas G,Bliznikas AM,Alencikiene G (2016) Effect of organic and inorganic iron in the diet on yolk iron content, fatty acids profile, malondialdehyde concentration, and sensory quality of chicken eggs. Eur Poult Sci 80: 139-151. Campbell TW (1988) Avian haematology and cytology. Iowa State Uni-
- versity Press, Ames, Iowa, pp:3-27.
- Chen YJ, ChoJH, YooJS, WangY, HuangY, KimIH (2008) Evaluation of  $\delta$ -aminolevulinic acid on serum iron status, blood characteristics, egg performance and quality in laying hens. Asian-Austral J Anim Sci 21:1355-1360.
- Choi JS, Koh IU, Lee HJ, Kim WH, Song J (2013) Effects of excess dietary iron and fat on glucose and lipid metabolism. J Nutr Biochem 24: 1634-1644
- Gibson RS, Bailey KB, Gibbs M, Ferguson EL (2010) A review of phytate, iron, zinc, and calcium concentrations in plant-based complementary foods used in low-income countries and implications for bioavailability. Food Nutr Bull 31:134-146.
- Gou Z, Fan Q, Li L, Jiang Z, Lin X, Cui X, Wang Y, Zheng C, Jiang S (2020) Effects of dietary iron on reproductive performance of Chinese Yellow broiler breeder hens during the egg-laying period. Poult Sci 99: 3921-3929.
- Gupta UC, WuK, Liang S (2008) Micronutrients in soils, crops, and livestock. Earth Sci Front 15:110-125.
- Leeson S, Summers JD (2001) Minerals. In: Lesson, S., Summers, J.D. (Eds.), Scott's Nutrition of the Chicken. Univ. Books, Guelph, Guelph, ON, Canada, pp. 331-428.
- Li L,Frei B (2006)Iron chelation inhibits NF-kB-mediated adhesion molecule expression by inhibiting p22phox protein expression and NA-DPH oxidase activity. Arterioscler Thromb Vasc Biol26: 2638-2643.
- Lin X, Gou Z, Wang Y, Li L, Fan Q, Ding F, Zheng C, Jiang, S (2020) Effects of dietary iron level on growth performance, immune organ indices and meat quality in Chinese yellow broilers. Animals 10: 670.
- Ma XY, Liu SB, Lu L, Li SF, Xie JJ, Zhang LY, ZhangJH, Luo XG(2014)

Relative bioavailability of iron proteinate for broilers fed a casein-dextrose diet. Poult Sci 93:556-563.

- Mackenzie EL, Iwasaki K, Tsuji Y(2008) Intracellular iron transport and storage: from molecular mechanisms to health implications. Antioxid Redox Signal 10:997-1030.
- McDowell LR (2003) Minerals in animal and human nutrition. Amsterdam: Elsevier Science BV. pp: 203-233.
- Paik I, LeeH, ParkS(2009) Effects of organic iron supplementation on the performance and iron content in the egg yolk of laying hens. J Poult Sci 46: 198-202
- Rajpathak SN, CrandallJP, Wylie-RosettJ, KobatGC, PohnTE(2009) The role of iron in type 2 diabetes in humans. Biochim Biophys Acta 1790: 671-681.
- Romero C, Brenes M, Garcia P, Garrido A (2002) Hydroxytyrosol 4-β-D-glucoside, an important phenolic compound in olive fruits and derived products. J Agric Food Chem 50: 3835-3839.
- Sarlak S, Tabeidian SA, Toghyani M, Shahraki ADF, Goli M, Habibian M (2021) Effects of replacing inorganic with organic iron on performance, egg quality, serum and egg yolk lipids, antioxidant status, and iron accumulation in eggs of laying hens. Biol Trace Elem Res 199: 1986-1999.
- Scott C, Arora G, Dickson K, Lehmann C (2021) Iron chelation in local infection. Molecules, 26: 189.
- Seo YM, Shin KS, Rhee AR, Chi YS, Han J, Paik IK (2010) Effects of dietary Fe-soy proteinate and MgO on egg production and quality of eggshell in laying hens. Asian-Austral J Anim Sci 23: 1043-1048.
- TaschettoD, VieiraSL, AngelCR, StefanelloC, KindleinL, EbbingMA, SimoesCT(2017) Iron requirements of broiler breeder hens. Poult Sci 96:3920-3927
- Vieira SL (2008) Chelated minerals for poultry. Rev Bras Cienc Avic 10: 73-79.
- Xie C, Elwan HAM, Elnesr SS, DongXY, Zou XT (2019a) Effect of iron glycine chelate supplementation on egg quality and egg iron enrichment in laying hens. PoultSci 98:7101-7109.
- Xie C, Elwan HA, Elnesr SS, Dong X, Feng J, Zou XT(2019b) Effects of iron glycine chelate on laying performance, antioxidant activities, serum biochemical indices, iron concentrations and transferrin mRNA expression in laying hens. J AnimPhysiolAnimNutr 103:547-554
- Yu B, Huang WJ, Chiou PW (2000) Bioavailability of iron from amino acid complex in weaning pigs. Anim Feed Sci Technol 86:39-52.

J HELLENIC VET MED SOC 2022, 73 (4) ПЕКЕ 2022, 73 (4)