

## Journal of the Hellenic Veterinary Medical Society

Vol 72, No 3 (2021)



### Differences in antimicrobial resistance of *Salmonella* spp. isolated from humans, animals and food products in Kazakhstan

R RYCHSHANOVA, M RUZAUSKAS, G CHUZHEBAYEVA, R MOCKELIUNAS, N MAMIYEV, M VIRGAILIS, P SHEVCHENKO, R SIUGZDINIENE, L ANSKIENE, A MENDYBAYEVA

doi: [10.12681/jhvms.28498](https://doi.org/10.12681/jhvms.28498)

Copyright © 2021, R RYCHSHANOVA, M RUZAUSKAS, G CHUZHEBAYEVA, R MOCKELIUNAS, N MAMIYEV, M VIRGAILIS, P SHEVCHENKO, R SIUGZDINIENE, L ANSKIENE, A MENDYBAYEVA



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

#### To cite this article:

RYCHSHANOVA, R., RUZAUSKAS, M., CHUZHEBAYEVA, G., MOCKELIUNAS, R., MAMIYEV, N., VIRGAILIS, M., SHEVCHENKO, P., SIUGZDINIENE, R., ANSKIENE, L., & MENDYBAYEVA, A. (2021). Differences in antimicrobial resistance of *Salmonella* spp. isolated from humans, animals and food products in Kazakhstan. *Journal of the Hellenic Veterinary Medical Society*, 72(3), 3091–3100. <https://doi.org/10.12681/jhvms.28498>

## Differences in antimicrobial resistance of *Salmonella* spp. isolated from humans, animals and food products in Kazakhstan

R. Rychshanova<sup>1</sup>, M. Ruzauskas<sup>2\*</sup>, G. Chuzhebayeva<sup>1</sup>, R. Mockeliunas<sup>2</sup>,  
N. Mamiyev<sup>1</sup>, M. Virgailis<sup>2</sup>, P. Shevchenko<sup>1</sup>, R. Siugzdiniene<sup>2</sup>, L. Anskiene<sup>2</sup>,  
A. Mendybayeva<sup>1</sup>

<sup>1</sup>Institute of Applied Biotechnology, Kostanay State University, Kazakhstan

<sup>2</sup>Veterinary Academy, Kaunas, Lithuanian University of Health Sciences

**ABSTRACT:** The aims of this study were twofold: 1. to isolate *Salmonella* spp. from different sources in Kazakhstan, to determine their serovars and resistance profiles and 2. to evaluate similarities in antimicrobial susceptibility of *Salmonella* spp. isolated from humans, animals and food products. From the 10 212 samples tested *Salmonella* spp. were isolated in 47 cases. The predominant serovar isolated from humans and food products was *S. Enteritidis*. Although different animal species were tested the *Salmonella* spp. were isolated mainly from chickens and ducks. *S. Enteritidis* and *S. Typhimurium* were the most prevalent serovars in raw poultry meat. The most frequent resistances were those to nalidixic acid, ampicillin and tetracycline. Human isolates demonstrated lower resistance compared to animal and food isolates. The genes encoding antimicrobial resistance in human isolates in most cases were absent except for some isolates which harboured *tet*, *sul*, *stre* and *bla* TEM genes encoding resistance to the oldest antimicrobial classes. The situation in animals and food products was less beneficial as besides different genes, integrons associated with the horizontal gene transfer were detected. The findings suggest that antimicrobials in animal sector should be used more strictly, paying attention to critically important antimicrobials for humans and possible horizontal transfer of pathogens and their genetic determinants.

**Keywords:** *Salmonella* Paratyphi, One Health, salmonellosis, zoonoses

*Corresponding Author:*  
M. Ruzauskas, Lithuanian University of Health Sciences, Mickeviciaus 9, Kaunas,  
LT-44307, Lithuania  
E-mail address: modestas.ruzauskas@lsmuni.lt

*Date of initial submission:* 26-03-2020  
*Date of revised submission:* 12-12-2020  
*Date of acceptance:* 13-01-2021

## INTRODUCTION

*Salmonella* spp. are one of the most frequently isolated foodborne pathogens. It is a major worldwide public health concern, accounting for 93.8 million foodborne illnesses and 155,000 deaths per year (Eng et al., 2015). Gastroenteritis is the most common manifestation of *Salmonella* infection, followed by bacteraemia and enteric fever (Majowicz et al. 2010). Non-typhoidal *Salmonella enterica* infection, caused by different serovars from contaminated food, is an important cause of both sporadic gastroenteritis and outbreaks internationally. In the European Union, over 91,000 human cases are reported each year (EFSA, 2018). In Australia, the incidence of infection due to *Salmonella* spp. in the community is estimated to be 185 infections per 100,000 population per year (Kirk et al., 2014). Enteric fever that is caused by *Salmonella enterica* Typhi causes illness in south-central Asia, Southeast Asia, and southern Africa by more than 100 cases per 100,000 person-years (Marks et al., 2017). It is also prevalent in many other parts of the world. For instance, approximately 200 to 300 cases of *S. Typhi* were reported in the United States each year (Lynch et al, 2009).

Epidemiological data and incidence cases of salmonellosis are registered in many countries including the USA, EU, China, Australia, India and others while some countries still are white spots in the map according to the prevalence of salmonellosis (Eng et al., 2015). In Kazakhstan the incidence of diagnosed salmonellosis in 1999 and 2000 was about 3500-4000 cases per year, while typhoid fever incidence was about 30-110 per year (WHO 2001).

Main factors associated with *Salmonella* outbreaks include incomplete cooking of food products, improper storage and direct contact with raw ingredients (Lynch et al. 2006). The food products that are predominantly associated with the outbreaks include animal products such as milk, poultry and eggs, as well as food products such as chocolate and peanut butter (Eng et al., 2015). Animals are the main carriers and reservoir of non-typhoid *Salmonella*. The transmission of *Salmonella* usually occurs through gastrointestinal tract by ingestion of food or water contaminated with infected animals. Direct contact with infected animals also constitutes a risk of infection (Swanson et al., 2007). The worldwide incidence rate of *Salmonella* infection is high as the strains can be found naturally in the environment and in both domestic and wild animals including cats, dogs, am-

phibians, reptiles and rodents (Eng et al., 2015). The diversity of possible reservoirs of infection results in significant challenges for public health authorities to control the infections (Swanson et al. 2007; Dione et al. 2011).

*Salmonella* resistance to at least 3 antibiotics started to be reported in 1960s (Dyson et al. 2019). Since then, the frequency of resistant isolates has increased in many countries; however, the spectrum of resistance or the rate of multidrug resistant isolates depends on country, source of isolation and *Salmonella* serovar. For instance, in the USA *Salmonella* spp. recovered from poultry were resistant to streptomycin (30.9%), gentamicin (12.6%), sulfadimethoxine (20.9%), tetracycline (13.9%), and trimethoprim-sulfamethoxazole combination (8.6%). Among these isolates, 20% were multi-drug resistant, showing resistance to three or more antibiotics; noteworthy is that 67% of *S. Heidelberg* and 54% of *S. Kentucky* isolates have shown resistance to five or more antibiotics (Liljebjelke et al., 2017). In China high rates of resistant *salmonellae* from chicken were found to sulfisoxazole (76.1%), tetracycline (75.3%), ampicillin (48.0%), and ofloxacin (44.7%). Notably, antimicrobial susceptibility tests have identified resistance to polymyxin B (2.0%) and imipenem (0.5%) (Zhang et al., 2018). In Poland the resistance of salmonella isolated from non-meat food products was low with the highest prevalence to nalidixic acid (35.2%), sulphonamides (6.6%), ampicillin (4.9%), amoxicillin/clavulanic acid (2.5%), streptomycin, cefoxitin, gentamicin and tetracycline (1.6%) (Mąka et al., 2015). Although an official data about salmonellosis prevalence in Kazakhstan is collected, there is no current systematic data about the situation of antimicrobial resistance of the strains that are prevalent in humans and in other sources including animals, food products or feedstuffs. Thus, the aims of this study were twofold: 1. to isolate *Salmonella* spp. strains from different sources in Kazakhstan, to determine their serovars and resistance profiles and 2. to evaluate similarities among antimicrobial susceptibility of *Salmonella* spp. isolated from humans, animals and food products.

## MATERIALS AND METHODS

### Place and sampling

Bacteriological investigations were performed at the Institute of Applied Biotechnology, Kostanay University during 2018-2019. The aim was to obtain wide spectrum of *Salmonella* isolates prevalent in Ka-

zakhstan. Animal carcasses and raw meat including poultry, lamb, beef were collected in supermarkets and markets at a retail level. Clinical and pathological material of animals was collected at slaughterhouses and animal farms. *Salmonella* spp. strains already isolated from humans (in hospitals and other health-care facilities) were delivered to the department of Microbiology of Kostanay University as isolates without the data about their concrete source (name of the patient) of isolation. Ready to use, raw and frozen food products were obtained at retail markets and supermarkets. In total 10212 samples were tested from which 1006 samples were from live animals, animal carcasses and milk, 383 food samples and 8823 human samples.

### Isolation and identification of *Salmonella*

Isolation of *Salmonella* was performed according to EN ISO 6579-1 (ISO, 2017) procedure for *Salmonella* detection. Xylose Lysine Deoxycholate (XLD) agar and Salmonella Shigella (SS) Agar (Oxoid, UK) were used as plating media after the enrichment procedure. Isolates obtained from health-care facilities were re-cultivated on XLD agar.

Identification of the isolates was performed using classical biochemical testing including fermentation of carbohydrates, production of hydrogen sulfide, indole, production of lysine decarboxylase, urease, oxidase, catalase, MR-VP as well as other conventional tests.

Serotyping was performed using slide agglutination test with sera to O and H antigens (Petsal, Russia) as well as Wellcolex Colour salmonella serogroup

identification latex test (Remel, UK) according to the manufacturer's instructions.

### Susceptibility testing

Antimicrobial susceptibility testing was performed using the disc diffusion method according to Kirby-Bauer (Hudzicki, 2009). The following discs with antimicrobials ( $\mu\text{g}$ ) (Thermo Fisher, UK) were used: ampicillin (10), cefoperazone (75), cefoxitin (30), streptomycin (10), kanamycin (30), gentamicin (10), tetracycline (30), doxycycline (30), chloramphenicol (30), ciprofloxacin (5), nalidixic acid (30), gemifloxacin (5), enrofloxacin (5), ofloxacin (5) and sulfamethoxazole-trimethoprim (25). Interpretation of the results was carried out using CLSI clinical breakpoints (CLSI, 2019) set for *Salmonella* except for ofloxacin, gemifloxacin and nalidixic acid for which breakpoints set for Enterobacteriaceae were used.

### Detection of the genes encoding antimicrobial resistance

Molecular testing of the isolates was performed at the Microbiology and Virology Institute, Lithuanian University of Health Sciences. DNA material for molecular testing was obtained after bacterial lysis according to the extraction protocol prepared by the EU Community Reference Laboratory for Antimicrobial Resistance with slight modifications (Ruzauskas et al., 2014). Detection of the genes encoding antimicrobial resistance was performed by PCR. Annealing temperatures and oligonucleotides used are presented in Table 1.

**Table 1.** Antimicrobial resistance genes tested and oligonucleotide primers used in the study

Primer name	Sequence (5'-3')	Size, bp and t (°C)	Target gene	Source
<i>bla</i> TEM-F	GAGTATTCAACATTTTCGT	857 (50)	<i>tem</i>	Maynard et al., 2003
<i>bla</i> TEM-R	ACCAATGCTTAATCAGTGA			
<i>bla</i> SHV-F	TCGCCTGTGTATTATCTCCC	768 (60)	<i>shv</i>	Ojdana et al., 2014
<i>bla</i> SHV-R	CGCAGATAAATCACCACAATG			
<i>oxa</i> 1-F	TCAACAAATCGCCAGAGAAG	276 (55)	<i>oxa</i> group I	Bert et al., 2002
<i>oxa</i> 1-R	TCCCACACCAGAAAACCAG			
<i>oxa</i> group 5-F	AGCCGCATATTTAGTTCTAG	644 (56)	<i>oxa</i> group V	Bert et al., 2002
<i>oxa</i> group 5-R	ACCTCAGTTCCTTTCTCTAC			
<i>ctx</i> M-F	ATGTGCAGYACCAGTAARGT	593 (50)	<i>ctx</i> M	Pagani et al., 2003
<i>ctx</i> M-R	TGGGTRAARTARGTSACCAGA			
<i>ctx</i> M group 2-F	ATGATGACTCAGAGCATTTCGCCGC	876 (56)	<i>ctx</i> M2	Celenza et al., 2006
<i>ctx</i> M group 2-R	TCAGAAACCGTGGGTTACGATTTT			
<i>cm</i> y2-F	GCACTTAGCCACCTATACGGCAG	758 (58)	<i>cm</i> y	Hasman et al., 2005
<i>cm</i> y2-R	GCTTTTCAAGAATGCGCCAGG			

<i>PER-1-F</i>	ATGAATGTCATTATAAAAGCT	927 (48)	<i>per</i>	Celenza et al., 2006
<i>PER-1-R</i>	TTAATTTGGGCTTAGGG			
<i>PER-2-F</i>	ATGAATGTCATCACAAAATG	927 (49)		Celenza et al., 2006
<i>PER-2-R</i>	TCAATCCGGACTCACT			
<i>tetA-F</i>	GTGAAACCCAACATACCCC	888 (55)	<i>tetA</i>	Maynard et al., 2003
<i>tetA-R</i>	GAAGGCAAGCAGGATGTAG			
<i>tetB-F</i>	CCTTATCATGCCAGTCTTGC	774 (55)	<i>tetB</i>	Maynard et al., 2003
<i>tetB-R</i>	ACTGCCGTTTTTTCGCC			
<i>aadB-F</i>	ATGGACACAACGCAGGTCGC	534 (55)	<i>aadB</i>	Asadollahi et al., 2012
<i>aadB-R</i>	TTAGGCCGCATATCGCGACC			
<i>aadA-F</i>	GTGGATGGCGGCCTGAAGCC	528 (68)	<i>aadA</i>	Asadollahi et al., 2012
<i>aadA-R</i>	AATGCCCAGTCGGCAGCG			
<i>rmtB-F</i>	ATGAACATCAACGATGCCCT	769 (55)	<i>rmtB</i>	Yan et al., 2004
<i>rmtB-R</i>	CCTTCTGATTGGCTTATCCA			
<i>armA-F</i>	CAAATGGATAAGAATGATGTT	774 (55)	<i>armA</i>	Galimand et al., 2003
<i>armA-R</i>	TTATTTCTGAAATCCACT			
<i>aphA1-F</i>	AAACGTCTTGCTCGAGGC	500 (55)	<i>aphA1</i>	Frana et al., 2001
<i>aphA1-R</i>	CAAACCGTTATTCATTCGTGA			
<i>aacA4-F</i>	ATGACTGACATGACCTTGCG	487 (55)	<i>aacA4</i>	Odumosu et al., 2015
<i>aacA4-R</i>	TTAGGCATCACTGCGTGTTCG			
<i>aac(3)II-F</i>	TGAAACGCTGACGGAGCCTC	369 (65)	<i>aac(3)II</i>	Sandvang et al., 2009
<i>aac(3)II-R</i>	GTCCAACAG GTAGCACTGAG			
<i>strA-F</i>	CCTGGTGATAACGGCAATTC	546 (55)	<i>strA</i>	Lanz et al., 2003
<i>strA-R</i>	CCAATCGCAGATAGAAGGC			
<i>strB-F</i>	ATCGTCAAGGGATTGAAACC	509 (55)	<i>strB</i>	Lanz et al., 2003
<i>strB-R</i>	GGATCGTAGAACATATTGGC			
<i>catII-F</i>	ACACTTTGCCCTTTATCGTC	495 (55)	<i>catII</i>	Vassort-Bruneau et al., 1996
<i>catII-R</i>	TGAAAGCCATCACATACTGC			
<i>cmlA-F</i>	TTGCAACAGTACGTGACAT	293 (55)	<i>cmlA</i>	Keyes et al., 2000
<i>cmlA-R</i>	ACACAACGTGTACAACCAG			
<i>sul1-F</i>	TTCGGCATTCTGAATCTCAC	822 (55)	<i>sul1-F</i>	Christabel et al., 2012
<i>sul1-R</i>	ATGATCTAACCCCTCGGTCTC			
<i>sul2-F</i>	CGGCATCGTCAACATAACC	722 (50)	<i>sul2-F</i>	Pereten et al., 2003
<i>sul2-R</i>	GTGTGCGGATGAAGTCAG			
<i>sul3-F</i>	GAGCAAGATTTTTGGAATCG	792 (51)	<i>sul3-F</i>	Pereten et al., 2003
<i>sul3-R</i>	CATCTGCAGCTAACCTAGGGCTTTGA			
<i>dfr1-F</i>	ACGGATCCTGGCTGTTGGTTGGACGC	254 (55)	<i>dfr1</i>	Gibreel et al., 1998
<i>dfr1-R</i>	CGGAATTCACCTTCCGGCTCGATGC			
<i>dfr5-F</i>	GCBAAGGDGARCAGCT	394 (44)	<i>dfr5</i>	Seputiene et al., 2010
<i>dfr5-R</i>	TTMCCAYATTGATAGC			
<i>dfrA7-F</i>	AAAATTTCAATTGATTTCTGCA	471 (44)	<i>dfr7</i>	Navia et al., 2003
<i>dfrA7-R</i>	TTAGCCTTTTTTCCAAATCT			
<i>mcr1-F</i>	CGGTCAGTCCGTTTGTTC	309 (58)	<i>mcr1</i>	Liu et al., 2015
<i>mcr1-R</i>	CTTGGTCGGTCTGTAGGG			
<i>mcr2-F</i>	TGTTGCTTGTGCCGATTGGA	567 (58)	<i>mcr2</i>	Xavier et al., 2016
<i>mcr2-R</i>	AGATGGTATTGTTGGTTGCTG			
<i>qnrA-F</i>	ATTTCTCACGCCAGGATTTG	516 (53)	<i>qnrA</i>	Robicsek et al., 2006
<i>qnrA-R</i>	GATCGGCAAAGGTTAGGTCA			
<i>qnrB-F</i>	GATCGTGAAAGCCAGAAAGG	469 (53)	<i>qnrB</i>	Robicsek et al., 2006
<i>qnrB-R</i>	ACGATGCCTGGTAGTTGTCC			
<i>qnrS-F</i>	ACGACATTCGTCAACTGCAA	417 (53)	<i>qnrS</i>	Robicsek et al., 2006
<i>qnrS-R</i>	TAAATTGGCACCCTGTAGGC			
<i>qepA-F</i>	CAGTGGACATAAGCCTGTTC	218 (60)	<i>qepA</i>	Liu et al., 2008
<i>qepA-R</i>	CCCAGGCATAGACTGTA			
<i>teg1-F</i>	TTATTGCTGGGATTAGGC	164 (55)	integrase I class	Chen et al., 2013
<i>teg1-R</i>	ACGGCTACCCTCTGTTATC			
<i>teg2-F</i>	ACGACATTCGTCAACTGCAA	233 (50)	integrase II class	Goldstein, 2001
<i>teg2-R</i>	TAAATTGGCACCCTGTAGGC			

**Table 2.** Characteristics of *Salmonella* isolated from humans in Kazakhstan

Isolate number	Serovar	Phenotypic Resistance	Genotypic Resistance	Integrans
733	<i>S. Tshiongwe</i>	CIP		-
735	<i>S. Enteritidis</i>	NAL, CIP		-
737	<i>S. Blegdam</i>			-
739	<i>S. Blegdam</i>			-
990	<i>S. Enteritidis</i>	NAL, ENR		-
991	<i>S. Enteritidis</i>	CIP		-
993	<i>S. Enteritidis</i>	TET		-
994	<i>S. Enteritidis</i>	STR, DOX	<i>strB</i>	-
998	<i>S. Enteritidis</i>	ENR, NAL, CIP		-
999	<i>S. Enteritidis</i>	NAL, ENR, CIP, GEMI,		-
1106	<i>S. Enteritidis</i>	NAL, CIP		-
1107	<i>S. Enteritidis</i>			-
1108	<i>S. Enteritidis</i>			-
1109	<i>S. Enteritidis</i>			-
1110	<i>S. Enteritidis</i>			-
1111	<i>S. Enteritidis</i>	NAL, CIP		-
1112	<i>S. Enteritidis</i>			-
1113	<i>S. Enteritidis</i>	TET	<i>tet B</i>	-
1205	<i>S. Enteritidis</i>	NAL, ENR, CIP, GEMI		-
1206	<i>S. Enteritidis</i>	AMP, TET, DOX, NAL, ENR, CIP, OFX, SXT	<i>blaTEM, tet A, tet B, sul3</i>	+ (class I)
1280	IIIc 4p	NAL, ENR, CIP		-

NAL - nalidixic acid; CIP - ciprofloxacin; ENR - enrofloxacin; TET - tetracycline, DOX - doxycycline; GEMI - gemifloxacin; AMP - ampicillin; OFX - ofloxacin; STRE - streptomycin; SXT - sulfamethoxazole/trimethoprim

**Table 3.** Characteristics of *Salmonella* isolated from animals in Kazakhstan

Serovar	Source of Isolation	Phenotypic Resistance	Genotypic resistance	Integrans
<i>S. Typhimurium</i>	duck	STR, KAN, ENR		-
<i>S. Typhimurium</i>	duck	STR, CHL, TET, ENR,		-
<i>S. Typhimurium</i>	chicken			-
<i>S. Typhimurium</i>	chicken			-
<i>S. Paratyphi C</i>	chicken	CHL, TET, DOX, NAL, CIP, SXT	<i>tetA, sul3, cmlA, catII</i>	+ (class I)
<i>S. Enteritidis</i>	chicken	NAL, CIP		-
<i>S. Enteritidis</i>	chicken	NAL, CIP		-
<i>S. Enteritidis</i>	chicken	AMP, STR, SXT	<i>Bla TEM, aadA, dfr1</i>	-
<i>S. Enteritidis</i>	chicken	STR, TET, DOX, ENR, CIP, GEMI, NAL, SXT	<i>tet A, sul 2, dfr1, strA, str B</i>	-
<i>S. Enteritidis</i>	chicken	ENR, NAL, CIP		-
<i>S. Enteritidis</i>	chicken	AMP, STR, ENR	<i>bla TEM, aacA4,</i>	-
<i>S. Enteritidis</i>	chicken	NAL, ENR, CIP		-
<i>S. Moscow</i>	chicken	NAL, ENR, CIP		-
<i>S. Paratyphi C</i>	chicken (imported)	FOX, STR, TET, DOX, NAL, ENR, CIP, OFX, GEMI	<i>aadA, tet A, qnr B</i>	+ (class I)

NAL - nalidixic acid; CIP - ciprofloxacin; ENR - enrofloxacin; TET - tetracycline, DOX - doxycycline; GEMI - gemifloxacin; AMP - ampicillin; OFX - ofloxacin; SXT - sulfamethoxazole/trimethoprim; CHL - chloramphenicol; STR - streptomycin; KAN - kanamycin

**Table 4.** Characteristics of *Salmonella* isolated from food products in Kazakhstan

Serovar	Source of isolation	Phenotypic Resistance	Genotypic resistance	Integrans
<i>S. Enteritidis</i>	Roasted chicken drumsticks			-
<i>S. Enteritidis</i>	Roasted chicken drumsticks	NAL, ENR, CIP		-
<i>S. Enteritidis</i>	Sausage	NAL, ENR, CIP		-
<i>S. Enteritidis</i>	Cake			-
<i>S. Enteritidis</i>	Ice-cream	NAL, ENR, CIP		-
<i>S. Enteritidis</i>	Cake	NAL, ENR, CIP		-
<i>S. Enteritidis</i>	cake		tet B	-
<i>S. Enteritidis</i>	Refrigerated chicken drumsticks	AMP, STR, KAN, TET, DOX, ENR		-
<i>S. Enteritidis</i>	Sausage	AMP, STR, KAN, CHL, DOX, ENR		-
<i>S. Enteritidis</i>	Refrigerated chicken drumsticks	AMP,STR, KAN, CN, TET, DOX		-
<i>Salmonella</i> group C	Force meat	AMP, STR, CHL, TET, DOX, NAL, ENR, CIP, OFX, GEMI, SXT	aadB, tet A, tet B, sul3, catII, qnrA	+ (Class I)
<i>S. Tennessee</i>	Meat in paste	AMP, CFP, KAN, CHL, TET, DOX, NAL, ENR, CIP, SXT	ctxM, aphA1, aadA, tet A, sul3, cmlA, catII	+ (Class I, Class II)

NAL - nalidixic acid; CIP - ciprofloxacin; ENR - enrofloxacin; TET - tetracycline, DOX - doxycycline; GEMI - gemifloxacin; AMP - ampicillin; CN - gentamicin; OFX - ofloxacin; SXT - sulfamethoxazole/trimethoprim; CHL - chloramphenicol; STR - streptomycin; KAN - kanamycin; CFP - cefpodoxime

### Data analysis

Occurrence of multi-resistant isolates was calculated by dividing the number of isolates resistant to at least 3 antimicrobial classes by the total number of obtained isolates. Statistical analysis was performed using IBM SPSS Statistics package, version 20.

Comparison between categorical variables was calculated using the chi-square test. Results were considered statistically significant if  $P \leq 0.05$ .

### RESULTS

From total 10 212 samples tested (1006 samples from live animals, animal carcasses and milk; 383 food and 8823 human samples) *Salmonella* were isolated in 47 cases (0.46%). Twenty one (21) isolates (0.24%) were obtained from humans, fourteen (14) (1.39%) - from live or slaughtered animals and twelve (12) isolates (3.13 %) - from ready to use, raw or refrigerated food products (Tables 2-4).

As it could be seen from Table 2, the predominant *Salmonella* serovar isolated from humans was *S. Enteritidis* (81% from all isolates). Susceptibility varied among the strains: seven isolates were susceptible to all tested antimicrobials while a single isolate was resistant to eight (8) of the tested antimicrobials. The most frequent resistance was towards nalidixic acid, fluoroquinolones and tetracyclines. The prevalence of the genes encoding resistance was low. Integrans

were detected in a single multi-resistant isolate of *S. Enteritidis*.

Although different animal species were tested the *Salmonella* were isolated only from chickens and ducks (Table 3). The serovar variety in poultry was higher to those of humans. The predominant serovar were *S. Typhimurium* and *S. Enteritidis*. Two isolates were identified as *S. Paratyphi* C. One of those isolates was resistant to tetracyclines, nalidixic acid, ciprofloxacin, chloramphenicol, sulfonamides and trimethoprim and harboured class I integrans while the other - to 2d generation cephalosporins, streptomycin, tetracyclines and all fluoroquinolones tested. Two (2) strains out of fourteen (14) isolated from animals were susceptible to all antimicrobials tested while six (6) isolates were multi-resistant.


*S. Enteritidis* was the predominant serovar in food products (81% from all isolates) however, two other serovars- (*Salmonella* group C and *S. Tennessee*) had much wider spectrum of resistance. Five (5) isolates out of twelve (12) of food origin were multi-resistant, two (2) strains were susceptible to all antimicrobials and four (4) had similar resistance patterns being resistant only to (fluoro)quinolones. The isolate of *S. Tennessee* was a single strain harboured integrans of both I and II class.

The frequency of antimicrobial resistance in *Sal-*

*monella* isolates obtained from different sources is presented in Fig 1.

The most frequent resistance among *Salmonella* isolates was demonstrated to fluoro(quinolones), ampicillin, streptomycin and tetracycline. Resistance to critically important antimicrobials for humans was also detected and varied depending on the source of isolation. Human isolates demonstrated lower resistance compared to animal and food isolates. Food iso-

lates were more frequently resistant to antimicrobials compared to animal isolates except for streptomycin and enrofloxacin which were more frequently non effective against animal isolates and for nalidixic acid for which the number of resistant isolates from animals and foods had the same range. The correlation between the strains isolated from different sources regarding their resistances to separate antimicrobials is presented in Table 5.



NAL - nalidixic acid; CIP - ciprofloxacin; ENR - enrofloxacin; TET - tetracycline, DOX - doxycycline; GEMI - gemifloxacin; AMP - ampicillin; CN - gentamicin; OFX - ofloxacin; SXT - sulfamethoxazole/trimethoprim; CHL - chloramphenicol; STR - streptomycin; KAN - kanamycin; FOX - cefoxitin; CFP - cefpodoxime

**Fig 1.** Antimicrobial resistance (% of resistant isolates) of *Salmonella* isolated from humans, animals and food products

**Table 5.** Statistical differences among resistances in *Salmonella* isolates from different sources

Antibiotics	Human vs Animals			Human vs Food			Animals vs Food		
	Value	df	P - value	Value	df	P - value	Value	df	P - value
Ampicillin	0.972	1	0.324	6.991	1	<b>0.008</b>	2.462	1	0.117
Cefoperazone	constant			2.414	1	0.120	1.400	1	0.237
Cefoxitin	1.805	1	0.179	constant			0.788	1	0.375
Streptomycin	10.862	1	<b>0.001</b>	7.966	1	<b>0.005</b>	0.248	1	0.619
Kanamycin	1.544	1	0.214	7.966	1	<b>0.005</b>	2.854	1	0.091
Gentamicin	constant			1.805	1	0.179	1.213	1	0.271
Levomycetin	3.182	1	0.074	5.775	1	<b>0.016</b>	0.478	1	0.490
Tetracycline	2.146	1	0.143	2.910	1	0.088	0.069	1	0.793
Doxycycline	1.176	1	0.278	4.721	1	<b>0.030</b>	0.991	1	0.319
Enrofloxacin	6.801	1	<b>0.009</b>	5.040	1	<b>0.025</b>	0.281	1	0.596
Ciprofloxacin	0.638	1	0.424	0.524	1	0.469	0.000	1	1.000
Ofloxacin	0.171	1	0.679	0.408	1	0.523	0.046	1	0.830
Gemifloxacin	0.366	1	0.545	0.879	1	0.348	0.103	1	0.748
Nalidixic acid	1.733	1	0.188	1.429	1	0.232	0.000	1	1.000
Sulfamethoxazole/ trimethoprim	2.305	1	0.129	1.310	1	0.252	0.094	1	0.759



## DISCUSSION

This study describes the prevalence of *Salmonella* spp. in different sources in Kazakhstan. Up to date only the data from humans were collected by national and international responsible institutions within the country. Salmonellosis is one of the leading causes of foodborne infections in humans with animals being reservoirs of *Salmonella*. *Salmonella* colonizes the gastrointestinal tract of food animals (Andino et al., 2015) and is shed via feces (Narváez-Bravo et al., 2013). Although the clinical manifestation of salmonellosis in different food-producing animal species can be unequally expressed, all animal species can be a source of foodborne infection (Chaney et al., 2017; Evangelopoulou et al., 2015; Pande et al., 2016).

Knowledge of the prevalence and diversity of *Salmonella* serovars in animals and food can provide important information necessary to develop preventive measures and strategies at different stages of the food chain such as application hazard analysis and critical control programs in meat production industries to ensure food safety (Tietjen and Fung, 1995; Gutema et al., 2019). According to this study the most prevalent serovar of *Salmonella* in Kazakhstan was *S. Enteritidis* which was the most frequently detected both in humans (81%) and food products (83%). According to the World Health Organization, *S. Enteritidis* is one of the most frequently isolated *Salmonella* serovars (along with *S. Typhimurium*) from countries involved in the Global Foodborne Infections Network (Hendriksen et al., 2011). The other serovars isolated from humans included *S. Blegdam* and *S. Tshiongwe*. *S. Blegdam* which is genetically similar to *S. Enteritidis* was isolated for the first time in the USA in the middle of last century from a patient suffering with the enteric type of fever and extensive erythema during the fifth week after onset (Holt and Newton, 1948). There is a lack of data about its prevalence during past decades, although *S. Blegdam* was recently isolated from broilers in Egypt (Ammar et al., 2019). *S. Tshiongwe* was described as a pathogenic serovar for both humans and animals and was also isolated from foods (Thong et al., 2004; Mshelbwala et al., 2017). This study demonstrated that the variety of serovars prevalent in animals in Kazakhstan was not wide; besides *S. Enteritidis*, which prevalence was 50% of all serovars, *S. Typhimurium* (29%) was also common. This serovar was recovered from poultry. The disease caused by *S. Typhimurium* is of public health significance, as it is associated with food poisoning in humans (Dar et al., 2017). The other highly pathogenic to man serovar-

*S. Paratyphi C* was isolated from chicken carcasses in two cases. This serovar is associated with human disease although the carriers can be different animal species (ECDC, 2019). Although paratyphoid fever is a rare disease in Europe and the USA it is still prevalent in Asia (Ekdahl et al., 2005).

Interesting data was also obtained according to the antimicrobial susceptibility testing: the isolates from humans were less frequently resistant to different antimicrobials comparing to animal and particularly food isolates, although it is known that salmonellosis is a foodborne human infection. All of the isolates obtained from humans, *in vitro* were susceptible to chloramphenicol, cephalosporins and aminoglycosides (except one isolate which was resistant to streptomycin), while the isolates from animals and foods had different susceptibility patterns. Although aminoglycosides are poorly effective against salmonellosis *in vivo* (Kihlstrom and Andaker, 1985), these antimicrobials are good indicators testing *in vitro* when comparing differences in antimicrobial susceptibility of different isolates, particularly isolated from different sectors. As it is important to understand the chain of salmonellosis “from farm to table” the data obtained in this study is interestingly enough demonstrating quite big differences among susceptibility in salmonella isolates from animals, foods and humans. The most frequent resistance in the isolates irrespective from the source of isolation was towards (fluoro)quinolones, aminopenicillins, streptomycin, tetracyclines and sulphamethoxazole/trimethoprim. These antimicrobials are frequently used both in humans and animals. Comparing the resistance between animal and food isolates the differences were less obvious except for beta-lactams and kanamycin with the highest resistance frequency being in food isolates. Isolates from animals had the highest resistance towards enrofloxacin compared to isolates from other sources. This is probably associated with the frequent use of enrofloxacin for animal treatment in Kazakhstan as it is known that broad use of fluoroquinolones has been followed by emergence of resistance to this class of antimicrobials (Hooper, 2001). Although all food products from which *Salmonella* were isolated contained ingredients of animal origin (meat, milk products or eggs) only a part of the products were originated from Kazakhstan while some of them were imported (data are not presented). Resistance to 3<sup>d</sup> generation cephalosporins was detected only in the isolates from foods but not from humans and animals, meaning that *salmonellae* circulating in Kazakh-

stan are quite specific and isolated. This fact is also proved by the low overall prevalence of *Salmonella* and by the poor variety of serovars circulating within the country. Imported food however, may change the situation in case of foodborne infections appear. The current situation regarding antimicrobial resistance in humans is beneficial with possibility to treat salmonellosis with antimicrobials of the 1<sup>st</sup> choice. The best known genes encoding antimicrobial resistance in human isolates mostly were absent except for some isolates which harboured *tet*, *sul*, *stre* and *bla*TEM genes encoding resistance to the oldest antibiotic classes. The situation in animals and food products was less beneficial as besides different genes, integrons associated with the horizontal gene transfer were detected more frequently. These findings suggest that antimicrobials in animal sector should be used more strictly particularly paying attention to critically important antimicrobials for humans and possible horizontal transfer of pathogens and their genetic determinants between environment (animals, food ingredients) and

humans. Since multi-resistant highly pathogenic to humans serovars, as *S. Paratyphi C*, were detected in animal carcasses, monitoring of *Salmonella* prevalence in animals should be implemented on a regular basis and measures for *Salmonella* eradication should be foreseen.

In conclusion, food import is one of the key factors changing the *Salmonella* variety and its antimicrobial resistance patterns in Kazakhstan. Usage of enrofloxacin in veterinary medicine however, is the factor of increasing salmonella resistance to fluoroquinolones that may have negative impact on treatment of salmonellosis infections.

## ACKNOWLEDGEMENTS

This study was supported by the Ministry of Education and Science of the Republic of Kazakhstan, Project number AP05131447.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## REFERENCES

- Ammar AM, Abdeen EE, Abo-Shauma UP, Fekry E, Kotb Elmahallawy E (2019) Molecular characterization of virulence and antibiotic resistance genes among *Salmonella* serovars isolated from broilers in Egypt. *Lett Appl Microbiol* 68:188-195.
- Andino A, Hanning I (2015) *Salmonella enterica*: survival, colonization, and virulence differences among serovars. *Sci World J* 2015:520179.
- Asadollahi P, Akbari M, Soroush S, Taherikalani M, Asadollahi K, Sayehmiri K (2012) Antimicrobial resistance patterns and their encoding genes among *Acinetobacter baumannii* strains isolated from burned patients. *Clin Infect Dis* 38:1198-1203.
- Bert F, Branger C, Lambert-Zechovsky N (2002) Identification of PSE and OXA  $\beta$ -lactamase genes in *Pseudomonas aeruginosa* using PCR-restriction fragment length polymorphism. *J Antimicrob Chemother* 50:11-18.
- Celenza G, Pellegrini C, Caccamo M, Segatore B, Amicosante G, Perilli M (2006) Spread of *bla*CTX-M-type and *bla*PER-2  $\beta$ -lactamase genes in clinical isolates from Bolivian hospitals. *J Antimicrob Chemother* 57:975-978.
- Chaney WE, Agga GE, Nguyen SV, Arthur TM, Bosilevac JM, Dreyling E, Rishi A, Brichta-Harhay D (2017) Rapid detection and classification of *Salmonella enterica* shedding in feedlot cattle utilizing the roka bioscience atlas *Salmonella* detection assay for the analysis of rectoanal mucosal swabs. *J Food Prot* 80:1760-1767.
- Chen CH, Huang CC (2013) Risk factor analysis for extended-spectrum  $\beta$ -lactamase-producing *Enterobacter cloacae* bloodstream infections in central Taiwan. *BMC Infect Dis* 13:417.
- Christabel M, Budambula N, Kiiru J, Kariuki S (2012) Characterization of antibiotic resistance in environmental enteric pathogens from Kibera slum in Nairobi-Kenya. *Afr J Bacteriol Res* 4:46-54.
- CLSI (2019) Performance Standards for Antimicrobial Susceptibility Testing. 29th ed. CLSI supplement M100. Wayne, P. A.: Clinical and Laboratory Standards Institute.
- Dar MA, Ahmad SM, Bhat SA, Ahmed R, Urwat U, Mumtaz PT, Bhat SA, Dar TA, Shah RA, Ganai NA (2017) *Salmonella typhimurium* in poultry: a review. *World Poultry Sci J* 73:345-354.
- Dione MM, Ikumapayi UN, Saha D, Mohammed NI, Geerts S, Ieven M, Adegbola RA, Antonio M (2011) Clonal differences between non-typhoidal *Salmonella* (NTS) recovered from children and animals living in close contact in the Gambia. *PLoS Negl Trop Dis* 5:e1148.
- Dyson ZA, Klemm EJ, Palmer S, Dougan G (2019) Antibiotic resistance and typhoid. *Clin Infect Dis* 68:165-170.
- Ekdahl K, de Jong B, Wollin R, Andersson Y (2005) Travel-associated non-typhoidal salmonellosis - geographical and seasonal differences and serotype distribution. *Clin Microbiol Infect* 11:138-144.
- Eng SK, Pusparajah P, Ab Mutalib NS, Ser HL, Chan KG, Lee LH (2015) *Salmonella*: A review on pathogenesis, epidemiology and antibiotic resistance. *Front Life Sci* 8:284-293.
- European Centre for Disease Prevention and Control. Facts about typhoid and paratyphoid fever. Interactive <https://www.ecdc.europa.eu/en/typhoid-and-paratyphoid-fever/facts>. Accessed: 26-03-2020.
- Evangelopoulou G, Kritas S, Christodoulopoulos G, Burriel A R (2015) The commercial impact of pig *Salmonella* spp. infections in border-free markets during an economic recession. *Vet World* 8:257-272.
- Frana TS, Carlson SA, Griffith RW (2001) Relative distribution and conservation of genes encoding aminoglycoside-modifying enzymes in *Salmonella enterica* serotype Typhimurium phage type DT104. *Appl Environ Microbiol* 67:445-448.
- Galimand M, Courvalin P, Lambert T (2003) Plasmid-Mediated High-Level Resistance to Aminoglycosides in *Enterobacteriaceae* Due to 16S rRNA Methylation. *Antimicrob Agents Chemother* 47:2565-2571.
- Gibreel A, Sködl O (1998) High-level resistance to trimethoprim in clinical isolates of *Campylobacter jejuni* by acquisition of foreign genes (*dfp1* and *dfp9*) expressing drug-insensitive dihydrofolate reductases. *Antimicrob Agents Chemother* 42:3059-3064.
- Goldstein C, Lee MD, Sanchez S, Hudson C, Phillips B, Register B, Grady M, Liebert C, Summers AO, White DG, Maurer JJ (2001) Incidence of class 1 and 2 integrons in clinical and commensal bacteria from livestock, companion animals, and exotics. *Antimicrob Agents Chemother* 45:723-726.
- Gutema FD, Agga GE, Abdi RD, De Zutter L, Duchateau L, Gabriël S (2019) Prevalence and serotype diversity of *Salmonella* in apparently healthy

- cattle: Systematic review and meta-analysis of published studies, 2000-2017. *Front Vet Sci*6:102.
- Hasman H, Mevius D, Veldman K, Olesen I, Aarestrup FM (2005)  $\beta$ -lactamases among extended-spectrum  $\beta$ -lactamase (ESBL)-resistant *Salmonella* from poultry, poultry products and human patients in The Netherlands. *J Antimicrob Chemother* 56:115-121.
- Hendriksen RS, Vieira AR, Karlsmose S, Lo Fo Wong DM, Jensen AB, Wegener HC, Aarestrup FM (2011) Global monitoring of *Salmonella* serovar distribution from the World Health Organization global foodborne infections network country data bank: results of quality assured laboratories from 2001 to 2007. *Foodborne Pathog Dis* 8:887-900.
- Holt RA, Newton H (1948) The occurrence of *Salmonella blegdam* in Louisiana. *J Lab Clin Med* 33:1115-1158.
- Hooper DC (2001) Emerging mechanisms of fluoroquinolone resistance. *Emerg Infect Dis* 7:337-341.
- Hudzicki J (2009) Kirby-Bauer disk diffusion susceptibility test protocol. ASM Microbe Libr, URL (<https://www.asm.org/getattachment/2594ce26-bd44-47f6-8287-0657aa9185ad/Kirby-Bauer-Disk-Diffusion-Susceptibility-Test-Protocol-pdf.pdf>). (accessed 26.03.2020).
- ISO 6579-1:2017. Microbiology of the food chain - Horizontal method for the detection, enumeration and serotyping of *Salmonella* - Part 1: Horizontal method for the detection of *Salmonella* spp. International Organization for Standardization, Geneva, Switzerland.
- Keyes K, Hudson C, Maurer JJ, Thayer S, White DG, Lee MD (2000) Detection of florfenicol resistance genes in *Escherichia coli* isolated from sick chickens. *Antimicrob Agents Chemother*44:421-424.
- Kihlstrom E, Andaker L (1985) Inability of gentamicin and fosfomicin to eliminate intracellular Enterobacteriaceae. *J Antimicrob Chemother* 15:723-728.
- Kirk M, Ford L, Glass K, Hall G (2014) Foodborne illness, Australia, circa 2000 and circa 2010. *Emerg Infect Dis* 20:1857-1864.
- Lanz R, Kuhnert P, Boerlin P (2003) Antimicrobial resistance and resistance genes determinants in clinical *Escherichia coli* from different animal species in Switzerland. *Vet Microbiol*91:73-84.
- Liljebjelke KA, Hofacre CL, White DG, Ayers S, Lee MD, Maurer JJ (2017) Diversity of antimicrobial resistance phenotypes in *Salmonella* isolated from commercial poultry farms. *Front Vet Sci* 4:96.
- Liu JH, Deng YT, Zeng ZL, Gao JH, Chen L, Arakawa Y, Chen ZL (2008) Copevalence of plasmid-mediated quinolone resistance determinants QepA, Qnr, and AAC(6)-Ib-cr among 16S rRNA methylase RmtB-producing *Escherichia coli* isolates from pigs. *Antimicrob Agents Chemother* 52:2992-2993.
- Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, Doi Y, Tian G, Dong B, Huang X, Yu LF, Gu D, Ren H, Chen X, Lv L, He D, Zhou H, Liang Z, Liu, JH, Shen J (2015) Emergence of plasmid mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis*, S1473-3099(15)00424-7.
- Lynch MF, Blanton EM, Bulens S, Polyak C, Vojdani J, Stevenson J, Medalla F, Barzilay E, Joyce, K, Barrett T, Mintz ED (2009) Typhoid fever in the United States, 1999-2006. *JAMA* 302:859-865.
- Lynch M, Painter J, Woodruff R, Braden C (2006) Surveillance for foodborne-disease outbreaks - United States, 1998-2002. *MMWR Surveill Summ*10:1-42.
- Majowicz SE, Musto J, Scallan E, Angulo FJ, Kirk M, O'Brien SJ, Jones TF, Fazil A, Hoekstra R M (2010) The global burden of nontyphoidal *Salmonella* gastroenteritis. *Clin Infect Dis* 50:882-889.
- Mąka Ł, Maćkiw E, Ścieżyńska H, Popowska M (2015) Occurrence and antimicrobial resistance of *Salmonella* spp. isolated from food other than meat in Poland. *Ann Agric Environ Med AAEM*22:403-408.
- Marks F, von Kalckreuth V, Aaby P, Adu-Sarkodie Y, El Tayeb MA, Ali M, Aseffa A, Baker S, Biggs HM, Bjerregaard-Andersen M, et al. (2017) Incidence of invasive salmonella disease in sub-Saharan Africa: a multi-centre population-based surveillance study. *Lancet Glob Health* 5:e310.
- Maynard C, Fairbrother JM, Bekal S, Sanschagrin F, Levesque RC, Brousseau R, Masson L, Larivière S, Harel J (2003) Antimicrobial resistance genes in enterotoxigenic *Escherichia coli* O149:K91 isolates obtained over a 23-year period from pigs. *Antimicrob Agents Chemother* 47:3214-3221.
- Mshelbwala FM, Ibrahim NDG, Saidu SNA, Azeed AA, Akinduti PA, Kwanashie CN, Kadiri AKF, Muhammed M, Fagdamila IO, Luka PD (2017) Motile *Salmonella* serotypes causing high mortality in poultry farms in three Southwestern States of Nigeria. *Vet Rec Open* 4:e000247.
- Narváez-Bravo C, Rodas-González A, Fuenmayor Y, Flores-Rondon C, Carruyo G, Moreno M, Perozo-Mena A, Hoet AE (2013) *Salmonella*. on feces, hides and carcasses in beef slaughter facilities in Venezuela. *Int J Food Microbiol* 166:226-230.
- Navia MM, Ruiz J, Cespedes SJ, Vila J (2003) Detection of dihydrofolate reductase genes by PCR and RFLP. *Diagn Microbiol Infect Dis* 46:295-298.
- Odumusu BT, Akintimehin AR (2015) Occurrence of extended-spectrum beta-lactamase producing *Enterobacteriaceae* isolates in communal water sources in Ogun State, Nigeria. *Afr J Clin Exp Microbiol*16:28-32.
- Ojdana D, Sacha PB, Wiczorek P, Czaban B, Michalska A, Jaworowska J, Jurczak A, Poniatowski B, Tryniszewska E (2014) The occurrence of *bla*CTX-M, *bla*SHV, and *bla*TEM genes in extended-spectrum  $\beta$ -Lactamase-positive strains of *Klebsiella pneumoniae*, *Escherichia coli*, and *Proteus mirabilis* in Poland. *Int J Antibiot* :93584.
- Pagani L, Dell'Amico E, Migliavacca R, D'Andrea MM, Giacobone E, Amicosante GER, Rossolini GM (2003) Multiple CTX-M-Type extended-spectrum  $\beta$ -lactamases in nosocomial isolates of *Enterobacteriaceae* from a hospital in northern Italy. *J Clin Microbiol* 41:4264-4269.
- Pande VV, Devon RL, Sharma P, McWhorter AR, Chousalkar KK (2016) Study of *Salmonella typhimurium* infection in laying hens. *Front Microbiol*7:203.
- Perreten V, Boerlin PA (2003) New sulfonamide resistance gene (*sul3*) in *Escherichia coli* is widespread in the pig population of Switzerland. *Antimicrob Agents Chemother*47:1169-1172.
- Robicsek A, Strahilevitz J, Sahn D, Jacoby GA, Hooper DC (2006) *qnr* prevalence in ceftazidime-resistant *Enterobacteriaceae* isolates from the United States. *Antimicrob Agents Chemother* 50:2872-2874.
- Ruzauskas M, Siugzdiniene R, Klimiene I, Virgailis M, Mockeliunas R, Vaskeviciute L, Zienius D (2014) Prevalence of methicillin-resistant *Staphylococcus haemolyticus* in companion animals: a cross-sectional study. *Ann Clin Microbiol Antimicrob*13:56.
- Sandvang D, Aarestrup FM (2009) Characterization of aminoglycoside resistance genes and Class 1 integrons in porcine and bovine gentamicin-resistant *Escherichia coli*. *Microb Drug Resist*6:19-27.
- Seputiene V, Povilonis J, Ruzauskas M, Pavilonis A, Suziedeliene E (2010) Prevalence of trimethoprim resistance genes in *Escherichia coli* isolates of human and animal origin in Lithuania. *J Med Microbiol*59:315-322.
- Swanson SJ, Snider C, Braden CR, Boxrud D, Wunschmann A, Rudroff JA, Lockett J, Smith KE (2007) Multidrug-resistant *Salmonella enterica* serotype *Typhimurium* associated with pet rodents. *N Engl J Med*356:21-28.
- Thong KL, Bakeri SA, Lai KS, Koh YT, Taib MZ, Lim VKE, Yasin RM (2004) Molecular subtyping of *Salmonella enterica* serovar Tshiongw recently isolated in Malaysia during 2001-2002. *Southeast Asian J Trop Med Public Health* 35:92-96.
- Tietjen M, Fung DYC (1995) *Salmonellae* and food safety. *Crit Rev Microbiol* 21:53-83.
- Tirado C, Schmidt K (2001) WHO surveillance programme for control of foodborne infections and intoxications: preliminary results and trends across greater Europe. *World Health Organization. J Infect*43:80-84.
- Vassort-Bruneau C, Lesage-Descauses M, Martel JL, Lafont JP, Chaslus-Dancla E (1996) CAT III chloramphenicol resistance in *Pasteurella haemolytica* and *Pasteurella multocida* isolated from calves. *J Antimicrob Chemother*38: 205-213.
- Xavier BB, Lammens C, Ruhel R, Kumar-Singh S, Butaye P, Goossens H, Malhotra-Kumar S (2016) Identification of a novel plasmid-mediated colistin-resistance gene, *mcr-2*, in *Escherichia coli*, Belgium. *Euro Surveill*.21:pil=30280.
- Yan JJ, Wu JJ, Ko WC, Tsai SH, Chuang CL, Wu HM, Lu YJ, Li JD (2004) Plasmid-mediated 16S rRNA methylases conferring high-level aminoglycoside resistance in *Escherichia coli* and *Klebsiella pneumoniae* isolates from two Taiwanese hospitals. *J Antimicrob Chemother* 54:1007-1012.
- Zhang L, Fu Y, Xiong Z, Ma Y, Wei Y, Qu X, Zhang H, Zhang J, Liao M (2018) Highly prevalent multidrug-resistant *Salmonella* from chicken and pork meat at retail markets in Guangdong, China. *Front Microbiol*9:2104.