

Journal of the Hellenic Veterinary Medical Society

Vol 72, No 4 (2021)



Epidemiology of digestive parasites in buffaloes from Romania

DA BĂRBURAȘ, A GYORKE, LM POP, R BĂRBURAȘ,
V MIRCEAN, V COZMA

doi: [10.12681/jhvms.29369](https://doi.org/10.12681/jhvms.29369)

Copyright © 2022, DA BĂRBURAȘ, A GYORKE, LM POP, R
BĂRBURAȘ, V MIRCEAN, V COZMA



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

To cite this article:

BĂRBURAȘ, D., GYORKE, A., POP, L., BĂRBURAȘ, R., MIRCEAN, V., & COZMA, V. (2022). Epidemiology of digestive parasites in buffaloes from Romania. *Journal of the Hellenic Veterinary Medical Society*, 72(4), 3329–3336. <https://doi.org/10.12681/jhvms.29369>

Epidemiology of digestive parasites in buffaloes from Romania

D.A. Bărburaș^a, A. Györke^a, L.M. Pop^{a,*}, R. Bărburaș^b, V. Mircean^a, V. Cozma^a

^aDepartment of Parasitology and Parasitic Diseases, Faculty of Veterinary Medicine, University of Agricultural Sciences and Veterinary Medicine, Cluj-Napoca, Cluj, Romania

^bS.C. MedivetBărburaș S.R.L, Românași, Salaj, Romania

ABSTRACT: Digestive parasites are a constraint on buffalo productivity. Data regarding the prevalence of parasitic diseases in buffaloes are reported in developed countries, but in Romania, they are missing. The aim of this study was to evaluate the prevalence of internal parasites in buffaloes raised in households and farms from Romania. Faeces samples were collected from 180 buffaloes of different ages (calves: 0-6 months; juveniles: 6-30 months; adults: > 30 months). The faecal samples were processed by coproparasitological methods: flotation, sedimentation, and McMaster. Digestive parasites were detected in 57.2% of buffaloes. Most buffaloes had single species infection (42.2%) ($p < 0.001$), and only 15.0% had infection with 2 (14.5%) or 3 (0.6%) parasites. The prevalence of digestive parasites was higher in calves (68.8%) and juveniles (76.9%) compared with adults (19.6%) ($p < 0.001$). *Eimeria* spp. (43.3%), *Buxtonella sulcata* (1.7%), *Fasciola hepatica* (4.4%), *Paramphistomum cervi* (2.8%), *Moniezia* spp. (0.6%), *Toxocara vitulorum* (11.7%), digestive strongyles (3.3%), *Strongyloides papillosus* (5.0%), and *Capillaria* spp. (0.6%) were identified. *Eimeria* spp. and *T. vitulorum* were the most prevalent parasites in calves and juveniles ($p < 0.001$). *F. hepatica* was the most prevalent parasite in adults. *Eimeria* spp. and *S. papillosus* were the only infections diagnosed in farmed buffaloes, with a higher prevalence ($p > 0.05$) than in household buffaloes. *B. sulcata*, *Moniezia* spp., *T. vitulorum* and *S. Papillosus* were diagnosed only in young buffaloes, while *Capillaria* spp. only in adults. The OPG of *Eimeria* spp. was higher in young buffaloes compared with adults ($p < 0.001$). Age was the risk factor identified for infections with *Eimeria* spp., *T. vitulorum* and *F. hepatica*.

Keywords: digestive parasites, buffaloes, coproparasitological examination, OPG.

Corresponding Author:

Loredana-Maria Pop, Department of Parasitology and Parasitic Diseases, Faculty of Veterinary Medicine, University of Agricultural Sciences and Veterinary Medicine, 3-5 Mănăștur Street, 400372, Cluj-Napoca, Romania
E-mail address: loredana-maria.pop@usamvcluj.ro

Date of initial submission: 26-07-2020
Date of acceptance: 10-06-2021

INTRODUCTION

The buffalo farming is continuously increasing worldwide, including in Western-European countries (Bodnar, 2017). Buffaloes are the second biggest milk-producing species in the world (Guo, 2010). Buffalo milk has a higher fat and protein content than cow's milk and it is used in the dairy industry to produce mozzarella, cream, yogurt or butter (Coroian, 2012). Romanian buffaloes are raised traditionally for milk and meat production, and to a lesser extent for draught. Romania has the second highest proportion of milk recorded buffaloes from the Mediterranean area, after Italy (Borghese, 2010). Romanian buffalo population was ranked 2nd in Europe in 2004 with 100,000 heads, but gradually decreased up to 25,000 heads in 2013 (Vidu et al., 2008; Borghese, 2013). Around 97% of buffaloes are raised in centre and north-western regions of Romania (Coroian, 2009).

Internal parasites are a constraint on productivity (Bodnar, 2017). Parasitic diseases are responsible of decreased weight gains and milk production, capricious appetite, anaemia, mortality, leading to significant economic losses (Fikruet al., 2006). Maintenance and climatic conditions are extrinsic factors that determine the presence of internal parasites (Sargison, 2016), but the intraspecific susceptibility is based on factors such as genetics, physiological, nutritional status, age and feeding mode (Bhutto et al., 2002).

Extensive data regarding parasite epidemiology is required in order to control parasitic diseases. Regular monitoring of parasites in animals is necessary to detect the early stages of infections (Gunathilaka et al., 2018).

Data on the epidemiology of internal parasites in buffaloes are reported in developed countries, however the information is scarce in developing countries (Mamun et al., 2011). The prevalence of parasites in buffaloes from Romania has not been studied so far.

Taking into consideration the importance of buffa-

lo rearing in Transylvania region, the purpose of this study was to evaluate the prevalence of internal parasites in buffaloes raised in households and farms from the North-West of Romania.

MATERIALS AND METHODS

Study area

Romania is a Southeastern European country that has a temperate continental climate, with four distinct seasons (spring, summer, autumn, and winter). The Carpathian Mountains split the territory into two groups: intra-Carpathian regions and extra-Carpathian regions. The Transylvania Depression and the western part of Romania is included in the first group and it is dominated by western moist air masses. Precipitation decreases on average from West to East and increases with altitude (A.N.M., 2008). However, in the last years, there is an increasing trend in extreme precipitation indices in north-western regions of Romania (Croitoru et al., 2016).

Animals

One-hundred-eighty buffaloes were sampled in this study. Their age varied from 2 weeks to 24.0 years with an average of 45.1 (± 5.3) months (3.8 years). The buffaloes were divided into three age groups as follows: calves ($n=101$) from 0 to 6 months old; juveniles ($n=28$) older than 6 months to 30 months (2.5 years) old; and adults ($n=58$) older than 2.5 years (> 30 months) (Table 1). The buffaloes came from five counties: Cluj, Sălaj, Bihor, Bistrița-Năsăud and Maramureș (Fig. 1). Most of the animals were raised in household system (160/180), while the rest of the animals originated from a farm in Cluj County.

The buffaloes from households were grazing on pastures nearby forests. These pastures were used by several species of animals, and all age groups. In the cold season, the animals were kept in small shelters. Milking was done manually; calves and young buffaloes were living in the same shelter as adults.


Table 1. Distribution of the study samples by age, gender, and origin

	Number	Age [mean (\pm sem)]		Gender		Origin	
		Months	Years	Females	Males	Household	Farm
Calves	77	4.0 (0.2)	-	44	33	67	10
Juveniles	52	11.7 (1.0)	-	32	20	42	10
Adults	51	141.4 (9.8)	11.8 (0.8)	46	5	51	0
Total	180	45.1 (± 5.3)	3.8 (0.5)	122	58	160	20

Table 2. Frequency, prevalence and its 95% confidence interval for identified parasites in buffaloes ($n=180$) from North-West of Romania

Infections	Frequency	Prevalence (%)	95% CI	Age Mean in months (95% CI)
<i>Eimeria</i> spp.	78	43.3***	36.3-50.6	11.7 (3.9-19.6)
<i>B. sulcata</i>	3	1.7	0.6-4.8	5.3 (1.5-9.1)
<i>F. hepatica</i>	8	4.4	2.3-8.5	93.0 (18.8-167.2)
<i>P. cervi</i>	5	2.8	1.2-6.3	52.8 (22.4-83.2)
<i>Moniezia</i> spp.	1	0.6	0.1-3.1	2
<i>T. vitulorum</i>	21	11.7	7.8-17.2	5.7 (4.3-7.1)
Strongyles	6	3.3	1.5-7.1	15.0 (-8.2-38.2)
<i>S. papillosus</i>	9	5.0	2.7-9.2	6.8 (4.1-9.4)
<i>Capillaria</i> spp.	1	0.6	0.1-3.1	60
Total	103	57.2	49.9-64.2	-

Chi-square $*p < 0.05$, $**p < 0.01$ and $***p < 0.001$

**Figure 2.**

The calves had a much higher prevalence of parasitic infections, than juvenile and adult buffaloes. Moreover, juveniles had a higher prevalence than adults. The infection with *Eimeria* spp. was the most prevalent in juveniles (59.6%) and in calves (57.2%) ($p = 0.0001$). In adults, the infection with *F. hepatica* had the highest prevalence (9.8 %) ($p = 0.004$) (Table 3).

The infection with *Eimeria* spp. was significantly more prevalent in males (56.9%) ($p = 0.02$) than in females (36.9%) (Table 3).

In both farm and household system, the infection with *Eimeria* spp. had the greatest prevalence, but without statistical significance ($p > 0.05$). Only the infections with *Eimeria* spp. and *S. papillosus* were detected in samples from buffaloes that were raised on farm (Table 3).

Single species infections with *Eimeria* spp., *F. hepatica*, *P. cervi*, *Moniezia* spp., *T. vitulorum*, strongyles and *S. papillosus* were identified. The infection with *Eimeria* spp. was more prevalent than the other single infections ($p = 0.0001$) (Table 4).

The most frequent parasitic association was that of *Eimeria* spp. and *T. vitulorum* (7.2%; 95% CI 3.9-12.03). Mixed infection with *Eimeria* spp. and *S. papillosus* was observed in 6 (3.3%) buffaloes. Mixed infection with *Eimeria* spp. and *F. hepatica* was detected in 2 (1.1%) buffaloes. Other parasitic associations identified were *Eimeria* spp. with digestive strongyles, *Eimeria* spp. with *Buxtonella sulcata*, *Eimeria* spp. with *Capillaria* spp. and *Eimeria* spp., *B. Sulcata* and *T. vitulorum*, each in a single buffalo (0.6%).



Table 3. Frequency and prevalence [$n(\%)$] of parasitic infections in buffaloes from North-West of Romania by age, gender and origin.

Infections	Age category			Gender		Origin	
	Calves <i>N</i> =77	Juveniles <i>N</i> = 52	Adults <i>N</i> = 51	Females <i>N</i> =122	Males <i>N</i> = 58	Household <i>N</i> =160	Farm <i>N</i> =20
<i>Eimeria</i> spp.	44 (57.2)***	31 (59.6)***	3 (5.9)	45 (36.9)	33 (56.9)*	66 (41.3)	12 (60.0)
<i>B. sulcata</i>	2 (2.6)	1 (1.9)	0	2 (1.6)	1 (1.7)	3 (1.9)	0
<i>F. hepatica</i>	0	3(5.8)	5 (9.8)**	6 (4.9)	2 (3.4)	8 (5.0)	0
<i>P. cervi</i>	0	2 (3.9)	3 (5.9)*	4 (3.3)	1 (1.7)	5 (3.1)	0
<i>Moniezia</i> spp.	1 (1.3)	0	0	1 (0.8)	0	1 (0.6)	0
<i>T. vitulorum</i>	14 (18.2)**	7 (13.5)**	0	13(10.7)	8 (13.8)	21 (13.1)	0
Strongyles	3 (3.9)	2 (3.9)	1 (2.0)	5 (4.1)	1 (1.7)	6 (3.8)	0
<i>S. papillosus</i>	4 (5.2)	5 (9.6)	0	7(5.7)	2 (3.4)	7 (4.4)	2 (10.0)
<i>Capillaria</i> spp.	0	0	1 (1.96)	0	1 (2.0)	1 (0.6)	0
Total	53 (68.8)***	40 (76.9)***	10 (19.6)	65 (53.3)	38 (65.5)	91 (56.9)	12 (60.0)

Chi-square * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$

Table 4. Frequency and prevalence of single and mixed parasitic infections in buffaloes from North-West of Romania.

	Frequency	Prevalence (%)	95% CI
Single infections	76	42.2	35.2-49.5
<i>Eimeria</i> spp.	53	29.44	22.9-36.68
<i>F. hepatica</i>	5	2.78	0.91-6.36
<i>P. cervi</i>	4	2.22	0.61-5.59
<i>Moniezia</i> spp.	1	0.56	0.01-3.06
<i>T. vitulorum</i>	5	2.78	0.91-6.36
Strongyles	5	2.78	0.91-6.36
<i>S. papillosus</i>	3	1.67	0.35-4.79
Mixed infections	27	15.0	10.5-20.9
<i>Eimeria</i> spp. + <i>B. sulcata</i>	1	0.56	0.01-3.06
<i>Eimeria</i> spp. + <i>Fasciola</i> spp.	2	1.11	0.13-3.96
<i>Eimeria</i> spp. + <i>Capillaria</i> spp.	1	0.56	0.01-3.06
<i>Eimeria</i> spp. + <i>T. vitulorum</i>	13	7.22	3.9-12.03
<i>Eimeria</i> spp. + <i>S. papillosus</i>	6	3.33	1.23-7.11
<i>Eimeria</i> spp. + digestive strongyles	1	0.56	0.01-3.06
<i>B. coli</i> + <i>T. vitulorum</i>	1	0.56	0.01-3.06
<i>F. hepatica</i> + <i>P. cervi</i>	1	0.56	0.01-3.06
<i>Eimeria</i> spp. + <i>B. coli</i> + <i>T. vitulorum</i>	1	0.56	0.01-3.06

**Figure 3.****Figure 4.**

OPG/EPG

The faecal egg and oocysts count per gram of faeces (EPG and OPG) are presented in Figure 3 and Figure 4: average (SE) *Eimeria* spp. 2176.6±395.1; *N. vitulorum* 4348.2 (±1643.9); strongyles 126.3 (±45.8). High values of *T. vitulorum* EPG and of *Eimeria* spp. OPG were recorded. *Eimeria* spp. OPG was significantly higher in calves (2679.8±631.8) and juveniles (1677.1±386.4) compared with adult buffaloes (132.7±54.9) ($p < 0.01$) (Fig. 3).

Risk factors

The age was identified as a risk factor for infection with digestive parasites (Table 3). The overall infection in young buffaloes was 2.7 (1.8-4.1) times more likely than in adult buffaloes. The same risk factor was identified for the infection with *Eimeria* spp. (young buffaloes OR 2.8; 95% CI: 1.8-4.3), *T. vitulorum* (young buffaloes OR 3.1; 95% CI: 1.5-6.5) and *F. hepatica* (adult buffaloes OR 3.8; 95% CI: 1.1-12.5).

DISCUSSION

In buffaloes, the parasites have a major economic impact through the high morbidity and mortality they cause. Thus, in order to design and apply effective surveillance programs, the prevalence of parasitic infections must be known (Sreedevi and Hafeez., 2014).

In the present study, the prevalence of internal parasites in buffaloes from North-Western Romania was 57.2%. In the study of Alam et al. (2016), 85.0% of examined animals were diagnosed with parasitic infections. Biswas et al. (2014) obtained a similar prevalence (84.9%). Jyoti et al. (2012) identified a prevalence of 73.6%, with coccidiosis the most prevalent infection detected.

Global prevalence of coccidiosis varies between 30 and 60%. The highest prevalence was reported in Iran (100%), Italy (100%) and Brazil (100%) (Bahrami and Alborzi, 2013; Fusco et al., 1997; Cringoli et al. 1998; Barbosa et al., 1992). In the present study, in buffaloes from North-West of Romania, the infection with *Eimeria* spp. had the highest prevalence (43.3%) from all detected infections, with significantly higher values in juveniles (59.6%) and calves (57.2%). In Brazil, a similar prevalence of coccidiosis was obtained (43.6%) (Rebouças et al., 1994). Coccidiosis mostly occurs in subclinical forms in bovines and its OPG value can vary considerably in different animals. An OPG greater than 1000 is considered as moderate infection, whilst in high infections the cattle shed

more than 1×10^4 oocysts per gram of faeces (Koutny, 2012). In the present study the OPG of buffaloes was greater than 2×10^3 , suggesting a moderate infection that can have an impact on health and productivity.

The data are scarce with respect to the infection with *B. sulcata*. Prevalence of *B. sulcata* infection was reported in Nepal (27.0%) and India (35.0%) (Adhikari et al., 2013; Kumar et al., 2017). In the present study *B. sulcata* was detected only in 1.7% of samples.

The prevalences of fasciolosis (4.4%) and paramphistomosis (2.8%) obtained in the present study are similar to other results obtained in Pakistan (4%), India (2.0%) and Italy (2.1%) (Bhutto et al., 2002; Condoleo et al., 2007; Sreedevi and Hafeez, 2014).

Similar results in terms of prevalence of infection with *Capillaria* spp. and *Moniezia* spp. were obtained in studies conducted in Bangladesh (*Capillaria* spp.: 0.4%; *Moniezia* spp.: 0.6%) (Biswas et al., 2014), Italy (*Moniezia* spp.: 0.2%) (Condoleo et al., 2007) and India (*Moniezia* spp.: 0.6%) (Sreedevi and Hafeez, 2014).

Global prevalence of *T. vitulorum* infection varies between 1.4% and 33.0%. The highest prevalence was reported in Laos (25.5%), Egypt (28.4%) and Pakistan (33.0%) (Rast et al., 2012; Abdel-Rahman and El-Ashmawy, 2013; Bhutto et al., 2002), but the prevalence value in the present study (11.7%) was much lower than these values. A faecal egg count greater than 1000 is considered highly infective for livestock (Leahy, 2017). In the present study, the buffaloes shed more than 4000 *T. vitulorum* eggs per gram, suggesting that this parasitic infection was highly damaging. Moreover, the zoonotic potential of *T. vitulorum* remains to be clarified, as *Toxocara canis* and *Toxocara cati*, parasites of carnivores, are responsible for visceral larva migrans in humans (Leahy, 2017).

Literature data on infestation with *S. papillosus* and digestive strongyles are numerous and different prevalence have been reported depending on the age category studied and geographical area. The prevalence of *S. papillosus* infection from the present study (5.0%) is similar to that obtained in Pakistan (5.9%) (Azam et al., 2002). The prevalence of infection with digestive strongyles (3.3%) is similar with that obtained in India (5.0%) (Das et al., 2017).

In the present study, calves had a much higher infection rate than juveniles and adults. Gunathilaka

found also that adults were less susceptible to parasitic infections, but the yearlings were more susceptible than calves (Gunathilaka et al., 2018).

The parasitism was higher in males than in females buffalo in the present study, but statistically significant only for the infection with *Eimeria* spp. Gunathilaka et al. observed a higher prevalence of gastro-intestinal parasites in males compared to females. According to these authors, this might be due to the attitude of farmers that focus more on female's health as they provide milk (Gunathilaka et al., 2018).

The prevalence of internal parasites was higher in buffaloes raised in household system because there is a permanent source of contamination (Radostits et al., 1994). Raising buffaloes on farm could be a method of prophylaxis that could lead to a decrease in the number of contaminated animals, and special atten-

tion should also be paid to deworming.

CONCLUSIONS

Internal parasites were detected in 57.2% of the samples of buffaloes from North-West of Romania. *Eimeria* spp., *B. sulcata*, *F. hepatica*, *P.cervi*, *Moniezia* spp., *T. vitulorum*, digestive strongyles, *S. papillosus*, and *Capillaria* spp. were the parasitic infections identified. The highest prevalence was recorded for *Eimeria* spp. and *T. vitulorum*. The buffaloes from households had greater infection rates. The calves had higher parasitic infection prevalence than juveniles and adults. Single infections were more prevalent than mixed infections.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

- Abdel-Rahman MAM., El-Ashmawy WR (2013) *Toxocara vitulorum* faeces, serum and milk of buffaloes in Giza Governorate. *Int. J. Livest. Res.* 3(2), 89-99.
- Adhikari BB, Rana HB, Sultan KMI, Devkota B, Nakao T, Kobayashi K, Sato H, Dhakal IP (2013) Prevalence of *Buxtonella sulcata* in water buffaloes and cows in Chitwan valley, Southern Nepal. *Jpn J Vet Parasitol* 11(2), 55-60.
- Administrația Națională de Meteorologie [National Meteorological Administration] (2008) *Clima României* [Climate of Romania]. Editura Academiei Române, București, Romania.
- Alam MdR., Erfan R., Sen AB, Das S, Rahman MdM., Nath SK (2016) Prevalence of gastrointestinal helminthiasis in naturally infested buffalo in Sylhet district. *Int. J. Adv. Multidiscip. Res.* 3(8), 52-58.
- Azam M, Siddiqui MM, Habib G (2002) Prevalence of parasitic infection in buffalo calves in Khadagzai, District Dir. *Pakistan Vet. J.* 22 (2).
- Bahrami S, Alborzi AR (2013) Prevalence of subclinical coccidiosis in river buffalo calves of southwest of Iran. *Acta Parasitol.* 58, 527-530.
- Barbosa MA, Blasi AC, de Oliveira MR, Corrêa FMA (1992) Natural parasitism of buffalo cows in Botucatu, SP, Brazil - III. Dynamics of gastro-intestinal parasitism in cows and calves. *Mem. Inst. Oswaldo Cruz.* 87(1), 37-41.
- Bhutto B, Phullan SM, Rind R, Soomro AH (2002) Prevalence of gastro-intestinal helminthes in buffalo calves. *J. Biol. Sci.* 2, 43-45.
- Biswas H, Dey AR, Begum N, Das PM (2014) Epidemiological aspects of gastro-intestinal parasites in buffalo in Bhola, Bangladesh. *Indian J. Anim. Sci.* 84 (3): 245-250.
- Bodnar K, Konyves T, Privoczki ZI, Jerney Z (2017) Globalization in buffalo milk production. *Scientific Research. Agricultural Management, Banat's University of Agricultural Sciences and Veterinary Medicine. Timișoara, Romania*, 19 (3).
- Borghese A (2010) Development and perspective of buffalo and buffalo market in Europe and Near East. *Proc. 9th World Buffalo Congress, Buenos Aires, 25-28 April.* In: *Revista Veterinaria.* 21 (1), 20-31.
- Borghese A (2013) Buffalo livestock and products in Europe. *Buffalo Bulletin.* 32 (1), 50-74.
- Condoleo RU, Veneziano V, Bruni G, Santaniello M, Carbone S, Pennacchio S, Rinaldi L, Cringoli G (2007) Distribution of helminths in buffalo farms from central Italy. *Ital. J. Anim. Sci.* 6(2), 920-922.
- Coroian A, Mireșan V, Coroian CO, Feștilă I, Cocan D, Răducu C, Dărăban S (2012) Production Parameters in a Romanian Buffalo Population. *Scientific Papers: Animal Science and Biotechnologies.* 45 (2).
- Coroian A (2009) Creșterea bubaline lor pentru producția de lapte. *Bioflux, Cluj-Napoca, Romania.*
- Cringoli G, Guarino A, Fusco G, Veneziano V, Rinaldi L (1998) Diffusion dynamics of *Eimeria* spp. in infected buffalo herds. *Parassitologia.* 40 (1), 38.
- Croitoru AE, Piticar A, Burada DC (2016) Changes in precipitation extremes in Romania. *Quatern. Int.* 415, 325-335.
- Das M, Deka DK, Sarmah AK, Sarmah PC, Islam S (2017) Gastrointestinal parasitic infections in cattle and swamp buffalo of Guwahati, Assam, India. *Indian J. Anim. Res.* B-3427.
- Fikru R, Teshale S, Reta D, Yosef K (2006) Epidemiology of gastrointestinal parasites of ruminants in Western Oromia, Ethiopia. *Int. J. Appl. Res. Vet. M.* 4, 51-57.
- Fusco G, Guarino A, Merola A, Veneziano V, Cringoli G (1997) Natural diffusion of *Eimeria* spp. in buffalo calves. *Proceedings of the 5th World Buffalo Congress. Royal Palace, Caserta, Italy, October 13-16.* 92, 569-573.
- Gunathilaka N, Niroshana D, Amarasinghe D, Udayanga L (2018) Prevalence of gastrointestinal parasitic infections and assessment of Deworming Program among cattle and buffaloes in Gampaha District, Sri Lanka. *Biomed Res Int.* 3048373.
- Guo M, Hendricks G (2010) Improving buffalo milk. In: Griffiths, M. (Ed.), *Improving the safety and quality of milk: improving quality in milk products.* Woodhead Publishing, Cambridge, UK, pp. 402-416.
- Jyoti N, Singh K, Juyal PD, Haque M, Rath SS (2012) Epidemiology of gastrointestinal parasites in buffalo calves of Punjab state. *J. Vet. Parasitol.* 26(1), 19-22.
- Koutny H, Joachim A, Tichy A, Baumgartner W (2012) Bovine *Eimeria* species in Austria. *Parasitol Res.* 110(5), 1893-1901.
- Kumar B, Biswa RM, Amit P, Joice PJ, Bhavika RP (2017) Incidence of *Buxtonella sulcata* in Jaffrabadi buffaloes of south-western Gujarat, India. *Buffalo Bulletin,* 36(4), 623-628.
- Leahy E, Bronsvoort B, Gamble L, Gibson A, Kaponda H, Mayer D, Mazeri S, Shervell K, Sargison N (2017) Proof of concept of faecal egg nematode counting as a practical means of veterinary engagement with planned livestock health management in a lower income country. *Ir. Vet. J.* 70, 16, doi:10.1186/s13620-017-0094-9.
- Mamun MAA, Begum N, Mondal MMH (2011) A coprological survey of gastro-intestinal parasites of water buffaloes (*Bubalus bubalis*)

- in Kurigram district of Bangladesh. J. Bangladesh Agril. Univ. 9(1), 103-109.
- Zajac A. M., Conboy G. A. (2012) Veterinary clinical parasitology. 8th ed. John Wiley & Sons, UK.
- Radostits OM, Blood DC, Gay CC (1994) Veterinary Medicine: A text book of disease of cattle, sheep, pigs, goats and horse. 8th ed. Baillere Tindall Publication. London, 1223-1225, 1237-1238.
- Rast L, Lee S, Nampanya S, Jenny-Ann LMLT, Khounsy S, Windsor PA (2012) Prevalence and clinical impact of *Toxocara vitulorum* in cattle and buffalo calves in northern Lao PDR. Trop Anim Health Prod. 45,539-546.
- Rebouças MM, Grasso LMPS, Spósito Filha E, do Amaral V, Santos SM, Silva DM (1994) Prevalência e distribuição de protozoários do gênero *Eimeria* (Apicomplexa: *Eimeriidae*) em bovinos nos municípios de Altinópolis, Taquaritinga, São Carlos e Guaira - estado de São Paulo, Brasil [in Portuguese]. Rev. Bras. Parasitol. Vet. 3, 125-130.
- Sargison ND (2016) Keys to solving health problems in small ruminants: Anthelmintic resistance as a threat to sustainable nematode control. Small Ruminant Res. 142, 11-15.
- Sergeant ESG (2018) Epitools Epidemiological Calculators. Aus vet. Available at: <http://epitools.ausvet.com.au>.
- Sreedevi C, Hafeez Md.(2014) Prevalence of gastrointestinal parasites in buffaloes (*Bubalus Bubalis*) in and around Tirupati, India. Buffalo Bulletin.33(3).
- Vidu L, Popa R, GrasM (2008) Stadiul, direcțiile și obiectivele ameliorării bubalinelor din România [The stage, directions and objectives of buffalo breeding improvement in Romania]. Alpha Mdn, Buzău, Romania.