Journal of the Hellenic Veterinary Medical Society

Vol 72, No 4 (2021)

Ameliorative effect of tocotrienol and selenium yeast against the adverse effect of florfenicol in broilers' liver

AI HOSNY, MH KHAIRY, AM ASY, EA ABOZEID

doi: [10.12681/jhvms.29399](https://doi.org/10.12681/jhvms.29399)

Copyright © 2022, AI HOSNY, MH KHAIRY, AM ASY, EA ABOZEID

This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

To cite this article:

HOSNY, A., KHAIRY, M., ASY, A., & ABOZEID, E. (2022). Ameliorative effect of tocotrienol and selenium yeast against the adverse effect of florfenicol in broilers' liver. *Journal of the Hellenic Veterinary Medical Society*, 72(4), 3481–3490. <https://doi.org/10.12681/jhvms.29399>

Ameliorative effect of tocotrienol and selenium yeast against the adverse effect of florfenicol in broilers' liver

A.I. Hosny¹, M.H. Khairy¹, A.M. Asy², E.A. Abozeid²

¹ Department of Pharmacology, Faculty of Veterinary Medicine, Zagazig University, Zagazig, Egypt

² Department of Biochemistry, Animal Health Research Institute, Benha-Branch, ARC, Dokki, Giza, Egypt

ABSTRACT: The prolonged use of florfenicol can lead to detrimental side effects in poultry. This work focuses on the role of tocotrienol and selenium yeast to mitigate the adverse effects of florfenicol in broilers' liver. One hundred and fifty, one-day-old Cobb broiler chicks were equally divided in 5 experimental groups according to the following experimental design: Group (1) control group chicks fed to a balanced diet only. Group (2) Chicks treated with florfenicol (20 mg/kg b.w.) per bird for 3 successive days and the florfenicol was administered in other groups by the same dose and for the same period of time. Group (3) chicks treated with florfenicol and tocotrienol (170 mg/kg b.w.) for 7 successive days. Group (4) Chicks treated with florfenicol and selenium yeast (0.15mg /kg b.w., on feed) for 7 successive days. Group (5) Chicks treated with a combination of florfenicol, tocotrienol and selenium yeast. Chickens treated with florfenicol exhibited an increased level in hepatic malondialdehyde (MDA), as well as a decreased level in hepatic superoxide dismutase (SOD) and reduced glutathione (GSH). Tocotrienol and selenium yeast decreased the MDA and increased SOD and GSH in hepatic tissue as well as return ALP, cholesterol, triglyceride and VLDL to their normal levels. Treated chicks with tocotrienol returned serum (ALT) to normal activity but serum total protein and albumin levels were increased. The selenium yeast treated groups showed an increase of serum total globulin. Histopathologically, florfenicol treated group had focal hepatic leukocytic infiltration and focal coagulative necrosis of hepatocytes but chickens with the combination of tocotrienol and selenium yeast had activated Kupffer cells and revealed less evident necrotic changes in liver. In conclusion, tocotrienol and selenium yeast administered alone or in combination highlighted improved antioxidant effects and mitigated the lipid peroxidation in broilers' liver treated with florfenicol. Thus, tocotrienol and selenium yeast can improve the safety of using florfenicol in broiler chickens under experimental conditions.

Keywords: Florfenicol, Tocotrienol, Selenium yeast, Liver, Antioxidants, Broilers.

Corresponding Author:

Eman Abd-El Moneim Abozeid, Department of Biochemistry, Toxicology and Nutritional Deficiency, Animal Health Research Institute, Benha-Branch, ARI, Giza, Dokki, Egypt
E-mail address: eman.abozeid89@gmail.com

Date of initial submission: 10-11-2020
Date of acceptance: 14-06-2021

INTRODUCTION

Florfenicol, a synthetic broad -spectrum antibiotic, contains a fluorine atom at the 3-carbon position, instead of the hydroxyl group found in thiamphenicol used in veterinary practice to treat most gram-positive and gram-negative bacteria (Chang et al., 2010). Florfenicol action is mainly bacteriostatic by inhibition protein synthesis of bacteria by binding to 50S and 70S subunits ribosome and abolishes the activity of peptidyl transferase (Khalil et al., 2012). Ronette (2012) reported that florfenicol is used for the treatment of many microorganisms which affect poultry such as *Escherichia coli*, *Klebsiella*, *Ornithobacterium rhinotracheale*, *Pasteurella multocida*, *Mannheimia haemolytica*, *Enterobacter cloacae*, *Haemophilus somnus*, *pneumonia*, *Salmonella typhi*, *Shigella dysenteriae* and *Staphylococcus aureus*. Florfenicol usually made biotransformation effect (more concentrations of reactive oxygen species) with biochemical changes after entering into cells because it is a highly lipophilic drug. Florfenicol damage changes differ according to the dosage of the drug, time of administration and animal (Ren et al., 2014). Florfenicol administration can produce oxidative stress in broiler's liver by inhibiting the expression of antioxidant proteins nuclear factor-erythroid 2-related factor 2 (Nrf2), hemeoxygenase-1 (HO-1) and NAD(P)H dehydrogenase quinone-1 (Han et al., 2020). In addition Wang et al. (2020) reported that administration of florfenicol by different doses (0.15, 0.3, 0.6, 1.2 and 1.8 g/L) in drinking water of broilers for five successive days produced oxidative stress effect on chicks through inhibition of the expression of related factors in Nrf2.

Antioxidants are substances that interact with unstable free radicals and prevent there damage effects (Sies, 1997; Olayinka et al., 2012). The antioxidants can be classified to natural and synthetic. Natural antioxidants are vitamin E, organic selenium, vitamin C and beta carotene. Synthetic antioxidants are butylated hydroxyanisole and butylatedhydroxytoluene (Hurrell, 2003).

Tocotrienols are the primary form of vitamin E which has α , β , γ , and δ forms. The main sources of tocotrienols are palm oil extract from *Elaeis guineensis* (African oil palm) contain more than 800 mg/kg of tocotrienol and cereal grains such as wheat, barley, rice and differ from tocopherol by the presence of three trans-double bonds in the hydrocarbon tail. Tocotrienols have hypo cholesterolemic, neuropro-

tection and anticancer effects (Sen et al., 2006). Tocotrienol rich fraction from palm oil can protect cellular membranes from damage effect and it can inhibit protein oxidation and lipid peroxidation in rat liver microsomes (Kamat et al., 1997). Palmitate oil (tocotrienol containing diet) displayed significant decrease in malondialdehyde MDA level with raises superoxide dismutase SOD activity and reduced glutathione GSH level (Khan et al., 2011). Selenium yeast is one of the best organic sources of selenium for poultry and other farmed animals which was approved for chicken consumption in June 2000 by the United States Food and Drug Administration US-FDA, and has a potent antioxidant effect (Wang and Xu, 2008). Elevation selenium level in the diet was led to lowering lipid peroxidation products, free radical elimination and protecting cell membranes (Fan et al., 2009). Diets supplemented with selenium yeast improve activity of hepatic glutathione peroxidase and help for production of oxidized glutathione then stimulate glutathione reductase, which prevent its deactivation by NADPH (Upton et al. 2009). The prolonged use of florfenicol can lead to detrimental side effects in poultry. This work focuses on the role of tocotrienol and selenium yeast alone or their combination to mitigate these side effects in broilers' liver.

MATERIAL AND METHODS

Animals

One hundred and fifty, one-day-old, unsexed Cobb broiler chicks were used and purchased from El-Watania Poultry Company - Cairo - Egypt. The birds were allocated in separate units of metal wire-floored battery for five successive weeks. The study was approved by the Ethical Committee for care and use of animals at Animal Health Research Institute Benha Branch, Egypt (25/3/2019).

Drugs

Three drugs were used in this experimental model. Firstly, florfenicol 10% (Floricol[®], PharmaSwede Co., Egypt) was administrated for 3 successive days at 20 mg/kg according to the instructions of manufacture. Each ml of the product contained 100 mg of florfenicol base. Secondly, tocotrienol 50% (Tocovid[®], Hovid Company, Malaysia) was administrated for 7 successive days, at the recommended dose of 170 mg/kg. Finally, selenium yeast 0.2% (Bio-SEL 2000[®], IBEX International Co. LTD, Egypt) was administrated at 0.15 mg/kg for 7 successive days.

Experimental design

One hundred and fifty (150), apparently healthy, one-day-old, unsexed, Cobb broiler chicks were used. The chicks were housed in clean and disinfected enclosure, with controlled environmental temperature, and fed with a well-balanced ration throughout the experimental period of five successive weeks. The chicks were equally divided into five groups of 30 chicks each. Group 1, was the negative control group, including chicks which fed with balanced diet only. Chicks in Group 2 were treated with florfenicol (20 mg/kg) for 3 successive days (15th - 17th day of age) by drinking water. Group 3 was treated with florfenicol and tocotrienol (170 mg/kg, per os) for 7 successive days (15th - 21st day of age). Group 4 was treated with florfenicol and selenium yeast (0.15 mg /kg, in feed) for 7 successive days (15th - 21st day of age). Group 5 was treated with the combination from florfenicol, tocotrienol and selenium yeast for 7 successive days (15th - 21st day of age).

Blood samples were collected from jugular vein of five birds of each group on the 18th, 25th and 35th days of age. The liver of chicks was divided into two parts. One part was obtained immediately for fixed with 10% formalin solution for 48h and the another part was stored in at -20°C for determination of anti-oxidant /oxidant status in the hepatic tissue on the 18th, 25th and 35th days of age.

Biochemical analysis (Liver function tests)

Blood samples were collected and sera were separated. The sera were stored -20°C until examination to evaluate alanine aminotransferase (ALT) and aspartate aminotransferase (AST) (Varli, 1974), alkaline Phosphatase ALP (Belfield and Goldberg, 1971), total protein (TP) (Domas, 1975), serum albumin (Dumas, 1971), serum total globulin (Coles, 1974), serum total cholesterol (Flegg, 1973), triglycerides (TG), high-density lipoprotein cholesterol (HDL-C) (Gordon et al., 1977), and low-density lipoprotein cholesterol (LDL-C) (Friedewald et al., 1972) were estimated. Very low-density lipoprotein cholesterol (VLDL-C) was calculated by dividing triglycerides value by 5 (Tietz, 1976). Low-density lipoprotein cholesterol (LDL-C) was calculated by the following equation: $LDL-C = (Total\ cholesterol) - (HDL-C) - (VLDL-C)$ (Ashayerizadeh et al., 2009).

Evaluation of antioxidant and oxidant status in hepatic tissue

Liver tissues from each chick were collected im-

mediately and stored in low temperatures for reserve. Prior to dissection, perfuse hepatic tissues with phosphate buffered saline solution (PH 7.4) containing heparin (0.16 mg/ml) to remove any blood cells and clots. Homogenize the tissue in 5 -10 ml cold buffer (I, e, 50mM potassium phosphate, pH 7.5.1 mM EDTA) per gram tissue, using tissue homogenizer. Centrifuge at 4000 rpm for 5 minutes at 4°C. Remove the supernatant for assay and store on ice. If not assaying on the same day, freeze the sample at - 80°C. The sample will be stable for at least one month. The prepared samples were ready to evaluate superoxide dismutase (SOD) activity (Nishikimi et al., 1972), reduced glutathione (GSH) concentration (Beutler et al., 1963) and malondialdehyde (MDA) level (Ohkawa et al., 1979).

Histopathological examination of liver tissue

Liver samples were fixed in 10% formalin solution for 48 hours for further histopathological examination following standard methodologies to obtain hematoxylin and eosin-stained slides (Bancroft and Stevens, 1977).

Statistical analysis

Statistical analysis was conducted with the Statistical Package for Social Science (SPSS Inc. Released, 2009) to determine if variables differed between groups, according to Snedecor and Cochran, 1989. The Shapiro-Wilk test was used to test the normal distribution of the data before statistical analysis was performed. Analysis of variance was conducted by one-way ANOVA and compare between means were conducted by Duncan's multiple range test (Duncan, 1955). Probability values of less than 5 % ($P < 0.05$) were considered a significant finding.

RESULTS

The blood biochemical parameters (liver enzymes) in broilers treated with florfenicol showed a significant ($P < 0.05$) increase in serum ALT on the 25th day of age and no significant ($P > 0.05$) differences on the 35th day of age while a significant ($P < 0.05$) increase in serum ALP activity when compared with normal group. Other treated group returned serum ALT and ALP activities to normal levels as shown in Table 1.

On the 25th day of age, florfenicol treated group showed significant ($P < 0.05$) decreases in serum TP, albumin and serum total globulin when compared with control group. Other groups displayed significant ($P < 0.05$) increases in serum TP and albumin

when compared with florfenicol treated group. Selenium yeast and florfenicol treated group alone and the group treated with its combination with tocotrienol displayed a significant ($P < 0.05$) increase in serum total globulin when compared with florfenicol treated group as shown in Table1.

Florfenicol and tocotrienol treated group showed a significant ($P < 0.05$) decrease in serum cholesterol and LDL levels on the 18th day of age with no significant ($P > 0.05$) differences on the 25th day of age but triglycerides returned to normal on the 18th and 25th day of age when compared with control group. Selenium yeast and florfenicol treated group alone and the group treated with its combination with tocotrienol returned to normal lipid profile levels as shown in Table1.

Florfenicol increased MDA level in hepatic tissue, but significantly ($P < 0.05$) decrease SOD activity and GSH concentration in hepatic tissue when compared with the control group. Florfenicol, tocotrienol and selenium yeast either alone or in combination showed a significant ($P < 0.05$) decrease in MDA as well as significant ($P < 0.05$) increases in SOD and GSH in hepatic tissue when compared with florfenicol treated

group as shown in Table2.

For histopathological examination five samples from 30 birds (5/30) per group were collected on the 18th, 25th and 35th days of age. As shown in Fig. 1A, the liver section of the negative control group revealed normal hepatic architecture. Liver sections of florfenicol treated group showed focal leukocytic infiltration in the hepatic tissue and focal coagulative necrosis of hepatocytes represented by deeply eosinophilic cytoplasm of the nuclei with karyorrhexis and karyolysis in the liver section of chicks (Fig. 1B). Liver sections of florfenicol with tocotrienol treated group displayed fatty degeneration of some hepatic cells, with eccentric location of the nuclei on the 35th day of age and focal mononuclear cell infiltration of the hepatic tissue and activation of the Kupffer cells on the 35th day of age (Fig. 1C). Liver sections of florfenicol and selenium yeast treated group revealed focal coagulation hepatocellular necrosis with karyolysis of the nuclei and ballooning degeneration represented by empty cytoplasm and centrally located nuclei in the adjacent hepatocytes (Fig. 1D). Liver sections of florfenicol, tocotrienol and selenium yeast treated group showed activation of Kupffer cells and less evident necrotic changes (Fig. 1E).

Figure 1. Histopathological analysis of liver tissue. Note: Liver tissues were fixed and stained with H&E. A) Control group 200X, B) Florfenicol treated group showed focal leukocytic infiltration in the hepatic tissue (star) focal coagulation necrosis of hepatocytes (black arrow) 400X, C) Florfenicol and tocotrienol treated group showed activations of Kupffer cells (black arrow) focal mononuclear cell infiltration of the hepatic tissue (star) 400X, D) Florfenicol and selenium yeast treated group showed focal coagulation necrosis (star) and ballooning degeneration (black arrow) 200X, E) Florfenicol, tocotrienol and selenium yeast treated group showed activation of Kupffer cells (black arrow) and less evident necrotic changes. 200X

Table 1: Effect of oral administration of tocotrienol and, selenium yeast on blood biochemical parameters of broilers exposed to adverse effect of florfenicol. (Mean $\pm$ SEM)

Parameters	Experimental Periods	Groups				
		1	2	3	4	5
ALT (U/L)	18 day	11.00 $\pm$.58 ^a	6.50 $\pm$.29 ^b	5.00 $\pm$.58 ^b	10.50 $\pm$.29 ^a	10.00 $\pm$.58 ^a
	25 day	4.50 $\pm$.29 ^c	10.00 $\pm$.29 ^{ab}	3.50 $\pm$.29 ^c	9.00 $\pm$ 1.15 ^b	11.50 $\pm$.29 ^a
	35 day	12.50 $\pm$.29 ^b	15.00 $\pm$ 1.15 ^{ab}	12.00 $\pm$ 1.73 ^b	16.50 $\pm$.87 ^{ab}	17.50 $\pm$ 2.02 ^a
AST (U/L)	18 day	12.50 $\pm$.87 ^a	12.00 $\pm$.00 ^a	12.00 $\pm$.58 ^a	12.00 $\pm$.58 ^a	12.50 $\pm$.29 ^a
	25 day	16.00 $\pm$ 1.73 ^{bc}	11.00 $\pm$ 1.15 ^c	14.00 $\pm$ 2.31 ^{bc}	21.00 $\pm$ 3.46 ^{ab}	25.00 $\pm$ 2.89 ^a
	35 day	21.5 $\pm$.87 ^a	22.00 $\pm$ 1.73 ^a	23.00 $\pm$.58 ^a	27.50 $\pm$ 3.75 ^a	27.50 $\pm$ 4.33 ^a
ALP (U/L)	18 day	522.50 $\pm$ 17.03 ^b	608.00 $\pm$ 23.67 ^a	600.00 $\pm$ 10.39 ^{ab}	595.00 $\pm$ 42.15 ^{ab}	672.00 $\pm$ 20.21 ^a
	25 day	660.00 $\pm$ 7.50 ^b	735.50 $\pm$ 18.76 ^a	733.50 $\pm$ 10.68 ^a	680.00 $\pm$ 1.15 ^b	680.00 $\pm$ 9.81 ^b
	35 day	618.50 $\pm$ 3.50 ^{ab}	708.50 $\pm$ 20.50 ^a	540.50 $\pm$ 22.50 ^b	594.00 $\pm$ 158.00 ^{ab}	519.00 $\pm$ 158.00 ^b
Total Protein (g/dl)	18 day	3.55 $\pm$.043 ^a	3.55 $\pm$.043 ^a	3.33 $\pm$.04 ^{ab}	3.25 $\pm$.07 ^b	3.49 $\pm$.14 ^{ab}
	25 day	4.27 $\pm$.12 ^b	3.25 $\pm$.12 ^d	3.62 $\pm$.05 ^c	3.99 $\pm$.08 ^b	4.79 $\pm$.04 ^a
	35 day	3.60 $\pm$.058 ^{ab}	3.35 $\pm$.09 ^b	3.45 $\pm$.09 ^b	4.00 $\pm$.23 ^a	4.00 $\pm$.17 ^a
Albumin (g/dl)	18 day	1.89 $\pm$.07 ^a	1.78 $\pm$.012 ^{ab}	1.70 $\pm$.017 ^{bc}	1.63 $\pm$.003 ^c	1.80 $\pm$.055 ^{ab}
	25 day	2.26 $\pm$.04 ^a	1.63 $\pm$.04 ^c	2.03 $\pm$.12 ^b	2.00 $\pm$.017 ^b	2.13 $\pm$.032 ^{ab}
	35 day	2.00 $\pm$.06 ^a	1.80 $\pm$.06 ^a	1.95 $\pm$.03 ^a	2.15 $\pm$.20 ^a	2.15 $\pm$.14 ^a
Globulin (g/dl)	18 day	1.65 $\pm$.02 ^a	1.75 $\pm$.01 ^a	1.63 $\pm$.02 ^a	1.62 $\pm$.07 ^a	1.69 $\pm$.08 ^a
	25 day	2.01 $\pm$.075 ^b	1.61 $\pm$.08 ^c	1.58 $\pm$.06 ^c	1.99 $\pm$.06 ^b	2.66 $\pm$.07 ^a
	35 day	1.60 $\pm$.00 ^b	1.55 $\pm$.03 ^b	1.50 $\pm$.11 ^b	1.85 $\pm$.03 ^a	1.85 $\pm$.03 ^a
A/G ratio	18 day	1.15 $\pm$.05 ^a	1.02 $\pm$.00 ^b	1.04 $\pm$.00 ^{ab}	1.00 $\pm$.05 ^b	1.07 $\pm$.02 ^{ab}
	25 day	1.12 $\pm$.02 ^{ab}	1.01 $\pm$.03 ^b	1.29 $\pm$.13 ^a	1.00 $\pm$.03 ^b	0.8 $\pm$.03 ^c
	35 day	1.25 $\pm$.03 ^a	1.16 $\pm$.02 ^a	1.32 $\pm$.12 ^a	1.16 $\pm$.09 ^a	1.16 $\pm$.05 ^a
Total Cholesterol (mg/dl)	18 day	129.00 $\pm$ 3.46 ^a	138.00 $\pm$ 9.23 ^a	102.50 $\pm$.29 ^b	127.00 $\pm$ 1.15 ^a	123.50 $\pm$ 2.02 ^a
	25 day	91.00 $\pm$ 10.97 ^{ab}	63.50 $\pm$.87 ^b	69.50 $\pm$ 3.17 ^b	91.00 $\pm$ 15.58 ^{ab}	95.00 $\pm$ 2.88 ^{ab}
	35 day	55.00 $\pm$ 1.73 ^c	61.00 $\pm$ 4.04 ^{bc}	75.50 $\pm$ 9.53 ^{ab}	73.00 $\pm$ 4.04 ^{ab}	69.50 $\pm$ 3.75 ^{abc}
Triglycerides (mg/dl)	18 day	101.50 $\pm$ 7.79 ^b	133.00 $\pm$ 5.77 ^a	100.50 $\pm$ 3.18 ^b	101.00 $\pm$ 1.15 ^b	94.50 $\pm$ 1.44 ^b
	25 day	77.50 $\pm$ 4.90 ^a	46.00 $\pm$ 1.73 ^b	67.50 $\pm$ 3.18 ^a	64.00 $\pm$ 7.51 ^a	63.50 $\pm$ 4.33 ^a
	35 day	75.50 $\pm$ 2.50 ^b	69.50 $\pm$ 1.44 ^b	71.50 $\pm$ 6.06 ^b	79.00 $\pm$ 2.30 ^{ab}	78.50 $\pm$ 2.02 ^b
HDL-C(mg/dl)	18 day	41.00 $\pm$.58 ^a	39.00 $\pm$ 1.15 ^{ab}	42.00 $\pm$ 1.73 ^a	41.50 $\pm$ 3.18 ^a	34.50 $\pm$.29 ^b
	25 day	38.50 $\pm$.87 ^a	33.00 $\pm$ 2.31 ^{ab}	34.50 $\pm$ 2.02 ^{ab}	32.00 $\pm$ 2.31 ^b	31.50 $\pm$.87 ^b
	35 day	38.00 $\pm$.58 ^b	41.00 $\pm$.58 ^{ab}	31.50 $\pm$ 1.44 ^c	42.50 $\pm$.29 ^a	37.00 $\pm$.58 ^b
LDL-C (mg/dl)	18 day	67.70 $\pm$ 2.30 ^a	72.40 $\pm$ 16.00 ^a	40.40 $\pm$ 4.600 ^b	65.30 $\pm$ 7.100 ^a	70.10 $\pm$ 3.500 ^a
	25 day	37.00 $\pm$ 15.80 ^{ab}	21.30 $\pm$ 6.10 ^b	21.50 $\pm$.90 ^b	46.20 $\pm$ 20.4 ^a	58.80 $\pm$ 8.00 ^a
	35 day	13.90 $\pm$ 1.50 ^b	15.10 $\pm$ 5.50 ^b	29.70 $\pm$ 11.90 ^a	14.50 $\pm$ 5.70 ^b	16.80 $\pm$ 4.80 ^b
VLDL-C (mg/dl)	18 day	20.30 $\pm$ 2.70 ^b	26.60 $\pm$ 2.00 ^a	20.10 $\pm$ 1.10 ^b	20.20 $\pm$.40 ^b	18.90 $\pm$.50 ^b
	25 day	15.50 $\pm$ 1.70 ^a	9.20 $\pm$.60 ^b	13.50 $\pm$ 1.10 ^a	12.80 $\pm$ 2.60 ^a	12.70 $\pm$ 1.50 ^a
	35 day	15.10 $\pm$.50 ^b	13.90 $\pm$.50 ^b	14.30 $\pm$ 2.10 ^b	17.00 $\pm$.80 ^{ab}	15.70 $\pm$.70 ^b

^{ab} Mean values within the same row with different superscript letter are statistically different at $P \leq 0.05$. SEM = Standard Error of Means. 1) Control group 2) Florfenicol treated group 3) Florfenicol and Tocotrienol treated group 4) Florfenicol and Selenium yeast treated group 5) Florfenicol and Tocotrienol with Selenium yeast treated group.

(ALT: Alanine aminotransferase, AST: Aspartate aminotransferase, ALP: Alkaline Phosphatase, A/G: Albumin/Globulin ratio, HDL: High-density lipoprotein, LDL: Low-density lipoprotein, VLDL: Very low-density lipoprotein).

Table 2: Effect of oral administration of tocotrienol and selenium yeast on hepatic antioxidant/oxidant Status in broilers exposed to adverse effect of florfenicol. (Mean $\pm$ SEM) (n=5)

Parameters	Experimental Periods	Groups				
		1	2	3	4	5
Hepatic SOD (mg/dl)	18 day	80.39 $\pm$ 7.33 ^a	46.72 $\pm$ 2.76 ^{cd}	62.57 $\pm$ 1.76 ^{bc}	60.71 $\pm$ 2.33 ^{bc}	68.79 $\pm$ 6.58 ^{ab}
	25 day	77.29 $\pm$ 8.75 ^a	47.78 $\pm$ 6.17 ^b	80.85 $\pm$ 15.67 ^a	82.55 $\pm$ 2.63 ^a	85.68 $\pm$ 4.24 ^a
	35 day	53.61 $\pm$.002 ^a	39.28 $\pm$ 5.18 ^b	61.42 $\pm$ 1.91 ^a	53.56 $\pm$.26 ^a	57.10 $\pm$ 4.41 ^a
Hepatic GSH (mg/dl)	18 day	4.76 $\pm$.06 ^a	3.88 $\pm$.069 ^c	4.68 $\pm$.037 ^a	4.62 $\pm$.133 ^a	4.18 $\pm$.032 ^b
	25 day	4.14 $\pm$.26 ^b	3.36 $\pm$.020 ^c	3.58 $\pm$.254 ^c	4.70 $\pm$.069 ^a	4.66 $\pm$.092 ^{ab}
	35 day	4.83 $\pm$.060 ^d	4.03 $\pm$.012 ^c	5.56 $\pm$.057 ^{bc}	5.66 $\pm$.10 ^a	5.32 $\pm$.008 ^b
Hepatic MDA (mg/dl)	18 day	7.82 $\pm$.10 ^b	9.95 $\pm$.32 ^a	7.95 $\pm$.40 ^b	5.95 $\pm$.46 ^c	7.00 $\pm$.72 ^{bc}
	25 day	6.25 $\pm$.144 ^b	7.50 $\pm$.69 ^a	7.67 $\pm$.043 ^a	5.83 $\pm$.13 ^b	6.85 $\pm$.086 ^{ab}
	35 day	5.20 $\pm$.12 ^c	6.18 $\pm$.10 ^a	5.56 $\pm$.057 ^{bc}	5.48 $\pm$.21 ^{bc}	5.97 $\pm$.30 ^{ab}

^{ab}Mean values within the same row with different superscript letter are statistically different at $P \leq 0.05$. SEM = Standard Error of Means. 1) Control group 2) Florfenicol treated group 3) Florfenicol and Tocotrienol treated group 4) Florfenicol and Selenium yeast treated group 5) Florfenicol and Tocotrienol with Selenium yeast treated group.

(SOD: Superoxide dismutase, GSH: Reduced glutathione, MDA: malondialdehyde).

DISCUSSION

Florfenicol administration has a damage effect in liver of broiler chickens so tocotrienol and selenium yeast used to mitigate these side effects. In serum biochemical examination, ALP activity returned to normal level in all groups except in the florfenicol treated group. This result was previously explained by Hasan et al. (2018) who stated that tocotrienol derived from seeds of *Bixa orellana* enhance bone structure and bone strength with decrease bone resorption and improve bone formation by increased activity of ALP. Moreover, Norazlina et al. (2002) reported that tocotrienol plays an important role in bone calcification. ALP has a critical role in bone calcification. Therefore, tocotrienol may improve the ALP activity. Our results related to selenium yeast were in accordance with those obtained by Yang et al. (2012) and Invernizzi et al. (2013) in maintaining ALP activity within normal. On the other hand, a significant increase in serum ALP activity of rats fed ontocotrienol (120-130 mg/kg b.w.) for 13 weeks was previously reported by Nakamura et al., 2001. This elevation may be caused by cholestasis or bone remodeling.

Albumin is one of main protein sources formed in the liver to maintain plasma osmotic pressure, providing energy and repairing tissue. Albumin is considered a carrier of nutrients to maintain the body tissue protein dynamic balance (Ahmed et al., 2002). In our results on the 25th day of age, all treated groups

showed significant increases in serum TP and serum albumin when compared with the florfenicol treated group. These results were agreed by who reported with Shi et al., (2018) stating that diets supplemented with selenium can improve metabolism of major elements due to elevation of total protein and globulin. Our results related totocotrienol disagreed with those previously described by Tasakia et al. (2008) and Shibata et al. (2012), while our results regarding selenium yeast agreed with those previously described by El-Demerdash and Nasr (2014) and Shi et al. (2018) and disagree with those previously reported by Attia et al. (2010), Invernizzi et al. (2013) and Liu et al.(2020) in serum albumin characterization.

On the 25th day of age, the florfenicol treated group showed significant decreases in serum total globulin when compared with the control group, which is in agreement with Shaheen and El-Far (2013). The immunosuppressive effects of florfenicol could be attributed to its protein inhibition. However, our results disagree with those previously published by Allam et al. (2014) who reported a significant increase in serum total globulin of Pekin ducklings.

The selenium yeast treated group alone and combined with tocotrienol displayed a significant increase in serum total globulin when compared with the florfenicol treated group. This result agree with that mentioned by Shi et al. (2018) who proved that elevation

of serum globulines enhance immune system of animals. In addition, these results were unconformity with those of Yang et al. (2012) who showed no significant difference in serum total globulin in broilers fed on 3 ppm selenium yeast from 0 to 3 weeks of age.

The tocotrienol treated group showed a significant decrease in serum cholesterol and LDL levels on the 18th day of age when compared with the control group. The selenium yeast treated group alone and combined with tocotrienol returned the lipid profile levels to normal. The hypocholesterolemic effect of tocotrienol may be attributed to the decrease activity of β -hydroxy- β -methylglutaryl coenzyme A reductase (Yuet al., 2006). These results referred to tocotrienol agreed with those previously described by Qureshi and Peterson (2001), Yu et al. (2006) and Budin et al. (2009). Our results involving selenium yeast are similar with those published by Sevcikova et al. (2008) and Yang et al. (2012). Our results were not in agreement with Hasselwander et al. (2001) who reported that tocotrienol had little effect on serum lipid levels. Attia et al. (2010) showed significant decreases in serum cholesterol level and triglycerides after selenium yeast treatment.

Florfenicol treated group showed a significant increase of MDA level and significant decrease of SOD activity and GSH concentration in hepatic tissues in comparison with the control group. These results attributed to the ability of florfenicol to inhibit the expression of antioxidant proteins Nrf2, HO-1 and NQO-1 resulted in decrease antioxidant factors SOD and GSH (Han et al., 2020). Firozian et al. (2020) reported that elevation of lipid peroxidation is usually joined by decrease GSH and SOD antioxidant factors. GSH depletion can cause oxidative damage but SOD can directly inhibit reactive oxygen species (ROS) formations which they are considered an important antioxidant defense enzymes. These results are in agreement with those described by Farombi et al. (2001) who reported that MDA level was elevated and glutathione was lowered in liver of rats, which received 28.6 mg/kg chloramphenicol. On the other hand, Elia and Pacini, (2016) reported an increase in liver glutathione levels of rainbow trout treated with a dose of 7.5 and 15 mg/kg b.w. florfenicol. Glutathione levels were 1.5 fold higher in elevated dose of florfenicol attributed to protect liver cells from oxidative damage of florfenicol.

Other treated groups such as the tocotrienol and selenium yeast alone and combination of them showed

significant decreases in MDA level while significant increases in SOD activity and GSH concentration in hepatic tissue when compared with the florfenicol treated group. The result of SOD activity in liver tissue with the addition of tocotrienol was attributed to the disturbance of defense mechanism of liver tissue which stimulates production of superoxide anion radicals which prevent lipid peroxide formations (Lee et al., 2009). Selenium yeast may improve the antioxidant status of broilers by increasing the activity of antioxidant enzymes and inhibiting lipid peroxidation (Jiang et al., 2009; Yang et al., 2012). Gladyshev and Hatfield, (1999) reported that selenium inter in the form of amino acid called selenocysteine (one of selenoprotein) which have important enzymatic functions associated with antioxidant activity. Selenium is important in sulphur amino acid metabolism. In this way, the sulphur amino acids methionine and cystine can spare selenium through their antioxidant role.

Our results referred to tocotrienol of liver tissue were in accordance to those explained by Khan et al. (2011). Moreover, Palozza et al. (2006) reported that tocotrienols decrease MDA in rat liver microsomes (obtained from tissue homogenization then *in vivo* added tocotrienols to suspension) by inhibition of 2,2'-azobis 2-amidinopropane (AAPH) which induced MDA production. This result related to in selenium yeast were also observed by several authors (Mahmoud and Edens, (2003); Petrovič et al., 2006; Bao-wei et al., 2011; Li et al., 2016; Hamidet al., 2018) who concluded that selenium yeast restored or increased the liver antioxidant defense. The antioxidant capacity of selenium is an integral component of glutathione peroxidase (GPx) which plays a crucial role to reduce cellular damage by ROS (Kong et al., 2017). Our results disagreed with those previously explained by Lee et al. (2005) who reported SOD activity in liver of rats was lowered with age when these rats were fed with palmvitee (palm oil). This decrease may be due to the compensating effect of palmvitee which replaces antioxidant enzyme activities. Our results in selenium yeast also disagreed with those described by Holovská et al. (2003) and Chen et al. (2013) who showed no significant differences in SOD activity of hepatic tissue in chickens fed on selenium yeast.

Liver sections of the florfenicol treated group displayed focal leukocytic infiltration in the hepatic tissue and focal coagulative necrotic foci represented by deeply eosinophilic cytoplasm of the nuclei with karyorrhexis and karyolysis in the liver section

of chicks. Our findings were in correlation to those described by Yue and Li-hai (2009) who reported fatty degeneration and necrosis in the livers of seventh day old broiler chicks, which were fed diet containing different doses of florfenicol (200, 400, 800 and 2000 mg/kg) for 14 days. In addition, Reda et al. (2013) noted diffused hydropic degeneration in the liver examined in Nile tilapia (*Oreochromis niloticus*) which received 5 mg/kg of florfenicol for 12 weeks, whereas Isa et al. (2020) described coagulative necrosis with karyopyknosis and karyorrhexis in livers of 8-week-broiler chicks which treated with 250 mg/kg chloramphenicol from the 1st day.

Liver sections of the tocotrienol treated group showed fatty degeneration of some hepatic cells with eccentric location of the nuclei on the 35th day of age and focal mononuclear cell infiltration of the hepatic tissue and activation of the Kupffer cells on 35th day of age. These findings are in agreement with those of Qureshi et al. (2011) who stated that mild chronic inflammation was observed in liver of chickens fed with diet containing 50 ppm δ -tocotrienol and Wong et al. (2012) who observed mild fatty changes in the liver of rats which fed a high carbohydrate diet, contained 240 mg TRF/mL palm olein, for 8 weeks. On the other hand, no histopathological changes were observed in the liver tissue of mice and rats treated with different doses of tocotrienol (Husain et al., 2009; Shibata et al., 2012).

In our study, treatment with selenium yeast revealed focal hepatocellular coagulative necrosis with karyolysis and ballooning degeneration represented by empty cytoplasm and centrally located nuclei in the adjacent hepatocytes.

Our results are in agreement with those previously published by Attia et al. (2010) who stated

that breeding hens fed on selenium yeast showed mild focal necrosis in hepatic tissue. Moreover, Hamid et al. (2018) concluded that rats treated with selenium-enriched yeast showed ballooning of liver hepatocytes, mild fatty changes, and mild degree of centrilobular necrosis with partial infiltration of inflammatory cells.

On other hand, Mirjana et al. (2004) reported that chickens orally fed on selenized yeast showed varying degree of intracellular edema and fatty changes in the liver of sacrificed chicks..

Broilers treated with the combination of tocotrienol and selenium yeast revealed focal mononuclear cell infiltration in the fibrous connective tissue of the portal area and activation of the Kupffer cells with little evidence of necrotic changes in the liver. This means that the addition of tocotrienol with selenium yeast mitigated the damage caused by florfenicol on liver tissue.

CONCLUSION

The addition of tocotrienol, selenium yeast alone or their combination can improve the antioxidant effect and mitigate lipid peroxidation in the livers of treated broiler. Therefore, tocotrienol and selenium yeast can improve the safety of using florfenicol in broilers under experimental conditions.

ACKNOWLEDGMENTS

We thank Prof. Dr. Fatma Darwish, head of the department of pathology of animal health research, for her valuable support and help during practical histopathology work.

CONFLICT OF INTEREST

None declared.

REFERENCES

- Ahmed A F, Consatable PD, Misk NA (2002) Effect of feeding frequency and route of administration on abomasal luminal PH in dairy calves fed milk replacer. *J Dairy Sci* 85, 1502-1509.
- Allam HH, Salah AAB, Zaki RH, Bahar NA (2014) Bacteriological and Biochemical Studies on Pekin Duckling Infected With *Pasteurella Multocida* with Trial for Treatment. *Zagazig Veterinary Journal* 42 (2):23-32.
- Ashayerizadeh A, Dabirini, Ashayerizadeh O, Mirzadeh KH, Roshanfekar H, Mamooee M (2009) Effect of dietary antibiotic, probiotic and prebiotic as growth promoters, on growth performance, carcass characteristics and hematological indices of broiler chickens. *Journal of Biological sciences* 12 (1): 52-57.
- Attia YA, Abdalrah AA, Zeweil HS, Bovera F, Tag-El-Din AA, Araft MA (2010) Effect of inorganic or organic selenium supplementation on productive performance, egg quality and some physiological traits of dual-purpose breeding hens. *Czech J. Anim. Sci.*, 55, (11): 505-519.
- Bancroft JD, Stevens AS, forward by Douson JMP (1977) Theory and practice of histologic techniques. Churchill Livingstone Edinburgh New York 222-278.
- Baowei W, Guoqing H, Qiaoli W, BinY (2011) Effects of selenium yeast supplementation on the growth performance, meat quality, immunity, and antioxidant capacity of goose *Animal Physiology and Animal Nutrition* 95: 440-448.
- Belfield A, Goldberg DM, (1971) Revised assay for serum phenyl phosphatase activity using 4-amino-antipyrine. *Enzyme J Clin Path* 12:561-573.
- Beutler E, Duron O, Kelly MJB (1963) *Lab clin. Med* 61: 882.
- Budin SB, Othman F, Louis SR., Bakar MA, Das S, Mohamed J (2009) The effects of palm oil tocotrienol-rich fraction supplementation on biochemical parameters, oxidative stress and the vascularwal diabetic rats. *CLINICS* 64(3):235-44.
- Chang SK, Davis JL, Cheng CN, Shien RH, Hsieh MK, Koh B W, Chou CC (2010) Pharmacokinetics and tissue depletion of florfenicol in Leghorn and Taiwan Native chickens. *Journal of veterinary Pharmacology Therapeutic* (33):471-479.
- Chen G, Wu J, Li C (2013) Effect of different selenium sources on production performance and biochemical parameters of broilers. *Journal of Animal Physiology and Animal Nutrition* 98:747-754.
- Coles EH (1974) Determination of globulin. *Vet Clin Path* 2nd Ed Saunders Company Philadelphia and London PP 560-568.
- Domas BL (1975) Colorimetric determination of total protein. *ClinChem* 21(1): 159-166.
- Doumas B (1971) Colorimetric method for albumin determination. *Clin-ChimActa* 31: 87-92.
- Duncan DB (1955) Multiple range and multiple F tests. *Biometrics* 11:1-42.
- El-Demerdash FM, Nasr HM (2014) Antioxidant effect of selenium on lipid peroxidation, hyperlipidemia and biochemical parameters in rats exposed to diazinon. *J Trace Elem Med Biol* 28, 89-93.
- Elia AC, Pacini N (2016) Assessment of Detoxifying Markers for Florfenicol in Rainbow Trout Liver. *Journal of Aquatic Animal Health* 28:258-265.
- Er A, Dik B (2014) The Effects of Florfenicol on the Values of Serum Tumor Necrosis Factor- α and Other Biochemical Markers in Lipopolysaccharide-Induced Endotoxemia in Brown Trout. *Mediators of Inflammation*. <http://dx.doi.org/10.1155/2014/464373>.
- Fan C, Yu B, Chen D (2009) Effects of Different Sources and Levels of Selenium on Performance, Thyroid Function and Antioxidant Status in Stressed Broiler Chickens. *International Journal of Poultry Science* 8 (6): 583-587.
- Farombi EO (2001) Antioxidant status and hepatic lipid peroxidation in chloramphenicol treated rats. *Tohoku J Exp Med* 194: 91-98.
- Firozian F, Karami S, Ranjbar A, Azandaryani MT, Nili-Ahmadabadi A (2020) Improvement of therapeutic potential N-acetylcysteine in acetaminophen hepatotoxicity by encapsulation in PEGylated nano-niosomes. *Life Science* 255:117832.
- Flegg HM (1973) Quantitative-enzymatic-colourimetric determination of total and HDL-C in serum or plasma. *Ann Clin Biochem* 10: 79-88.
- Friedewald WT, Levy RI, Fredrickson DS (1972) Estimation of the concentration of LDL cholesterol in plasma, without use of the preparative ultracentrifuge. *ClinChem* 18: 499-502.
- Gladyshev VN, Hatfield DL (1999) Selenocysteine-containing proteins in mammals. *J Biomed Sci* 1999;6:151-60.
- Gordon T, Castelli WP, Hjortland MC, Kannel WB, Dawber TR (1977) High density lipoprotein as protective factor against coronary heart disease: The Framingham study. *Am J Med* 62: 707-714.
- Hamid A A, Aiyelaagbe O O, Usman L A, Ameen O M, Lawal A (2010) Antioxidants: Its medicinal and pharmacological applications. *Afr J Pure Appl Chem.*, 4 (8):142-151.
- Hamid M, Abdulrahim Y, Liu D, Qian G, Khan A, Huang K (2018) The hepatoprotective effect of selenium-enriched yeast and gum arabic combination on carbon tetrachloride-induced chronic liver injury in rats. *Journal of Food Science* 83(2):525-534.
- Han C, Wei Y, Cui Y, Geng Y, Bao Y, Shi W (2020) Florfenicol induces oxidative stress and hepatocyte apoptosis in broilers via Nrf2 pathway. *Ecotoxicology and Environmental Safety* 191:110239.
- Hasan WNW, Abd-Ghafar N, Chin K, Ima-Nirwana S (2018) Annatto-derived tocotrienol stimulates osteogenic activity in preosteoblastic MC3T3-E1 cells: a temporal sequential study. *Drug Des Devel Ther* 12: 1715-1726.
- Hasselwander O, Krämer K, Oberfrank U, Baldeus K, Schroder H, Kaufmann W, Bahnemann R, Nowakowsky B (2001) Effects of feeding various tocotrienol sources on plasma lipids and aortic atherosclerotic lesions in cholesterol-fed rabbits. *Food Research International* 35:245-251.
- Holovská K Jr, Holovská K, Boldižárová K, Čekonová S, Lenártová V, Levkut M, Javorský M, Leng L (2003) Antioxidant enzyme activities in liver tissue of chickens fed diets supplemented with various forms and amounts of selenium. *Journal of Animal and Feed Sciences* 12: 143-152.
- Hurrell R (2003) Influence of vegetable protein sources on trace element and mineral bioavailability. *Journal of nutrition* 133(9): 2973S-2977S.
- Husain K, Francois RA, Hutchinson S Z, Neuger A M, Lush R, Coppola D, Sebt S, Malafa MP (2009) Vitamin E δ -Tocotrienol Levels in Tumor and Pancreatic Tissue of Mice after Oral Administration. *Pharmacology* 83:157-163.
- Invernizzi G, Agazzi A, Ferroni M, Rebucci R, Fanelli A, Baldi A, Dell'Orto V, Savoini G (2013) Effects of Inclusion of Selenium-Enriched Yeast in the Diet of Laying Hens on Performance, Eggshell Quality, and Selenium Tissue Deposition. *Italian Journal of Animal Science* 12:1-8.
- Isa MM, Bukar M, Saheed Y, Anka BA (2020) The Effects of Prolonged Chloramphenicol Administration on Hematological Parameters and Histopathology of Liver, Kidney and Spleen in Broiler Chicken. *Asian Journal of Research in Infectious Diseases* 3(3): 34-40.
- Jiang ZY, Lin YC, Zhou GL, Luo LH, Jiang SQ, Chen F (2009) Effects of dietary selenomethionine supplementation on growth performance, meat quality and antioxidant property in yellow broilers. *J Agric Food.Chem* 57: 9769-9772.
- Kamat JP, Sarma HD, Devasagayam TPA, Nesaretnam K, Basiron Y (1997) Tocotrienols from palm oil as effective inhibitors of protein oxidation and lipid peroxidation in rat liver microsomes. *Mol Cell Biochem* 170(1-2):131-8, <http://dx.doi.org/10.1023/A:1006853419214>.
- Khalil HS, Mansourb AT, Godaa AM, Omar EA (2019) Effect of selenium yeast supplementation on growth performance, feed utilization, lipid profile, liver and intestine histological changes, and economic benefit in meagre, *Argyrosomus regius*, fingerlings. *Aquaculture* 501:135-143.
- Khalil S, Hamed E, Hassanin O (2012) Residue Withdrawal of Florfenicol from the Serum and Edible Tissues of Broiler Chickens. *Journal of American science* 8(12): 112-123.
- Khan MS, Khan MK A, Siddiqui MH, Arif JM (2011) An in vivo and in silico approach to elucidate the tocotrienol-mediated fortification against infection and inflammation induced alterations in antioxidant

- defense system. *European Review for Medical and Pharmacological Sciences* 15: 916-930.
- Kong Y, Ding Z, Zhang Z, Ye J, Du Z (2017) Dietary selenium requirement of juvenile oriental river prawn *Macrobrachium nipponense*. *Aquaculture* 476: 72-78.
- Lee LM, Hamid NA, Yusof Y (2005) Effects of Palmvitee on Status of Superoxide Dismutase and Glutathione Peroxidase in Rat Liver during Aging. *Malaysian Journal of Biochemistry and Molecular Biology* 12:21-24.
- Lee SP, Mar GY, Lean-Teik Ng (2009) Effects of tocotrienol-rich fraction on exercise endurance capacity and oxidative stress in forced swimming rats. *Eur J Appl Physiol* 107:587-595.
- Li X, Zhang Y, Yuan Y, Sun Y, Qin Y, Deng Z, Li H (2016) Protective Effects of Selenium, Vitamin E, and Purple Carrot Anthocyanins on D-Galactose-Induced Oxidative Damage in Blood, Liver, Heart and Kidney Rats. *Biol Trace Elem Res.*, 173:433-442.
- Liu H, Yu Q, Fang C, Chen S, Tang X, Ajuwon K, Fang R (2020) Effect of Selenium Source and Level on Performance, Egg Quality, Egg Selenium Content, and Serum Biochemical Parameters in Laying Hens. *Foods* 9, 68; doi:10.3390/foods9010068.
- Mahmoud K, Edens FW (2003) Influence of selenium sources on age-related and mild heat stress related changes of blood and liver glutathione redox cycle in broiler chickens (*Gallus domesticus*). *Comparative Biochemistry and Physiology Part B* 136:921-934.
- Mirjana T, Jovanovic M, Jokic Z, Hristov S, Vesna D (2004) Alterations in liver and kidneys of chickens fed with high levels of sodium selenite or selenized yeast. *Acta Veterinaria (Beograd)* 54 (2-3):191-200.
- Nakamura H, Furukawa F, Nishikawa A, Miyauchi M, Son HY, Imazawa T, Hirose M, (2001) Oral toxicity of a tocotrienol preparation in rats *Food Chem. Toxicology* 39:799-805.
- Nishikimi M, Roa NA, Yogi K (1972) The occurrence of superoxide anion in the reaction of reduced phenazinemetosulfate and molecular oxygen. *Biochem Biophys Res Commun* 46:849-854.
- Norazlina M, Ima-Nirwana S, Abdul Gapor MT, Khalid BA (2002) Tocotrienols are needed for normal bone calcification in growing female rats. *Asia Pac J Clin Nutr* 11(3):194-9.
- Ohkawa H, Ohishi W, Yagi K (1979) Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analytical Biochemistry* 95 (2):351-358.
- Olayink OO, Kareem AM, Ariyo IB, Omotugba SK, Oyebanji AO (2012) Antioxidant Contents (Vitamin C) of Raw and Blanched Different Fresh Vegetable Samples. *Food and Nutrition Sciences* 3: 18-21.
- Palozza P, Verdecchia S, Avanzi L, Vertuani S, Serini S, Iannone A, Manfredini S (2006) Comparative antioxidant activity of tocotrienols and the novel chromanyl-polyisoprenyl molecule FeAox-6 in isolated membranes and intact cells. *Molecular and Cellular Biochemistry* 287: 21-32.
- Petrovič V, Boldižárová K, Faix S, Mellen M, Arpášová H, Leng L (2006) Antioxidant and selenium status of laying hens fed with diets supplemented with selenite or Se-yeast. *Journal of Animal and Feed Sciences* 15, 435-444.
- Qureshi A, Reis C, Qureshi N, Papisian J, Morrison C, Schaefer M (2011) δ -Tocotrienol and quercetin reduce serum levels of nitric oxide and lipid parameters in female chickens. *Lipids in Health and Disease* 10: 2-22.
- Qureshi AA, Peterson DM (2001) The combined effects of novel tocotrienols and lovastatin on lipid metabolism in chickens. *Atherosclerosis* 156: 39-47.
- Reda RM, Ibrahim R E, Ahmed EG, El-Bouhy ZM (2013) Effect of oxytetracycline and florfenicol as growth promoters on the health status of cultured *Oreochromis niloticus*. *Egyptian Journal of Aquatic Research* 39: 241-248.
- Ren X, Pan L, Wang L (2014) Effect of florfenicol on selected parameters of immune and antioxidant systems, and damage indexes of juvenile *Litopenaeus vannamei* following oral administration. *Aquaculture* 432:106-113.
- Ronette G, (2012) Pharmacokinetics and Bioequivalence of Florfenicol Oral Solution formulations (Flonicol® and Veterin® 10%) in Broiler Chickens. *J. Bioequivalence and Bioavailability* 4(1): 1-5.
- Sen CK, Khanna S, Roy S (2006) Tocotrienols: vitamin E beyond tocopherols. *Life Science* 78: 2088-2098.
- Sevcikova S, Skrivan M, Dlouha G (2008) The effect of lycopene supplementation on lipid profile and meat quality of broiler chickens. *Czech J Anim Sci* 53(10): 431-440.
- Shaheen H, El-Far A (2013) Evaluation of the Therapeutic Efficacy of Pefloxacin and Florfenicol Combination in Broilers Experimentally Challenged by *Escherichia coli*. *Int J Pharm Sci Rev Res* 23 (64): 396-404.
- Shi L, Ren Y, Zhang C, Yue W, Lei F (2018) Effects of organic selenium (Se-enriched yeast) supplementation in gestation diet on antioxidant status, hormone profile and haematobiochemical parameters in Taihang Black Goats. *Animal Feed Science and Technology* 238:57-65.
- Shibata A, Nakagawa K, Shirakawa H, Kobayashi T, Kawakami Y, Takashima R, Ohashi A, Sato S, Ohsaki Y, Kimura F, Kimura T, Tsuduki T, Komai M, Miyazawa T (2012) Physiological Effects and Tissue Distribution from Large Doses of Tocotrienol in Rats. *Biosci-Biotechnol Biochem* 76 (9):1805-1808.
- Sies H (1997) Oxidative stress: oxidants and antioxidants. *The physiological society Experimental physiology* 82:291-295.
- Snedecor GW, Cochran WC (1989) *Statistical methods*. The eighth Edition Iowa University Press Ames Iowa USA.
- SPSS Inc. Released (2009) *PASW Statistics for Windows Version 18.0*. Chicago: SPSS Inc.
- Tasakia M, Umemura T, Inoue T, Okamura T, Kuroiwa Y, Ishii Y, Maeda M, Hirose M, Nishikawa A (2008) Induction of characteristic hepatocyte proliferative lesion with dietary exposure of Wistar Hanover rats to tocotrienol for 1 year. *Toxicology* 250: 143-150.
- Tietz NW (1976) *Fundamentals of clinical chemistry*. WB Saunders Company Washington D.C.
- Upton JR, Edens F W, Ferket PR (2009) The effects of dietary oxidized fat and selenium source on performance, glutathione peroxidase, and glutathione reductase activity in broiler chickens. *J. Appl Poult Res* 18:193-202.
- Varly H (1974) *Clinical chemistry methodology, past and present*. Ann Clin Chem 11: 161-63.
- Wang Y, Xu B (2008) Effect of different selenium source (sodium selenite and selenium yeast) on broiler chickens. *Animal Feed Science and Technology* 144: 306-314.
- Wang X, Han C, Cui Y, Geng Y, Wei Y, Shi W, Bao Y, (2020): Florfenicol induces renal toxicity in chicks by promoting oxidative stress and apoptosis. *Environmental Science and Pollution Research* <https://doi.org/10.1007/s11356-020-10550-4>.
- Yue QU, Li-hai GAO (2009): *Pathological Study of Florfenicol Poisoning in Experimental Broiler*. China Animal Husbandry and Veterinary Medicine. en.cnki.com.cn.