

Journal of the Hellenic Veterinary Medical Society

Vol 74, No 2 (2023)



Prevalence of Haemoproteus and Leucocytozoon spp. in Wild Birds in Hatay, Turkey

A Zerek, I Erdem, H Alakus, M Yaman, ME Altuğ

doi: [10.12681/jhvms.29714](https://doi.org/10.12681/jhvms.29714)

Copyright © 2023, I Erdem, A Zerek, M Yaman, H Alakus, ME Altuğ



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

To cite this article:

Zerek, A., Erdem, I., Alakus, H., Yaman, M., & Altuğ, M. (2023). Prevalence of Haemoproteus and Leucocytozoon spp. in Wild Birds in Hatay, Turkey . *Journal of the Hellenic Veterinary Medical Society*, 74(2), 5651–5656.
<https://doi.org/10.12681/jhvms.29714> (Original work published July 6, 2023)

Prevalence of *Haemoproteus* and *Leucocytozoon* spp. in Wild Birds in Hatay, Turkey

A. Zerek¹ , İ. Erdem^{1*} , H. Alakuş² , M. Yaman¹ , M.E. Altuğ² 

¹ Hatay Mustafa Kemal University, Faculty of Veterinary Medicine, Parasitology Department, TR-30001 Hatay-Türkiye

² Hatay Mustafa Kemal University, Faculty of Veterinary Medicine, Surgery Department, TR- 31001 Hatay-Türkiye

ABSTRACT: Hemosporidian parasites that cause infections in poultry and death in susceptible animals show a global distribution. Infected wild birds play an important role in this distribution due to seasonal migration among different regions. Hatay, where migratory birds enter Turkey, is an active region regarding vectors, disease agents and hosts. This study aimed to investigate the prevalence of *Haemoproteus* and *Leucocytozoon* parasites in wild birds of Hatay Province by microscopic and molecular methods. Blood samples were taken from a total of 50 wild birds belonging to three orders, eight species. In the microscopic examination, *Haemoproteus* spp. infection 18% (9/50) and coinfections (*Haemoproteus* spp. and *Leucocytozoon* spp.) 4% (2/50) were detected while in the PCR examination, coinfections 4% (2/50), *H. columbae* 20% (10/50) and *Leucocytozoon* spp. 2% (1/50) were detected. As a result, haemosporidian parasites were detected in approximately one of every four wild birds (microscopic 22%, molecular 26%).

Keywords: *Haemoproteus*; Hatay; *Leucocytozoon*; Turkey; wild bird

Corresponding Author:
İ. Erdem, Hatay Mustafa Kemal University, Faculty of Veterinary Medicine,
Parasitology Department
E-mail address: ipekerdem@mku.edu.tr

Date of initial submission: 23-02-2022
Date of acceptance: 19-03-2023

INTRODUCTION

Worldwide, haemosporidian blood parasites are vector-borne parasites that infect mammals, amphibians, reptiles, and poultry. More than 200 haemosporidian parasites have been identified in the strains of *Plasmodium*, *Haemoproteus* and *Leucocytozoon* that infect domestic and wild birds (Valkiunas, 2004). The vectors of the *Plasmodium* species are mosquitoes; vectors of the *Haemoproteus* species are *Pseudolynchia*, *Microlynchia* and *Culicoides*; vectors of the *Leucocytozoon* species are *Culicoides* and *Simulium*, which are hematophagous flies. Species belonging to the *Haemoproteus* and *Leucocytozoon* strains are prevalent in poultry. *Haemoproteus meleagridis* in domestic and wild turkeys; *H. nettionis* in domestic and wild ducks, geese and a variety of poultry; *H. iophoryx* in quails; *H. sacharovi* in pigeons and doves; *H. columbae* in domestic and wild pigeons, doves and other wild birds are parasitized. *Leucocytozoon simondi* from waterfowl, *L. marchouxi* from pigeons and doves, *L. toddi* from birds of prey have been reported (Derinbay Ekici, 2017). Although it is stated that haemosporidian parasites are not very harmful to poultry, it has been stated that they cause severe infections resulting in death in wild birds when high parasitemia occurs (Palinauskas et al., 2016). These parasites can cause anemia, fever, weakening, loss of appetite, difficulty in breathing, limping and decrease in lifespan, loss of productivity, changes in metabolic activities such as migration and reproduction, impairments in thermoregulation, and even deaths in poultry (Atkinson et al., 2000). In pathological findings, liver, spleen and kidney enlargement has been encountered (Valkiunas, 2004).

Microscopic and molecular studies on the presence of haemosporidian parasites have been carried out in domestic (Öz and Turut, 2007; Sürsal et al., 2017; Tasci et al., 2018) and wild birds (Balkaya et al., 2016; Ciloglu et al., 2016; Ciloglu et al., 2020; Marzal and Albayrak, 2012; Özmen et al., 2006; Özmen et al., 2009; Yildirim et al., 2013) in different provinces of Turkey.

Turkey is a crucial Europe region due to its diverse ecosystems and its location on the migration routes of migratory birds. They constitute approximately 20% of the recorded bird species in Turkey. The province of Hatay is a critical region where migratory birds enter Turkey in the spring period and leave in the autumn period (Yaman et al., 2018). The prevalence of haemosporidian parasites in migratory birds that have

the potential to encounter more vectors is higher than in those that do not migrate (Møller and Erritzøe, 1998). Hatay Province has a crucial position in migratory birds, yet no study on this subject is up to date.

This study aimed to investigate the prevalence of *Haemoproteus* and *Leucocytozoon* parasites in wild birds in Hatay by microscopic and molecular methods and contribute to the literature in this field.

MATERIALS AND METHODS

Ethical Statement

This study was performed with the Hatay Mustafa Kemal University Animal Experiments Local Ethics Committee decision. It was decided that ethics committee approval was not required. (Approval no: 2019/06-4, date: 31.05.2019).

Features of the Work Areas

The province of Hatay, where the temperate Mediterranean climate prevails, has an area of approximately 6,000 km² at 36 ° 14 North and 36 ° 10 East coordinates (Muz et al., 2015).

Collection of Blood Samples

In the study, 1 ml of blood samples was taken from vena subcutanea ulnaris to blood tubes with anticoagulant substance from a total of 50 wild birds belonging to three orders, eight species (determined with the help of the relevant literature) (Kiziroğlu, 2013; Porter et al., 2009). They were brought to the HMKU Veterinary Health, Practice and Research Center as wounded between May 2018 and January 2020.

Microscopic Method

The smears prepared from each of the blood samples brought to the laboratory of Parasitology were stained with the Giemsa solution and examined under the light microscope with the help of the relevant literature (Valkiunas, 2004). The remaining parts of the blood samples were stored at -20 °C for molecular analysis.

Molecular Method

The genomic DNA was extracted using the protocol of the commercial kit of Roche. The extracted DNA was stored at -20 °C until it was used in PCR. Then, PCR was performed using primers specific to the mitochondrial *cytochrome b* (mt-cytb) gene region of *Haemoproteus* and *Leucocytozoon* parasites.

The primers of PCR reaction for detection of *Haemoproteus columbae* were as follows: *H. clom-F*: 5'-TTA GAT ACA TGC ATG CAA CTG GTG-3' and *H. clom-R*: 5'-TAG TAA TAA CAG TTG CAC CCC AG-3'. PCR amplification was carried out in 20 µl reaction mixtures containing 4 µl 5x Master Mix (Solis BioDyne), 1 µl of each primer (10 pmol), 1 µl DNA (100 ng/µl) and 13 µl water (PCR grade). PCR was performed as follows: initial denaturation at 94 °C for 5 min followed by 30 cycles consisting of denaturation at 94 °C for 1 min, annealing at 60 °C for 1 min, extension at 72 °C for 1 min and final extension at 72 °C for 5 min. PCR products were separated by electrophoresis on 1% agarose gels and were visualized under UV light in terms of the presence of specific band. The size of *H. columbae* applicant was 207 base pairs (bp) (Doosti et al., 2014).

The extracted DNA was used in nested PCR reactions to amplify *Leucocytozoon* spp. For the first amplification, primers were used DW2: 5'-TAA TGC CTA GAC GTA TTC CTG ATT ATC CAG-3', and DW4: 5'-TGT TTG CTT GGG AGC TGT AAT CAT AAT GTG-3'. The primers were described by Perkins and Schall (Perkins and Schall, 2002). PCR amplification was carried out in 20 µl reaction mixtures containing 4 µl Master Mix (5x, Solis BioDyne), 1 µl of each primer (10 pmol), 1 µl DNA (100 ng/µl) and 13 µl water (PCR grade). The first PCR reaction was performed as follows: initial denaturation at 94 °C for 3 min followed by 35 cycles consisting of denaturation at 94 °C for 30 sec, annealing at 52 °C for 30 sec, extension at 72 °C for 1 min and final extension at 72

°C for 10 min. For the second PCR reaction, primers were used belonging to the mitochondrial *cytochrome b* (mt-cytb) gene region specific to *Leucocytozoon* spp. These primers were Leuco-F: 5'-TCT TAC TGG TGT ATT ATT AGC AAC-3' and Leuco R: 5'-AGC ATA GAA TGT GCAAAT AAA CC-3'. PCR amplification steps were carried out to be the same as the first PCR reaction. PCR products were separated by electrophoresis on 1% agarose gels and were visualized under UV light in terms of the presence of specific band. The size of *Leucocytozoon* spp. applicant was 865 bp (Sehgal et al., 2006).

RESULTS

According to the microscopic results, only *Haemoproteus* spp. infection 18% (9/50) and coinfections (*Haemoproteus* spp. and *Leucocytozoon* spp.) 4% (2/50) were detected (Table 1). *Haemoproteus* spp. and *Leucocytozoon* spp. gametocytes were observed in the blood smears (Figure 1). In the PCR examination, *H. columbae* 20% (10/50), *Leucocytozoon* spp. 2% (1/50), coinfections (*H. columbae* and *Leucocytozoon* spp.) 4% (2/50) were determined (Table 1).

As a result, these parasites could not be detected in the Ciconiiformes (*Ciconia ciconia*) and Strigiformes (*Tyto alba*) orders and the *Pernis apivorus* species of the Accipitriformes order by molecular and microscopic methods. Haemosporidian parasites were detected in approximately one of every four wild birds of the Accipitriformes order (microscopic 22%, molecular 26%) (Table 1).

Table1: Prevalence of haemosporidian parasites in wild birds in Hatay by microscopic and molecular methods

Orders	Species	Number of samples (n)	Number of positive samples (Microscope)				Number of positive samples (PCR)			
			H	L	Coinfections	Total	H*	L	Coinfections	Total
Accipitriformes	<i>Buteo buteo</i> (common buzzard)	17	4	-	-	4	4	-	-	4
Accipitriformes	<i>Clanga pomarina</i> (lesser spotted eagle)	11	2	-	2	4	2	-	2	4
Accipitriformes	<i>Circaetus gallicus</i> (Short-toed snake eagle)	9	1	-	-	1	2	1	-	3
Accipitriformes	<i>Buteo rufinus</i> (Long-legged Buzzards)	5	1	-	-	1	1	-	-	1
Accipitriformes	<i>Milvus migrans</i> (black kite)	3	1	-	-	1	1	-	-	1
Accipitriformes	<i>Pernis apivorus</i> (European honey buzzard)	1	-	-	-	-	-	-	-	-
Ciconiiformes	<i>Ciconia ciconia</i> (white stork)	3	-	-	-	-	-	-	-	-
Strigiformes	<i>Tyto alba</i> (western barn owl)	1	-	-	-	-	-	-	-	-
Total	8	50	9	-	2	11	10	1	2	13

H - *Haemoproteus* spp.; H* - *H. columbae*; L - *Leucocytozoon* spp.




Figure 1. Thin smears from blood samples stained with Giemsa; A-B) *Haemoproteus* spp. gametocyte (arrowed), C-D) *Leucocytozoon* spp. gametocyte (arrowed) (x1000 magnification). Scale bar = 10 µm.

DISCUSSION

The prevalence of haemosporidian parasites in wild birds in the world has been reported in various studies, and the reasons, such as the difference in bird species, haemosporidian parasites and geographical regions, make it difficult to compare the studies with each other (Inumaru et al., 2017). However, in some molecular studies conducted worldwide, the prevalence rate of *Haemoproteus* and *Leucocytozoon* parasites in wild birds is between 9% and 80.5%. In the Philippine Islands, in 215 wild birds, the infection was detected in a total of 26.4%, and the prevalence rates of *Haemoproteus*, *Leucocytozoon* and coinfections were 14.4%, 8.3% and 3.7%, respectively (Silva-Iturriza et al., 2012). In Slovakia, the rates of *Haemoproteus* and *Leucocytozoon* infections in 251 wild birds were 46.6% and 33.9%, respectively, and a total infection was 80.5% (Berthová et al., 2012). In 475 wild birds in Tokyo, Japan, the total infection rate was determined at 20%, the prevalence of *Haemoproteus* and *Leucocytozoon* was the same ratio as 8.8%, and coinfections were 2.4% (Inumaru et al., 2017). In another study, the total infection rate was found to be 9% in 55 wild birds, the *Haemoproteus*

prevalence rate was 3.6%, and *Leucocytozoon* prevalence rate was 5.4% (Tanigawa et al., 2012). In Indonesia, in 112 wild birds, *Haemoproteus* and *Leucocytozoon* the total infection rate was detected at 8% and 1.8%, respectively, and the total infection rate was 9.8% (Yuda, 2019). The prevalence of the species in the *Haemoproteus* lineage is generally higher than those of the *Leucocytozoon* and *Plasmodium* species in poultry (Atkinson and Van Riper, 1991). In this study, in 50 wild birds, *Haemoproteus columbae* and *Leucocytozoon* spp. were detected by PCR method at 20% and 2%, respectively; coinfections were 4%, and the prevalence rate of total infection was 26%. The *Haemoproteus* lineage was detected as higher than the *Leucocytozoon* lineage and was parallel to some studies conducted in different countries around the world, except for Japan (Inumaru et al., 2017; Tanigawa et al., 2012). In our study, wild birds infected with this blood parasite belong to the Accipitriformes order and generally prefer to live in a habitat of forestry, agriculture and open lands, in which *Leucocytozoon* spp. infection may have low prevalence. According to the literature that draws attention to the vector-host interaction (Krams et al., 2012), the risk of infection

with the *Leucocytozoon* lineage increase significantly with increasing proximity to rivers where blackflies (Diptera: Simuliidae) breed.

In this study, haemosporidian parasites were not detected in *Ciconia ciconia* (white stork) from the order Ciconiiformes, *Tyto alba* (Barn Owl) in the order Strigiformes, and *Pernis apivorus* (Bee hawk) in the order Accipitriformes. Although *Haemoproteus ciconiae* has been reported from the white stork (Valkiūnas et al., 2016), there are few studies about the diagnosis of blood parasites in these birds due to the wide living area and the difficulty of catching them (Valkiūnas, 2004; Valkiūnas et al., 2002; Jovani et al., 2002). It has been notified that these blood parasites were detected in some of the studies conducted on Barn Owls and Bee hawks (Salakij et al., 2018; Walther et al., 2016), however they were not detected in some of the studies (Krone et al., 2001; Sacchi and Prigioni, 1984). It is believed that these parasites were not found in this study due to examining only a few of the blood samples of wild birds in this order.

Turkey is on the migration routes of wild birds (Yaman et al., 2018), but there are few studies about the prevalence of haemosporidian parasites in wild birds, two of which are case reports (Özmen et al., 2009; Yildirim et al., 2013). While some of the studies used the microscopic method (Balkaya et al., 2016; Özmen et al., 2006; Özmen et al., 2009), some of them were conducted molecularly (Ciloglu et al., 2016; Ciloglu et al., 2020; Marzal and Albayrak, 2012; Yildirim et al., 2013). When looking at the molecular studies conducted in Turkey, *Haemoproteus* infection was detected in one Tawny Owl (*Strix aluco*) (Yildirim et al., 2013). The another study was conducted in 67 Anatolian nuthatches (*Sitta krueperi*), and the prevalence of *Haemoproteus* was 10.44% (7/67) while it was 13.43% (9/67) in *Leucocytozoon*. Coinfections had the same ratio as the last one. The total prevalence was 37,3% (25/67) (Marzal and Albayrak, 2012). *Leucocytozoon* infection was detected in 5 out of 22 (22.7%) predatory bird species in 6 different species (Ciloglu et al., 2016). In a study conducted in Sultan Sazlığı, there were 565 migratory birds from 39 species belonging to 23 families. A total of 193 (34.2%) birds were infected. Birds infected with *Haemoproteus* and *Leucocytozoon* were 168 and 6, respectively. Coinfections were 3.36% (19/565) (Ciloglu et al., 2020). Between our study and conducted the other studies, could not the relationship except for one (Ciloglu et al., 2020). The reason is that bird species, their localities and the

number of blood samples were different. Birds may not have encountered vectors or disease agents.

H. columbae parasitizes domestic and wild pigeons, doves and other wild birds (Derinbay Ekici, 2017). Although it is stated that infections with this species are apathogenic for birds, they can cause severe and fatal infections in young individuals whose immune system is not sufficiently developed (Sol et al., 2003). Based on the microscopic methods, *H. columbae* has been reported only in pigeons in Turkey (Karacaoğlu and Karatepe, 2017; Karatepe and Karatepe, 2018). The presence of *H. columbae* has not been reported in wild birds in the world except for pigeons (Al-Rubaie et al., 2020; Doosti et al., 2014; Nematollahi et al., 2012). With this study, the presence and prevalence of *H. columbae* were reported for the first time by PCR method in wild birds worldwide and in Turkey.

Although it is stated that microscopic diagnosis based on its prevalence and morphology in the host can be overlooked at low intensity (Valkiūnas et al., 2008), it has been claimed that it is a reliable tool even at low density and that the PCR test does not make a significant difference compared to microscopy (Krams et al., 2012). However, since the microscopic detection of parasites requires taxonomic expertise to identify the parasite species and stages of development accurately (Krams et al., 2012; Valkiūnas et al., 2008), it is recommended to verify the microscopic methods with molecular studies in today's conditions (Ciloglu et al., 2016; Valkiūnas et al., 2008; Valkiūnas et al., 2014). In this study, there was no significant difference between the microscopic findings and molecular findings. This situation supports the view of Krams et al. (2012).

CONCLUSION

This study investigates the prevalence of *Haemoproteus* and *Leucocytozoon* parasites in wild birds in Hatay Province, which is located on the crucial migratory bird routes in Turkey. During their migration, wild birds cause to be transport haemosporidian parasites and spread to different parts of the world. For this reason, it is beneficial to carry out more detailed studies on the subject of wild birds in Turkey.

ACKNOWLEDGEMENTS

This study was supported by the Hatay Mustafa Kemal University Scientific Research Project Coordination with the number 19.M.042.

REFERENCES

- Al-Rubaie HMA, Al-Biatee ST, Al-Saffar NSJ (2020) Molecular Diagnosis Of Haemoproteus Columbæ In Local Domestic Pigeons (Columba Livia Domestica) In Baghdad City. *Plant Arch*, 20(1): 195-198.
- Atkinson CT, Dusek RJ, Woods KL, Iko WM (2000) Pathogenicity of avian malaria in experimentally-infected Hawaii Amakihi. *J Wildl Dis*, 36(2): 197-201.
- Atkinson CT, Van Riper C (1991) Pathogenicity And Epizootiology Of Avian Haematozoa: Plasmodium, Leucocytozoon And Haemoproteus. In: *Bird-Parasite Interactions: Ecology, Evolution And Behaviour*. Oxford Ornithology Series, London: pp 19-48.
- Balkaya I, Oruc E, Guven E, Avcioglu H (2016) Haemosporidian parasites in blood smears of Sparrowhawks (Accipiter nisus, Falconiformes: Accipitridae) in Northeastern Turkey. *Isr J Vet Med*, 71(3): 26-30.
- Berthová L, Valkiūnas G, Országhová Z (2012) The first report of nine species of haemosporidian parasites (Haemosporida: Haemoproteus, Plasmodium and Leucocytozoon) in wild birds from Slovakia. *Biologia*, 67(5): 931-933.
- Ciloglu A, Ergen AG, Inci A, Dik B, Duzlu O, Onder Z, Yetismis G, Bensch S, Valkiūnas G, Yildirim A (2020) Prevalence and genetic diversity of avian haemosporidian parasites at an intersection point of bird migration routes: Sultan Marshes National Park, Turkey. *Acta Trop*, 210: 105465.
- Ciloglu A, Yildirim A, Duzlu O, Onder Z, Dogan Z, Inci A (2016) Investigation of avian haemosporidian parasites from raptor birds in Turkey, with molecular characterisation and microscopic confirmation. *Folia parasitol*, 63: 1.
- Derinbay Ekici Ö (2017) Protozoon Hastalıkları ve Tedavisi. In: Güvercin El Kitabı. Güneş Tıp Kitabevi, Ankara: pp 87-97.
- Doosti A, Ahmadi R, Mohammadalipour Z, Zohoor A (2014) Detection of Haemoproteus columbae in Iranian pigeons using PCR. In: *International Conference on Biological, Civil and Environmental Engineering*, Dubai: pp 36-38.
- Inumaru M, Murata K, Sato Y (2017) Prevalence of avian haemosporidia among injured wild birds in Tokyo and environs, Japan. *Int J Parasitol Parasites Wildl*, 6(3): 299-309.
- Jovani R, Tella JL, Blanco G, Bertellotti M (2002) Absence of haematozoa on colonial white storks Ciconia ciconia throughout their distribution range in Spain.
- Karacaoğlu M, Karatepe M (2017) Adana yöresi evcıl güvercinlerinde (Columba livia domestica) kan parazitlerinin araştırılması. *FU Vet J Health Sci*, 31(1): 33-38.
- Karatepe B, Karatepe M (2018) Niğde Yöresi Güvercinlerinde (Columba livia) Kan Protozoonlarının Yaygınlığı. *Türkiye Parazit Derg*, 42(4): 258-61.
- Kızıroğlu İ (2013) Türkiye Kuşları Cep Kitabı, Geoturka, İstanbul: pp 1-541.
- Krams I, Suraka V, Cirule D, Hukkanen M, Tummeleht L, Mierauskas P, Rytönen S, Rantala MJ, Vrublevska J, Orell M (2012) A comparison of microscopy and PCR diagnostics for low intensity infections of haemosporidian parasites in the Siberian tit Poecile cinctus. In: *Ann Zool Fenn*, 49(5): 331-340.
- Krone O, Priemer J, Streich J, Sommer P, Langgemach T, Lessow O (2001) Haemosporida of birds of prey and owls from Germany. *Acta Protozool*, 40(4): 281-290.
- Marzal A, Albayrak T (2012) Geographical variation of haemosporidian parasites in Turkish populations of Krüper's Nuthatch Sitta krueperi. *J Ornithol*, 153(4): 1225-1231.
- Møller AP, Erritzøe J (1998) Host immune defence and migration in birds. *Evol Ecol*, 12(8): 945-953.
- Muz BN, Oruç Kılınç Ö, İşler CT, Altuğ E, Karakavuk M (2015) Bazı yabani kuşların beyin dokularında Toxoplasma gondii ve Neospora caninum'un moleküler tanısı. *Kafkas Univ Vet Fak Derg*, 21 (2): 173-178.
- Nematollahi A, Ebrahimi M, Ahmadi A, Himan M (2012) Prevalence of Haemoproteus columbae and Trichomonas gallinae in pigeons (Columba domestica) in Isfahan, Iran. *J Parasit Dis*, 36(1): 141-142.
- Öz İ, Turut N (2007) Adana Yöresinde Evcıl Güvercinlerde Haemoproteus Columbæ'nın Yaygınlığı. *Bornova Vet Kont Araşt Enst Derg*, 29(43): 25-29.
- Özmen Ö, Haligür M, Adanir R (2009) Identification of different protozoa species from a common buzzard (Buteo buteo). *Turkish J Vet Anim Sci*, 33(3): 257-260.
- Özmen Ö, Haligür M, Yukari BA (2006) A study on the presence of leucocytozoonosis in wild birds of Burdur district. *Turkish J Vet Anim Sci*, 29(6): 1273-1278.
- Palinauskas V, Žiegėytė R, Iezhova TA, Ilgūnas M, Bernotienė R, Valkiūnas G (2016) Description, molecular characterisation, diagnostics and life cycle of Plasmodium elongatum (lineage pERIRUB01), the virulent avian malaria parasite. *Int J Parasitol*, 46(11): 697-707.
- Perkins SL, Schall J (2002) A molecular phylogeny of malarial parasites recovered from cytochrome b gene sequences. *J Parasitol*, 88(5): 972-978.
- Porter R, Christensen S, Schiermacker Hansen P (2009) Türkiye ve Orta Doğu'nun Kuşları, Uzerler Matbaası, Ankara: pp 1-455.
- Sacchi L, & Prigioni, C. (1984) Occurrence of leucocytozoon and haemoproteus (Apicomplexa, Haemosporina) in Falconiformes and Strigiformes of Italy. *Ann Parasitol Hum Comp*, 59(3): 219-226.
- Salakij C, Pompanom P, Lertwatcharasarakul P, Kasomdorkbua C, Salakij J (2018) Haemoproteus in barn and collared scops owls from Thailand. *J Vet Sci*, 19(2): 280-289.
- Sehgal RN, Hull AC, Anderson NL, Valkiūnas G, Markovets MJ, Kawamura S, Tell LA (2006) Evidence for cryptic speciation of Leucocytozoon spp. (Haemosporida, Leucocytozoidae) in diurnal raptors. *J Parasitol*, 92(2): 375-379.
- Silva-Iturriza A, Ketmaier V, Tiedemann R (2012) Prevalence of avian haemosporidian parasites and their host fidelity in the central Philippine islands. *Parasitol Int*, 61(4): 650-657.
- Sol D, Jovani R, Torres J (2003) Parasite mediated mortality and host immune response explain age-related differences in blood parasitism in birds. *Oecologia*, 135(4): 542-547.
- Sürsal N, Atan P, Gökpinar S, Duru Ö, Çakmak A, Yıldız K (2017) Kirikkale Yöresi'nde Yetistirilen Taklaci Güvercinlerde (Columba livia domestica) Haemoproteus spp.'nin Yaygınlığı/Prevalence of Haemoproteus spp. in Tumbler Pigeons (Columba livia domestica) in Kirikkale Province, Turkey. *Türkiye Parazit Derg*, 41(2): 71-76.
- Tanigawa M, Sato Y, Ejiri H, Imura T, Chiba R, Yamamoto H, Kawaguchi M, Tsuda Y, Murata K, Yukawa M (2012) Molecular identification of avian haemosporidia in wild birds and mosquitoes on Tsushima Island, Japan. *J Vet Med Sci*, 75(3): 319-326.
- Tasci G, Olmez N, Aydın PN, Akca A, Sari B, Arslan M, Mor N, Vatansever Z (2018) Prevalence and molecular characterization of haemosporidians in domestic geese: a new focus of haemosporidian parasites, kars province, northeastern Turkey. *Isr J Vet Med*, 73: 33-42.
- Valkiūnas G (2004) *Avian malaria parasites and other haemosporidia*, CRC press, Florida: pp 1-946.
- Valkiūnas G, Bolshakov C, Chernetsov N, Kosarev V (2002) Lack of haematozoa in the White Stork Ciconia ciconia in the Kaliningrad Region of Russia. *Avian Ecol Behav*, 9: 63.
- Valkiūnas G, Iezhova TA, Krizanauskienė A, Palinauskas V, Sehgal RN, Bensch S (2008) A comparative analysis of microscopy and PCR-based detection methods for blood parasites. *J Parasitol*, 94(6): 1395-1401.
- Valkiūnas G, Ilgūnas M, Bukauskaitė D, Iezhova TA (2016) Description of Haemoproteus ciconiae sp. nov. (Haemoproteidae, Haemosporida) from the white stork Ciconia ciconia, with remarks on insensitivity of established polymerase chain reaction assays to detect this infection. *Parasitol Res*, 115(7): 2609-2616.
- Valkiūnas G, Palinauskas V, Ilgūnas M, Bukauskaitė D, Dimitrov D, Bernotienė R, Zehntindjiev P, Ilieva M, Iezhova TA (2014) Molecular characterization of five widespread avian haemosporidian parasites (Haemosporida), with perspectives on the PCR-based detection of haemosporidians in wildlife. *Parasitol Res*, 113(6): 2251-2263.
- Walther E, Valkiūnas G, Wommack EA, Bowie RC, Iezhova TA, Sehgal RN (2016) Description and molecular characterization of a new Leucocytozoon parasite (Haemosporida: Leucocytozoidae), Leucocytozoon californicus sp. nov., found in American kestrels (Falco sparverius sparverius). *Parasitol Res*, 115(5): 1853-1862.
- Yaman M, Zerek A, Akkücüç Ş (2018) A case of traumatic myiasis in a white stork (Ciconia ciconia) caused by Lucilia sericata (Diptera: Calliphoridae). *Eurasian J Vet Sci*, 34(3): 201-203.
- Yildirim A, Aysul N, Bayramli G, Abdullah I, Hasan E, Aypak S, Duzlu O, Ciloglu A, Onder Z (2013) Detection and molecular characterization of a Haemoproteus lineage in a Tawny Owl (Strix aluco) in Turkey. *Ankara Univ Vet Fak Derg*, 60(3): 179-183.
- Yuda P (2019) Detection of avian malaria in wild birds at Trisik Beach of Yogyakarta, Java (Indonesia). *Ann Parasitol*, 65(2): 171-175.