



# Journal of the Hellenic Veterinary Medical Society

Vol 74, No 4 (2023)



# To cite this article:

Ozguden-Akkoc, C., & Akkoç, A. (2024). Immunohistochemical Demonstration of Mast Cells in Bovine Papillomatous Digital Dermatitis of Holstein-Friesian Dairy Cattle. *Journal of the Hellenic Veterinary Medical Society*, *74*(4), 6369–6376. https://doi.org/10.12681/jhvms.29927

# Immunohistochemical Demonstration of Mast Cells in Bovine Papillomatous Digital Dermatitis of Holstein-Friesian Dairy Cattle

C.G. Ozguden-Akkoc<sup>1</sup>, A. Akkoc<sup>2</sup>

<sup>1</sup>Bursa Uludag University, Faculty of Veterinary Medicine, Department of Histology & Embryology, Bursa, Turkey

<sup>2</sup>Bursa Uludag University, Faculty of Veterinary Medicine, Department of Pathology, Bursa, Turkey

ABSTRACT: Mast cells (MC) are unique members of immune system; their location and functions are of great importance in health and disease status. This study aims to evaluate MCs distribution and heterogeneity in healthy and digital dermatitis lesions of dairy cattle for the first time. A total of 50 skin samples, 25 healthy and 25 with digital dermatitis (DD) lesions were sampled in Holstein-Friesian dairy cattle. All samples were stained with hematoxylin-eosin (H&E) for microscopic examination and also stained with Toluidine blue for MCs demonstration. Epidermal acanthosis and hyperkeratosis with dermal inflammatory cell infiltrations were in consistent with digital dermatitis lesions. Spirochetal agents were successfully demonstrated with Warthin-Starry staining in 22 out of 25 DD samples. Increased number of mast cell was observed in digital dermatitis samples when compared with healthy skin samples. The average number of intact MCs was  $4.6 \pm 2$  and  $9.3 \pm 1$  in healthy and digital dermatitis samples, respectively. Twofold increase in the number of intact MCs in digital dermatitis samples was observed. Degranulated mast cell numbers in tissue sections were also higher in digital dermatitis samples and 4.25-fold increase was recorded in affected skin samples. Immunophenotype of MCs in skin samples were identified by immunohistochemical staining with anti-tryptase and chymase antibodies. Only tryptase positive MCs were observed in healthy and DD samples. In the statistical analysis, differences in the mean intact and degranulated MCs were found to be significant (mean intact: p<0.05, mean degranulated: p<0.01). Our results possibly suggest that MCs may have important roles in the pathogenesis of bovine digital dermatitis.

Keywords: Bovine papillomatous Digital dermatitis; Holstein-Friesian cattle; Immunohistochemistry; Mast cells

Corresponding Author: Ahmet Akkoc, Bursa Uludag University, Faculty of Veterinary Medicine, Department of Pathology, 16059-Bursa, Turkey. E-mail address: aakkoc@uludag.edu.tr

Date of initial submission: 18-03-2022 Date of acceptance: 27-07-2023

#### **INTRODUCTION**

**B** ovine digital dermatitis (BDD) is one of the most common infectious diseases of the foot skin in dairy cattle and causes serious economic losses worldwide. BDD was first described in Italy in 1974 as a disease affecting 70% of adult dairy cattle. The prevalence of the disease in the herd can reach up to 74% (Cheli and Mortellaro, 1974; Orsel et al., 2017). Similarly, there are data showing that the prevalence among herds can vary between 0% and 83% (Holzhaue r et al., 2006). Serious financial losses due to decreased milk production, treatment costs and animal culling are major drawbacks of BDD. Annual cost of BDD has been reported to be around 190 million dollars in the United States (Losinger, 2006).

BDD is characterized macroscopically by papillomatous tissue growths on the plantar surface of the foot over the coronary bands. (Cheli and Mortellaro, 1974; Evans et al., 2016). In histopathological examination, acanthosis, parakeratosis, hyperkeratosis, intraepidermal pustules, infiltrations of inflammatory cells in dermis are major lesions. Although neutrophil leukocytes constitute big portion of these inflammatory cells (Görgül et al., 2002; Demirkan et al., 2018) other mononuclear leukocytes such as lymphocytes, plasma cells and macrophages are also present in dermis. BDD is of multifactorial etiology; large number of anaerobic bacteria and spiral shaped Treponema species are commonly isolated from clinical cases (Evans et al., 2009; Klitgaard et al., 2013; Demirkan et al., 2018; Caddeyand De Buck, 2021). The exact pathogenesis of BDD is still obscure. Hypoxic changes (Gomez et al., 2012) and various bacteria are major actors in the etiology of BDD. Recently, some studies have been conducted on the interaction between causative spirochete species and the immune system to understand the pathogenesis of the disease (Elliott and Alt, 2009; Moe et al., 2010; Refaai et al., 2013). In a study, inflammatory responses of foot skin fibroblasts and keratinocytes against to treponeme wholecell sonicates have been investigated in the context of RANTES/CCL5, MMP-12, TNF-α, TGF-β, and TIMP3 molecules (Evans et al., 2014). They speculated that significantly upregulated expression of target molecules (RANTES/CCL5, MMP-12, TNF-a, TGF- $\beta$ , and TIMP3) in dermal fibroblasts may involve BDD lesion formation. In another study, it was documented that neutrophils that encounter causative agents at inflammation area synthesize more cytokines and chemokines for modulation of inflammation and tissue regeneration (Refaai et al., 2013). In

the same study, T and B lymphocyte distributions and strong upregulation of IL-8 in keratinocytes have been identified in clinical BDD cases. Until now; however, presence, activity and immunophenotypes of mast cells have been omitted in the understanding of the pathogenesis of clinical BDD cases. The present study was designed to reveal the presence, activity, and immunophenotypes of mast cells in clinical BDD cases to fill the aforementioned gap.

Mast cells (MCs) are found in large numbers in certain body parts such as respiratory tract mucosa, gastrointestinal mucosa and skin which are in close contact with the external environment. Therefore, MCs are basically divided into two main subgroups as mucosal and cutaneous mast cells. These cells can also be classified according to their cytoplasmic enzyme contents; chymase (+), tryptase (+) and both tryptase and chymase (+) mast cells. Localizations of bovine MC send their enzyme contents in various organs have been documented, but there is no data regarding their presence in the foot skin. MCs are principal members in allergic reactions; moreover, very recent studies demonstrate they play active roles in the pathogenesis of some infectious diseases. MCs with phagocytic properties engulf infectious disease agents, so it should be kept in mind that mast cells may have local phagocytic functions against infectious agents responsible for the formation of digital dermatitis. On the other aspect, mast cells are known to induce proliferation of various cell types such as endothelial, smooth muscle, mesenchymal and epithelial cells (Marks et al., 1986; Cairns et al., 1996).

The role of MCs in BDD pathogenesis is still obscure. We believe that presence of MCs in bovine foot skin which is very vulnerable area due to constant contact with external environment (feces and urine) may be important in the pathogenesis of BDD. The proliferative effects of MCs granules on various cell types are known, they may contribute to formation of proliferative papillomatous skin lesions which are cardinal clinical and histopathological findings in BDD. According to the authors' knowledge, there is no study on the presence of mast cells in BDD lesions. Since the significance of MCs and their immunophenotypes are not yet clear in bovine foot, in the present study, demonstration, number, distribution and immunophenotyping of mast cells and their activation in healthy and digital dermatitis lesions of dairy cattle was aimed.

#### MATERIALS AND METHODS

# Tissue Sampling and Histopathological Examination

Skin samples were obtained from local slaughterhouses in Bursa, Turkey. A total of 50 skin samples, 25 healthy and 25 with BDD specific lesions were sampled in Holstein-Friesian dairy cattle. The samples were placed in 10% buffered formalin solution and fixed for 48 hours at room temperature. Following fixation, tissues were routinely processed, and 4 micrometer serial sections were cut and adhered to poly-L-lysine-coated slides. All samples were stained with hematoxylin-eosin for microscopic examination and stained with Toluidine blue for demonstration of Mast cells. Tissue samples displaying papillomatous digital dermatitis lesions were also stained with Warthin-Starry silver stain to demonstrate the causative agents (Warthin and Chronister, 1920). BDD was confirmed by gross and histopathological findings.

#### Warthin-Starry Staining for Spirochetes

Sections from healthy and digital dermatitis samples were stained with Warthin-Starry method to demonstrate spirochetal agents. Commercially available Warthin-Starry staining kit from Bio-Optica (04-040903, Italy) was used according to the instructions of manufacturer.

# Demonstration of Mast Cells with Toluidine Blue Staining

Tissue sections were stained with toluidine blue to identify mast cells. Briefly, toluidine blue solution was prepared as follows; 0.5 g of toluidine blue (198161, Sigma Aldrich, St. Louis, MO, USA) was dissolved in 70% ethanol, 5 ml of this solution was taken and mixed with 45 ml of 1% sodium chloride solution, and the sections were incubated in toluidine blue for 30 minutes at room temperature. After washing with tap water, the sections were dehydrated through graded alcohol solutions and cleared in xylene. Images obtained from 10 randomly selected areas at x400 magnification. Intact and degranulated mast cell numbers in healthy and bovine digital dermatitis samples were recorded. Results were expressed as the average number of mast cells.

## **Immunohistochemical Staining**

To investigate mast cells in tissue samples according to their enzyme content, tissues were stained with commercially available anti-chymase (sc-59586, Santa Cruz, USA) and anti-tryptase (sc-59587, Santa Cruz, USA) antibodies, which are known to work in bovine tissues. Deparaffinized tissues were boiled in citrate buffer (pH:6) at 600 watts for 3 times for 2 minutes to reveal antigens. When the temperature of the solution was equalized with room temperature, 3% hydrogen peroxide solution was added on the slides and endogenous peroxidase activity was suppressed. Then, protein block solution was placed on the slides and incubated for 10 minutes. After decanting the protein block solution without any washing, primary antibodies were applied to the tissues and incubated overnight at +4°C. At the end of the period, the tissues were washed 3 times in phosphate buffer solution (PBS) for 5 minutes and secondary antibody (TL015-HDJ, Thermo Scientific, USA) was applied on the tissues and incubated for 20 minutes. At the end of the period, streptavidin biotin solution (TL015-HDJ, Thermo Scientific, USA) was applied to the tissues washed in the same way with PBS and waited for 15 minutes. After a similar washing process, the reaction was completed by applying diaminobenzidine (DAB) on the tissues for 2 minutes. All tissues were counter stained with Harris' hematoxylin solution (HHS16, Sigma Aldrich, USA). During immunohistochemical staining, bovine udder tissue is used as positive control since all phenotypes are present.

#### Statistical analysis

Mean intact and degranulated mast cell counts obtained from healthy and digital dermatitis samples were compared with T-test using SPSS program.

#### RESULTS

#### **Macroscopic Findings**

Severe proliferative skin lesions were noted in the sampled tissues. Just above the heels, nodular, cauli-flower-like, vegetative appearance hyperplastic papillomatous tissue growths with an average size of 3.5 cm  $\pm$  0.4 cm were noted. In the cross-sections applied to the tissues, it was seen that the border of epidermis and dermis could be easily distinguished. Irregular growths in the epidermis were also evident (Figure-1 and 2).

#### **Histopathological Findings**

In light microscopic examination, severe hyperkeratosis (25 cases), parakeratotic hyperkeratosis (21 cases), sub corneal pustular epidermitis (19 cases), balloon-like degeneration of epithelial cells (25 cases), dermal neutrophil leukocyte infiltrates (25 cases), lymphocytic plasmocytic mononuclear cell infil-



Figure 1. Epithelial overgrowths (arrow) and wart formation in the digital skin of affected cattle.



Figure 2. Intraepidermal pustular changes, infiltration of polymorphonuclear leukocytes (arrow), acanthosis and hyperkeratosis (A), hematoxylin-eosin staining, X200 magnification.

trations were observed in tissue samples (25 cases). Hyperemia in dermal blood vessels and endothelial hypertrophy were evident. Increased amount of collagenous matrix and fibroblastic proliferation were noted in the dermis (Figure-2). In healthy animals, normal architecture of the skin was preserved. Keratinized stratified squamous epithelial cells and the dermis rich in collagen tissue were observed within normal limits. None of the skin samples in healthy animals displayed prominent inflammatory changes.

## Warthin-Starry Staining

Twenty-two out of 25 (88%) digital dermatitis

samples were positively stained with the kit. Spiral-shaped black bacteria on a yellowish background were demonstrated. Most of the agents were located within deeper acanthotic epidermal layer, in some cases (3 out of 22, 13%) spirochetal bacteria were also observed just beneath the stratum basale of epidermis.

#### **Toluidine Staining**

In animals in the control and digital dermatitis groups, mast cells with granules showing metachromasia in toluidine staining were found just below the epidermis and around the blood vessels located in the dermis (Figure-3 and 4). It was observed that the mean number of mast cells in skin samples with digital dermatitis was higher than in the control groups. An average of  $4.6 \pm 2$  mast cells were found in each objective area in the foot skin sections of the healthy animals, while this number was  $9.3 \pm 1$  in digital dermatitis. When the number of mast cells in healthy samples was normalized to 1, digital dermatitis samples were displayed 2.02-fold increase in number of mast cells. Average degranulated mast cell numbers in healthy and digital dermatitis samples were 1.2and 5.1, respectively. Again, when we normalized the number of degranulated mast cells in healthy samples to 1, digital dermatitis samples had 4.2-fold increase in the number of degranulated mast cells. In the statistical analysis, differences in the mean intact and degranulated mast cells were found to be significant (mean intact: p<0.05, mean degranulated: p<0.01).

#### **Immunohistochemical Staining**

To demonstrate the phenotype of mast cells in healthy and digital dermatitis samples we used commercially available primary anti-chymase and anti-tryptase antibodies. Staining with anti-chymase and anti-tryptase antibodies showed only tryptase (+) mast cells were positively reacted with antibody (Figure-5 and 6). No chymase (+) mast cells were iden-



Figure 3. Mast cells with metachromatic granules (arrows) in healthy skin samples, Toluidine blue staining, X400 magnification.



Figure 4. Intact mast cells with metachromatic granules in their cytoplasms(arrows) and degranulated mast cells (open arrows) in skin sample with digital dermatitis, Toluidine blue staining, X400 magnification.

tified. Chymase (+) mast cells were demonstrated in bovine mammary tissue samples which were previously known to have both chymase (+) and tryptase (+) phenotype MCs.

## DISCUSSION

In this study, presence of mast cells and their immunophenotypes in the foot skin of dairy cattle with digital dermatitis were investigated for the first time. Dairy cattle breeding is extremely important for the economy and human health. Foot diseases are serious welfare and health problems, and they are resulted with production losses in dairy cattle. BDD is indisputably one of the most important foot diseases of cattle due to the severity of lesions and higher prevalence in herds. The foot skin is constantly subjected to external environmental stimuli such as physical trauma, urine, and feces, and depending on the stage of BDD, ulcerative wounds and/or papillomatous growths is formed on the heels. Determination of the presence and immunophenotype of cutaneous mast cells of foot skin may provide useful data for BDD.



Figure 5. Tryptase positive mast cells (arrows) in the dermis of healthy skin samples. Streptavidin-biotin peroxidase, DAB chromogen, X400 magnification.



Figure 6. Tryptase positive mast cells (arrows) in the dermis of skin samples with digital dermatitis. Streptavidin-biotin peroxidase, DAB chromogen, X400 magnification.

J HELLENIC VET MED SOC 2023, 74 (4) ПЕКЕ 2023, 74 (4) Histopathologically, severe acanthosis, infiltrations of large numbers of polymorphonuclear (especially neutrophils) and mononuclear leukocytes in skin layers have been reported, but no data were found regarding the presence of MCs among those inflammatory cells. To fill this gap and shed light on future studies on the pathogenesis and treatment options of the disease, mast cells and their immunophenotypes in bovine foot skin were investigated in the submitted study. For the confirmation of BDD, in addition to unique gross findings, Warthin-Starry staining procedure was followed to demonstrate causative agents on suspected tissue samples. Most of digital dermatitis samples (22 out of 25) were stained positively; black spiral bacteria on a yellowish background were evident. The rest three samples did not show any positive reaction with Warthin-Starry staining; however, those samples were having characteristic gross and microscopic changes with no hesitation.

Mast cells are unique cells of the immune system and play important roles in allergic reactions and infectious diseases. MCs contain multifunctional cellular granules which are primarily responsible for modulation and regulation of inflammation. Their functions in allergy, asthma, anaphylaxis, and various immune responses are known in detail, but results of some studies on the function of mast cells direct attention into the infectious diseases (Siebenhaar et al., 2007; DiNardo et al., 2008; Nakamura et al., 2013; Zimmermann et al., 2019). In addition to classical functions of MCs, their phagocytic properties have been documented (Lima et al., 2013). Moreover, upon bacterial challenge, MCs quickly responds and secrete many inflammatory cytokines like TNF- $\alpha$ , IL-3, IL-4 and IL-13 to orchestrate other immune cells. Zuerneret al. (2007) have reported that innate immune functions of bovine macrophages were hampered by treponemal cell wall fragments in vitro. In their study, they exposed macrophages to treponemal cell sonicates and evaluated many inflammatory genes. According to their data, IL-1, -6, -11 genes were downregulated or unchanged and genes encoding proteins responsible for antigen presentation were downregulated. While mast cells are known with their undeniable functions in innate immunity and they serve as a pool of cytokines (Mukai et al., 2018), it is intriguing that their possible roles in BDD have not been investigated.

Albeit in small numbers, there are data on the localization of mast cells in various organs of cattle, (Küther et al., 1998; Özen et al., 2007; Ertuğrul and Kurtdede, 2017). In a study, the presence of mast cells in urinary system tissues of cattle were demonstrated histochemically (Ertuğrul and Kurtdede, 2017) Özenet al. (2007) successfully demonstrated the presence of mast cells in the ovarian tissues of healthy cows. However, these studies did not provide any information on immunophenotypes of mast cells on various locations. Kütheret al. (1998) showed that mast cells with different immunophenotypes were found in forestomach, duodenum, lung, lymph node, uterus and skin. According to their findings, mast cells were primarily located in the dermal layer of skin, only tryptase (+) mast cells were noticed immunophenotypically. In the submitted study, mast cells were successfully identified in the dermis of healthy and digital dermatitis samples. We noticed that MCsin dermis were only positive for tryptase phenotype both in healthy and digital dermatitis lesions. No positivity was found in staining with anti-chymase antibody. To confirm the specificity of the chymase antibody, we used bovine mammary tissue as positive control and chymase positive mast cells were successfully demonstrated in udder tissue.

The submitted study includes comparison of intact and degranulated mast cells and mast cell immunophenotypes in healthy and papillomatous digital dermatitis lesions in cattle. We observed 2.02 and 4.2fold increase in number of intact and degranulated MCs in healthy and digital dermatitis samples, respectively. The presence of higher number of cutaneous intact and degranulated mast cells in animals with digital dermatitis when compared with healthy animals strongly suggest that MCs may play important roles in the pathogenesis of BDD. Increased amount of fibrotic tissue and acanthotic changes in keratinocytes were noticed in all digital dermatitis samples. Since mast cell tryptase is known as a potent mitogen for epithelial cells and fibroblasts, we believe that increased number of tryptase (+) mast cell in dermis may contribute to the proliferation of keratinocytes and fibroblasts inBDD. It is known that different immunophenotypes of mast cells are predominanttype in certain inflammatory reactions (Rozniecki et al., 1995; Knight et al., 2000; Huang et al., 2001; Caughey, 2007). Moreover, heterogeneous population of MCs have been reported in lung tissue even in different microanatomical positions (Bradding, 2009). Some in vitro studies showed that MCs with one phenotype can shift to another MC phenotype in special culture circumstances (Mierke et al., 2000; Hsieh et al., 2005).

#### CONCLUSION

We successfully demonstrated MCs in bovine digital dermatitis lesions for the first time. Presence of MCs in bovine foot and determination of their phenotypes may bring new perspectives into BDD management. There are many chemical compounds capable of inhibition and stabilization of MCs, these chemicals may be useful in the alleviation of hyperplastic epidermal lesions. Further clinical studies are strongly needed to test the action of MC stabilizers or inhibitors in the treatment of BDD.

#### ACKNOWLEDGMENTS

This study was partly supported financially by The Scientific and *Technological Research Council of Türkiye* (TUBITAK/TOVAG 1180893).

#### **CONFLICT OF INTEREST**

None declared by the authors.

#### REFERENCES

- Bradding P (2009)Human lung mast cell heterogeneity. Thorax64:278-280. Caddey B, De Buck J (2021)Meta-analysis of bovine digital dermatitis
- microbiota reveals distinct microbial community structures associated with lesions.Front Cell Infect Microbiol 16:685-861.
- Cairns JA, Walls AF (1996) Mast cell tryptase is a mitogen for epithelial cells. Stimulation of IL-8 production and intercellular adhesion molecule-1 expression. J Immunol 156:275-283.
- Caughey GH (2007)Mast cell tryptases and chymases in inflammation and host defense.Immunol Rev 217:141-154.
- CheliR, Mortellaro C (1974) La dermatite digitale del bovino. In: 8th International Meeting on Diseases of Cattle, Milan, Italy: pp 208-213.
- Di Nardo A, Yamasaki K, Dorschner RA, Lai Y, Gallo RL (2008) Mast cell cathelicidin antimicrobial peptide prevents invasive group A Streptococcus infection of the skin.J Immunol180:7565-7573.
- Elliott MK, Alt DP (2009) Bovine immune response to papillomatous digital dermatitis (PDD)-associated spirochetes is skewed in isolate reactivity and subclass elicitation.Vet Immunol Immunopathol 130:256-261.
- Evans NJ, Brown JM, Demirkan I, Murray RD, Birtles RJ, Hart CA, Carter SD (2009) Treponema pedis sp.nov., a spirochaeta isolated from bovine digital dermatitis lesions. Int J Syst Evol Microbiol59:987-991.
- Evans NJ, Brown JM, Scholey R, Murray RD, Birtles RJ, Hart CA, Carter SD (2014)Differential inflammatory responses of bovine foot skin fibroblasts and keratinocytes to digital dermatitis treponemes. Vet Immunol Immunopathol 161:12-20.
- Evans NJ, Murray RD, Carter SD (2015) Bovine digital dermatitis: Current concepts from laboratory to farm. Vet J211:3-13.
- Gomez A, Cook NB, Bernardoni ND, Rieman J, Dusick AF, Hartshorn R, Socha MT, Read DH, Döpfer D (2012) An experimental infection model to induce digital dermatitis infection in cattle.J Dairy Sci95:1821-1830.
- Görgül OS, Kahraman MM, Çeçen G, Akkoç A, Gül NY, Sevimli A (2002) Sığırlarda digital ve interdigital dermatitis'lerdekliniktanı, sağaltımvehistopatolojik bulgular. J Res Vet Med21:115-124.
- Hsieh FH, Sharma P,Gibbons A, Goggans T, Erzurum SC, Haque SJ (2005) Human airway epithelial cell determinants of survival and functional phenotype for primary human mast cells.Proc Natl Acad Sci USA102:14380-14385.
- Holzhauer M, Hardenberg C, Bartels CJ, Frankena KJ (2006)Herdand cow- level prevalence of digital dermatitis in the Netherlands and associated risk factors. Dairy Sci89:580-588.
- Huang C, De Sanctis GT, O'Brien PJ, Mizgerd JP, Friend DS, Drazen JM, Brass LF, Stevens RL (2001) Evaluation of the substrate specificity of human mast cell tryptase beta I and demonstration of its importance in bacterial infections of the lung.J Biol Chem276:26276-26284.
- Klitgaard K, Foix Bretó A, Boye M, Jensen TK (2013) Targeting the treponemal microbiome of digital dermatitis infections by high-resolution phylogenetic analyses and comparison with fluorescent in situ hybridization. J Clin Microbiol51:2212-2219.
- Knight PA, Wright SH, Lawrence CE, Paterson YY, Miller HR(2000) De-

layed expulsion of the nematode Trichinella spiralis in mice lacking the mucosal mast cell-specific granule chymase, mouse mast cell protease-1.J Exp Med192:1849-1856.

- Küther K, Audigé L, Kube P, Welle M (1998) Bovine mast cells: distribution, density, heterogeneity, and influence of fixation techniques. Cell Tissue Res293:111-119.
- Lima HG, Pinke KH, Gardizani TP, Souza-Júnior DA, Carlos D, Avila-Campos MJ, Lara VS (2013) Mast cells act as phagocytes against the periodontopathogen Aggregatibacter Actinomycetemcomitans.J Periodontol84:265-272.
- Losinger WCJ (2006)Economic impacts of reduced milk production associated with papillomatous digital dermatitis in dairy cows in the USA. Dairy Res73:244-256.
- Marks RM, Roche WR, Czerniecki M, Penny R, Nelson DS (1986) Mast cell granules cause proliferation of human microvascular endothelial cells.Lab Invest 55:289-294.
- Mierke CT, Ballmaier M, Werner U, Manns MP, Welte K, Bischoff SC (2000)Human endothelial cells regulate survival and proliferation of human mast cells.J Exp Med192:801-811.
- Moe KK, Yano T, Misumi K,Kubato C, Yamazaki W, Muguruma M, Misawa N (2010) Analysis of the IgG immune response to Treponema phagedenis-like spirochetes in individual dairy cattle with papillomatous digital dermatitis.Clin Vaccine Immunol17:376-83.
- Mukai K, Tsai M, Saito H, Galli SJ (2018) Mast cells as sources of cytokines, chemokines, and growth factors. Immunol Rev282:121-150.
- Nakamura Y, Oscherwitz J, Cease KB (2013)Staphylococcus δ-toxin induces allergic skin disease by activating mast cells.Nature503:397-401.
- Orsel K, Plummer P, Shearer J, De Buck J, Carter SD, Guatteo R, Barkema HW (2018) Missing pieces of the puzzle to effectively control digital dermatitis.TransboundEmerg Dis65:186-198.
- Refaai W, Ducatelle R, Geldhof P, Mihi B, El-shair M, Opsomer G (2013) Digital dermatitis in cattle is associated with an excessive innate immune response triggered by the keratinocytes.BMC Vet Res9:193.
- Rozniecki JJ, Hauser SL, Stein M, Lincoln R, Theoharides TC (1995) Elevated mast cell tryptase in cerebrospinal fluid of multiple sclerosis patients. Ann Neurol37:63-66.
- Siebenhaar F, Syska W, Weller K, Magerl M, Zuberbier T, Metz M, Maurer M (2007) Control of Pseudomonas aeruginosa skin infections in mice is mast cell-dependent. Am J Pathol170:1910-1916.
- Warthin AS, Chronister AC (1920) A more rapid and improved method of demonstrating spirochetes in tissues (Warthin and Starry's cover-glass method). AmJSyp4:97-103.
- Zimmermann C, Troeltzsch D, Giménez-Rivera VA, Galli SJ, Metz M, Maurer M, Siebenhaar F (2019)Mast cells are critical for controlling the bacterial burden and the healing of infected wounds. Proc Natl Acad Sci USA116:20500-20504.
- Zuerner RL, Heidari M, Elliott MK, Alt DP, Neill JD (2007) Papillomatous digital dermatitis spirochetes suppress the bovine macrophage innate immune response.Vet Microbiol125:256-264.

J HELLENIC VET MED SOC 2023, 74 (4) ПЕКЕ 2023, 74 (4)