

## Journal of the Hellenic Veterinary Medical Society

Vol 74, No 3 (2023)



### Effects of Antioxidants Supplemented Feed in Coccidiosis Treatment, Blood Antioxidative Status, and Enzymatic Activity of Domestic Cats

*B Bizhga, S Cocoli, A Stevanović, S Bajić, E Lika, T Shtylla Kika, N Puvača*

doi: [10.12681/jhvms.30849](https://doi.org/10.12681/jhvms.30849)

Copyright © 2023, B Bizhga, S Cocoli, A Stevanović, S Bajić, E Lika, T Shtylla Kika, N Puvača



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

### To cite this article:

Bizhga, B., Cocoli, S., Stevanović, A., Bajić, S., Lika, E., Shtylla Kika, T., & Puvača, N. (2023). Effects of Antioxidants Supplemented Feed in Coccidiosis Treatment, Blood Antioxidative Status, and Enzymatic Activity of Domestic Cats: Natural antioxidants against endoparasites. *Journal of the Hellenic Veterinary Medical Society*, 74(3), 6099–6106. <https://doi.org/10.12681/jhvms.30849>

## Effects of Antioxidants Supplemented Feed in Coccidiosis Treatment, Blood Antioxidative Status, and Enzymatic Activity of Domestic Cats

B. Bizhga<sup>1</sup>, S. Cocoli<sup>1</sup>, A. Stevanović<sup>2</sup>, S. Bajić<sup>3</sup>, E. Lika<sup>1</sup>, T. Shtylla Kika<sup>1</sup>,  
N.Puvača<sup>4,\*</sup>

<sup>1</sup>Faculty of Veterinary Medicine, Agricultural University of Tirana, Tirana, Albania

<sup>2</sup>High Medical - Sanitary School of Professional Studies, Visan", Belgrade-Zemun, Serbia

<sup>3</sup>Faculty of Technical Sciences, University of Novi Sad, Novi Sad, Serbia

<sup>4</sup>Department of Engineering Management in Biotechnology, Faculty of Economics and Engineering Management in Novi Sad, University Business Academy in Novi Sad, Novi Sad, Serbia

**ABSTRACT:** As coccidia become increasingly resistant to anticoccidial drugs, efforts have been made to find alternatives. In recent years, botanicals have been reported as potential alternatives to anticoccidials since they are effective against protozoa, arthropods, and helminths. In this study, different doses of dried pomegranate fruit (*Punica granatum* L.) were evaluated for their effectiveness in reducing the number of oocysts in domestic cats and their antioxidant properties. Under *in vivo* conditions, 24 six-month-old domestic cats of both genders naturally infected with *Cytosporafelis* were tested. Four equal groups of infected cats were formed. Six cats made up each group. The control group (C) was on a basal diet, and one group received the chemical coccidiostat robenidine (CR), supplemented in the amount of 0.5%. Two levels of whole dried pomegranate fruit as a natural antioxidant were applied in a concentration of 0.5% (P1) and 1.0% (P2) on top of the basic diet. With the McMaster technique, oocysts number and eggs per gram of feces were determined. From each cat, 6 per group, blood samples were taken from a jugular vein at the end of the experimental period to investigate the influence of dried pomegranate fruit on blood enzymatic activity and lipid oxidation. In conclusion, supplementing cats' diets with dried pomegranate fruit reduced the number of oocysts per gram of feces significantly, but it is important to carry out further and more detailed studies to prove the anticoccidial and antioxidant properties of dried pomegranate fruit in cats' diets.

**Keywords:** cats; coccidiosis; parasites; pomegranate, nutrition.

*Corresponding Author:*

Prof. dr Nikola Puvača, DVM, PhD, Department of Engineering Management in Biotechnology, Faculty of Economics and Engineering Management Novi Sad, University Business Academy, 8 Cvečarska 2, 21000, Novi Sad, Serbia.  
E-mail address: nikola.puvaca@fimek.edu.rs

*Date of initial submission:* 18-07-2022  
*Date of acceptance:* 27-07-2022

## INTRODUCTION

Cats' and dogs' feces have been harboring coccidian parasites for more than a century, but their clinical and public health significance was unknown until 1970 (Cox, 2002). The feces of cats and dogs were thought to contain only one species of coccidia before Wenyon's report in 1923 (Wenyon, 1923). It was determined that three coccidian parasites exist based on their oocyst sizes, the parasite with large oocysts (*Cystoisosporafelis*), medium-sized oocysts (*Cystoisosporarivolta*), and small-sized oocysts (*Cystoisosporabigemina*), and in addition, it was believed that they are non-host specific (Duszynski, 2021). The parasites were later demonstrated to be host-specific with extraintestinal stages as well. Rodents, dogs, cattle, pigs, camels, albino mice, and rabbits can serve as paratenic hosts according to bioassay studies (Dubey et al., 2007; Gajadhar et al., 2015; Koutsoumanis et al., 2018). Cats infected with *C. felis* excrete *Toxoplasma gondii* oocysts without showing any clinical signs, while *C. rivolta* was pathogenic for a newborn but not for weaned cats (Dubey, 2018). Diarrhea usually begins 3-4 days after the newborn cat has been injected with 100,000 sporocysts or infected mice. Cats are reported to be at risk of infection with *C. felis*, but there is conflicting information regarding its pathogenicity (Shaw et al., 2004). There have been reports of cats infected with enteric viruses experiencing severe diarrhea, especially before prophylactic antibiotics were available (Truyen et al., 2009). The treatment of *Cystoisospora* infection in cats has been made possible by a variety of anticoccidial drugs. When cats are administered toltrazuril during the prepatent period, the number of oocysts excreted is significantly reduced (Litster et al., 2014; Petry et al., 2011).

In the last decade, there have been questions raised about the use of antibiotics, such as sulfonamides as well as salinomycin and robenidine, in poultry nutrition for the treatment of coccidiosis. As avian coccidia (protozoa) become increasingly resistant to anticoccidial drugs, efforts have been made to find alternatives.

In recent years, botanicals have been reported as potential alternatives to anticoccidials since they are effective against protozoa, arthropods, and helminths (Puvača et al., 2022). Pomegranate fruit (*Punica granatum* L.) is a nutrient-dense fruit rich in phytochemical compounds (Juhaimi et al., 2017). As a mechanism of defense, plants produce low molecular weight

compounds called phytochemicals. About 50% of the total fruit weight corresponds to the peel, which is an important source of bioactive compounds such as phenolics, flavonoids, ellagitannins, and proanthocyanidin compounds. Due to their biological and free radical scavenging activities, phenolic compounds, flavonoids, anthocyanins, and tannins are the most important antioxidant phytochemicals (Elfalleh et al., 2011). Many studies have demonstrated that combinations of pomegranate extracts from different parts of the fruit are more effective than single extracts of the fruit because different parts of the fruit contain different phenolic acids, flavonoids, and tannins (Husain et al., 2021).

As a result of normal cellular metabolism, reactive oxygen species (ROS) are produced by living organisms, leading to cell oxidative destruction, also known as oxidative stress. Humans, as well as animals, may suffer from numerous pathological changes caused by oxidative stress, including membrane dysfunction, DNA damage, protein inactivation, cancer, arthritis, and neurodegenerative diseases (Islam, 2017). In contrast, aerobic organisms possess highly effective antioxidants that are usually effective in blocking the negative effects of ROS. These antioxidants are both enzymatic and nonenzymatic, therefore, defensive processes involve a variety of protective mechanisms (Uttara et al., 2009).

Activated oxygen is eliminated through the action of antioxidant enzymes. Several enzymes, such as superoxide dismutase (SOD), convert superoxide anion radicals into  $H_2O_2$  and reduce  $H_2O_2$  to water, and glutathione peroxidase (GSHPx) reduces  $H_2O_2$  and terminates lipid peroxidation. The enzyme glutathione reductase (GR) can also regenerate glutathione (GSH), which plays an important role in the reduction of acute toxicity of xenobiotics and lipid peroxidation products (Bailly, 2004). By stealing electrons from cell membrane lipids, free radicals cause lipid peroxidation, which damages the cells. Polyunsaturated fatty acids are particularly susceptible to this since they contain multiple double bonds, between which there are methylene bridges (-CH<sub>2</sub>-) with highly reactive hydrogens, resulting in lipid peroxides such as malondialdehyde (MDA) and hydroxyl, which damage tissues (Tsikas, 2017). Consequently, MDA content is a significant indicator of the level of lipid peroxidation and an indirect indicator of cell damage (Bhattacharjee, 2014).

In this study, different doses of dried pomegranate

fruit (*Punica granatum* L.) were evaluated for their effectiveness in reducing the number of oocysts in domestic cats and their antioxidant properties.

## MATERIALS AND METHODS

The animal study protocol was approved by the Ethics Committee of University BA in Novi Sad, Serbia (EC-01/1/22).

### Experimental design and pre-experimental investigation

Domestic cats were observed and documented daily for natural infection before the experimental trial. All cats were found to be infected with *Cystoisospora felis* given the mild bloody diarrhea which was confirmed based on fecal examination. The direct smear and the saturated sugar floatation technique were

used for determining the presence and identification of oocysts. Using the McMaster technique, coccidia oocysts per gram (OPG) of feces were determined.

Under *in vivo* conditions, 24 six-month-old domestic cats of both genders naturally infected with *C. felis* were tested. All pet cats were observed in their home environment; therefore, four equal groups of infected cats were formed. Six cats made up each group. The control group (C) was on a basal diet, and one group received the chemical coccidiostat robenidine (CR), supplemented in the amount of 0.5%. Two levels of whole dried pomegranate fruit as a natural antioxidant were applied in a concentration of 0.5% (P1) and 1.0% (P2) on top of the basic diet (Tables 1 and 2). From 6 to 8 months of age, the cats were monitored. A standard basal diet was fed to the cats, with access to water and food as needed.

**Table 1.** Ingredient composition of domestic cats' diets.

Ingredients, %	C	CR	P1	P2
Boneless chicken	30.0	30.0	30.0	30.0
Dehydrated chicken protein	28.0	28.0	28.0	28.0
Sweet potato	4.3	4.3	4.3	4.3
Chicken fat	3.0	3.0	3.0	3.0
Dehydrated eggs	9.0	9.0	9.0	9.0
Dehydrated herring protein	9.0	9.0	9.0	9.0
Fish oil	8.0	8.0	8.0	8.0
Alfalfa flour	2.5	2.5	2.5	2.5
Sodium chloride	0.2	0.2	0.2	0.2
Dry beer yeast	6.0	6.0	6.0	6.0
Coccidiostat robenidine	0.0	0.5	0.0	0.0
Dried pomegranate fruit	0.0	0.0	0.5	1.0

C - control group; CR - group received chemical coccidiostat robenidine; P1 - group received dried pomegranate fruit in the concentration of 0.5%; P2 - group received dried pomegranate fruit in the concentration of 1.0%.

**Table 2.** Chemical composition of domestic cats' diets.

Nutrients	%
Crude protein	44.0
Crude fat	20.0
Crude fiber	1.8
Crude moisture	8.0
Crude ash	8.5
Calcium	1.1
Phosphorus	0.9
Magnesium	0.08
Omega 6 fatty acids	3.3
Omega 3 fatty acids	0.9
ME, MJ/kg	17.59

Additives per 1kg of feed: Vitamin A 18000IU; vitamin D3 1200IU; vitamin E 600mg; vitamin C 300mg; niacin 150mg; calcium D-pantothenate 50mg; vitamin B2 20mg; vitamin B6 8.1mg; vitamin B1 10mg; biotin 1.5mg; folic acid 1.5mg; vitamin B12 0.1mg; choline chloride 250mg; beta-carotene 1.5 mg; zinc (zinc chelate methionine hydrate) : 163.8mg; manganese (manganese chelate methionine hydrate) : 64.6mg; iron (iron- (II) -chelate glycine hydrate) : 58.3mg; copper (copper chelate of methionine hydrate) : 15.8mg; DL-methionine 5000mg; taurine 4000 mg; L-carnitine 300 mg.

### Oocyst count (OPG index)

A fecal sample was taken from each cat before natural antioxidants and chemical coccidiostat were introduced into the diets. An oocyst count per gram of feces was measured at the end of six, seven, and eight months of the cat's life. The number of oocysts in ten samples per gram of feces from each group was determined by collecting fecal samples of around 30 g from each group using mosquito net mesh. Refrigeration was used to keep the samples before analysis. With the McMaster technique, OPG and eggs per gram of feces were determined.

### Preparation of blood hemolysates

From each cat, 6 per group, blood samples were taken from a jugular vein after fasting overnight at the end of the eighth month and the end of the experimental period. The plasma was separated by centrifugation (3500 g, 10 min) and stored at -70 °C for analysis of biochemical parameters. The remaining red blood cells (RBCs) were washed three times with isotonic NaCl (0.15 M) before use in the biochemical assay. The RBC hemolysates were prepared by diluting RBCs 1:10 with ice-cold double distilled water, shaken vigorously to force hemolysis, and stored at -70 °C (Fébel et al., 2008).

### Determination of enzymatic activity

The glutathione peroxidase (GSHPx) activity was determined by spectrophotometric measurement of absorbance at 412 nm with cumene hydroperoxide as the substrate (Chiu et al., 1976), while malondialdehyde (MDA) concentration in the serum was measured by the 2-thiobarbituric acid assay. Further on,


superoxide-dismutase (SOD) activity was determined by the spectrophotometric method based on the inhibition of adrenalin reduction to adrenochrome at pH 10.2 (L. M. Kostadinović et al., 2011), and activity of the glutathione-reductase (GR) was determined from the rate of NADPH oxidation and it was monitored by measuring the absorbance at 340 nm (Łukasiewicz-Hussain & Moniuszko-Jakoniuk, 2004).

### Statistical analysis

Significant effects were evaluated using ANOVA. Fisher's l.s.d. post-hoc multiple-range test was used to ascertain differences among groups. A significance level of  $P = 0.05$  was used. Statistical analyses were conducted using the statistical software program Statistica 13 for Windows (StatSoft, Inc., Tulsa, OK, USA).

## RESULTS AND DISCUSSION

The addition of dried pomegranate fruit to the diet of domestic cats led to statistically significant ( $P < 0.05$ ) differences in the number of oocysts per gram of feces (Figure 1). A statistically significant difference in OPG between the cats was not observed at the beginning of the study ( $P > 0.05$ ). Later, the level of contamination by oocysts in the C group was systematically higher compared to the other groups ( $P < 0.05$ ), while there was no significant difference between groups P1 and P2. Finally, at the end of the experimental period, the lowest number of oocysts achieved was with group P1 (36998 oocysts per g of feces) followed by groups P2 (37510 oocysts per g of feces) and CR (38123 oocysts per g of feces).



**Figure 1.** Coccidiosis oocyst counts in domestic cats' fecal samples (number of oocyst per gram of feces). C - control group; CR - group received chemical coccidiostat robenidine; P1 - group received dried pomegranate fruit at the concentration of 0.5%; P2 - group received dried pomegranate fruit at the concentration of 1.0%.

In the present study the antioxidant defense system (both enzymatic and nonenzymatic) in blood hemolysates (Table 3) of the cats, was investigated. As can be seen from Table 3 and Figure 2, the content of glutathione (GSH) and malondialdehyde (MDA) in the blood hemolysates of the cats treated with dried pomegranate fruit showed a decrease in comparison with the control (C) and robenidine (CR) groups ( $P < 0.05$ ), while glutathione peroxidase, superoxide dismutase, and glutathione reductase activities significantly increased in the P1 group compared to the other diets.

Finally, it was noticed that the content of GSH in the blood samples of cats treated with dried pomegranate fruit showed a significant decrease ( $P < 0.05$ ) in comparison with the control and robenidine groups. The same tendency was recorded for MDA content. The GSHPx activity in samples was the highest in the P1 group and was statistically significantly different ( $P < 0.05$ ) from the other groups. A similar tendency was observed in the activities of SOD and GR in the blood samples of cats, where the highest values were


recorded in the group receiving 0.5% of dried pomegranate fruit with a significant difference ( $P < 0.05$ ) in comparison with the control and robenidine groups.

To our knowledge, there are no relevant studies in the literature on the effects of pomegranate fruit (*Punica granatum* L.) on cats in the fight against coccidiosis. Consequently, our findings will be compared with other published studies about coccidiosis in companion and food animals, respectively. The results obtained with other animal species with the addition of medicinal plants as alternatives to coccidiosis indicate numerous positive effects on animals' health (Hashemi & Davoodi, 2011; L. Kostadinović & Lević, 2018). When it comes to cats, animal shelter kittens are often afflicted with intestinal coccidiosis, and its pathogenesis remains unclear (Dubey et al., 2009). In mammals, coccidia regularly inhabits the intestinal tracts as obligate intracellular parasites (Mundt et al., 2006). The most common species of coccidia in cats, *C. felis* and *C. rivolta*, appear to be influenced by age and living conditions (Dubey, 2018). The rate of nat-

**Table 3.** The effect of dried pomegranate fruit on the antioxidative systems of cats' blood.

Parameters	C	CR	P1	P2
GSH (nmol/mg protein)	7.24±0.6 <sup>a</sup>	7.02±0.2 <sup>a</sup>	5.98±0.2 <sup>b</sup>	5.64±0.4 <sup>b</sup>
MDA (nmol/mL)	0.81±0.5 <sup>b</sup>	0.69±0.11 <sup>a</sup>	0.57±0.3 <sup>a</sup>	0.58±0.8 <sup>a</sup>
GSHPx (nmol/mg protein min)	8.51±0.4 <sup>c</sup>	9.31±0.1 <sup>b</sup>	9.66±0.1 <sup>a</sup>	9.54±0.2 <sup>a</sup>
SOD (nmol/mg protein min)	33.24±1.2 <sup>c</sup>	78.70±1.9 <sup>b</sup>	93.61±0.8 <sup>a</sup>	81.00±2.4 <sup>b</sup>
GR (nmol/mg protein min)	15.87±0.4 <sup>c</sup>	18.20±0.3 <sup>b</sup>	21.96±0.1 <sup>a</sup>	18.28±0.2 <sup>b</sup>

Means within a row followed by the different letters are significantly different ( $P < 0.05$ ). C - control group; CR - group received chemical coccidiostat robenidine; P1 - group received dried pomegranate fruit in a concentration of 0.5%; P2 - group received dried pomegranate fruit in a concentration of 1.0%.



**Figure 2.** Positive differences of dried pomegranate fruit on the antioxidative systems of cats' blood compared to control group of cats, %.

ural infection is highest in neonates and kittens less than three months old. Synthetic drugs are common in the treatment of coccidiosis in cats. Different drugs such as sulfadimethoxine, toltrazuril, and ponazuril have been regularly used. Dosing recommendations have ranged from only one dose, one dose followed by a second dose five days later, to daily doses for three or five days, also depending on drug availability in certain countries and on the approval of drug usage by the Food and Drug Administration (FDA). Resistance of coccidia to currently used coccidiostats to treat coccidiosis represents a serious problem in veterinary practice (Zhang et al., 2013). The use of whole plants or plant extracts to treat coccidiosis is not a new approach (Abbas et al., 2012; Naidoo et al., 2008). When searching for the best natural extract to treat coccidiosis, it is necessary to take into account that the extract needs to be at least partially soluble in lipids to penetrate the cellular membrane, because coccidia is located inside the cells. Besides, herbs and spices can protect the feed against oxidative deterioration during storage. This is a widely used practice in the pet food and human food industry (Vapa Tankosić et al., 2022). Naidoo et al. (2008) studied the capacity of four African plants which would be appropriate to treat coccidiosis: leaves of *Combretum woodii*, leaves and stem of *Artemisia afra*, a whole plant, and seeds of *Vitis vinifera*. Extracts of all chosen plants improved the feed conversion in poultry to the same extent as coccidiostats toltrazuril used in cats' treatment against coccidiosis. The best effect was seen with *Tulbaghiavioleacea*, which also partially lowered the shedding of oocysts. Furthermore, Popović et al. (2017) have investigated wormwood (*Artemisia absinthium*) as a potential anticoccidial drug in the rabbit diet, as well as a growth promoter and antioxidant promoter, and obtained similar positive results as in our present study. In the experiment with tea tree essential oil Puvača et al. (2020) showed that this essential oil can be usefully used in the treatment of coccidiosis of mature laying hens. Nowadays, significantly higher attention is being paid to coccidia in companion and food animals than earlier (Pritt et al., 2012). A variety of parasites have been successfully treated with medicinal herbs in a variety of studies, and enviable success has been achieved against a variety of parasites (Culliney, 2005). The most active components in such as trans-sabinene,  $\alpha$ - and  $\beta$ -thujon, trans-sabinylac-

etate, linalyl acetate, and cis-chrysanthenyl acetate, have been responsible for the anticoccidial activity of plants in animals (Ismail et al., 2021; Puvača et al., 2022).

On the other side, in a chain reaction, antioxidant enzymes like glutathione peroxidase and superoxide dismutase can break down free radicals. As a substrate for glutathione peroxidase, glutathione serves as an antioxidant protector, reducing the acute toxic effects of xenobiotics and lipid peroxidation products (Agarwal et al., 2012; Ighodaro & Akinloye, 2018). Lipid oxidation contributes to the deterioration of physiological functions such as growth, reproduction, and immunity, making individuals more susceptible to infectious diseases (Sultana et al., 2017). To detect the level of oxidative stress in the body, malondialdehyde is used as a sensitive indicator (Beaulieu & Costantini, 2014). As a simple way to assess lipid peroxidation in biological materials, determining malondialdehyde has attracted widespread interest. Thus, the increased activity of all antioxidant enzymes in our study may be attributed to the environmental pathogens of *C. felis* in domestic cats. The results indicate that pomegranate fruit (*Punica granatum* L.) could improve antioxidant activities and reduce the damage caused by free radicals in domestic cats.

## CONCLUSION

According to the results obtained, cats can benefit greatly from supplementing their diet with dried pomegranate fruit, without experiencing any adverse effects. Due to its ability to activate the enzymes responsible for antioxidant protection in animals, dried pomegranate fruit may be beneficial in terms of antioxidant protection in cats. In addition, supplementing cats' diets with dried pomegranate fruit reduced the number of oocysts per gram of feces significantly. It is important to carry out further and more detailed studies to prove the anticoccidial and antioxidant properties of dried pomegranate fruit in cats' diets.

## CONFLICT OF INTEREST

The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

## REFERENCES

- Abbas, R. Z., Colwell, D. D., & Gilleard, J. (2012). Botanicals: An alternative approach for the control of avian coccidiosis. *World's Poultry Science Journal*, 68 (2), 203-215. <https://doi.org/10.1017/S0043933912000268>
- Agarwal, A., Aponte-Mellado, A., Premkumar, B. J., Shaman, A., & Gupta, S. (2012). The effects of oxidative stress on female reproduction: A review. *Reproductive Biology and Endocrinology*, 10 (1), 49. <https://doi.org/10.1186/1477-7827-10-49>
- Bailly, C. (2004). Active oxygen species and antioxidants in seed biology. *Seed Science Research*, 14 (2), 93-107. <https://doi.org/10.1079/SSR2004159>
- Beaulieu, M., & Costantini, D. (2014). Biomarkers of oxidative status: Missing tools in conservation physiology. *Conservation Physiology*, 2 (1), cou014. <https://doi.org/10.1093/conphys/cou014>
- Bhattacharjee, S. (2014). Membrane lipid peroxidation and its conflict of interest: The two faces of oxidative stress. *Current Science*, 107 (11), 1811-1823.
- Chiu, D. T. Y., Stults, F. H., & Tappel, A. L. (1976). Purification and properties of rat lung soluble glutathione peroxidase. *Biochimica et Biophysica Acta (BBA) - Enzymology*, 445 (3), 558-566. [https://doi.org/10.1016/0005-2744\(76\)90110-8](https://doi.org/10.1016/0005-2744(76)90110-8)
- Cox, F. E. G. (2002). History of Human Parasitology. *Clinical Microbiology Reviews*, 15 (4), 595-612. <https://doi.org/10.1128/CMR.15.4.595-612.2002>
- Culliney, T. W. (2005). Benefits of Classical Biological Control for Managing Invasive Plants. *Critical Reviews in Plant Sciences*, 24 (2), 131-150. <https://doi.org/10.1080/07352680590961649>
- Dubey, J. P. (2018). A review of Cystoisosporafelis and C. rivolta-induced coccidiosis in cats. *Veterinary Parasitology*, 263, 34-48. <https://doi.org/10.1016/j.vetpar.2018.09.016>
- Dubey, J. P., Lindsay, D. S., & Lappin, M. R. (2009). Toxoplasmosis and Other Intestinal Coccidial Infections in Cats and Dogs. *Veterinary Clinics of North America: Small Animal Practice*, 39 (6), 1009-1034. <https://doi.org/10.1016/j.cvsm.2009.08.001>
- Dubey, J. P., Schares, G., & Ortega-Mora, L. M. (2007). Epidemiology and Control of Neosporosis and Neospora caninum. *Clinical Microbiology Reviews*, 20 (2), 323-367. <https://doi.org/10.1128/CMR.00031-06>
- Duszynski, D. W. (2021). Biodiversity of the Coccidia (Apicomplexa: Conoidasida) in vertebrates: what we know, what we do not know, and what needs to be done. *Folia Parasitologica*, 68. <https://doi.org/10.14411/fp.2021.001>
- Elfalleh, W., Tlili, N., Nasri, N., Yahia, Y., Hannachi, H., Chaira, N., Ying, M., & Ferchichi, A. (2011). Antioxidant Capacities of Phenolic Compounds and Tocopherols from Tunisian Pomegranate (*Punica granatum*) Fruits. *Journal of Food Science*, 76 (5), C707-C713. <https://doi.org/10.1111/j.1750-3841.2011.02179.x>
- Fébel, H., Mézes, M., Pálffy, T., Hermán, A., Gundel, J., Lugasi, A., Balogh, K., Kocsis, I., & Blázovics, A. (2008). Effect of dietary fatty acid pattern on growth, body fat composition and antioxidant parameters in broilers. *Journal of Animal Physiology and Animal Nutrition*, 92 (3), 369-376. <https://doi.org/10.1111/j.1439-0396.2008.00803.x>
- Gajadhar, A. A., Lalonde, L. F., Al-Adhami, B., Singh, B. B., & Lobanov, V. (2015). Foodborne apicomplexan protozoa: Coccidia. In A. A. Gajadhar (Ed.), *Foodborne Parasites in the Food Supply Web* (pp. 101-147). Woodhead Publishing. <https://doi.org/10.1016/B978-1-78242-332-4.00006-0>
- Hashemi, S. R., & Davoodi, H. (2011). Herbal plants and their derivatives as growth and health promoters in animal nutrition. *Veterinary Research Communications*, 35 (3), 169-180. <https://doi.org/10.1007/s11259-010-9458-2>
- Hussain, S. Z., Naseer, B., Qadri, T., Fatima, T., & Bhat, T. A. (2021). Pomegranate (*Punica granatum*) —Morphology, Taxonomy, Composition and Health Benefits. In S. Z. Hussain, B. Naseer, T. Qadri, T. Fatima, & T. A. Bhat (Eds.), *Fruits Grown in Highland Regions of the Himalayas: Nutritional and Health Benefits* (pp. 131-144). Springer International Publishing. [https://doi.org/10.1007/978-3-030-75502-7\\_10](https://doi.org/10.1007/978-3-030-75502-7_10)
- Ighodaro, O. M., & Akinloye, O. A. (2018). First line defence antioxidants-superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX): Their fundamental role in the entire antioxidant defence grid. *Alexandria Journal of Medicine*, 54 (4), 287-293. <https://doi.org/10.4314/bafm.v54i4>
- Islam, Md. T. (2017). Oxidative stress and mitochondrial dysfunction-linked neurodegenerative disorders. *Neurological Research*, 39 (1), 73-82. <https://doi.org/10.1080/01616412.2016.1251711>
- Ismail, I. E., Alagawany, M., Taha, A. E., Puvaca, N., Laudadio, V., & Tuffarelli, V. (2021). Effect of dietary supplementation of garlic powder and phenyl acetic acid on productive performance, blood haematology, immunity and antioxidant status of broiler chickens. *Animal Bioscience*, 34 (3), 363-370. <https://doi.org/10.5713/ajas.20.0140>
- Juhaimi, F. A., Özcan, M. M., & Ghafoor, K. (2017). Characterization of pomegranate (*Punica granatum* L.) seed and oils. *European Journal of Lipid Science and Technology*, 119 (10), 1700074. <https://doi.org/10.1002/ejlt.201700074>
- Kostadinović, L., & Lević, J. (2018). Effects of phytoadditives in poultry and pigs diseases. *Journal of Agronomy, Technology and Engineering Management*, 1 (1), 1-7.
- Kostadinović, L. M., Pavkov, S. T., Lević, J. D., Galonja-Coghil, T. A., Dozet, G. K., & Bojat, N. Č. (2011). Effect of sulphachloropyrazine on antioxidant systems in blood and liver of broilers. *Acta Veterinaria Brno*, 80 (2), 165-170. <https://doi.org/10.2754/avb201180020165>
- Koutsoumanis, K., Allende, A., Alvarez-Ordóñez, A., Bolton, D., Bover-Cid, S., Chemaly, M., Davies, R., De Cesare, A., Herman, L., Hilbert, F., Lindqvist, R., Nauta, M., Peixe, L., Ru, G., Simmons, M., Skandamis, P., Suffredini, E., Cacciò, S., Chalmers, R., ... Robertson, L. (2018). Public health risks associated with food-borne parasites. *EFSA Journal*, 16 (12), e05495. <https://doi.org/10.2903/j.efsa.2018.5495>
- Litster, A. L., Nichols, J., Hall, K., Camp, J., & Mohamed, A. S. (2014). Use of ponazuril paste to treat coccidiosis in shelter-housed cats and dogs. *Veterinary Parasitology*, 202 (3), 319-325. <https://doi.org/10.1016/j.vetpar.2014.03.003>
- Lukaszewicz-Hussain, A., & Moniuszko-Jakoniuk, J. (2004). Liver Catalase, Glutathione Peroxidase and Reductase Activity, Reduced Glutathione and Hydrogen Peroxide Levels in Acute Intoxication with Chlorfenvinphos, an Organophosphate Insecticide. *Polish Journal of Environmental Studies*, 13 (3), 303-309.
- Mundt, H.-C., Joachim, A., Becka, M., & Dausgchies, A. (2006). Isosporasuis: An experimental model for mammalian intestinal coccidiosis. *Parasitology Research*, 98 (2), 167-175. <https://doi.org/10.1007/s00436-005-0030-x>
- Naidoo, V., McGaw, L. J., Bisschop, S. P. R., Duncan, N., & Eloff, J. N. (2008). The value of plant extracts with antioxidant activity in attenuating coccidiosis in broiler chickens. *Veterinary Parasitology*, 153 (3-4), 214-219. <https://doi.org/10.1016/j.vetpar.2008.02.013>
- Petry, G., Kruedewagen, E., Kampkoetter, A., & Krieger, K. (2011). Efficacy of Emodepside/Toltrazuril Suspension (Procox® Oral Suspension for Dogs) against Mixed Experimental Isosporafelis/Isosporarivolta Infection in Cats. *Parasitology Research*, 109 (1), 29-36. <https://doi.org/10.1007/s00436-011-2400-x>
- Popović, S. J., Kostadinović, L. M., Puvaca, N. M., Kokić, B. M., Čabarkapa, I. S., & Đuragić, O. M. (2017). Potential of wormwood (*Artemisia absinthium*) as a feed supplement in rabbit diet: Effect on controlling rabbit coccidiosis, antioxidative systems and growth performance. *Veterinarski Arhiv*, 87 (6), 769-782. <https://doi.org/10.24099/vet.arhiv.160704a>
- Pritt, S., Cohen, K., & Sedlacek, H. (2012). Parasitic Diseases. In M. A. Suckow, K. A. Stevens, & R. P. Wilson (Eds.), *The Laboratory Rabbit, Guinea Pig, Hamster, and Other Rodents* (pp. 415-446). Academic Press. <https://doi.org/10.1016/B978-0-12-380920-9.00015-8>
- Puvaca, N., Horvatek Tomić, D., Popović, S., Đorđević, S., Brkić, I., Lalić,



- N., ShtyllaKika, T., & Lika, E. (2020) . Influence of tea tree (*Melaleuca alternifolia*) essential oil as feed supplement on production traits, blood oxidative status and treatment of coccidiosis in laying hens. *Veterinary Archives*, 90 (4) , 331-340.
- Puvača, N., Tufarelli, V., & Giannenas, I. (2022) . Essential Oils in Broiler Chicken Production, Immunity and Meat Quality: Review of *Thymus vulgaris*, *Origanum vulgare*, and *Rosmarinus officinalis*. *Agriculture*, 12 (6) , 874. <https://doi.org/10.3390/agriculture12060874>
- Shaw, S. E., Kenny, M. J., Tasker, S., & Birtles, R. J. (2004) . Pathogen carriage by the cat flea *Ctenocephalides felis* (Bouché) in the United Kingdom. *Veterinary Microbiology*, 102 (3) , 183-188. <https://doi.org/10.1016/j.vetmic.2004.06.013>
- Sultana, Z., Maiti, K., Aitken, J., Morris, J., Dedman, L., & Smith, R. (2017) . Oxidative stress, placental ageing-related pathologies and adverse pregnancy outcomes. *American Journal of Reproductive Immunology*, 77 (5) , e12653. <https://doi.org/10.1111/aji.12653>
- Truyen, U., Addie, D., Belák, S., Boucraut-Baralon, C., Egberink, H., Frymus, T., Gruffydd-Jones, T., Hartmann, K., Hosie, M. J., Lloret, A., Lutz, H., Marsilio, F., Pennisi, M. G., Radford, A. D., Thiry, E., & Horzinek, M. C. (2009) . Feline panleukopenia. ABCD guidelines on prevention and management. *Journal of Feline Medicine & Surgery*, 11 (7) , 538-546. <https://doi.org/10.1016/j.jfms.2009.05.002>
- Tsikias, D. (2017) . Assessment of lipid peroxidation by measuring malondialdehyde (MDA) and relatives in biological samples: Analytical and biological challenges. *Analytical Biochemistry*, 524, 13-30. <https://doi.org/10.1016/j.ab.2016.10.021>
- Uttara, B., Singh, A. V., Zamboni, P., & Mahajan, R. T. (2009) . Oxidative Stress and Neurodegenerative Diseases: A Review of Upstream and Downstream Antioxidant Therapeutic Options. *Current Neuropharmacology*, 7 (1) , 65-74.
- VapaTankosić, J., Puvača, N., Giannenas, I., Tufarelli, V., & Ignjatijević, S. (2022) . Food Safety Policy in the European Union. *Journal of Agronomy, Technology and Engineering Management (JATEM)* , 5 (2) , 712-717. <https://doi.org/10.55817/EMRK6646>
- Wenyon, C. M. (1923) . Coccidiosis of Cats and Dogs and the Status of the Isospora of Man. *Annals of Tropical Medicine & Parasitology*, 17 (2) , 231-289. <https://doi.org/10.1080/00034983.1923.11684357>
- Zhang, J. J., Wang, L. X., Ruan, W. K., & An, J. (2013) . Investigation into the prevalence of coccidiosis and maduramycin drug resistance in chickens in China. *Veterinary Parasitology*, 191 (1) , 29-34. <https://doi.org/10.1016/j.vetpar.2012.07.027>