



## Journal of the Hellenic Veterinary Medical Society

Vol 74, No 4 (2023)



### To cite this article:

Aghdashi, M., Nobakht, A., & Mehmannavaz, Y. (2024). Effects of different levels of iron, iodine and vitamin B12supplementation on egg quality and antioxidant capacity of eggs at different times of storage in laying hens. *Journal of the Hellenic Veterinary Medical Society*, *74*(4), 6501–6510. https://doi.org/10.12681/jhvms.31449

# Effects of different levels of iron, iodine and vitamin B<sub>12</sub>supplementation on egg quality and antioxidant capacity of eggs at different times of storage in laying hens

M.N. Aghdashi<sup>®</sup>, A. Nobakht<sup>\*</sup><sup>®</sup>, Y. Mehmannavaz<sup>®</sup>

Department of Animal Science, Maragheh Branch, Islamic Azad University, Maragheh Iran

ABSTRACT: The experiment was carried out to evaluate the effects of adding different levels of iron salts, iodine and vitamin B12 on yield, quality traits and antioxidant capacity of eggs in laying hens. A total of 320 laying hens (HyLine Variety White, 28wk old) in a 2×2×2 factorial experiment were randomly assigned to 8 groups, and each group consisting of 40 hens (5 replicates of 8 hens each) for a period of approximately 13 weeks in a completely randomized design. Hens in the control group received a basal diet with no supplementation whereas hens in treat 2 received basal diet supplemented with 400 mg Fe/kg as FeSO<sub>4</sub>, treat 3 (basal diet + 450 mg I/kg as KI), treat 4 (basal diet + 0.1 mg B<sub>12</sub>vit/ kg), treat 5 (basal diet + 400 mg Fe/kg as FeSO<sub>4</sub> + 450 mg I/kg as KI), treat 6 (basal diet + 400 mg Fe/kg as FeSO<sub>4</sub> +  $0.1 \text{ mg B}_{12}$ Vit/kg), treat 7 (basal diet + 450 mg I/kg as KI + 0.1 mg B $_{12}$ Vit/kg), treat 8 (basal diet + 400 mg Fe/kg as  $FeSO_4 + 450 \text{ mg I/kg}$  as KI + 0.1 mg  $B_{12}$ Vit/kg), respectively. The results showed that the use of iodine and vitamin B<sub>12</sub> increased the shell thickness of eggs (p<0.05). The use of iron, iodine and vitamin B12 supplements increased the antioxidant capacity of eggs compared to the control group (p < 0.05). The highest antioxidant capacity of eggs (0.77) was observed when these three supplements were used together in the diet of laying hens. The length of storage had significant effects on the weight of albumin and yolk and antioxidant content of eggs (p < 0.05). The highest weight of albumin and antioxidant content was at the beginning of storage, while the highest weight of yolk was obtained in 4 weeks after storage, with increasing of storage time to 8 weeks, all three attributes of albumin weight, yolk weight and antioxidant content of eggs decreased (p < 0.05). The related effects of interval of the storage and the experimental supplementation, the use of iron, iodine and vitamin B<sub>12</sub> supplements increased the antioxidant capacity of eggs, so that the highest amount of antioxidants in eggs in 8 weeks after storage and by supplementing the diets with all three additives (p<0.05). According to the present experiment, the use of iron, iodine and vitamin  $B_{1,2}$  additives in the diet of laying hens, improves the performance, the content of iron and iodine elements in eggs and maintains the suitable condition of eggs during storage.

Keywords: Antioxidants; Eggs quality; Minerals; Storage time; Vitamins

Corresponding Author: A. Nobakht, Department of Animal Science, Maragheh Branch, Islamic Azad University, Maragheh Iran E-mail address: anobakht20@yahoo.com

Date of initial submission: 19-09-2022 Date of acceptance: 27-12-2022

#### **INTRODUCTION**

S tudies show that the efficiency of transferring vitamins and minerals to egg yolk is divided into 4 levels (low, medium, high and very high). The efficiency of transferring vitamins and minerals to egg yolk is around 50% to 80%. Although the antagonistic interactions between some vitamins have been observed, however, normal use of these vitamins does not seem to cause problems in the nutritional performance of laying hens (Gan, 2020).

The use of rations containing high levels of vitamins has increased the vitamin content of ordinary eggs daily recommended intake (DRI) up to 50%. Vitamin  $B_{12}$  showed the highest efficiency among all vitamins and when the amount of vitamin  $B_{12}$  in the diet of laying hens had increased up to 11 fold its DRI value increased from36 to 139 (Leeson and Caston, 2003). Vitamins K and biotin provide 50% of the DRI (Leeson and Caston, 2003).

Due to antagonism and synergism effects among minerals and vitamins, finding the best and most suitable level of these nutrients seems very important and necessary. In this regard, the interaction between vitamin D3 and calcium can be mentioned as an example. Vitamin D gives the ability to the body to obtain calcium from food sources that passing through the digestive tract instead of taking it from the bones. Also vitamin C helps iron absorption. Furthermore, zinc can be involved in the absorption, transfer, physiological and biochemical efficiency of vitamin B9 (folate). However, nutrients interactions are not always be positive. For example, vitamin C blocks the absorption of copper by the body, or the lowest magnesium deposition causes iron deficiency (Bagheri et al., 2019). The experiments have shown that iodine storage in eggs is mostly done in the yolk (Damaziak et al., 2018). Kaufmann et al. (1998) stated that the consumption of iodine from zero, 0.5, 1, 2, and 5 mg/kg diet cannot improve feed efficiency and egg production.

They also concluded that the amount of iodine in eggs increases significantly by supplementing the diet of laying hens. There is no significant difference in body weight, feed consumption, egg production, egg shell index, egg thickness and yolk index with consumption of zero, 3, 6, 12 and 24 mg/kg iodinein diet of laying hens that were fed with calcium iodide (Yalçinet al., 2004).Christensen et al. (1991) reported that in a study with two commercial strains of turkey the amount of iodine increased from 0.7 to 4.2 mg/kg, in which the hatchability in mother flocks statistically increased. Also Damaziak et al. (2018) stated that the use of a laying hen diet supplemented with iodine up to 10ppmcompletely inhibit the Salmonella growth until day 10 of incubation and impart high level of resistance to the vitelline membrane against the growth of Salmonella in case of eggs stored at 30°C.Shah and Dwivedi (2022) reported that, Actinobacteria such as Streptomyces and Propionibacteria are among the efficient producers of B vitamins and can serve as probiotics to increase the productivity in the poultry industry through improving growth performance and immune response of poultry. Research has shown that the utilization coefficient of inorganic trace elements was about 30% (Behroozlak et al., 2019). With these aspects the effects of different levels of iron, iodine and vitamin B<sub>12</sub> supplementation on egg quality and antioxidant capacity of eggs at different times of storage in laying hens evaluated in this study.

#### MATERIALS AND METHODS

#### **Birds and Housing**

Three hundred twenty white Leghorn hens of w-36 strains from the age of 28 to 41 weeks for approximately 13 weeks in the form of a completely randomized design with a factorial arrangement of 2×2×2 with food rations including 3 factors. Which are 2 levels of iodine (zero and 450 mg/kg) and 2 levels of iron (zero and 400 mg/kg) and 2 levels of vitamin B12 (zero and 0.1 mg/kg). Each experimental group consisted of 5 replications, where 2 pens together were considered as one experimental unit, which included a total of 40 experimental units. The chickens were kept with 17 hours of artificial lighting and optimal temperature and proper ventilation. Food rations were balanced according to the guidelines of the target strain and using the UFFDA ration software. The experimental hens was freely access to food and water. The study was carried out in an industrial poultry farm located in Khoy city called Aqlari in the months of November and December of 2018 and January of 2019 which are among the coldest months of the year. This farm has 2 halls that each hall has 4 rows of 3-story cages with cage dimensions of  $40 \times 35 \times 30$ . Experimental hens were selected from the third row in the middle of the hall and from the cages in the middle row. After 10 days of habituation, data collection started.

#### **Experimental Diets and Treatments**

The diet used for experimental hens is shown in Table 1. The target ration is balanced according to the 2016 version of the white leghorn strain manual using

	1		
Ingredients	Content (%)	Nutrients	
Corn	58.1	Crude protein, %	15.88
Soybean Meal	22.18	Calcium, %	4.42
Oyster Shell	10	Available phosphorus, %	0.52
Wheat Bran	6	Sodium, %	0.31
Soybean Oil	2	Methionine, %	0.58
Dical. Phos	1.6	Lysine, %	0.94
Mineral Premix <sup>1</sup>	0.03	Methionine + cysteine, %	1.36
Vitamin Premix <sup>2</sup>	0.03	Argenine %	1.06
L-lysine HCL	0.025	Linoleic acid, %	1.91
Common Salt	0.025	Threonine, %	0.68
DL-Methionine	0.013	Tryptophan, %	0.31
Total	100	Metabolizable energy, kcal/kg	2700

Table 1. Feed ingredients and nutrient composition of the basal diet

<sup>1</sup>Mineral supplement per kg of diet: 2.4 mg of copper, 0.34 mg of iodine, 30 mg of iron, 29.76 mg of manganese, 0.08 mg of selenium, 25.87 mg of zinc. <sup>2</sup>Vitamin supplement per kg of diet: 3520Iu Vitamin A, 0.59 mg vitamin B1, 1.6 mg vitamin B2, 13.86 mg niacin, 13.3 mg pantothenic acid, 1 mg B6, 0.06 mg biotin, 80 mg choline, 0.004 mg vitamin B12, 0.19 mg B9, 1000IU vitamin D3, 8.8 IU vitamin E, 0.88 mg k3, all calculations are evaluated based on the requirements of (1994) NRC.

#### the UFFDA ration software.

Furthermore, the experimental treatments were contained different dietary supplementations. Hens in control group received a basal diet with no supplementation whereas hens in treat 2 received basal diet supplemented with 400 mg Fe/kg as FeSO<sub>4</sub>, treat 3 included basal diet + 450 mg I/kg as KI, treat 4 contained basal diet + 0.1 mg B<sub>12</sub>vit/kg, treat 5 were basal diet + 400 mg Fe/kg as FeSO<sub>4</sub> + 450 mg I/kg as KI, treat 6 included basal diet + 400 mg Fe/kg as FeSO<sub>4</sub> + 450 mg I/kg as KI, treat 6 included basal diet + 400 mg Fe/kg as FeSO<sub>4</sub> + 450 mg I/kg as KI + 0.1 mg B<sub>12</sub>Vit/kg, treat 7 contained basal diet + 400 mg Fe/kg as KI + 0.1 mg B<sub>12</sub>Vit/kg as KI + 0.1 mg B<sub>12</sub>Vit/kg, respectively.

#### Albumen quality

Height meter was used to determine the height of albumen. In order to determine the haugh (HU) unit, 5 eggs from each replicate were randomly selected and after numbering, they were taken to the laboratory and first weighed with a digital scale with an accuracy of  $\pm 0.001$  grams and then broken on a glass surface. The height of albumen from a certain point was calculated for all eggs using a height measuring device and the HU unit using the following formula (haugh, 1937):

$$HU = 100 \times Log (H + 7.57 - 1.7 W^{0.37})$$

In this equation, H is the height of the albumen (mm) and W is the weight of the egg (grams).

#### **Albumen Index**

It was calculated as the average between the max-

imum and minimum diameter of albumin.

#### Shell Quality

To determine the quality of the egg shell, two factors of the shell contain shell thickness and shell strength were measured. To determine the shell thickness, 5 eggs from each replicate were selected, numbered and broken. Then, the inside contents were completely washed with water and dried well with a paper towel, and then the thickness of the shell was calculated using a micrometer. A strength tester was used to determine the shell strength (Mertens et al., 2006).

#### Yolk color

Color fan method is a common method to measure yolk color and is accepted as a standard in most countries of the world. Color fan consists of 15 color stencils that are fan-shaped. Each stencil has a color that is measured visually. The range of colors is from pale yellow (number 1) to reddish orange (number 15). By comparing the color of egg yolks with a color template, scores were given according to the observed colors.

#### Yolk index and egg shape index

To calculate the yolk index, the height of the yolk was measured first using a height meter, and then its diameter was measured with a caliper (in centimeters). The result of dividing the height of the yolk by the diameter of the yolk represents the yolk index. For the egg shape index, like the yolk, the length and width of the egg were measured with a caliper. Egg shape index=  $100 \times W/L$ 

W is the width and L is the length of the egg

After 13 weeks, 4 eggs were randomly taken from each replicate to determine the concentration of iron, iodine and cobalt inside the eggs and were taken to the laboratory. From the 3 treatments and the pure treatments, 4 eggs were transferred to the laboratory in the first days, 28 days and 56 days with an interval of 28 days to measure the antioxidant capacity and the effect of storage on the eggs. The storage conditions of the eggs were kept in a refrigerator at 4 degrees Celsius and 60% humidity for the effect of storage. The laboratory chosen to measure the internal content of eggs, including the amount of iodine, iron, cobalt and antioxidants, is the University Jihad Laboratory of West Azarbaijan Province, located in Urmia city, which is equipped with equipment including a furnace, an atomic absorption device for measuring substances. Mineralogy and spectrophotometry were used to measure the amount of antioxidants using a laboratory kit.

#### Egg quality

The height of yolk and Albumen was measured using a stand-based device (made in Japan, model. Alimet OSK3470) with an accuracy of 0.01 mm. The haugh unit was calculated using the formula previously presented by haugh (1937). The shell strength (kg/ cm2) was obtained using an egg shell strength measuring device (made in Japan, model. OSK-13473). The thickness of the shell was evaluated using a micrometer with an accuracy of 0.01 mm (F.K.H model, made in Japan) in the back part of the egg from three different points (Nobakht et al., 2022).

#### **Minerals Preparation**

The experimental diet with the addition of ferro sulfate, potassium iodide and synthetic vitamin  $B_{12}$  to the basic diet in 2 levels of iron (zero and 400 mg/kg) and 2 levels of iodine (zero and 450 mg/kg) and 2 levels of vitamin B12 (zero and 0.1 mg/kg) were prepared. In the NRC table, the amount of iron, iodine, and vitamin  $B_{12}$  needed by laying poultry is 55 mg per kg, iodine 1.7 mg per kg, and vitamin  $B_{12}$  0.2 mg per kg of diet. Ferro sulfate with the chemical formula FeSo<sub>4</sub>.7H<sub>2</sub>O in the form of green crystals and soluble in water with the grade of industrial, food and agricultural products was purchased from Iran's Pars Oxid Company, which contains 70% iron. Iodide potassium with the chemical formula KI and the appearance of

white solid crystals and molecular weight of 166 g/ mol and solubility in water and for use in the food industry, which was purchased from Barnard Company with a purity of 99%. Vitamin  $B_{12}$  with the name of cyanocobalamin with 99% purity and food grade was purchased from the representative of Germany's Lohmann Company in Iran with the trade name Microvit and with the chemical formula of  $C_{63}$ H-<sub>88</sub>CoN<sub>14</sub>O<sub>14</sub>P.

#### **Statistical analysis**

At the end of the experiment, statistical analysis of data was done using SAS software version 3.9 and GLM procedure. The experiment was carried out in the form of a completely randomized design with a factorial arrangement of  $2 \times 2 \times 2$  with 2 levels of iodine, 2 levels of iron and 2 levels of vitamin B<sub>12</sub>. The results related to antioxidants and storage conditions were completely randomized with a factorial arrangement including the first factor (control, iron, iodine, vitamin B<sub>12</sub> and the combination of all 3) and the second factor (zero, 4 weeks and 8 weeks storage time). Before analyzing the data, the normality of the data was evaluated using the Kolmogorov-Smirnov test. In this model:

$$\begin{split} Y_{ijkl} = & \mu + A_i + B_j + C_k + AB_{ij} + AC_{ik} + BC_{jk} + ABC_{ijk} \\ + & \epsilon_{ijkl} \end{split}$$

 $Y_{ijkl}$  = measured value for each piece;  $\mu$  = community average;  $A_i$  = effect of iron;  $B_j$  = effect of iodine;  $C_k$  = effect of vitamin  $B_{12}$ ;  $AB_{ij}$  = interaction effect of iron and iodine;  $AC_{ik}$  = interaction effect of iron and vitamin  $B_{12}$ ;  $BC_{jk}$  = interaction between iodine and vitamin  $B_{12}$ ;  $ABC_{ijk}$  = interaction effect of iron, iodine and vitamin  $B_{12}$ ;  $\epsilon_{iikl}$  = experimental error;

Data were analyzed using GLM procedure of SAS software version 9.3, and the means were compared with Tukey test at the level of 0.05.

#### **RESULTS AND DISCUSSION**

The effects of using different levels of iron, iodine, and vitamin  $B_{12}$  on egg parameters are presented in Tables 2, 3, and 4 during the first, second, and third 20 days of the experiment, respectively.

As it shown in table 2, the treatment containing 0.1 mg vitamin  $B_{12}$  with 450 mg of iodine per kg of the diet of laying hens had significant effects on the shell thickness (p<0.05), but the other treatments had no effect on egg indices in the first 20 days of the experiment. Also, it can be seen in the table 2 that the index

of shell thickness in which with addition of iodine to the diet contained vitamin  $B_{12}$ , the thickness of the shell had increased, which is 7% more than treatment without iodine, and it can be indicated that vitamin

 $B_{12}$  with iodine could increase the thickness of the egg shell. Also, the addition of vitamin  $B_{12}$  increases es the shape index, but this increase is not significant (p>0.05).

<b>Table 2.</b> Effects of treatments on egg indices in the first 20 days of the experiment.											
	large	Small	Shape	Yolk	shell	Egg	Albumen	Yolk	shell	Volk	
Treatments	diameter	diameter	Index	diameter	strength	weight	height	height	thickness	color	HU
	(cm)	(cm)		(cm)	(gr/cm <sup>2</sup> )	gr	(mm)	(mm)	(mm)		
Fe											
0	5.303	4.081	0.766	4.314	2.468	60.441	7.015	16.882	0.3498	3.7	82.495
400	5.396	4.109	0.761	5.572	1.892	61.672	6.876	15.664	0.3523	4	81.362
SEM	0.0243	0.012	0.0034	0.5531	0.5732	0.5146	0.2481	0.5447	0.0041	0.1497	1.5316
P-value	0.0643	0.1176	0.3417	0.1179	0.4829	0.1005	0.6944	0.1238	0.6722	0.166	0.6043
Iodine											
0	5.337	4.106	0.766	4.342	1.731	61.33	6.903	16.945	0.347	3.94	81.492
450	5.335	4.08	0.761	5.544	2.63	60.78	6.987	15.602	0.355	3.76	82.365
SEM	0.0243	0.0122	0.0034	0.553	0.5732	0.5146	0.248	0.5447	0.004	0.1497	1.5316
P-alue	0.9655	0.2206	0.3417	0.1341	0.2757	0.4578	0.8117	0.0909	0.1415	0.4014	0.6897
B <sub>12</sub>											
0	5.355	4.094	0.762	4.316	2.602	60.291	6.938	16.78	0.352	3.875	81.661
0.1	5.32	4.096	0.766	5.57	1.759	60.822	6.952	15767	0.35	3.825	82.196
SEM	0.0243	0.0122	0.0034	0.553	0.5732	0.5146	0.2481	0.5447	0.0041	0.1497	1.5316
P-value	0.3695	0.8995	0.3838	0.1187	0.3067	0.5239	0.9677	0.1977	0.7755	0.8148	0.8066
Fe + Iodine											
0 + 0	5.294	4.082	0.768	4.331	1.757	60.586	6.974	16.771	0.351	3.69	82.084
400 + 0	5.312	4.081	0.764	4.298	3.18	60.296	7.055	16.993	0.349	3.71	82.906
0 + 450	5.379	4.13	0.764	4.353	1.706	62.074	6.832	17.119	0.342	4.19	80.9
450 + 400	3.358	4.088	0.759	6.79	2.08	61.27	6.92	14.21	0.362	3.81	81.823
SEM	0.0344	0.0173	0.0049	0.7821	0.8106	0.7278	0.3509	0.7703	0.0057	0.2117	2.167
P-value	0.5785	0.2606	0.8227	0.1241	0.5222	0.7263	0.9926	0.0505	0.0807	0.3518	0.9815
$Fe + B_{12}$											
0 + 0	5.294	4.091	0.769	4.311	3.139	60.634	6.966	16.683	0.352	3.54	82.146
0.1 + 0	5.312	4.072	0.762	4.318	1.797	60.348	7.063	17.081	0.348	3.86	82.845
400 + 0	5.409	4.098	0.754	4.321	2.064	61.948	6.91	16.876	0.352	4.21	81.177
400 + 0.1	5.328	4.121	0.769	6.822	1.721	61.396	6.842	14.452	0.353	3.79	81.547
SEM	0.0344	0.0172	0.0049	0.7821	0.8106	0.7278	0.3509	0.7703	0.0057	0.2117	2.1661
P-value	0.1565	0.2377	0.1207	0.1208	0.542	0.9099	0.8152	0.0764	0.6473	0.09	0.9398
Iodine + B <sub>12</sub>											
0 + 0	5.381	4.114	0.762	4.345	1.737	62.308	7.044	16.846	0.355ª	4.04	81.939
0 + 0.1	5.295	4.098	0.77	4.34	1.725	60.352	6.763	17.043	0.338 <sup>b</sup>	3.84	81.045
450 + 0	5.322	4.075	0.761	4.288	3.466	60.274	6.833	16.714	0.349 <sup>ab</sup>	3.71	81.383
0.1 + 450	5.349	4.095	0.762	6.801	1.794	61.292	7.142	14.49	0.362ª	3.81	83.347
SEM	0.0344	0.0173	0.0049	0.7821	0.8106	0.7278	0.3509	0.7703	0.0057	0.2116	2.166
P-value	0.1019	0.3109	0.4061	0.1174	0.3136	0.2181	0.4059	0.1259	0.0122	0.4837	0.5142
$Fe + Iodine + B_{12}$											
0 + 0 + 0	5.334	4.109	0.768	4.325	1.642	61.864	7.217	16.621	0.364	3.66	83.27
0 + 0 + 0.1	5.255	4.055	0.767	4.337	1.871	59.308	6.732	16.921	0.337	3.72	80.898
0 + 450 + 0	5.254	4.072	0.771	4.297	4.636	59.404	6.716	16.746	0.34	3.42	81.021
0.1 + 450 + 0	5.37	4.089	0.758	4.299	1.724	61.188	7.394	17.241	0.358	4	84.792
0 + 0 + 400	5.429	4.118	0.756	4.364	1.832	62.752	6.871	17.071	0.346	4.42	80.608
0.1 + 0 + 400	5.329	4.142	0.773	4.342	1.579	61.396	6.793	17.166	0.339	3.96	81.192
0 + 450 + 400	5.39	4.077	0.752	4.279	2.296	61.144	6.949	16.681	0.357	4	81.745
0.1 + 450 + 400	5.327	4.1	0.766	4.172	1.864	61.396	6.89	16.802	0.366	3.62	81.901
SEM	0.0486	0.0244	0.0069	1.1061	1.1463	1.0293	0.4962	1.0894	0.0081	0.2993	3.063
P-value	0.2592	0.3056	0.6399	0.1161	0.3679	0.3551	0.4205	0.0992	0.2281	0.6068	0.4537

Table 3. Effects of treatments on egg indices in the second 20 days of the experiment.											
	large	Small	C1	Yolk	shell	Egg	Albumen	Yolk	shell	V-11-	
Treatments	diameter	diameter	Snape	diameter	strength	weight	height	height	thickness	rolk	HU
	(cm)	(cm)	mucx	(cm)	(gr/cm <sup>2</sup> )	gr	(mm)	(mm)	(mm)		
Fe											
0	5.673	4.384	0.77	4.527	2.584	61.529	7.832	17.515	0.583	3.35	87.447
400	5.690	4.369	0.769	4.509	2.245	62.150	8.163	17.693	0.310	3.6175	89.121
SEM	0.02216	0.01671	0.00345	0.02550	0.16569	0.8851	0.19700	0.14531	0.19571	0.0793	1.1099
P-value	0.5749	0.6811	0.8709	0.6097	0.1583	0.4034	0.2435	0.395	0.3326	0.0233	0.2941
Iodine											
0	5.686	4.385	0.767	4.510	2.541	61.627	7.679	17.497	0.306	3.447	86.504
450	5.676	4.394	0.771	4.526	2.288	62.052	8.315	17.711	0.5876	3.52	90.064
SEM	0.02216	0.01671	0.00345	0.02550	0.16569	0.5188	0.19700	0.14531	0.19571	0.0793	1.1099
P-alue	0.7494	0.7121	0.4914	0.6683	0.2893	0.5667	0.0291	0.3051	0.3171	0.523	0.0302
B <sub>12</sub>											
0	5.696	4.394	0.768	4.5011	2.364	61.975	8.043	17.617	0.306	3.475	88.504
0.1	5.667	4.385	0.770	4.5355	2.465	61.704	7.952	17.591	0.5874	3.492	88.064
SEM	0.02216	0.01671	0.00345	0.02550	0.16569	0.5188	0.19700	0.14531	0.19571	0.0793	1.1099
P-value	0.3602	0.7058	0.7454	0.3474	0.6709	0.7146	0.7461	0.8978	0.3178	0.8771	0.7814
Fe + Iodine											
0 + 0	5.673	4.389	0.771	4.512	2.712	61.22	7.605	17.441	0.301	3.3	87.511
450 + 0	5.672	4.379	0.769	4.542	2.456	61.838	8.059	17.590	0.864	3.4	87.383
400 + 0	5.7	4.380	0.764	4.508	2.37	62.035	7.753	17.553	0.311	3.595	89.496
450 + 400	5.680	4.408	0.773	4.509	2.121	62.266	8.572	17.832	0.310	3.64	88.746
SEM	0.03134	0.02363	0.00488	0.03607	0.23432	0.73369	0.27860	0.20550	0.27678	0.1122	1.5696
P-value	0.7735	0.4321	0.2772	0.6924	0.9882	0.7935	0.5164	0.7546	0.3157	0.808	0.4487
$Fe + B_{12}$											
0 + 0	5.684	4.394	0.770	4.494	2.57	61.706	7.848	17.5742	0.302	3.15 <sup>b</sup>	87.511
400 + 0	5.661	4.375	0.769	4.561	2.598	61.352	7.815	17.4576	0.863	3.55 <sup>b</sup>	87.383
0 + 0.1	5.708	4.394	0.767	4.508	2.159	62.244	7.237	17.6614	0.310	3.8ª	89.496
400 + 0.1	5.672	4.394	0.771	4.509	2.332	62.057	8.088	17.7248	0.311	3.435 <sup>ab</sup>	88.746
SEM	0.03134	0.02363	0.0488	0.03607	0.23432	0.73369	0.27860	0.20550	0.27678	0.1122	1.5696
P-value	0.8371	0.6934	0.5705	0.37	0.759	0.9099	0.8353	0.6644	0.3191	0.0018	0.8443
Iodine + B <sub>12</sub>											
0 + 0	5.701	4.366	0.763	4.527	2.466	61.602	7.725	17.5094	0.304	3.61	86.891
0 + 0.1	5.672	4.404	0.772	4.493	2.616	61.653	7.633	17.4854	0.307	3.285	86.118
450 + 0	5.691	4.421	0.774	4.474	2.263	62.348	8.360	17.7262	0.308	3.34	90.117
0.1 + 450	5.662	4.366	0.768	4.577	2.314	61.756	8.271	17.697	0.867	3.7	90.011
SEM	0.03134	0.02363	0.00488	0.03607	0.23432	0.73369	0.27860	0.20550	0.27678	0.1122	1.5696
P-value	0.9975	0.0583	0.1029	0.0648	0.834	0.664	0.9955	0.99	0.3229	0.0046	0.8333
Fe + Iodine +B12											
0 + 0 + 0	5.693	4.384	0.768	4.539	2.66	61.424	7.659	17.507	0.298	3.22 <sup>b</sup>	86.642
0 + 0 + 0.1	5.653	4.395	0.773	4.486	2.764	61.016	7.550	17.374	0.304	3.38 <sup>b</sup>	85.89
0 + 450 + 0	5.675	4.403	0.772	4.448	2.48	61.988	8.037	17.640	0.307	3.08°	88.380
0.1 + 450 + 0	5.67	4.356	0.765	4.636	2.432	61.688	8.080	17.540	0.422	3.72ª	88.869
0 + 0 + 400	5.709	4.348	0.757	4.516	2.272	61.78	7.791	17.511	0.311	4 <sup>a</sup>	87.139
0.1 + 0 + 400	5.690	4.412	0.772	4.500	2.468	62.29	7.715	17.596	0.310	3.19 <sup>b</sup>	86.340
0 + 450 + 400	5.707	4.44	0.776	4.500	2.046	62.708	8.684	17.811	0.309	3.6ª	91.853
0.1 + 450 + 400	5.654	4.376	0.770	4.519	2.196	61.824	8.461	17.853	0.311	3.68ª	91.152
SEM	0.04433	0.03344	0.00690	0.05101	0.33138	1.03759	0.39400	0.29063	0.39142	0.1587	2.219
P-value	0.584	0.472	0.7149	0.1639	0.9107	0.6125	0.7906	0.9277	0.3256	0.368	0.857

As it cleared in Table 3, the treatment containing 400 mg Fe/kg of diet of laying hens and the treatment containing all three supplemented substances (iron, iodine and vitamin B12) improved egg yolk color index significantly (p<0.05), but the other treatments had no effect on the egg indices at the second 20 days of the experiment (p>0.05).

Also, as it shown in table 4, the treatment containing 400 mg Fe/kg with 0.1 mg vit  $B_{12}$ /kg of diet of laying hens and the treatment containing all three supplemented substances (iron, iodine and vitamin  $B_{12}$ ) improved egg yolk color index significantly (p<0.05), but the other treatments had no effect on the egg indices at the third 20 days of the experiment (p>0.05).

Table 4. Effects of treatments on egg indices in the third 20 days of the experiment.											
	large	Small	Shape	Yolk	shell	Egg	Albumen	Yolk	shell	Yolk	
Treatments	diameter	diameter	Index	diameter	strength	weight	height	height	thickness	color	HU
E-	(cm)	(cm)		(cm)	(gr/cm <sup>2</sup> )	gr	(mm)	(mm)	(mm)		
Fe	5 710	4 410	0 7(7	4.011	2 217	(2, 172	7 200	17 (70)	0.004	2 (1	04.262
0	5./18	4.410	0.767	4.911	2.21/	62.4/3	7.308	17.502	0.284	3.61	84.363
400	5.693	4.40/	0.772	4.548	0.1229	0.4592	/.329	17.583	0.290	3.65	84.518
SEM	0.0192	0.0118	0.0032	0.2556	0.1228	0.4582	0.1130	0.1138	0.0037	0.08003	0.74213
P-value	0.3624	0.82/3	0.2929	0.3230	0.0981	0.41/8	0.8948	0.5581	0.217	0./261	0.8835
lodine	5 702	4 412	0 772	4.570	2 007	(2.2(0)	7.245	17 (0)	0.2057	2 515	04 57(
0	5.703	4.413	0.772	4.572	2.087	62.368	7.345	17.686	0.2857	3.515	84.576
450 SEM	5.709	4.404	0.768	4.88/	0.052	0.4592	/.292	0.1129	0.2889	3./45	84.306
SEM D alara	0.0192	0.0118	0.0032	0.2556	0.1228	0.4582	0.1130	0.1138	0.0037	0.08003	0.74213
P-alue	0.8351	0.5762	0.4/66	0.3905	0.8439	0.6227	0.4/3/	0.4952	0.5460	0.0505	0.7985
B <sub>12</sub>	5 711	4 410	0 7702	4.005	2.126	(1.221	7 4 4 0	177(1	0.201	2 (5	05 104
0	5./11	4.410	0.7703	4.895	2.126	61.331	7.199	17.701	0.291	3.65	85.124
0.1	5.700	4.407	0.7704	4.304	2.013	02.083	/.188	0.1120	0.283	3.01	83./5/
SEM	0.0192	0.0118	0.0032	0.2556	0.1228	0.4582	0.1130	0.1158	0.0037	0.08003	0.74213
P-value	0.6748	0.8365	0.9827	0.36//	0.5206	0.7045	0.1129	0.1150	0.146/	0./261	0.2022
Fe + Iodine	5 (00	4 412	0 771	4.5.4.4	2.2(1	(2.449	7 277	17 (15	0.004	2.46	94160
0 + 0	5.699	4.413	0.771	4.544	2.261	62.448	7.277	17.015	0.284	3.46	84.160
$400 \pm 0$	5./384	4.408	0.746	5.278	2.1/4	62.498	7.338	17.741	0.283	3.76	84.566
0 + 450	5.7076	4.414	0.773	4.600	1.912	62.288	7.412	17.757	0.287	3.57	84.992
450 + 400	5.6769	4.400	0.772	4.496	1.931	61.594	/.246	17.409	0.294	3./3	84.045
SEM	0.02/1	0.0168	0.0045	0.3614	1.1/38	0.6480	0.1598	0.1610	0.0052	0.01131	1.04953
P-value	0.2237	0./999	0.4/00	0.2555	0.7620	0.5700	0.4812	0.1504	0.4512	0.5407	0.5257
$Fe + D_{12}$	5 715	4 4124	0.770	5 206	2 261	62 462	7 522	17 120	0.204	2 1 1 b	95 650
0 + 0 $400 \pm 0$	5 722	4.4124	0.770	J.200 4.616	2.301	62.402	7.002	17.120	0.304	2.79a	82.059
$400 \pm 0.1$	5 709	4.409	0.703	4.010	2.074	62 200	7.095	17.209	0.290	2.96a	83.007
0 + 0.1 $400 \pm 0.1$	5.708	4.409	0.775	4.504	1.051	61.682	7.575	16.099	0.308	2.01a	04.309 94.449
400 + 0.1	0.0271	0.01668	0.0045	0.2614	0.1728	01.082	0.1508	0.2870	0.270	0.1121	1 04052
D value	0.0271	0.01008	0.2550	0.3014	0.1750	0.0480	0.1398	0.2079	0.1131	0.0020	0.2516
I - value	0.4910	0.9859	0.3350	0.4787	0.3237	0.0797	0.2979	0.0432	0.4739	0.0020	0.2310
0 + 0	5 738	1 121	0.770	1 136	2 211	63 012	7 183	17 177	0 306	3 01	85 264
0 + 0 0 + 0.1	5.668	4.424	0.773	4.430	1.0626	61 724	7.405	17.177	0.300	3.91	83.204
0 + 0.1 $450 \pm 0$	5.685	4.405	0.775	4.410	2 041	61 650	7.200	17.204	0.264	3.82	81 081
$430 \pm 0$ 0.1 + 450	5 732	4.397	0.767	5 689	2.041	62 442	7.170	16.093	0.204	3.70	83 627
SEM	0.0271	0.0168	0.0045	0.4091	0.1738	0.6481	0.1598	0.0052	0.1131	0.0495	1 04953
P_value	0.0271	0.2952	0.6169	0.9011	0.1750	0.0401	0.1398	0.3653	1	0.3384	0.0020
Fe + Iodine + B12	0.0505	0.2752	0.0107	0.0711	0.3430	0.7700	0.1104	0.5055	1	0.5504	0.7727
0 + 0 + 0	5 722	4 4 3 1	0 774	4 484	2 580	63 060	7 429	17 932	0 294	3 18 <sup>b</sup>	84 986
0 + 0 + 0 1	5 676	4 3956	0.767	4 605	1 943	61.836	7.125	17.299	0.274	3.74ª	83 334
0 + 450 + 0	5 708	4 3936	0.765	5 928	2 142	61.864	7.616	17.847	0.285	3.70ª	86 333
0.1 + 450 + 0	5,768	4,422	0.764	4.628	2.206	63,132	7.060	17.636	0.282	3.82ª	82.800
0+0+400	5 754	4 417	0.766	4 644	1 842	62 964	7 537	17.850	0.284	Δa	85 542
0 + 0 + 400	5 660	4 4 1 0	0.778	4 556	1 982	61 612	7 287	17.655	0.289	3 14b	84 441
0+450+400	5.662	4 400	0 774	4 523	1.962	61 436	7 212	17.416	0 3004	3.17 3.72ª	83 636
0.1 + 450 + 400	5.602	4 401	0 7712	4 469	1 922	61 52	7 279	17 401	0.288	3.72 3.74a	84 454
SEM	0.0384	0.0237	0.0064	0.5112	0.2457	0.9165	0.2260	0 2277	0.0074	0 16007	1 48426
P-value	0.8465	0.3951	0.2560	0.3214	0.2261	0.7526	0.3799	0.6978	0 1190	0.0064	0.3720

In our study, adding minerals and vitamins to diet has increased the egg yolk color index and these findings in agreement with other reports (Zurak et al. 2022). Also, it has been reported that the addition of some minerals such as selenium to the diet of laying hens increases the activity of enzymes related to antioxidants, especially the glutathione peroxidase enzyme, which reduces the potentials of cell damage of egg liquid and egg shell, and protects it against free radicals damages (Wakebe, 1998; Dvorskaet al., 2003).Vitamins as natural antioxidants protect the yolk pigments by protecting them against oxidation and preserving the color of the yolk (Omri et al. 2019;Kim and Shin, 2022). Guoet al. (2021) observed that adding vitamin A and K3 to the diet resulted an improvement in egg shell quality and yolk color, as well as the antioxidant status of the egg shell gland of old laying hens.

#### Egg quality in storage conditions

The effects of using iron and iodine salts, and vitamin  $B_{12}$  on albumen and yolk weight indices and antioxidant content in the process of 8 weeks of egg storage are shown in Table 5. As can be seen in table 5, the effect of two factors of supplementation and storage time on egg albumen and yolk weight and the amount of total antioxidant content characteristics has been investigated.

Regarding the albumen weight, the highest amount of this parameter is observed in control group and this

difference is significant (p<0.05), while no significant difference was seen in other treatments (p>0.05). Also, concerning the yolk weight trait, a significant difference be observed between the treatment containing Fe and the treatment containing a mixture of 3 supplements (p < 0.05), while the other treatments did not show a significant difference (p>0.05). Moreover, the results of this experiment showed that by increasing the levels of iron, iodine and vitamin  $B_{12}$  (alone and with combination) the antioxidant content of eggs will be improved significantly (p<0.05). About the storage time and the effect of different times of storage on egg quality and egg antioxidant content, the results indicated that increasing the storage time caused a decrease in albumen and yolk weight and this decrease was statistically significant (p < 0.05)and also the antioxidant content of the egg increased significantly with increasing days of storage in the

Table 5:	The effects of Fe,	I and vit B <sub>12</sub> on	the albumen and y	olk weight and	antioxidant c	content in the p	process of egg storage.
----------	--------------------	------------------------------	-------------------	----------------	---------------	------------------	-------------------------

Trantments	Items							
	Albumen weight	Yolk weight	Antioxidant content					
Control	62ª	19 <sup>ab</sup>	0.38 <sup>d</sup>					
Fe	55 <sup>b</sup>	20ª	0.68 <sup>b</sup>					
Ι	53 <sup>b</sup>	19 <sup>ab</sup>	0.67 <sup>b</sup>					
Vit B <sub>12</sub>	55 <sup>b</sup>	20ª	0.61°					
$Fe + I + Vit B_{12}$	52 <sup>b</sup>	18 <sup>b</sup>	0.77ª					
SEM	1.0880	0.389	0.0123					
P-Value	< 0.0001	0.0180	< 0.0001					
Time								
Storage start time	60°	19 <sup>b</sup>	0.68ª					
4 weeks storage	57 <sup>b</sup>	20ª	0.63 <sup>b</sup>					
8 weeks storage	49ª	19 <sup>b</sup>	0.56 <sup>b</sup>					
SEM	0.8427	0.3014	0.0095					
P-Value	< 0.0001	0.0008	< 0.0001					
Treatments + Time								
Control + Storage start time	65 <sup>ab</sup>	20 <sup>ab</sup>	0.42 <sup>d</sup>					
Control + 4 weeks storage	61ª	19 <sup>ab</sup>	$0.37^{d}$					
Control + 8 weeks storage	59ª	19 <sup>ab</sup>	$0.36^{d}$					
Fe + Storage start time	58ª	21 <sup>ab</sup>	$0.74^{\mathrm{ab}}$					
Fe + 4 weeks storage	54 <sup>ab</sup>	$20^{ab}$	0.68 <sup>b</sup>					
Fe + 8 weeks storage	52 <sup>ab</sup>	19 <sup>ab</sup>	0.62 <sup>b</sup>					
I + Storage start time	61ª	18 <sup>bc</sup>	$0.76^{ab}$					
I + 4 weeks storage	56 <sup>ab</sup>	21 <sup>ab</sup>	0.71 <sup>b</sup>					
I + 8 weeks storage	42°	$20^{ab}$	0.56°					
Vit $B_{12}$ + Storage start time	59ª	19 <sup>ab</sup>	$0.66^{\mathrm{bc}}$					
Vit $B_{12}^{+}$ + 4 weeks storage	55 <sup>ab</sup>	22ª	0.60 <sup>bc</sup>					
Vit $B_{12}^{12}$ + 8 weeks storage	52 <sup>ab</sup>	19 <sup>ab</sup>	0.57°					
$Fe + I + Vit B_{12} + Storage start time$	56 <sup>ab</sup>	17c	0.83ª					
$Fe + I + Vit B_{12} + 4$ weeks storage	58ª	$20^{ab}$	$0.77^{ab}$					
$Fe + I + Vit B_{12} + 8$ weeks storage	41°	18 <sup>bc</sup>	$0.70^{a}$					
SEM	1.8844	0.6740	0.0213					
P-Value	0.0004	0.0080	0.0058					

J HELLENIC VET MED SOC 2023, 74 (4) ПЕКЕ 2023, 74 (4) refrigerator condition (4 degrees of Celsius and 60% humidity) (p>0.05).

About the mutual effects of supplements and storage time which there was observed a significant difference in albumen weight in the treatment containing 450 mg I/kg of diet of laying hens, (p<0.05) and also in treatment mixture of Fe, I and vitamin  $B_{12}$  (p<0.05), while other treatments did not show significant difference in albumen weight (p>0.05). Besides, the effect of treatment include three supplements (Fe + I + vit  $B_{12}$ ) showed a significant difference on yolk weight (p<0.05) during egg storage time, while other treatments did not show a significant difference (p>0.05). Furthermore, the mutual effects of mineral supplements and vitamin  $B_{12}$  on the antioxidant content of eggs of laying hens there were significant improvements (p<0.05) in treatments containing 450 mg I/ kg and a mixture of 3 supplements. Meanwhile, there was no significant difference in the interaction effects between treatments and time in other experimental groups (p<0.05).

According to the findings obtained in this experiment, minerals and vitamins did not have much effect on the weight of egg yolk and albumen, and the most important factor could be storage conditions and the loss of moisture and Co<sub>2</sub> through the egg shell and our findings in agreement with other reports (Eke et al., 2013). Therefore, the amount of antioxidants in eggs did not affect the weight of egg yolk and albumen, and it is just the storage conditions that affect the external and internal quality of eggs. Also, according to the decreasing trend of antioxidants in 8 weeks of storage and for treatments without mineral and vitamin supplements, it is a constant trend and covers approximately 12% and numerically it is 0.05 g/kg. But then for the treatments that have two minerals and vitamin  $B_{12}$  supplements, this reduction trend was 7% of the initial amount of antioxidants, so therefore, the antioxidants capacity of these eggs increases in storage conditions.

Antioxidants play an important role in maintaining the balance between free radicals produced by the metabolic system or taken from environmental sources and the body's antioxidant system. Damage to the endogenous antioxidant system contributes to the accumulation of free radicals and stimulates lipid peroxidation processes (Singh et al., 2004). Therefore, supplementing the diet with antioxidants is a suitable method to reduce oxidative stress, and in this field, the role of low-requirement elements and vitamins cannot be ignored and it was bold with other researchers (Khan et al., 2014).On the other hand, trace elements and vitamins have an effect on various physiological functions and have a major role in bone and egg shell formation and internal quality of eggs in poultry (Obianwuna et al., 2022). Subsequently deficiency of these elements can have a negative effect on egg quality (Nestor et al., 1985; Lilburn, 1994; Bessei, 2006; Angel, 2007). These factors can reduce production efficiency and ultimately lead to economic losses (Cook, 2000; Cotoet al., 2008). Based on previous studies, the mineral elements and vitamins can increase the antioxidant capacity of eggs and our findings cover that (Ghasemi et al., 2022).

The internal quality of the egg, even at the moment of laying, is affected by some chemical changes in the egg, and the deterioration process begins with the production of  $CO_2$  gas. During the storage period, some physical and chemical changes, which include the increase in the area of the air bag, the decrease in the height of the egg yolk and albumen, which is affected by the loss of water and  $Co_2$  from the shell, cause a decrease in the quality of the yolk and albumen (Arivazhaganet al., 2013). The amount of albumen reducing process during the storage, is highly dependent on the storage conditions (temperature and relative humidity) and shell characteristics (Obianwuna, 2022).

Storage time is usually used to identify eggs that are fresh and suitable for consumer use (according to above, it is not possible to rely only on the number of days after laying) (Rossi et al., 2010) and the supplemental treatments of this experiment could be helpful in stability of egg quality in 4 and 8 weeks of egg storage time.

#### CONCLUSION

Since our knowledge is still limited in practical usages of mineral and vitamin additives, totally the writers suggest that the proposed levels of Iron, Iodine and Vit  $B_{12}$  in this study maybe beneficial to enrichment the eggs of laying hens and therefore it could improve the storage condition of eggs but it needs more studies.

#### **CONFLICT OF INTEREST DECLARATION**

The authors declare that they have no known competing financial interests or personal relationship that could have appeared to influence the work reported in this paper.

#### ACKNOWLEDGEMENTS

The article was extracted from the PhD. thesis pre-

pared by Mir Nima Aghdashi, under the guidance of Prof. Ali Nobakht.

#### REFERENCES

- Angel, R., 2007. Metabolic disorders: limitations to growth of and mineral deposition into the broiler skeleton after hatch and potential implications for leg problems. *Journal of Applied Poultry Research*, 16(1), pp.138-149.
- Arivazhagan, S., Shebiah, R.N., Sudharsan, H., Kannan, R.R. and Ramesh, R., 2013. External and internal defect detection of egg using machine vision. *Journal of Emerging Trends in Computing and Information Sciences*, 4(3), pp.257-262.
- Bagheri, S., Janmohammadi, H., Maleki, R., Ostadrahimi, A. and Kianfar, R., 2019. Laying hen performance, egg quality improved and yolk 5-methyltetrahydrofolate content increased by dietary supplementation of folic acid. *Animal nutrition*, 5(2), pp.130-133.
- Behroozlak, M. A., Daneshyar, M., Farhoomand, P., & Nikoo, A. (2019). Effect of Replacing Dietary FeSO4 with FeHPO4 Nanoparticles on Growth Performance, Carcass Characteristics and Tissue Iron Content in Broilers. *Iranian Journal of Applied Animal Science*, 9(4), 709-716.
- Bessei, W., 2006. Welfare of broilers: a review. *World's Poultry Science Journal*, 62(3), pp.455-466.
- Christensen, V.L. and Ort, J.F., 1991. Iodine toxicity in large white turkey breeder hens. *Poultry Science*, 70(12), pp.2402-2410.
- Cook, M.E., 2000. Skeletal deformities and their causes: introduction. *Poultry Science*, 79(7), pp.982-984.
- Coto, C., Yan, F., Cerrate, S., Wang, Z., Sacakli, P., Halley, J.T., Wiernusz, C.J., Martinez, A. and Waldroup, P.W., 2008. Effects of dietary levels of calcium and nonphytate phosphorus in broiler starter diets on live performance, bone development and growth plate conditions in male chicks fed a corn-based diet. *International Journal of Poultry Science*, 7(2), pp.101-109.
- Damaziak, K., Marzec, A., Riedel, J., Szeliga, J., Koczywąs, E., Cisneros, F., Michalczuk, M., Łukasiewicz, M., Gozdowski, D., Siennicka, A. and Kowalska, H., 2018. Effect of dietary canthaxanthin and iodine on the production performance and egg quality of laying hens. *Poultry Science*, 97(11), pp.4008-4019.
- Dvorska, J.E., Yaroshenko, F.A., Surai, P.F. and Sparks, N.H.C., 2003. Selenium-enriched eggs: quality evaluation. In *Proc. 14 th European Symp. Poult. Nutr. Lillehammer, Norway* (pp. 23-24).
- Eke, M.O., Olaitan, N.I. and Ochefu, J.H., 2013. Effect of storage conditions on the quality attributes of shell (table) eggs. *Nigerian Food Journal*, 31(2), pp.18-24.
- Gan, L., Zhao, Y., Mahmood, T., & Guo, Y. 2020. Effects of dietary vitamins supplementation level on the production performance and intestinal microbiota of aged laying hens. *Poultry Science*, 99(7), 3594-3605.
- Ghasemi, H.A., Hajkhodadadi, I., Hafizi, M., Fakharzadeh, S., Abbasi, M., Kalanaky, S. and Nazaran, M.H., 2022. Effect of advanced chelate compounds-based mineral supplement in laying hen diet on the performance, egg quality, yolk mineral content, fatty acid composition, and oxidative status. *Food Chemistry*, 366, p.130636.
- Guo, S., Niu, J., Xv, J., Fang, B., Zhang, Z., Zhao, D., Wang, L. and Ding, B., 2021. Interactive effects of vitamins A and K3 on laying performance, egg quality, tibia attributes and antioxidative status of aged Roman Pink laying hens. *Animal*, 15(6), p.100242.
- Haugh, R.R., 1937. A new method for determining the quality of an egg. US Egg Poultry Mag, 43, 552-555.
- Kaufmann, S., Wolfram, G., Delange, F. and Rambeck, W.A., 1998. Iodine supplementation of laying hen feed: A supplementary measure

to eliminate iodine deficiency in humans?. Zeitschrift für Ernährungswissenschaft, 37(3), pp.288-293.

- Khan, R.U., Zia-ur-Rahman, Javed, I. and Muhammad, F., 2014. Serum antioxidants and trace minerals as influenced by vitamins, probiotics and proteins in broiler breeders. *Journal of Applied Animal Research*, 42(3), pp.249-255.
- Kim, M.J. and Shin, W.S., 2022. Stability of zeaxanthin/lutein in yolk oil obtained from microalgae-supplemented egg under various storage conditions. *LWT*, 155, p.112899.
- Leeson, S. and Caston, L.J., 2003. Vitamin enrichment of eggs. Journal of Applied Poultry Research, 12(1), pp.24-26.
- Lilburn, M.S., 1994. Skeletal growth of commercial poultry species. *Poultry science*, 73(6), pp.897-903.
- Mertens, K., Bamelis, F., Kemps, B., Kamers, B., Verhoelst, E., De Ketelaere, B., Bain, M., Decuypere, E. and De Baerdemaeker, J., 2006. Monitoring of eggshell breakage and eggshell strength in different production chains of consumption eggs. *Poultry Science*, 85(9), pp.1670-1677.
- National Research Council, 1994. Nutrient requirements of poultry: 1994. National Academies Press.
- Nestor, K.E., Bacon, W.L., Saif, Y.M. and Renner, P.A., 1985. The influence of genetic increases in shank width on body weight, walking ability, and reproduction of turkeys. *Poultry Science*, 64(12), pp.2248-2255.
- Nobakht, A., Palangi, V., Ayaşan, T., &Coçkun, I. (2022). Efficacy of Tragopogóngraminifolius medicinal powder as an inulin source for laying hens. South African Journal of Animal Science, 52(3), 252-258.
- Obianwuna, U.E., Oleforuh-Okoleh, V.U., Wang, J., Zhang, H.J., Qi, G.H., Qiu, K. and Wu, S.G., 2022. Potential Implications of Natural Antioxidants of Plant Origin on Oxidative Stability of Chicken Albumen during Storage: A Review. *Antioxidants*, 11(4), p.630.
- Omri, B., Alloui, N., Durazzo, A., Lucarini, M., Aiello, A., Romano, R., Santini, A. and Abdouli, H., 2019. Egg yolk antioxidants profiles: Effect of diet supplementation with linseeds and tomato-red pepper mixture before and after storage. *Foods*, 8(8), p.320.
- Rossi, M., Casiraghi, E., Primavesi, L., Pompei, C. and Hidalgo, A., 2010. Functional properties of pasteurised liquid whole egg products as affected by the hygienic quality of the raw eggs. *LWT-Food Science and Technology*, 43(3), pp.436-441.
- Shah, F., & Dwivedi, M. 2022. Vitamin B12 Producing Actinobacteria as Probiotics for Poultry Production. *In Methods in Actinobacteriology* (pp. 571-588). Humana, New York, NY.
- Singh, R.P., Sharad, S. and Kapur, S., 2004. Free radicals and oxidative stress in neurodegenerative diseases: relevance of dietary antioxidants. *J Indian Acad Clin Med*, 5(3), pp.218-225.
- Wakebe, M., 1998. Organic selenium and egg freshness. Patent# 10-23864. Feed for meat chickens and feed for laying hens. Japanese Patent Office, Application Heisei, pp.8-179629.
- Yalçin, S.I.D.D.I.K.A., Kahraman, Z., Yalcin, S., Yalcin, S.S. and Dedeoğlu, H.E., 2004. Effects of supplementary iodine on the performance and egg traits of laying hens. *British poultry science*, 45(4), pp.499-503.
- Zurak, D., Slovenec, P., Janječić, Z., Bedeković, X.D., Pintar, J. and Kljak, K., 2022. Overview on recent findings of nutritional and non-nutritional factors affecting egg yolk pigmentation. *World's Poultry Science Journal*, 78(2), pp.531-560.

J HELLENIC VET MED SOC 2023, 74 (4) ПЕКЕ 2023, 74 (4)