

Journal of the Hellenic Veterinary Medical Society

Vol 76, No 3 (2025)



Acute phase proteins and oxidative stress before and after treatment in neonatal calves with sepsis caused by *Escherichia coli* K99

E Akyüz, O Merhan, M Sezer, MR Coşkun, EE Erkiçiç, YU Batı, K Bozukluhan, M Kuru, M Şahin, G Gökce

doi: [10.12681/jhvms.32294](https://doi.org/10.12681/jhvms.32294)

Copyright © 2025, E Akyüz, O Merhan, M Sezer, MR Coşkun, EE Erkiçiç, YU Batı, K Bozukluhan, M Kuru, M Şahin, G Gökce



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

To cite this article:

Akyüz, E., Merhan, O., Sezer, M., Coşkun, M., Erkiçiç, E., Batı, Y., Bozukluhan, K., Kuru, M., Şahin, M., & Gökce, G. (2025). Acute phase proteins and oxidative stress before and after treatment in neonatal calves with sepsis caused by *Escherichia coli* K99. *Journal of the Hellenic Veterinary Medical Society*, 76(3), 9419–9428. <https://doi.org/10.12681/jhvms.32294>

Acute phase proteins and oxidative stress before and after treatment in neonatal calves with sepsis caused by *Escherichia coli* K99

E. Akyüz,^{*1,a} O. Merhan,^{2,b} M. Sezer,^{1,c} MR. Coşkun,^{3,d} EE. Erkiş, ^{1,e} YU. Batı,^{1,f}
K. Bozukluhan,^{4,g} M. Kuru,^{5,h} M. Şahin,^{3,i} G. Gökçe^{1,j}

¹Kafkas University, Faculty of Veterinary Medicine, Department of Internal Medicine, 36100 Kars, Türkiye

²Kafkas University, Faculty of Veterinary Medicine, Department of Biochemistry, 36100 Kars, Türkiye

³Kafkas University, Faculty of Veterinary Medicine, Department of Microbiology, 36100 Kars, Türkiye

⁴Kafkas University, Kars Vocational School, Department of Veterinarian and Laboratory, 36100 Kars, Türkiye

⁵Kafkas University, Faculty of Veterinary Medicine, Department of Obstetrics and Gynecology, 36100 Kars, Türkiye

ABSTRACT: This study aimed to detect changes in acute-phase proteins endothelin-1 (ET-1), pentraxin-3 (P-3), ceruloplasmin (Cp), haptoglobin (Hp), and albumin, in calves with diarrhea caused by *Escherichia coli* K99 neonatal sepsis before and after treatment. Diagnostic and prognostic clinical, biochemical, and hematological parameters were evaluated. The sepsis group was composed of 20 newborn calves which met neonatal sepsis criteria, did not receive any treatment, and were referred to the clinics within 24 hours after the detection of clinical findings. The control group consisted of 10 healthy calves for comparison. Clinical examinations were performed before (0th h) and after (7th d) treatment. Blood samples were drawn via the jugular vein from sick calves before and after treatment and once from control animals. The P-3, ET-1, Cp, Hp, and malondialdehyde (MDA) concentrations in the sepsis group before treatment were statistically higher than post-treatment and control group values ($P<0.05$); while the glutathione (GSH) concentration was statistically lower ($P<0.05$). In conclusion, the high pre-treatment concentrations of P-3, ET-1, Cp, Hp, and MDA in the sepsis group were of diagnostic importance. The approximation of these parameters after treatment to the mean values of the control group was also of prognostic value.

Keyword: Calf; Ceruloplasmin; Endothelin-1; Haptoglobin; Pentraxin-3.

Correspondence author:

E. Akyüz,
Kafkas University, Faculty of Veterinary Medicine,
Department of Internal Medicine, 36100 Kars, Türkiye
E-mail address: enesakuyuz_44@hotmail.com

Date of initial submission: 7-12-2022

Date of acceptance: 16-5-2023

INTRODUCTION

Neonatal sepsis is a serious cause of mortality in calves (Radostits et al., 2006; Basoglu et al., 2018; Akyüz, 2020; Akyüz and Gökce, 2021), and *E. coli* is responsible for the majority of economic loss related to neonatal calves in the first 5 days after birth (Ok et al., 2009; Ok et al., 2020). The etiology of sepsis is very complex. Death usually occurs due to diarrhea, respiratory system problems, and multiple organ failure (Asadi et al., 2015; Merhan et al., 2016; Akyüz et al., 2016; Chatre et al., 2017). Neonatal calf diseases can be caused by a single factor or due to a combination of many factors (Ok et al., 2009; Erkılıç et al., 2019; Akyüz, 2020), including viral, bacterial, and protozoal components (Ok et al., 2009; Lorenz et al., 2011; Akyüz and Gökce, 2021; Akyüz and Kükürt, 2021). Although the transmission route of these agents is usually via the oral route from contaminated equipment that does not meet hygienic conditions, spread can also occur by omphalogen or aerogenous routes (Trefz et al., 2013). Inflammatory situations created by infectious agents lead to activation of the body's defense mechanisms. As a result, systemic inflammatory response syndrome (SIRS) and sepsis may develop. Affected animals may experience decreased interest in the environment, anorexia, diarrhea, weakening of the sucking reflex, depression, lethargy, drop in body temperature, increase in respiratory rate, prolongation of capillary refill time (CRT), and dehydration (House et al., 2015; Bonelli et al., 2018; Akyüz, 2020). Bacterial infections such as *E. coli*, viral infections such as rotavirus and coronavirus, and parasitic diseases such as cryptosporidiosis that cause sepsis are responsible for many newborn calf deaths (Chatre et al., 2017; Basoglu et al., 2018; Akyüz and Gökce, 2021). Although different treatment options have been tried, sepsis severity and mortality rates remain high (Novelli et al., 2010).

The importance of using biomarkers to diagnose sepsis is increasing in both human and veterinary medicine (Moore et al., 2007; Köse and Maden, 2013). Acute-phase proteins are blood proteins used to evaluate the body's response to infection, inflammation, and trauma (Murata et al., 2004; Petersen et al., 2004; Coşkun and Şen, 2011). These proteins play a significant role as biomarkers in the evaluation and diagnosis of sepsis and in monitoring the response to treatment (Meisner, 2005; Dupuy et al., 2013; Sönmezer and Tülek, 2015). Haptoglobin (Hp) and ceruloplasmin (Cp) are positive acute-phase pro-

teins (Murata et al., 2004; Coşkun and Şen, 2011; Bozukluhan et al., 2021; Ceron et al., 2005; Erkılıç, 2019). The role of Hp is to prevent iron loss after forming complex structures by binding hemoglobin (Cray, 2012; Tuna and Ulutaş, 2015; Erkılıç, 2019). Cp, synthesized from the liver (Murata et al., 2004; Erkılıç, 2019), is a plasma antioxidant that binds free copper in the serum and provides protection against the damage that can be caused by iron during inflammatory events (Cray, 2012; Tuna and Ulutaş, 2015; Erkılıç, 2019).

Pentraxin-3 (P-3) is an acute-phase protein released from macrophages, dendritic cells, leukocytes, and endothelial cells during the inflammatory response (Libby et al., 2009). Studies have found a positive correlation between disease severity and plasma P-3 levels (Kao et al., 2013; Sönmezer and Tülek, 2015). Endothelin-1 (ET-1) is a very potent vasoconstrictor released by endothelial cells. In endotoxemias, prepro-ET-1 is released from the heart and lungs and increases in direct proportion to disease mortality rates (Shah, 2007; Tschakowsky et al., 2000; Sönmezer and Tülek, 2015). Glutathione (GSH), a substance synthesized from the amino acids glycine, cysteine, and glutamic acid, is intended to protect against oxidative tissue damage (Parlakpınar et al., 2013; Akyüz et al., 2021). The organic compound malondialdehyde (MDA) is widely used to evaluate oxidative stress and determine cellular damage (Akyüz et al., 2021; Bozukluhan et al., 2021).

This study aimed to evaluate the diagnostic and prognostic value of some acute-phase proteins, oxidative stress indexes, as well as clinical, biochemical, and hematological parameters (measured before and after treatment) in calves with diarrhea and neonatal sepsis caused by *E. coli* K99. We hypothesized that neonatal calves with sepsis caused by *E. coli* would have higher concentrations of ET-1 and P-3 than normal calves.

MATERIAL AND METHODS

Ethical Approval

This study was conducted with approval number "KAU-HADYK/2020-078" from the local ethics committee of Kafkas University, Kars, Türkiye. Signed informed consent document was obtained from the calf owners.

Animals

The animals evaluated in this study were 30 calves divided into two groups. The sepsis group was com-

posed of 20 newborn calves (0-10 days old) which met neonatal sepsis criteria but did not receive any treatment, referred to the clinics of the Department of Internal Medicine, Faculty of Veterinary Medicine, Kafkas University within 24 hours after clinical findings were detected. A control group consisting of 10 healthy calves (0-10 days old) housed in the closed barn system in the University of Veterinary Medicine Education, Research, and Application Farm were used for comparison.

Procedures

Agent Detection from Feces

A rapid test kit (BoviD-5 Ag Test Kit[®], Bionote Inc., Korea) was used to detect etiological factors in the feces of calves with neonatal sepsis. Only samples with positive *E. coli* confirmed by feces rapid test kit results were included in the study. Mixed samples were not included. Positive feces samples were sent to Kafkas University Veterinary Faculty Microbiology Laboratory for confirmation.

Obtaining and Measuring Blood Samples

Blood was collected in gel-containing vacuum tubes (BD Vacutainer[®], BD, UK) for biochemical measurements and K₂EDTA-containing tubes for hematological measurements (BD Vacutainer[®], BD, UK). Blood samples were drawn via the jugular vein from sick calves before (0th h) and after (7th d) treatment and once from control animals. Blood samples were obtained from patients brought to Kafkas University Veterinary Faculty Clinics for treatment and healthy calves brought for control purposes. Total leukocyte count (WBC, x10³/µL) and other hematological parameters were determined using a complete blood count device (VG-MS4e[®], Melet Schloesing Labs., Osny, France) within 30 minutes of sample collection. For analysis requiring sera, after approximately 1 hour at room temperature, blood samples were centrifuged for 10 min at 3000 rpm (Hettich Rotina 380R[®], Hettich, Germany) and stored at -20°C until measured. Alanine aminotransferase (ALT), aspartate aminotransferase (AST), gamma-glutamyl transferase (GGT), alkaline phosphatase (ALP), lactate dehydro-

genase (LDH) activities, total protein (TP), cholesterol, triglycerides, glucose, creatinine, urea, and creatine kinase (CK) were assessed using a fully automated biochemistry device (Mindray BS120[®], Mindray Medical Technology İstanbul, Türkiye). Albumin was measured colorimetrically (Epoch[®], Biotek, USA) with the methods reported for Hp by Skinner et al. (1991) and for Cp by Colombo and Richterich (1964). Cattle P-3 (PTX3[®], Cat: ELK8840, ELK Biotechnology, Wuhan, China) and cattle ET-1 (EDN-1[®], Cat: ELK8839, ELK Biotechnology, Wuhan, China) were measured colorimetrically with ELISA kits. Serum measurements of MDA were performed according to Yoshioka et al., (1979) and GSH according to Beutler et al., (1963). The obtained data were processed using a spectrophotometric microplate reader (Spectramax Plus[®], Marshall Scientific, Product Code: MD-SMP, NH, USA).

Polymerase Chain Reaction Procedure

Polymerase chain reaction (PCR) was performed to confirm the *E. coli* K99 strain detected in the feces samples using the rapid test kit. For PCR, 1 g of feces was diluted with 3 ml of phosphate-buffered saline (PBS), from which 1 mL was taken and added to 9 mL of buffered peptone water and incubated at 37°C for 18 hours. At the end of the incubation, 1 mL of the culture was centrifuged at 8000 rpm for 10 minutes. The supernatant was removed, and the pellet was washed 3 times with PBS. At the end of washing, DNA extraction was performed by adding 100 µL of nuclease-free water and boiling the pellet at 99°C for 10 minutes (Franck et al. 1998). The primer pair in Table 1 was used to determine the K99 gene region of the obtained DNA. For the 50 µL PCR mix, 5 µL of 10x PCR buffer (Thermo Fisher EP0402[®], Thermo Fisher Scientific, USA), 1.5 mM MgCl₂ (Thermo Fisher EP0402[®], Thermo Fisher Scientific, USA), 1 mM dNTP (Ampliqon A502004[®], 40 mM mix, Ampliqon A/S, Denmark), 0.5 µM from each primer sequence (Sentebiolab[®], Türkiye), 1.25 U Taq polymerase (Thermo Fisher EP0402[®], 5 U/µL, Thermo Fisher Scientific, USA), and 5 µL of the obtained bacterial DNA were added

Table 1. Study primer pair

Virulence factor	Primary Sequence 5'- 3'	Reference	Product size
K99 (F)	TATTATCTTAGGTGGTATGG	(Roosendaal et al. 1984)	314
K99 (R)	GGTATCCTTTAGCAGCAGTATTTTC		

to nuclease-free water. Amplification of the prepared mixture consisted of 25 cycles of denaturation at 94°C for 30 seconds, bonding at 50°C for 45 seconds, elongation at 70°C for 90 seconds, with 3 seconds per cycle added. The final bonding was conducted at 70°C for 10 minutes (Franck et al. 1998). The K99 strain found in the collection of the laboratory of the Microbiology Department of the Faculty of Veterinary Medicine of Kafkas University was used as a positive control. The primer pair used was per Roosendaal et al., (1984) (Table 1).

SIRS and Sepsis Evaluation

There are specific criteria that should be met when evaluating for SIRS and sepsis. The presence of at least two of the following criteria is required for diagnosis of SIRS: Hypothermia or hyperthermia, tachycardia, tachypnea, increased arterial partial carbon dioxide pressure, leukopenia or leukocytosis. If infection accompanies SIRS, this is considered sepsis (Fecteau et al., 2009; Sen and Constable, 2013; Yıldız et al., 2018; Beydilli and Gökçe, 2019; Akyüz and Gökçe, 2021). Calves with symptoms of depression, diarrhea, low/lack of sucking reflex, dehydration, and constant urge to lie down were examined and evaluated for sepsis criteria. The SIRS criteria for neonatal calves used were: body temperature >39.5°C or <37°C, pulse rate per minute <100 or >160, respiratory rate per minute >45, leukocyte count >12×10³/µL or <4×10³/µL (Fecteau et al., 1997, 2009; Yıldız et al., 2018; Akyüz and Gökçe, 2021). Calves exhibiting at least two of the specified criteria and with proven infection (sepsis) were included in the study.

Treatment

Calves with sepsis were kept under observation in separate compartments during treatment. Fluid therapy was provided to these animals according to the severity of dehydration. For this purpose, the calves were intravenously administered 0.9% NaCl (60-80 mL/kg, PVC[®], 1000 mL Eczacıbaşı Baxter, İstanbul, Türkiye), 1.3% NaHCO₃ (5 mL/kg, Bikarvil[®], Vilsan, Ankara, Türkiye), and 5% dextrose (1 gr/kg, Polifleks[®], Polifarma, Tekirdağ, Türkiye). For infection control, they were intramuscularly administered enrofloxacin (Baytril %10[®], Bayer, Germany) 2.5-5 mg/kg, once a day, for 7 days. They were orally administered antipyretic neomycin sulfate and bismuth subcarbonate-containing powder (Cesamol[®], Topkim, Türkiye) 10-20 mg/kg for 3 days. For vitamin supplementation, the calves were parenterally administered vitamin B complex (Berovit B12[®],

Ceva, Australia) at a dose of 8-10 mL/calf for 7 days, vitamin C (Maxivit-C[®], Bavette, Türkiye) 4-6 mg/kg for 7 days, and vitamin ADE in a single subcutaneous dose of 1 mL/50 kg (Ademin[®], Ceva, Australia). For mineral supplementation, they were subcutaneously administered a solution of Ca, P, and Mg (Kalsimin[®], Vilsan, Germany) at a single dose of 10 mL/50 kg. The calves were also given a single 4-6 mg/kg dose of meloxicam (Bavet Meloxicam[®], Bavet, Türkiye). Sick calves were treated for 7 days. No application was made to the control group calves.

Statistical analysis

Normal distribution of the data was evaluated using visual methods (histogram graph and Q-Q graph) and the Shapiro-Wilk test. The group with sepsis was compared with the Paired t-test before and after treatment. Independent sample t-test was used to compare the group with sepsis before and after treatment with the control group. Pearson correlation coefficients were calculated to define the correlation between variables. The data obtained in the study were reported as mean ± standard error (SEM). All analyses were performed with the SPSS[®] software program (SPSS Statistics 26.0, Chicago, IL, USA). Differences obtained in group comparisons were considered significant at P<0.05.

RESULTS

Clinical examinations of the calves that met the sepsis criteria revealed different degrees of diarrhea, no or reduced sucking reflex, depression, little interest in the environment, and inability to stand. The K99 gene region was detected by PCR in all stool samples of the 20 calves with sepsis (Fig. 1). All of the calves with sepsis that were treated survived. Physical examination of the calves with sepsis found the mean values of CRT and heart and respiratory rates per minute before treatment to be statistically significantly higher than post-treatment and control calf values (P<0.001, Table 2).

Before treatment, WBC, lymphocyte, and granulocyte counts were higher in calves with sepsis than after treatment and controls (P<0.05, Table 2).

In calves with sepsis, pre-treatment ALT, AST, ALP, urea, and creatinine activities/concentrations were statistically significantly higher than after treatment and control concentrations (P<0.05, Table 2). On the other hand, serum glucose was at its lowest level before treatment in the sepsis group, but increased after treatment, approaching control animal values. Albumin, and triglyceride concen-

Table 2. Clinical, hematological, and biochemical parameters of calves

Parameters	Sepsis Group		Control Group	P-values		
	Before Treatment	After Treatment		BT-AT	BT-C	AT-C
	Mean ± SEM					
Rectal temperature (°C)	38.52±0.43	38.42±0.07	38.38±0.06	0.822	0.756	0.725
Breaths/min	59.05±3.05	35.16±0.67	28.60±0.89	<0.001	<0.001	<0.001
Heartbeats/min	158.20±8.66	93.90±3.99	80.40±1.80	<0.001	<0.001	0.005
Capillary refill time/sec	3.00±0.14	1.40±0.06	1.50±0.04	<0.001	<0.001	0.161
Total leukocytes count (×10 ³ /µL)	15.85±1.60	9.15±0.92	7.66±0.28	0.005	<0.001	0.273
Lymphocytes count (x10 ³ /µL)	5.10±0.73	3.15±0.35	3.04±0.43	0.060	0.009	0.856
Monocytes count (x10 ³ /µL)	0.34±0.05	0.42±0.03	0.56±0.13	0.218	0.158	0.325
Granulocytes count (x10 ³ /µL)	9.43±0.74	5.58±0.77	3.85±0.65	0.004	<0.001	0.158
Red blood cell count (x10 ⁶ /µL)	7.08±0.37	7.84±0.27	8.63±0.23	0.108	0.002	0.073
Mean red cell volume (fL)	43.54±0.67	42.43±0.56	41.50±0.94	0.080	0.090	0.380
Hematocrit (%)	36.82±2.07	30.85±1.86	35.99±2.28	<0.001	0.805	0.108
Hemoglobin (g/dL)	8.78±0.60	9.22±0.55	10.51±0.42	0.547	0.070	0.140
Platelet count (x10 ³ /µL)	453.52±73.12	497.05±41.58	288.91±25.46	0.238	0.036	<0.001
Alanine aminotransferase (IU/L)	35.23±2.89	26.14±2.41	15.04±0.68	0.005	<0.001	<0.001
Aspartate aminotransferase (IU/L)	84.04±6.70	64.36±4.22	42.23±3.62	0.017	0.001	<0.001
Alkaline phosphatase (IU/L)	275.66±22.52	193.72±16.26	162.43±11.06	0.025	0.007	<0.001
Gamma-glutamyl transferase (IU/L)	383.82±72.21	250.26±86.17	302.98±52.97	0.002	0.466	0.670
Creatine (mg/dL)	3.17±0.20	1.87±0.10	1.46±0.10 ^a	<0.001	<0.001	0.003
Urea (mg/dL)	66.59±7.94	42.82±3.65	23.81±2.65	0.011	<0.001	0.002
Lactate dehydrogenase (IU/L)	742.30±34.72	760.48±56.31	771.71±60.70	0.705	0.655	0.902
Glucose (mg/dL)	50.70±5.37	82.80±4.12	97.15±2.31	<0.001	<0.001	0.005
Total protein (g/dL)	6.99±0.41	6.72±0.28	6.47±0.27	0.385	0.412	0.573
Albumin (g/dL)	2.69±0.09	2.92±0.13	2.96±0.07	0.163	0.005	0.952
Cholesterol (mg/dL)	37.75±2.77	28.89±1.96	39.79±2.65	0.003	0.581	0.002
Creatine kinase (IU/L)	416.49±84.52	498.64±95.83	360.49±74.70	0.354	0.673	0.466
Triglyceride (mg/dL)	23.80±4.19	16.15±1.71	42.60±7.64	0.068	0.026	0.007
Ceruloplasmin (mg/dL)	17.30±0.60	12.72±0.70	11.26±1.09	<0.001	<0.001	0.257
Haptoglobin (g/L)	0.63±0.03	0.11±0.007	0.06±0.005	<0.001	<0.001	0.008
Glutathione (umol/mL)	2.28±0.14	2.89±0.18	2.85±0.19	0.015	0.002	0.721
Malondialdehyde (nmol/mL)	1.53±0.05	1.31±0.02	1.27±0.02	0.005	<0.001	0.388
Pentraxin-3 (ng/mL)	9.81±0.35	4.85±0.16	3.11±0.09	<0.001	<0.001	<0.001
Endothelin-1 (pg/mL)	38.79±1.36	28.30±1.33	23.33±1.60	<0.001	<0.001	0.057

SEM: standard error of the mean. BT: Before treatment, AT: After treatment, C: Control.

trations were statistically significantly lower in the sepsis group before treatment compared to the control group ($P<0.05$, Table 2).

Haptoglobin and Cp acute-phase protein concentrations were considerably higher in the sepsis group (before treatment) compared to the controls; how-

ever, after treatment, concentrations reached values close to the controls ($P<0.001$, Table 2). While the GSH concentration was lower in the sepsis group compared to the control group before treatment ($P=0.002$, Table 2), the MDA concentration was higher ($P<0.001$, Table 2). ET-1 and P-3 concentrations before treatment were higher in the group




Figure 1. *E. coli* K99 PCR image obtained from a stool sample of a calf with sepsis. M: 100bp Marker 1: Positive control, 2: *E. coli* K99 positive sample.

with sepsis compared to post-treatment and controls ($P < 0.001$, Table 2).

Haptoglobin, Cp, GSH, MDA, P-3, and ET-1 were compared with all other study data (Table 3). A positive correlation was determined with Hp, Cp, P-3, ET-1, and total leukocytes. Hp, P-3, and ET-1 negatively correlated with erythrocyte count. Hp, Cp, and P-3 positively correlated with ALT and AST. Hp, Cp, GSH, MDA, P-3, and ET-1 parameters correlated differently with physical findings, hematology, clinical, and biochemical parameters (Table 3). While a significant positive correlation was determined between Hp and Cp, MDA, P-3, and ET-1, a negative correlation was determined with GSH. Cp showed similar correlations to Hp in ET-1, P-3, and MDA comparisons among themselves (Table 3). GSH, on the other hand, showed a significant negative correlation with Hp, MDA, P-3, and ET-1 (Table 3).

DISCUSSION

Sepsis is an *in vivo* response to infection associated with a high mortality rate. Early diagnosis and initiation of appropriate treatment are essential in the battle against sepsis (Dupuy et al., 2013; Bachmann et al., 2005; Sönmezer and Tülek, 2015). In light of this information, in our study, a rapid test kit for feces examination was used for early diagnosis in calves, and treatment was started within 24 hours at the latest after sepsis symptoms appeared.

Studies have reported that clinical findings such as depression, decreased interest in the environ-

ment, increase in respiratory and pulse rates, rise or fall in body temperature, and diminished appetite occur with sepsis (Aldridge et al., 1993; Fecteau et al., 1997; Çitil and Gökçe, 2013; Yıldız et al., 2018; Beydilli and Gökçe, 2019; Akyüz et al., 2016; Akyüz, 2020; Akyüz and Gökçe, 2021). In the present study, the physical examination findings before treatment in the sepsis group were similar to the results of other studies. These effects are most likely due to the deterioration of hemostasis due to SIRS and the inability of organs and systems to function properly. In animals with sepsis, tachycardia develops due to fluid loss and SIRS (Fecteau et al., 2009; Naseri, 2017; Yıldız et al., 2018; Akyüz and Gökçe, 2021). In this study, tachycardia detected before treatment in calves with sepsis may have been due to dehydration/hypovolemia and SIRS. Tachypnea occurs as a result of activation of compensation mechanisms associated with sepsis and infectious conditions (Fecteau et al., 1997; Fecteau et al., 2009; Naseri, 2017). Similarly, in our study, tachypnea occurred in septic animals. CRT prolongation in sepsis results from multiple organ dysfunction, adversely affecting the cardiovascular system (Fecteau et al., 2009; Akyüz and Gökçe, 2021). CRT was prolonged in our sepsis group compared to the control group prior to treatment.

Leukocytosis may occur in calves with sepsis (Dellinger et al., 2013; Naseri, 2017; Yıldız et al., 2018; Beydilli and Gökçe, 2019). In our study, leukocytosis associated with the increase in lymphocytes and granulocytes before treatment in the sepsis group was similar to the results of other studies. This leukocytosis may be due to SIRS-induced inflammation. Depending on disease severity, a decrease in the number of erythrocytes can be observed in calves with sepsis (Pardon and Deprez, 2018; Beydilli and Gökçe, 2019). Consistent with the literature, the low number of erythrocytes found in our study may be due to the absence of or low level of sucking reflex, the disruption of erythrocyte production, and the deterioration of organ function due to sepsis.

Increases in serum ALT, AST, and ALP activities have been reported in neonatal calves with diarrhea (Baser and Civelek, 2013; Naseri, 2017; Bozukluhan et al., 2017). We found serum ALT, AST, and ALP to be higher in calves with sepsis before treatment than after treatment and the control group, which may be a result of hepatic cell damage due to sepsis. Dehydration/hypovolemia is the most important reason for increased serum urea and creatinine concentrations in calves with diarrhea (Dratwa-Chalupnik et al., 2012).

Table 3. Pearson correlation of measured parameters

Parameters	Hp	Cp	GSH	MDA	P-3	ET-1
	Pearson Correlation Coefficient (R)					
Total leukocytes count ($\times 10^3/\mu\text{L}$)	0.481**	0.445**	-0.176	0.126	0.556**	0.579**
Lymphocytes count ($\times 10^3/\mu\text{L}$)	0.369**	0.136	-0.170	0.048	0.383**	0.440**
Monocytes count ($\times 10^3/\mu\text{L}$)	-0.189	-0.341*	0.144	-0.135	-0.267	-0.272
Granulocytes count ($\times 10^3/\mu\text{L}$)	0.554**	0.453**	-0.280*	0.282*	0.493**	0.531**
Red blood cell count ($\times 10^6/\mu\text{L}$)	-0.342*	-0.276	0.182	-0.195	-0.429**	-0.341*
Mean red cell volume (fL)	0.206	0.305*	-0.166	-0.095	0.288*	0.167
Hematocrit (%)	0.200	0.293*	-0.073	-0.031	0.108	0.109
Alanine aminotransferase (IU/L)	0.475**	0.446**	-0.056	0.214	0.618**	0.277
Aspartate aminotransferase (IU/L)	0.555**	0.281*	-0.381**	0.223	0.512**	0.294*
Glucose (mg/dL)	-0.636**	-0.328*	0.344*	-0.440**	-0.574**	-0.342*
Creatine (mg/dL)	0.734**	0.422**	-0.364**	0.253	0.755**	0.541**
Urea (mg/dL)	0.522**	0.244	-0.251	0.354*	0.578**	0.428**
Hp (g/L)	-	0.609**	-0.481**	0.577**	0.891**	0.647**
Cp (mg/dL)	0.609**	-	-0.254	0.269	0.616**	0.542**
Albumin (g/dL)	-0.383**	-0.309*	0.176	-0.186	-0.387**	-0.427**
GSH (umol/mL)	-0.481**	-0.254	-	-0.399**	-0.441**	-0.434**
MDA (nmol/mL)	0.577**	0.269	-0.399**	-	0.467**	0.286*
P-3 (ng/mL)	0.891**	0.616**	-0.441**	0.467**	-	0.712**
ET-1 (pg/mL)	0.647**	0.542**	-0.434**	0.286*	0.712**	-
Heartbeats/min	0.681**	0.523**	-0.421**	0.278	0.675**	0.560**
Breaths/min	0.841**	0.570**	-0.503**	0.501**	0.743**	0.677**
Capillary refill time/sec	0.761**	0.541**	-0.320*	0.459**	0.758**	0.539**

*: Correlation is significant at the 0.05 level (2-tailed), **: Correlation is significant at the 0.01 level (2-tailed). Hp: Haptoglobin, Cp: Ceruloplasmin, GSH: Glutathione, MDA: Malondialdehyde, P-3: Pentraxin-3, ET-1: Endothelin-1

It has been reported that serum creatinine and urea concentrations are higher in calves with sepsis than in healthy calves (Ercan et al., 2016; Akyüz and Gökce, 2021). In our study, the high pre-treatment serum urea and creatinine concentrations in calves with sepsis compared to healthy calves may be the result of dehydration/hypovolemia. This opinion is supported by the improvement of renal function as a result of fluid therapy in the sepsis group and the fact that serum urea and creatinine values reached the levels of the control group after treatment. In calves with diarrhea, fluid and electrolyte losses impair hemostasis, and carbohydrate stores are depleted, resulting in hypoglycemia (Dratwa-Chalupnik et al., 2012). Depletion and loss of carbohydrates, and the cessation of food intake, are known to cause hypoglycemia (Seifi et al., 2006). Serum glucose concentrations in calves with sepsis were reported to be lower than in healthy

calves (Ercan et al., 2016). In our study, the blood glucose concentration before treatment was also lower in the sepsis group than in the control group. This result is likely due to lack of food intake, glucose loss through diarrhea, and depletion of carbohydrate stores due to sepsis. There may be a decrease in the concentration of albumin, a negative acute-phase protein associated with inflammation and diarrhea (Uzlu et al., 2010; Marcato et al., 2018; Akyüz and Gökce, 2021). The results of our study are consistent with the literature in this regard. Low albumin levels in the sepsis group before treatment compared to the control group may be due to sepsis-induced inflammation. In many acute cases, plasma triglyceride levels were normal or low (Akgün et al., 1998). In our study, triglyceride levels were lower in calves with sepsis compared to controls. While cholesterol levels increase in nephrotic syndrome or protein-los-

ing nephropathy, hypothyroidism, acute pancreatitis, and cholestasis, the levels decrease in protein-losing enteropathy, hypoadrenocorticism, and acute inflammatory response (Carpenter and Scruel, 2002). In our study, the low cholesterol concentration in the sepsis group was most likely due to an inflammatory reaction and enteritis.

Acute-phase proteins, blood proteins generally synthesized in the liver, are used to evaluate the body's response in cases of inflammation or infection (Pradeep, 2014). One of the acute-phase proteins, Hp, which is important for ruminants, is typically at a low level in healthy animals, but increases in cases of inflammation, trauma, and infection (Petersen et al., 2004; McGrotty et al., 2003; Erkiliç, 2019). Thus, Hp is a clinically useful parameter to determine the severity of the inflammatory response in ruminants (Murata et al., 2004; Çitil, 2003; Heegaard et al., 2000; Coşkun and Şen, 2011). Merhan et al. (2016) reported that Hp and Cp levels increased in a study of calves with diarrhea. Pourjafar et al. (2011) also reported an increase in Hp concentration. This increase in Hp was probably due to tissue destruction. In addition, the Hp acute-phase protein increase may be the result of activation of the body's defense mechanisms against infection and the severe inflammation that develops in sepsis. After treatment, Hp concentration decreased. Cp is also a positive acute-phase protein (Ceron et al., 2005; Erkiliç, 2019) synthesized in the liver (Murata et al., 2004; Erkiliç, 2019). Cp, which is used less commonly in cattle than Hp, is reported as an indicator of infection (Gruys et al., 2005; Petersen et al., 2004). Cp acts on defense system cells and causes an increase in the antimicrobial power of these cells (Ceron et al., 2000). In a study conducted in calves, the level of Cp increased in cases of enteritis (Murata et al., 2004; Coşkun and Şen, 2011). In this study, the increase in Cp may be due to its effect on the defense system. In addition, the reason for this increase is thought to be due to the acute-phase response, infection, and plasma antioxidant properties.

Inflammation and infection activate many cells with phagocytic activity such as monocytes and macrophages (Tokoyuni, 1999) and cause the formation of free radicals (such as hydrogen peroxide, superoxide anion) due to excessive oxygen consumption. Studies have reported that the oxidant-antioxidant balance is disrupted, and oxidative stress occurs in bacterial, viral, and parasitic diseases such as traumatic reticuloperitonitis (Atakişi et al., 2010), sheeppox in sheep (Bozukluhan et al., 2018), and

hypodermosis in cattle (Merhan et al., 2017). In this study, the oxidant-antioxidant balance was disturbed, the MDA concentration increased, and the GSH concentration decreased, which may be due to free radicals formed by phagocytes, which play an important role in host defense. As a result of treatment, the MDA concentration decreased, while the GSH concentration increased.

Pentraxin-3 is an acute phase protein released from macrophages, dendritic cells, leukocytes, and endothelial cells during the inflammatory response (Libby et al., 2009). Studies have found a positive correlation between disease severity and the plasma P-3 level (Kao et al., 2013; Sönmezer and Tülek, 2015). Consistent with the literature, pre-treatment P-3 was higher in the sepsis group than in the controls in our study. This result may be due to the presence of leukocytosis in the sepsis group, the widespread inflammation caused by SIRS, and the increase in acute-phase proteins associated with infection. ET-1 is secreted by endothelial cells (Shah, 2007; Tschakowsky et al., 2000; Sönmezer and Tülek, 2015). The higher ET-1 values in the sepsis group in our study compared to the control group may be the result of endothelial damage caused by sepsis. In light of the data presented, it was determined that the oxidative stress load increased, the antioxidant capacity decreased, concentrations of Hp, Cp, and P-3 acute phase proteins increased, and the negative acute phase protein albumin decreased in calves with sepsis. In addition, it was determined that endothelial damage occurred. Accordingly, there were changes in the related enzymes and some organ tissue dysfunction.

CONCLUSION

Significantly higher levels of Hp, Cp, MDA, ET-1 and P-3 in calves with sepsis before and after treatment and controls are important for diagnosis. After treatment, these parameters decreased and reached close to control levels, which is important for prognosis evaluation. In addition, Hp, Cp, GSH, MDA, ET-1, and P-3 had significant correlations; therefore, evaluation of these parameters together is essential to determine a sepsis diagnosis swiftly and accurately. It is vital to start treating sick calves as soon as possible after clinical findings appear and a diagnosis of sepsis is made. The survival of all calves with sepsis in this study indicates that early diagnosis and treatment are critical. In conclusion, Hp, Cp, GSH, MDA, ET-1, and P-3 are important diagnostically and prognostically in neonatal calves with sepsis due to *E. coli* K99.

Author contributions: EA conceived and supervised the study. EA, OM, MS, and YUB collected and analyzed data. OM analyzed biochemical measurements. MRC and MŞ performed the PCR procedure. EA, GG, EEE, MK, and KB contributed to writing the article. All authors contributed to the critical revision of the manuscript and have read and approved the final version.

Funding: The authors declare that no funds, grants, or other support were received during the preparation of this manuscript.

Data availability: The datasets generated and analyzed during the current study are not publicly available due to all the results in the form of means and statistics are presented in this paper, but are available from the corresponding author on reasonable request.

Declarations

Competing interest The authors declare no conflicts of interest associated with this study or its results.

Consent to participate All authors participated and helped voluntarily in the research.

Consent for publication All authors read and approved the final manuscript.

REFERENCES

- Akgün, S., Ertel, N.H., Mosenthal, A., Oser, W., 1998. Postsurgical reduction of serum lipoproteins: Interleukin-6 and the acute-phase response. *The Journal of Laboratory and Clinical Medicine* 131, 103-108.
- Akyüz, E., 2020. Kars bölgesindeki neonatal buzağlarda sıklıkla karşılaşılan ishal kaynaklı sepsis nedenleri. In: *'Tüm Yönleri ile Kuzey Doğu Anadolu Bölgesinde Hayvancılık'*, (Ed.): Ayvazoğlu Demir, P. İksad Yayınevi, Ankara, Türkiye, 401-423.
- Akyüz, E., Kükürt, A., 2021. Evaluation of oxidative stress index and some biochemical parameters in neonatal calves with diarrhea. *Acta Scientiae Veterinariae* 3, 58-63.
- Akyüz, E., Makav, M., Kuru, M., Bayram, P., 2021. The effect of tarantula cubensis extract on gentamicin-induced acute kidney injury in ovariectomized rats. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi* 27, 559-566.
- Akyüz, E., Coşkun, A., Şen, İ., 2016. Deneysel endotoksemi oluşturulan buzağlarda sıvı tedavisinin hemodinamik parametreler üzerine etkisi. *Eurasian Journal of Veterinary Sciences* 32, 246-254.
- Akyüz, E., Gökçe, G., 2021. Neopterin, procalcitonin, clinical biochemistry, and hematology in calves with neonatal sepsis. *Tropical Animal Health and Production* 53, 354.
- Aldridge, B.M., Garry, F.B., Adams, R., 1993. Neonatal septicemia in calves: 25 cases (1985-1990). *Journal of the American Veterinary Medical Association* 203, 1324-1329.
- Asadi, A.H., Baghinezhad, M., Asadi, H., 2015. Neonatal calf diarrhea induced by rotavirus and coronavirus: a review. *International Journal of Biosciences* 6, 230-236.
- Atakisi, E., Bozukluhan, K., Atakisi, O., Gökçe, H.I., 2010. Total oxidant and antioxidant capacities and nitric oxide levels in cattle with traumatic reticuloperitonitis. *Veterinary Record* 167, 908-909.
- Bachmann, L.M., Juni, P., Reichenbach, S., Ziswiler, H.R., Kessels, A.G., Vogelin, E., 2005. Consequences of different diagnostic "gold standards" in test accuracy research: Carpal tunnel syndrome as an example. *International journal of epidemiology* 34, 953-955.
- Başer, D.F., Civelek, T., 2013. Akut ishallerde neonatal buzağlarda venöz asit baz durumu ve renal fonksiyon arası korelasyon. *Kocatepe Veteriner Journal* 6, 25-31.
- Basoglu, A., Sen, I., Meoni, G., Tenori, L., Naseri, A., 2018. NMR-Based plasma metabolomics at set intervals in newborn dairy calves with severe sepsis. *Mediators of Inflammation* 2018, 1-10.
- Beutler, E., Duron, O., Kelly, B., 1963. Improved method for the determination of blood glutathione. *The Journal of Laboratory and Clinical Medicine* 61, 882-888.
- Beydilli, Y., Gökçe, H., 2019. Sepsisli neonatal buzağlarda bazı hematolojik ve biyokimyasal parametrelerin araştırılması. *Mehmet Akif Ersoy Üniversitesi Sağlık Bilimleri Enstitüsü Dergisi* 7, 55-67.
- Bonelli, F., Meucci, V., Divers, T.J., Boccardo, A., Pravettoni, D., Meylan, M., Belloli, A.G., Sgorbini, M., 2018. Plasma procalcitonin concentration in healthy calves and those with septic systemic inflammatory response syndrome. *Veterinary Journal* 234, 61-65.
- Bozukluhan, K., Merhan, O., Gökçe, H.I., Devenci, H.A., Gökçe, G., Ogun, M., Marasli, S., 2017. Alterations in lipid profile in neonatal calves affected by diarrhea. *Veterinary world* 10, 786-789.
- Bozukluhan, K., Merhan, O., Gökçe, H.I., Ögün, M., Atakışı, E., Kiziltepe, Ş., Gökçe, G., 2018. Determination of some acute phase proteins, biochemical parameters and oxidative stress in sheep with naturally infected Sheeppox virus. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi* 24, 437-441.
- Bozukluhan, K., Merhan, O., Kiziltepe, Ş., Egritag, H.E., Akyuz, E., Gökçe, H.I., 2021. Determination of haptoglobin, some biochemical and oxidative stress parameters in calves with pneumonia. *Fresenius Environmental Bulletin* 30, 9492-9496.
- Carpentier, Y.A., Scruel, O., 2002. Changes in the concentration and composition of plasma lipoproteins during the acute phase response. *Current Opinion in Clinical Nutrition & Metabolic Care* 5, 153-158.
- Ceron, J.J., Eckersall, P.D., Martınez-Subiela, S., 2005. Acute phase proteins in dogs and cats: current knowledge and future perspectives. *Veterinary Clinical Pathology* 34, 85-99.
- Cerone, S.I., Sansinanea, A.S., Streitenberger, S.A., Garcia, M.C., Auza, N.J., 2000. Cytochrome C oxidase, Cu, Zn-superoxide dismutase, and ceruloplasmin activities in copper-deficient bovines. *Biological trace element research* 73, 269-278.
- Chatre, L., Verdonk, F., Rocheteau, P., Crochemore, C., Chrétien, F., Ricchetti, M., 2017. A novel paradigm links mitochondrial dysfunction with muscle stem cell impairment in sepsis. *Biochimica et Biophysica Acta - Molecular Basis of Disease* 1863, 2546-2553.
- Çitil, M., 2003. Puerperal enfeksiyonlu ve abomasum deplasmanlı ineklerde serum amiloid A ve haptoglobin düzeyleri. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi* 9, 147-51.
- Çitil, M., Gökçe, E., 2013. Neonatal sepsismi. *Türkiye Klinikleri Journal of Veterinary Sciences* 4, 62-70.
- Colombo, J.P., Richterich, R., 1964. Zur bestimmung des caeruloplasmin im plasma [on the determination of ceruloplasmin in plasma]. *Schweizerische Medizinische Wochenschrift* 23, 715-720.
- Coşkun, A., Şen, İ., 2011. Sığırlarda akut faz proteinleri ve klinik kullanılm alanları. *Sağlık Bilimleri Dergisi* 20, 240-246.
- Cray, C., 2012. Acute phase proteins in animals. *Progress in Molecular Biology and Translational Science* 105, 113-150.
- Dellinger, R.P., Levy, M.M., Rhodes, A., Annane, D., Gerlach, H., Opal, S.M., Sevransky, J.E., Sprung, C.L., Douglas, I.S., Jaeschke, R., Osborn, T.M., 2013. Surviving sepsis campaign: International guidelines for management of severe sepsis and septic shock. *Critical Care Medicine* 41, 580-637.
- Dratwa-Chalupnik, A., Herosimczyk, A., Lepczyński, A., Skrzypczak, W.F., 2012. Calves with diarrhea and a water-electrolyte balance. *Medycyna weterynaryjna* 68, 5-8.
- Dupuy, A.M., Philippart, F., Pean, Y., Lasocki, S., Charles, P.E., Chalumeau, M., Claessens, Y.E., Quenot, J.P., Gras-Le Guen, C., Ruiz, S., Luyt, C.E., Roche, N., Stahl, J.P., Bedos, J.P., Pugin, J., Gauzit, R., Misset, B., Brun-Buisson, C., 2013. Role of biomarkers in the

- management of antibiotic therapy: an expert panel review: I - currently available biomarkers for clinical use in acute infections. *Annals of Intensive Care* 3, 22.
- Ercan, N., Tuzcu, N., Başbug, O., Tuzcu, M., Alim, A., 2016. Diagnostic value of serum procalcitonin, neopterin, and gamma interferon in neonatal calves with septicemic colibacillosis. *Journal of Veterinary Diagnostic Investigation* 28, 180-183.
- Erkılıç, E.E., Merhan, O., Kırmızıgül, A.H., Ögün, M., Akyüz, E., Çitil, M., 2019. Salivary and serum levels of serum amyloid A, haptoglobin, seruloplasmin and albumin in neonatal calves with diarrhoea. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi* 25, 583-586.
- Erkılıç, E.E., 2019. Babesiosis'li köpeklerde tedavi öncesi ve sonrası haptoglobin, seruloplazmin ve bazı biyokimyasal parametrelerin seviyelerinin belirlenmesi. *Harran Üniversitesi Veteriner Fakültesi Dergisi* 8, 77-80.
- Fecteau, G., Paré, J., Van, Metre, D.C., Smith, B.P., Holmberg, C.A., Guterbock, W., Jang, S., 1997. Use of a clinical sepsis score for predicting bacteremia in neonatal dairy calves on a calf rearing farm. *Van Veterinary Journal* 38, 101-104.
- Fecteau, G., Smith, P.B., George, L.W., 2009. Septicemia and meningitis in newborn calf. *Veterinary Clinics of North America - Food Animal Practice* 25, 195-208.
- Franck, S.M., Brad, T.B., Harley, W.M., 1998. Multiplex PCR for enterotoxigenic, attaching and effacing, and shiga toxin-producing *Escherichia coli* strains from calves. *Journal of Clinical Microbiology* 36, 1795-1797.
- Gruys, E., Toussaint, M.J.M., Niewald, T.A., 2005. Acute phase reaction and acute phase proteins. *Journal of Zhejiang University: Science* 11, 1045-1056.
- Heegaard, P.M.H., Godson, D.L., Toussaint, M.J.M., Tjørnehoj, K., Larsen, L.E., Viuff, B., Ronsholt, L., 2000. The acute phase response of haptoglobin and serum amyloid A (SAA) in cattle undergoing experimental infection with respiratory syncytial virus. *Veterinary Immunology and Immunopathology* 77, 151-159.
- House, J.K., Smith, G.W., McQuirk, S.M., Gunn, A.A., Izzo, M., 2015. Manifestations and management of disease in neonatal ruminants. In: *'Large Animal Internal Medicine'*, (Ed): Smith BP, 5th edition, Elsevier Saunders, St. Louis, U.S.A., 302-338.
- Kao, S.J., Yang, H.W., Tsao, S.M., Cheng, C.W., Bien, M.Y., Yu, M.C., Bai, K.J., Yang, S.F., Chien, M.H., 2013. Plasma long pentraxin 3 (PTX3) concentration is a novel marker of disease activity in patients with community-acquired pneumonia. *Clinical Chemistry and Laboratory Medicine* 51, 907-913.
- Köse, S.İ., Maden, M., 2013. Biyobelirteçler ve klinik kullanımları. *Dicle Üniversitesi Veteriner Fakültesi Dergisi* 2, 1-8.
- Libby, P., Ridker, P.M., Hansson, G.K., 2009. Inflammation in atherosclerosis: from pathophysiology to practice. *Journal of the American College of Cardiology* 54, 2129-38.
- Lorenz, I., Fagan, J., More, S.J., 2011. Calf health from birth to weaning. II. management of diarrhoea in pre-weaned calves. *Irish veterinary journal* 64, 9.
- Marcato, F., Van den Brand, H., Kemp, B., van Reenen, K., 2018. Evaluating potential biomarkers of health and performance in veal calves. *Frontiers in Veterinary Science* 5, 133.
- McGrotty, Y.L., Knottenbelt, C.M., Ramsey, I.K., Reid, S.W.J., Eckersall, P.D., 2003. Haptoglobin concentrations in a canine hospital population. *Veterinary Record* 152, 562-564.
- Meisner, M., 2005. Biomarkers of sepsis: clinically useful? *Current Opinion in Critical Care* 11, 473-480.
- Merhan, O., Bozukluhan, K., Gökçe, G., Yılmaz, O., 2016. İshalli buzağılarda haptoglobin, seruloplazmin ve bazı biyokimyasal parametre düzeylerinin araştırılması. *Fırat Üniversitesi Sağlık Bilimleri Veteriner Dergisi* 30, 195-198.
- Merhan, O., Bozukluhan, K., Gökçe, H.I., 2017. Acute phase proteins and biochemical and oxidative stress parameters in Hypoderma spp. infested cattle. *The Journal of the Hellenic Veterinary Medical Society* 68, 535-540.
- Moore, R.E., Kirwan, J., Doherty, M.K., Whitfield, P.D., 2007. Biomarker discovery in animal health and disease: The application of post-genomic technologies. *Biomarker Insights* 2, 185-196.
- Murata, H., Shimada, N., Yoshioka, M., 2004. Current research on acute phase proteins in veterinary diagnosis: an overview. *Veterinary Journal* 168, 28-40.
- Naseri, A., 2017. Echocardiographic assesment of left ventricular systolic function in calves with naturally occurring severe sepsis and septic shock and changes in these functions related to applied treatment; longitudinal study. Selçuk University Health Sciences Institute, Konya, Türkiye.
- Novelli, G., Ferretti, G., Poli, L., Pretagostini, R., Ruberto, F., Perrella, S.M., Levi, S., Morabito, V., Berloco, P.B., 2010. Clinical results of treatment of postsurgical endotoxin-mediated sepsis with polymyxin-b direct hemoperfusion. *Transplantation Proceedings* 42, 1021-1024.
- Ok, M., Güler, L., Turgut, K., Ok, U., Sen, I., Gündüz, I.K., 2009. The studies on the etiology of diarrhoea in neonatal calves and determination of virulence gene markers of *Escherichia coli* strains by multiplex PCR. *Zoonoses Public Health* 56, 94-101.
- Ok, M., Yıldız, R., Hatipoğlu, F., Baspınar, N., Ider, M., Uney, K., Ertürk, A., Durgut, M., Terzi, F., 2020. Use of intestine-related biomarkers for detecting intestinal epithelial damage in neonatal calves with diarrhoea. *American journal of veterinary research* 81, 139-146.
- Pardon, B., Deprez, P., 2018. Rational antimicrobial therapy for sepsis in cattle in face of the new legislation on critically important antimicrobials. *Vlaams Diergeneeskundig Tijdschrift* 87, 37-46.
- Parlakpınar, H., Örum, M.H., Acet, A., 2013. İlaça bağlı nefrotoksisitede serbest oksijen radikalleri. *Fırat Üniversitesi Sağlık Bilimleri Tıp Dergisi* 27, 51-56.
- Petersen, H.H., Nielsen, J.P., Heegard, P.M.H., 2004. Application of acute phase protein measurements in veterinary clinical chemistry. *Veterinary Research* 35, 163-187.
- Pourjafar, M., Badiei, K., Nazifi, S., Naghib, S.M., 2011. Acute phase response in holstein dairy calves affected with diarrhoea. *Bhikhabhai Jivabhai Vanijya Mahavidyalaya* 14, 142-149.
- Pradeep, M., 2014. Application of acute phase proteins as biomarkers in modern veterinary practice. *Indian Journal of Veterinary and Animal Sciences Research* 43, 1-13.
- Radostits, O.M., Gay, C.C., Hinchcliff, K.W., Constable, P.D., 2006. *Veterinary Medicine, A textbook of the diseases of cattle, sheep, goats, pigs and horses*, 10th edition, Saunders Co, London.
- Roosendaal, B., Wim, G., De Graaf, F.K., 1984. The nucleotide sequence of the gene encoding the K99 subunit of enterotoxigenic *Escherichia coli*. *FEMS Microbiology Letters* 22, 253-258.
- Seifi, H.A., Mohri, M., Shoorei, E., Farzaneh, N., 2006. Using haematological and serum biochemical findings as prognostic indicators in calf diarrhoea. *Comparative Clinical Pathology* 15, 143-147.
- Sen, I., Constable, P.D., 2013. General overview to treatment of strong ion (metabolic) acidosis in neonatal calves with diarrhoea. *Eurasian Journal of Veterinary Sciences* 29, 114-120.
- Shah, R., 2007. Endothelins in health and disease. *European Journal of Internal Medicine* 18, 272-82.
- Skinner, J.G., Brown, R.A., Roberts, L., 1991. Bovine haptoglobin response in clinically defined field conditions. *Veterinary Record* 128, 147-149.
- Sönmez, Ç.M., Tülek, N., 2015. Biomarkers in bacterial infections and sepsis. *Klinik Dergisi* 28, 96-102.
- Tokoyuni, S., 1999. Reactive oxygen species-induced molecular damage and its application in pathology. *Pathology International* 49, 91-102.
- Trefz, F.M., Lorch, A., Feist, M., Sauter-Louis, C., Lorenz, I., 2013. The prevalence and clinical relevance of hyperkalaemia in calves with neonatal diarrhoea. *Veterinary Record* 195, 350-356.
- Tschaikowsky, K., Sagner, S., Lehnert, N., Kaul, M., Ritter, J., 2000. Endothelin in septic patients: effects on cardiovascular and renal function and its relationship to proinflammatory cytokines. *Critical Care Medicine* 28, 1854-1860.
- Tuna, G.E., Ulutaş, B., 2015. Hastalıkların biyobelirteçleri olarak akut faz proteinleri. *Türkiye Klinikleri Veteriner Sciences- Internal Medicine Special Topics* 1, 8-19.
- Uzlu, E., Karapehlivan, M., Çitil, M., Özçelik, E., Erdoğan, H.M., 2010. İshal semptomu belirlenen buzağılarda serum sialik asit ile bazı biyokimyasal parametrelerin araştırılması. *Yüzüncü Yıl Üniversitesi Veteriner Fakültesi Dergisi* 21, 83-86.
- Yıldız, R., Beslek, M., Beydilli, Y., Özçelik, M.M., Biçici, Ö., 2018. Evaluation of platelet activating factor in neonatal calves with sepsis. *Veteriner Hekimler Derneği Dergisi* 89, 66-73.
- Yoshioka, T., Kawada, K., Shimada, T., Mori, M., 1979. Lipid peroxidation in maternal and cord blood and protective mechanisms. *American Journal of Obstetrics and Gynecology* 135, 972-976.