

Journal of the Hellenic Veterinary Medical Society

Vol 75, No 4 (2024)



Investigation of the relationship between apelin hormone response and some physiological parameters in enzootic bovine leukosis virus infected cattle

A Sait, B Bayraktar, S Korkmaz, M Genç, S Bayraktar, GC Kotan, Ş Aydemir, Ç Takma, AS Bülbül, V Aksakal, E Tekce, GB Büyük Özcan, H Kaya, M Odabaşı

doi: [10.12681/jhvms.34664](https://doi.org/10.12681/jhvms.34664)

Copyright © 2025, A Sait, B Bayraktar, S Korkmaz, M Genç, S Bayraktar, GC Kotan, Ş Aydemir, Ç Takma, AS Bülbül, V Aksakal, E Tekce, GB Büyük Özcan, H Kaya, M Odabaşı



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

To cite this article:

Sait, A., Bayraktar, B., Korkmaz, S., Genç, M., Bayraktar, S., Kotan, G., Aydemir, Ş, Takma, Ç, Bülbül, A., Aksakal, V., Tekce, E., Büyük Özcan, G., Kaya, H., & Odabaşı, M. (2025). Investigation of the relationship between apelin hormone response and some physiological parameters in enzootic bovine leukosis virus infected cattle . *Journal of the Hellenic Veterinary Medical Society*, 75(4), 8191–8196. <https://doi.org/10.12681/jhvms.34664>

Investigation of the relationship between apelin hormone response and some physiological parameters in enzootic bovine leukosis virus infected cattle

A. Sait^{1*}, B. Bayraktar², S. Korkmaz¹, M. Genç³, S. Bayraktar⁴, G.C. Kotan⁵,
Ş. Aydemir⁵, Ç. Takma⁶, A.S. Bülbül⁷, V. Aksakal⁷, E. Tekce⁷, G.B. Büyük Özcan⁸,
H. Kaya⁹, M. Odabaşı¹⁰

¹ Istanbul Pendik Veterinary Control Institute, Virology Laboratory, İstanbul, Türkiye

² Faculty of Health Sciences, Bayburt University, Bayburt, Türkiye

³ Faculty of Veterinary Medicine, Atatürk University, Erzurum, Türkiye

⁴ Faculty of Veterinary Medicine, Ondokuz Mayıs University, Samsun, Türkiye

⁵ Alaca Vocational School, Hitit University, Çorum, Türkiye

⁶ Ege University, Faculty of Agriculture, Department of Biometry and Genetics, İzmir, Türkiye

⁷ Faculty of Applied Sciences, Bayburt University,

⁸ Ankara Medipol University, Faculty of Medicine, Department of Physiology, Ankara, Türkiye

⁹ Kelkit Vocational School, Gümüşhane University, Gümüşhane, Türkiye

¹⁰ Istanbul Food Control Laboratory Directorate, İstanbul, Türkiye

ABSTRACT: Apelin is a hormone with lymphangiogenesis potential, associated with tumor growth and lymph node metastasis, and is reported to stimulate tumor vascularization. This study aimed to investigate the response of the apelin hormone in cattle with and without Enzootic Bovine Leukosis (EBL) infection, based on physiological parameters such as breed, age, and body condition score (BCS). A total of 120 Simmental and Montafon cattle of varying ages (<2, 2-8, and >8) and BCS categories (<2, 3-3.5, and ≥4) were included in the study. Serum apelin hormone levels were measured using the ELISA technique in blood samples collected from the neck vein (*Vena jugularis*) of the animals. The analysis of mean serum apelin (ng/mL) values revealed that, among EBL-positive cattle, the highest value was 2.77 ng/mL in Simmental cattle aged 2-8 years with a BCS of 3-3.5, while a value of 2.69 ng/mL was observed in Montafon cattle aged >8 years with a BCS of 3-3.5. For EBL-negative cattle, the highest mean serum apelin value was 2.24 ng/mL, recorded in Simmental cattle aged 2-8 years with a BCS of 3-3.5, whereas it was 1.45 ng/mL in Montafon cattle aged >8 years with the same BCS. In conclusion, this study demonstrated for the first time the effect of apelin hormone levels in cattle with or without EBL, depending on physiological parameters such as age, breed, and BCS. It is suggested that the apelin hormone, which is involved in neoplastic lymphocyte cell infiltration in the lymphoreticular tissues of cattle, and plays a role in regulating tumor growth in EBL, a malignant tumor disease, may contribute to eradication programs. Moreover, the evaluated parameters may prove valuable for disease prevention and control strategies.

Keywords: Enzootic Bovine Leukosis; apelin; bovine; physiological parameters

Corresponding Author:

A. Sait, Istanbul Pendik Veterinary Control Institute, Viral Diagnostics Laboratory,
İstanbul, Türkiye
E-mail address: ahmet.sait@tarimorman.gov.tr

Date of initial submission: 6-8-2023

Date of acceptance: 29-8-2024

INTRODUCTION

Enzootic bovine leukosis (EBL) is a disease caused by the bovine leukemia virus (BLV), an oncovirus belonging to the Retroviridae family that affects cattle, leading to significant economic losses worldwide (Camargos et al., 2002). BLV, also known as bovine lymphosarcoma, primarily targets the bone marrow, skin, lymph node, blood cells (lymphocytes) and thymus, causing lymphocyte proliferation and tumoral formations (Murakami et al., 2013). In animals with EBL, there is a decline in body condition score, live weight, and production parameters (Murtaugh et al., 1991). Clinical signs in cattle include fever, anorexia, weight loss, increased heart rate, elevated respiratory frequency, swelling of all superficial lymph nodes, and firm tumoral formations in the prescapular, mandibular, and inguinal lymph nodes. BLV infection is challenging to control, as most infected cattle (approximately 70%) are asymptomatic or subclinically infected (Juliarena et al., 2017). Permanent lymphocytosis occurs in 30% of BLV-infected cattle, while malignant B-cell lymphosarcoma tumors may develop after a prolonged latent form (1-8 years) in 1-5% of cases (Aida et al., 2013; Pandey et al., 2017). The EBL virus is reportedly absent from semen and is not considered a transmission route for artificial insemination (Monkey, 1986). However, semen may act as a transmission medium if contaminated with blood during collection via massage technique (Lucas et al., 1980). Age plays a crucial role in the seroprevalence of the disease, with BLV commonly occurring between 5 and 8 years of age. The seroprevalence rate increases with age, as animals face prolonged exposure to infected cattle (Maezawa et al., 1998).

Apelin is a hormone secreted from adipose tissue, derived from a 77 amino-acids precursor (Tatemoto et al., 2001). Among its isoforms, apelin-13 exhibits biological activity that is 8 times stronger than apelin-17 and 60 times stronger than apelin-36 (Tatemoto et al., 1998). The Apelin/APJ system plays a crucial role in various physiological processes, including the endocrine stress response (Taheri et al., 2002), cardiovascular functions (Szokodi et al., 2002), regulation of blood pressure (Tatemoto et al., 2001), angiogenesis (Zhang et al., 2016), thermoregulation (Reaux et al., 2001), and energy metabolism regulation (Bertrand et al., 2015). Apelin and APJ are expressed in endothelial cells of newly developing blood vessels during angiogenesis (Kidoya et al., 2008). Apelin promotes tumor growth by stimulating tumor vascularization, which has lymphangiogenic potential and is associat-

ed with tumor progression and lymph node metastasis (Sorli et al., 2006; Sorli et al., 2007). Recognized as a therapeutic agent, apelin plays a significant role in tumor initiation and progression, making it a potential target for the treatment of tumors and cancers (Masoumi et al., 2020).

The findings of this study are expected to contribute to research on Enzootic bovine leukosis (EBL), a disease for which no treatment or vaccine currently exists. Moreover, no existing studies have examined the apelin hormone response in cattle with EBL concerning breed, age and body condition score. The current study aimed to investigate the relationship between EBL infection and the apelin hormone response in cattle, alongside some physiological parameters.

MATERIALS AND METHOD

Selection of the animals and sampling

This study was approved by the Veterinary Control Institute Animal Care and Use Committee Ethics Committee (Decision No: 08/2022). The study was carried out in cattle farms located in the Marmara region of Türkiye, which were raised under intensive conditions and whose records were followed regularly. To categorize the groups based on body condition score (BCS), palpation of the waist region was performed as described by Khan (1993). In accordance with the principles outlined by Thompson and Meyer (2006) scoring was conducted based on muscle and fat accumulation in the waist region, as follows: BCS <2 (thin), BCS 3-3.5 (normal), and BCS ≥4 (obese). The animal material for the study consisted 120 Simmental and Montafon cattle of varying ages (<2, 2-8, and >8) and body condition scores (≤2, 3-3.5 and ≥4). Those with questionable ELISA test results, age information, and body condition scores were not included in the study. Detailed information on the animal material and sample numbers is presented in Table 1.

During the research period, care was taken to ensure that the environment and feed factor (Table 2) remained consistent. The feed content in this study was analyzed using standard AOAC methods.

Collection of serum samples:

A 5 ml blood sample was collected from the neck vein (*vena jugularis*) of the cattle in the study groups into tubes without anticoagulant (VACUETTE® TUBE 9 ml Z Serum Clot Activator). After centrifuging the blood samples at 3000 rpm for 10 minutes in a refrigerated centrifuge (NF 1200R, NUVE, Ankara,

Table 1. Number of cattle included in the study across different ages and body condition scores

BCS	Age																	
	<2						2-8						>8					
	Montafon			Simmental			Montafon			Simmental			Montafon			Simmental		
	Neg	Pos	Total	Neg	Pos	Total	Neg	Pos	Total	Neg	Pos	Total	Neg	Pos	Total	Neg	Pos	Total
≤2	3	5	8	2	3	5	3	5	8	2	3	5	3	5	8	2	4	6
3-3.5	3	4	7	5	3	8	3	4	7	4	6	10	3	4	7	3	5	8
≥4	3	2	5	4	3	7	2	3	5	3	2	5	2	3	5	3	3	6
Total	9	11	20	11	9	20	8	12	20	9	11	20	8	12	20	8	12	20

*BCS: Body Condition Score, Neg: Negative, Pos: Positive

Table 2. Nutrient composition of feed used in the study (%)

Ingredient	% DM
Alfalfa haylage	31,35
Corn Silage	26,7
Soybean meal (30% CP)	15,2
Barley Straw	26,3
Mineral and vitamin mixture*	0,15
Sodium bicarbonate	0,3
Nutrient composition	
CP %	12,3
NDF	39,95
ADF	26,98
NFC	33,12
Ca	0,72
P	0,46
Energy (Mcal/kg)	1,41

*1 kg vit.-min. Premix contains vitamin A, 7,000,000 IU; vitamin D3, 1,000,000; vitamin E, 30,000; Mn, 50,000 mg; Zn, 50,000 mg; Fe, 50,000; Cu, 10,000 mg; I, 8,000 mg; Co, 200 mg; Se, 150 mg; and Mg, 100 mg

TÜRKİYE) in the laboratory, the obtained blood serums were transferred to sterile tubes and stored at -80°C until analysis.

Antibody ELISA test

The Enzootic Bovine Leukosis Virus (BLV) Antibody ELISA Test Kit (IDEXX, Lot SN 19090, France) was used to detect BLV antibodies in the serum samples obtained in this study. The test was performed following the manufacturer's instructions, and the optical density (OD) values were measured using an ELISA reader at a wavelength of 450 nm. According to the test procedure, the degree of positivity of each serum was determined, and any sample with a result equal to or greater than a plus (+) value was considered positive.

Measurement of apelin hormone levels in serum:

The minimum detectable concentration of the apelin hormone kit used for measuring of apelin levels in the blood serums obtained in this study was reported as 16pg/ml. The race-specific Bovine Apelin (AP)

ELISA Kit (Apelin, SinoGeneClon, Product code: SG-60825, CHINA) was used in accordance with the procedure described in the manufacturer's catalog, with a detection limit of < 35 pg/ml, an intra-assay coefficient of 8.0% and an inter-assay coefficient of 10.0%.

Statistical analysis:

In this study, apelin hormone levels in the serum of Simmental and Montafon cattle breeds during lactation and pregnancy were analyzed as the dependent variable; while gender, body score, and breed were analyzed as fixed effects. To investigate the effect of gender, body score and breed on apelin hormone levels in serum during lactation and pregnancy, a normal distribution test was first applied to the apelin measurements. Logarithmic transformation was applied to the data as a result of which the Shapiro-Wilk test turned out to be significant. It was determined by the Levene test ($P=0.33$), which showed a normal distribution, and the group variances were homogeneous. Then, analyses were performed according to the experiment

pattern of coincidence plots in factorial order (2x3x3) using a single-variable procedure in linear models generalized to logarithmic apelin measurements, and the interaction effects of body score and breed, gender and body score, race and body score, gender and breed factors were also investigated. Tukey multiple comparison tests was applied to compare the mean differences of Apelin hormone. IBM SPSS v25 statistical package program was used for all analyses in this study. All significant differences were evaluated by testing at the $P < 0.05$ level.

RESULTS

In the current study, cattle were categorized based on body condition scores (BCS) and age groups: <2 years, 2-8 years, and >8 years. BCS classifications were defined as low (≤ 2), optimal (3-3.5), and high (≥ 4). A total of 120 female cattle were included in the study, with 20 cattle in each age group (<2 years, 2-8

years, and >8 years) for each BCS category. The cattle were also evaluated for Enzootic Bovine Leukosis (EBL) infection status, as detailed in Table 3. As a result of the analyzes, when the mean serum Apelin (ng/mL) values for the BCS*EBL*Age*Race interaction were examined, the highest values for EBL positive cattle was 2.77 ng/mL in Simmental breed aged 2-8 years with BCS 3- 3.5, while it was 2.69 ng/mL in Montafon breed with BCS 3-3.5 and aged >8 years. For EBL negative cattle, the highest mean serum apelin value was 2.24 ng/mL in Simmental breed aged 2-8 years with BCS 3-3.5, and 1.45 ng/mL in Montafon breed aged >8 years with BCS 3-3.5. When analyzed for the BCS*EBL*Age* Breed interaction, the highest apelin hormone levels were observed in EBL positive Montafon and Simmental cattle aged ≥ 8 years, regardless of BCS, and in Simmental cattle aged 2-8 years with BCS 3-3.5 and EBL-positive status ($p < 0.01$) (Table 4).

Table 3. The comparisons of apelin (ng/mL) levels based on physiological parameters and EBL status

Parameters		n	Apelin	±	SEM
Race	Montofon	60	1,479b	±	0,05
	Simental	60	1,925a	±	0,05
BCS	≤ 2	40	1,551b	±	0,062
	3-3.5	47	1,863a	±	0,055
	≥ 4	33	1,692ab	±	0,066
Age	< 2	40	1,449c	±	0,061
	2-8	40	1,625b	±	0,062
	≥ 8	40	2,031a	±	0,061
EBL +/-	Positive	67	2,111a	±	0,047
	Negative	53	1,292b	±	0,053

Source of variation (P-values)

Race	0,003
BCS	<0,01
Age	<0,01
EBL	<0,01

Table 4. Interaction of apelin hormone level (ng/mL) according to enzootic bovine leukosis, age, races and body condition score.

BCS	EBL+						EBL-					
	Age						Age					
	<2		2-8		>8		<2		2-8		>8	
	Montafon	Simmental	Montafon	Simmental	Montafon	Simmental	Montafon	Simmental	Montafon	Simmental	Montafon	Simmental
≤ 2	1,71±0,165jmr	1,713±0,214r	2,132±0,165jn	1,31±0,214dm	2,232±0,165hj	2,433±0,185h	0,97±0,214cefg	0,590±0,262adf	0,847±0,214ce	1,785±0,262ad	0,99±0,214c	1,895±0,262a
3-3.5	2,218±0,185jmr	2,19±0,214fr	2,043±0,214ejn	2,773±0,151hk	2,693±0,185hj	2,626±0,165h	0,643±0,214eg	1,674±0,165adf	0,803±0,214be	1,385±0,185ad	1,453±0,185ab	1,86±0,214a
≥ 4	1,72±0,262rt	1,81±0,214fr	0,567±0,214p	2,695±0,262dhk	2,433±0,185hj	2,707±0,214h	1,007±0,214befg	1,142±0,185df	0,92±0,262be	2,243±0,214ad	1,24±0,262abc	1,817±0,214a

Source of variation (P-values)

BCS*Age*Race	<0,01
BCS*EBL*Race	0,016
BCS*EBL*Age	0,029
Age*Race*EBL	0,024
BCS*EBL*Age*Race	<0,01

DISCUSSION

Enzootic Bovine Leukosis is a malignant tumoral disease characterized by neoplastic lymphocyte infiltrations in the lymphoreticular tissues of various organs and systems. The clinical symptoms very depending on the organs where the lesions are located (Barlet et al., 2020). The disease manifests in five forms: enzootic viral leukosis in adult cattle, sporadic juvenile leukosis in calves under six 6 months old, sporadic thymic leukosis in young cattle under two years old, sporadic skin leukosis in young cattle, and benign lymphocytosis (Donnik et al., 2021). EBL-infected animals are persistently infected and spread the disease without displaying clinical symptoms (Asfaw et al., 2005). The absence of a vaccine or treatment makes the disease one of the major livestock problems globally. Pregnancy and lactation involve physiological and hormonal changes that can influence disease transmission. The disease can be transmitted to the fetus via blood, and to offspring through infected milk and colostrum after birth. The secretion of the apelin hormone, produced by adipose tissue, varies with physiological conditions such as body mass index, pregnancy, and lactation, resulting in fluctuating levels in circulation (Hughes & Watson, 2018; Lérias et al., 2014). Apelin is a hormone with a well-defined role in tumor progression and is easily measurable in blood serum (Zhang et al., 2016). Apelin/APJ system has been reported to inhibit tumor growth and regulate tumor formation in an autocrine manner through its role in tumor angiogenesis. Endothelial

cell-derived Apelin is also known to regulate tumor growth via the Apelin/APJ axis (Hu et al., 2021). It was determined that BCS*EBL*Age*Breed was effective on the mean serum apelin hormone level in cattle with positive EBL disease with different body condition scores, which is the target of our current study ($p<0.05$). In addition, the results of our current research are similar to the results of studies in humans and different animals in which serum apelin levels are examined in tumor patients (Picalt et al., 2014; Feng et al., 2016; Podgórska et al., 2019).

CONCLUSION

In conclusion, we believe that the apelin hormone may serve a useful biomarker for evaluating tumor progression in cattle infected with EBL, a disease characterized by tumoral formations in lymph nodes. However, further and more comprehensive studies are necessary.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

Authors' Contribution Statements: Design and idea stage of the article AS, BB, Apelin analysis part BB, SB, GCK, GBÖ, MO; in statistical analysis; SK, ET, VA, HK; In the fieldwork, AS, BB, SB, SK, ŞA, GCK, ASB, VA has taken part. All authors read and approved the final manuscript.

REFERENCES

- Aida, Y., H. Murakami, M. Takagashi and S.N. Takeshima.2013. Mechanisms of pathogenesis induced by bovine leukemia virus as a model for human T-cell leukemia virus. *Front. Microbiol.* 4:328-339.
- Asfaw, Y., Tsuduku, S., Konishi, M., Murakami, K., Tsuboi, T., Wu, D. and H. Sentsui. 2005. Distribution and superinfection of bovine leukemia virus genotypes in Japan. *Arch. Viro.* 150: 493-505.
- Bartlett, P. C., Ruggiero, V. J., Hutchinson, H. C., Droscha, C. J., Norby, B., Sporer, K. R and T.M. Taxis. 2020. Current developments in the epidemiology and control of enzootic bovine leukosis as caused by bovine leukemia virus. *Pathogens*.9(12):1058.
- Bertrand, C., P. Valet and I. Castan-Laurell. 2015. Apelin and energy metabolism. *Frontiers in physiology.* 6: 115-120.
- Blackburn, E. H. 1991. Structure and function of telomeres. *Nature.* 350(6319): 569-573.
- Buzala, E. and W. Dereń. 2003. Comparison of PLA with AGID and ELISA results in serology diagnosis of bovine leukosis. *Polish Journal of Veterinary Sciences*, 6: 9-11.
- Camargos, M. F., D. Stancek, M.A. Rocha, L.M. Lessa, J.K.P. Reis and R.C. Leite. 2002. Partial sequencing of env gene of bovine leukaemia virus from Brazilian samples and phylogenetic analysis. *Journal of Veterinary Medicine, Series B.* 49(7): 325-331.
- Diakowska, D., K. Markocka-Mączka, M. Nienartowicz, J. Rosińczuk and M. Krzystek-Korpacka. 2019. Assessment of apelin, apelin receptor, resistin, and adiponectin levels in the primary tumor and serum of patients with esophageal squamous cell carcinoma. *Advances in Clinical and Experimental Medicine.* 28(5): 671-678.
- Donnik, I. M., Gulyukin, M. I., Busol, V. A., Kovalenko, L. V. A. M. Kovalenko. 2021.Enzootic Bovine Leukosis—Diagnostics, Eradication, And Anthroponotic Potential *Biol. Ag.* 56 (2):230-244.
- Feng, M., G. Yao, H. Yu, Y. Qing and K. Wang.2016. Tumor apelin, not serum apelin, is associated with the clinical features and prognosis of gastric cancer. *BMC Cancer.* 16(1): 1-8.
- Hu, L., Y. Hayashi, H. Kidoya and N. Takakura. 2021. Endothelial cell-derived Apelin inhibits tumor growth by altering immune cell localization. *Scientific Reports*, 11(1), 1-12.
- Hughes, K. And C.J. Watson. 2018.The mammary microenvironment in mastitis in humans, dairy ruminants, rabbits and rodents: a one health focus. *Journal of Mammary Gland Biology and Neoplasia*, 23: 27-41.
- Juliarena, M. A., C.N. Barrios, C.M. Lützelshwab, E.N. Esteban and S.E. Gutierrez.2017. Bovine leukemia virus: current perspectives. *Dove Press. Manchester, UK.* pp 1-9.
- Kidoya, H., M. Ueno, Y. Yamada, N. Mochizuki, M. Nakata, T. Yano and N. Takakura.2008. Spatial and temporal role of the apelin/APJ system in the caliber size regulation of blood vessels during angiogenesis. *The EMBO Journal*, 27(3): 522-534.
- Khan, K. 1993. Effects of Body Condition and PreLambing Supplementa-

- tion on Ewe Productivity. MS Thesis, Oregon State University, 102p, Oregon, USA.
- Lérias, J. R., L.E. Hernández-Castellano, A. Suárez-Trujillo, N. Castro, Pourlis A. and A.M. Almeida. 2014. The mammary gland in small ruminants: major morphological and functional events underlying milk production-a review. *Journal of Dairy Research*, 81(3):304-318.
- Lucas, M.H., M. Dawson, D. Chasey, G. Wibberley, Roberts D.H. and R. Saunders. 1980. Enzootic Bovine Leukosis Virus in Semen. *Veterinary Record*. 1980, 106, 128.
- Ma, W. Y., T.Y. Yu, J.N. Wei, C.S. Hung, M.S. Lin, Liao Y.J. and H.Y. Li. 2014. Plasma apelin: a novel biomarker for predicting diabetes. *Clinica Chimica Acta*. 435:18-23.
- Maezawa M., K.I. Watanabe, N. Horiuchi, K. Matsumoto, Y. Kobayashi and Inokuma, H.A. 2018. clinical case of enzootic bovine leukosis in a 13-month-old Holstein heifer. *Japanese Journal of Veterinary Research*. 66: 209-213.
- Masoumi, J., A. Jafarzadeh, H. Khorramdelazad, M. Abbasloui, J. Abdolizadeh and Jamali, N. 2020. Role of Apelin/APJ axis in cancer development and progression. *Advances in Medical Sciences*. 65(1): 202-213.
- Monke, D.R. 1986. Noninfectivity of semen from bulls infected with bovine leukosis virus. *Journal of the American Veterinary Medical Association*. 188: 823-826.
- Murakami, K., S. M. Kobayashi, K. Konishi Kameyama and Tsutsui, T. 2013. Nationwide survey of bovine leukemia virus infection among dairy and beef breeding cattle in Japan from 2009-2011. *The Journal of Veterinary Medical Science*. 75:1123-1126.
- Murtaugh, M. P., G.F. Lin, D.L. Haggard, A.F. Weber and Meiske, J. C. 1991. Detection of bovine leukemia virus in cattle by the polymerase chain reaction. *Journal of Virological Methods*. 33(1-2):73-85.
- Mutlak, S. S., M.S. Kadhim and Hussein, R. M. 2018. Apelin and some biomarkers in females with metabolic syndrome. *Biomedical and Pharmacology Journal*, 11(1): 247-253.
- Norvezh, F., M.R. Jalali, M.R. Tabandeh, M.R.H. Hajikolaie and Gooranjad, S. 2019. Serum Apelin-36 alteration in late pregnancy and early lactation of dairy cows and its association with negative energy balance markers. *Research in Veterinary Science*, 125: 285-289.
- Pandey, G. S., E. Simulundu, D. Mwiinga, K.L. Samui, A.S. Mweene, M. Kajihara and Takada, A. 2017. Clinical and subclinical bovine leukemia virus infection in a dairy cattle herd in Zambia. *Archives of Virology*. 162: 1051-1056.
- Picault, F. X., C. Chaves-Almagro, F. Progetti, H. Prats, B. Masri and Audigier, Y. 2014. Tumour co-expression of apelin and its receptor is the basis of an autocrine loop involved in the growth of colon adenocarcinomas. *European Journal of Cancer*. 50(3): 663-674.
- Podgórska, M., D. Diakowska, K. Pietraszek-Gremplewicz, M. Nie-nartowicz and Nowak, D. 2019. Evaluation of Apelin and Apelin receptor level in the primary tumor and serum of colorectal cancer patients. *Journal of Clinical Medicine*. 8(10): 1513.
- Reaux, A., N. De Mota, I. Skultetyova, Z. Lenkei, S. El Messari, K. Gallatz, and Llorens-Cortès, C. 2001. Physiological role of a novel neuropeptide, apelin, and its receptor in the rat brain. *Journal of Neurochemistry*. 77(4):1085-1096.
- Sorli, S. C., S. Le Gonidec, B. Knibiehler and Audigier, Y. 2007. Apelin is a potent activator of tumour neoangiogenesis. *Oncogene*, 26(55):7692-7699.
- Sorli, S. C., L. Van den Berghe, B. Masri, B. Knibiehler and Audigier, Y. (2006). Therapeutic potential of interfering with apelin signaling. *Drug Discovery Today*, 11(23-24):1100-1106.
- Szokodi, I., P. Tavi, G. Földes, S. Voutilainen-Myllylä, M. Ilves, H. Tokola and Ruskoaho, H. 2002. Apelin, the novel endogenous ligand of the orphan receptor APJ, regulates cardiac contractility. *Circulation Research*. 91(5):434-440.
- Taheri, S., K. Murphy, M. Cohen, E. Sujkovic, A. Kennedy, W. Dhillon and Bloom, S. 2002. The effects of centrally administered apelin-13 on food intake, water intake and pituitary hormone release in rats. *Biochemical and Biophysical Research Communications*, 291(5): 1208-1212.
- Tatemoto, K., M. Hosoya, Y. Habata, R. Fujii, T. Kakegawa, M.X. Zou and Kurokawa, T. 1998. Isolation and characterization of a novel endogenous peptide ligand for the human APJ receptor. *Biochemical and biophysical research communications*. 251(2):471-476.
- Thompson, J., H. Meyer. 1994. Body Condition Scoring of Sheep. *Oregon State University Extension Service*; 1-4.
- Wrathall, A. E., H.A. Simmons and Van Soom, A. 2006. Evaluation of risks of viral transmission to recipients of bovine embryos arising from fertilisation with virus-infected semen. *Theriogenology*. 65(2): 247-274.
- Wysocka, M. B., K. Pietraszek-Gremplewicz and Nowak, D. 2018. The role of apelin in cardiovascular diseases, obesity and cancer. *Frontiers in Physiology*. 9:1-15.
- Zhang, J., Q. Liu, Z. Fang, X. Hu, F. Huang, L. Tang and Zhou, S. 2016. Hypoxia induces the proliferation of endothelial progenitor cells via upregulation of Apelin/APLNR/MAPK signaling. *Molecular Medicine Reports*. 13(2): 1801-1806.