

Journal of the Hellenic Veterinary Medical Society

Vol 76, No 2 (2025)



New insights into polymorphisms in candidate genes associated with incidence to repeat breeder in Italian buffaloes (*Bubalus bubalis*)

A Ateya, FA Safhi , AR Alhimaidi , AA Ammari , RA Amran , B Marghani , SI Rakha , S Sakr, H El-Emam

doi: [10.12681/jhvms.36239](https://doi.org/10.12681/jhvms.36239)

Copyright © 2025, A Ateya, FA Safhi , AR Alhimaidi , AA Ammari , RA Amran , B Marghani , SI Rakha , S Sakr, H El-Emam



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

To cite this article:

Ateya, A., Safhi, F., Alhimaidi, A., Ammari, A., Amran, R., Marghani, B., Rakha, S., Sakr, S., & El-Emam, H. (2025). New insights into polymorphisms in candidate genes associated with incidence to repeat breeder in Italian buffaloes (*Bubalus bubalis*). *Journal of the Hellenic Veterinary Medical Society*, 76(2), 9055–9072. <https://doi.org/10.12681/jhvms.36239>

New insights into polymorphisms incandidate genes associated with incidence to repeat breeder in Italian buffaloes (*Bubalus bubalis*)

Ahmed Ateya,^{1*} Fatmah A. Safhi,² Ahmad R. Alhimaidi,³ Aiman A. Ammari,³
Ramzi A. Amran,³ Basma Marghani,^{4,5} Shima I. Rakha,⁶ Shima A. Sakr¹
and Huda El-Emam¹

¹Department of Development of Animal Wealth, Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt.

²Department of Biology, College of Science, Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia.

³Zoology Department, College of Science, King Saud University, Riyadh, Saudi Arabia

⁴Department of Physiology, Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt.

⁵Department of Biochemistry, Physiology, and Pharmacology, Faculty of Veterinary Medicine, King Salman International University, Egypt.

⁶Department of Theriogenology, Faculty of Veterinary Medicine, Mansoura University, Mansoura 35516, Egypt.

ABSTRACT: This study looked at the reproductive genes that could interact with repeat breeder incidence in Italian buffaloes. Two hundred and forty female Italian buffaloes (120 repeat breeder and 120 apparently normal) were used. Blood samples were collected from each buffalo into tubes encompassing EDTA anticoagulant to isolate DNA. PCR-DNA sequencing elicited nucleotide sequence differences between normal and repeat breeder buffaloes for the reproductive (*MFSD14A*, *PALMD*, *VPS13B*, *BMP*, *MTNR1A*, *TSHR*, *CSGALNACT1*, *CADM2*, *ZNF503*, *KRR1*, *MTPN*, *IGFBP7*, *LEP*, and *ABCC4*) genes. The likelihood of dissemination of all notable nucleotide variations varied noticeably between buffalo groups with and without repeat breeders, depending on Fisher's exact test ($p < 0.01$). All of the reproductive markers under research had the exonic region mutations in repeat breeder buffaloes versus normal ones. The results may confirm the importance of these markers' nucleotide variations as candidates for repeat breeder occurrence and offer a practical buffalo management strategy.

Keyword: Italian buffaloes; candidate gene; repeat breeder.

Correspondence author:

Ahmed Ateya
Department of Development of Animal Wealth, Faculty of
Veterinary Medicine, Mansoura University, Mansoura, Egypt
dr_ahmedismail@mans.edu.eg; ahmed_ismail888@yahoo.com

Date of initial submission: 18-12-2023

Date of acceptance: 18-11-2024

INTRODUCTION

Economically important for long-term food production are reproductive features, especially in monotocous animals like cattle and buffalo (Shao et al., 2021). Repeat breeding, or an extended interval between two calves, may be indicative of poor reproduction or sterility. More inseminations, veterinary care, and hormone therapies are required to address this illness, all of which have an impact on current and upcoming lactations (Biochard, 1990). Additionally, extra expenses are spent due to the culling and replacement of animals with reproduction problems (Roxström & Strandberg, 2002). The best option for reducing culling expenses, maintaining significant genetic features, and boosting farm profits is fertility improvement (Dekkers, 1991). With relation to statistical distribution, binary, interval, and continuous attributes have previously been used to classify reproductive characteristics (Berry & Evans 2014). Reproductive characteristics have been classified to make them simpler for understanding and utilizing in livestock and breeding programs (Cammack et al., 2009). These categories include features relating to ovulation, mating, and calving.

Because buffalo are mono-ovulatory, only one follicle typically grows and ovulates during the oestrus phase, which significantly reduces fertility and limits the growth of the buffalo business (Li et al., 2018). The reproductive activity of the buffaloes follows a seasonal pattern, with compromised activity in the summer and regular reproductive cyclicality and conception in the autumn and winter (Das & Khan 2010). Untangling the reproductive endocrinology of buffaloes will help researchers determine the cause of this reproductive restriction (Warriach et al., 2015). Quantitative features that affect reproduction are influenced by a variety of circumstances. Repeat breeding's occurrence in dairy farms can be reduced, and with it, the associated financial loss, by being aware of its underlying causes (Regmi&Dhakal, 2020). Repeat breeding can be caused by a wide range of factors, including insufficient fertilization, premature embryonic death, genital tract congenital or genetic faults, spermatozoa, ova or first zygote defects, microbial or traumatic inflammation, endocrine/hormonal problems, and management flaws (such as failing to identify estrus) (Yusuf et al., 2010). It is anticipated that genetic variants will influence gene expression, which will change the amount of one or more proteins, changing the phenotype of attributes (Musunuru et al., 2010; Lap-

palainen et al., 2013). A way to locate inherited components which affect susceptibility for common economic problems is marker-assisted selection (MAS) (Ashwell et al., 1997; Bishop et al., 1995). In the era of genomic assortment, excellent cow training groups that combine attributes with high-throughput genomic SNP marker data allow the use of selection strategies for emerging functional characteristics, such as disease resistance (Buch et al., 2012). Another significant advantage of genetic selection is shortening of generational intervals (Schaeffer, 2006). A novel understanding of the identity of underlying genes and change in the genetic material and how they might affect the characteristic has been claimed to be gained by integrating genetic markers to find significant genes linked with target traits (Li et al., 2014). The term "transcriptome" denotes all the genes in the genome that are accurately and at high throughput during particular physiological and pathological states (Kukurba & Montgomery, 2015).

Recent genome-wide association analysis studies have focused on novel genes related to bovine reproductive performance; however, no studies have yet looked at the relationship between the SNPs in these genes and the frequency of repeat breeders (Li et al., 2018; Mahrouset al., 2022). It is important to note that more research is necessary to follow potential symptoms that can imply reproductive failure from repeat breeder because there is limited knowledge on the reproduction pathways that determine buffalo resistance to or sensitivity to repeat breeder (Li et al., 2018). The molecular alterations may also help to identify repeat breeders and provide crucial information about the interactions between the physiologies of the various reproductive pathways (Sammad et al., 2022; Heidari et al., 2019). The effectiveness of putative reproductive (*MFS14A*, *PALMD*, *VPS13B*, *BMP*, *MTN1A*, *TSHR*, *CSGALNACT1*, *CADM2*, *ZNF503*, *KRR1*, *MTPN*, *IGFBP7*, *LEP*, and *ABCC4*) genes as candidates for repeat breeder incidence prediction and tracking in Italian buffaloes was examined in this study employing PCR-DNA sequencing methods.

MATERIALS AND METHODS

Dairy buffaloes and research samples

The current study was carried out on 240 pure Italian buffaloes aged 4.5 years. Buffaloes upraised on a private farm in the province of Delta region of northern Egypt of which 120 were repeat breeder and 120 appeared to be normal. Based on breeding history and clinical indications seen by skilled vets,

both groups were selected. All of the animals were raised in stalls with free access to water. It was determined that none of the study animals had ovarian pathologies like ovarian cysts or uterine, cervical, or vaginal inflammation or infection. The buffalo cows in the control group appeared to be in good condition and had a regular estrous cycle, nonetheless, no pregnancies were achieved in the repeat breeder group notwithstanding at least three successful attempts to artificially inseminate or mate with a bull in estrus (Devkota et al., 2022). Visual observation was used to identify estrus in both groups twice daily in the morning and in the evening. All animals were checked for foetal membrane slip and/or fremitus, and rectal palpation was used to confirm pregnancy diagnoses.

Five milliliters of blood were obtained by puncturing the jugular vein of each buffalo. To obtain whole blood and recover DNA samples were placed in tubes containing EDTA as anticoagulant. The Ethical Committee approved the sample collection and animal care techniques utilized in this study, and they complied with Mansoura University's regulations. The Mansoura University Animal Care and Use Committee (MU-ACUC) gave its approval to the study's protocol (code VM.R.23.11.26).

DNA extraction and PCR amplification

DNA from the genome was extracted using total blood by means of the genetic material JET full blood genomic DNA isolation kit and the producer's directions (Thermo scientific, Vilnius, Lithuania). Using Nanodrop, DNA of good purity and concentration was considered. The following reproductive (*MFS14A*, *PALMD*, *VPS13B*, *BMP*, *MTNR1A*, *TSHR*, *CSGALNACT1*, *CADM2*, *ZNF503*, *KRR1*, *MTPN*, *IGFBP7*, *LEP*, and *ABCC4*) genes' parts of coding sites (CDS) were amplified. The oligonucleotide sequences for amplification were constructed using the PubMed *Bubalus bubalis* genome. Table 1 provides the primers utilized in the PCR.

The polymerase chain cloning mixture had been treated in a thermal cycler with a final volume of 50 μ L. The succeeding elements were present in every reaction container: 19 microliters d.d. water, 4 microliters of genetic material, 1 microliters of each primer pair, and 25 microliters of the master combination (Jena Bioscience, Jena, Germany). The PCR combinations remained in use for four minutes at a starting temperature of 95 °C for unwinding. The 34 cycles included one-minute denaturation rounds

at 95 °C, one-minute annealing cycles based on the Table 1 temperature range, and 30-second for elongation at 72 °C, followed by ten minutes of final elongation. The samples were retained at 4 °C. Using agarose gel electrophoresis to obtain demonstrable outcomes and viewing PCR segment configurations underneath UV light, a gel certification technique was used.

Discovering polymorphism

Before DNA sequencing, Hamburg, Germany-based Jena Bioscience # pp-201s/Munich provided methods to purify PCR for getting rid of primer dimers, non-specific bands, and other impurities and create the predicted scope's planned amplified product (Boom et al., 1990). Using a Nanodrop (Waltham, Massachusetts, USA, UV-Vis spectrometer Q5000), adequate quality and good concentrations were obtained while measuring PCR output (Boesenberg-Smith et al., 2012). Sequence analysis of the PCR-produced amplification results has been utilized to find SNPs using normal and repeat breeder buffaloes. The PCR yields were sequenced using the Sanger et al., (1977) outlined enzyme chain terminator technique on an ABI 3730XL DNA sequencer (United States: Applied Biosystems, Waltham).

Chromas 1.45 and BLAST 2.0, the tools to evaluate results of DNA analysis (Altschul et al., 1990). Comparing the reproductive gene PCR results to the GenBank-provided reference gene sequences has revealed polymorphisms. The MEGA6 tool can detect differences in the amino acid categorizations among the examined genes according to sequence matching among the buffaloes under investigation (Tamura et al., 2007).

Statistical analysis

H_0 : Polymorphisms of reproductive genes could not interact in the incidence to repeat breeder in Italian buffaloes.

H_A : Polymorphisms of reproductive genes could interact in the incidence to repeat breeder in Italian buffaloes.

Observing the significant distribution of SNPs for the discovered genes between the examined buffaloes was conducted using Fisher's exact test analysis ($p < 0.01$). Graphpad statistical software was used to conduct the statistical analysis (Graphpad prism for Windows version 5.1, Graphpad software, Inc., San Diego, CA, USA).

Table 1. Oligonucleotide primers for reproductive genes used to study genetic polymorphisms.

Investigated marker	Sense	Antisense	Accession number	Annealing Temperature (°C)	Size of PCR product (bp)
<i>MFSDI4A</i>	5'-GCCGAACCGCAGTATCATGCTG-3'	5'-CCAGGCCATACGCCATACTTC-3'	XM_055584984.1	60	457
<i>PALMD</i>	5'-TGGCTTCTAGATGGAATCAGC-3'	5'-CTATTCCTTCATTTCTTCC-3'	XM_006045619.1	58	343
<i>VPSI3B</i>	5'-AATGAGCTATGTGAATCGCT-3'	5'-ACAACCTCGTCTGATCAGACTCT-3'	XM_055546577.1	60	414
<i>BMP</i>	5'-ATGGTCCCTTCTGAGCATCCTTA-3'	5'-GTGGAGCCCTCTTCCCTGAAGAAG-3'	EF375880.1	60	480
<i>MTNRI4</i>	5'-GTGAGCCCTGGCAGTTGCAGAC-3'	5'-CACTCTCCATCTGACCTGAAG-3'	MF173058.1	56	465
<i>TSHR</i>	5'-ATCCGCCCTCTGGCACGCCCTAC-3'	5'-TGGCATAAGGAATGGATTGGC-3'	KC415275.1	58	442
<i>CSGALNACT1</i>	5'-CTGCTCGCCCTGGGCCCTCCCCGG-3'	5'-CGGTGGCCTGCTTGACGCCCGGC-3'	XM_025275645.3	58	385
<i>CADM2</i>	5'-CGCAGCGCCGTTCTCCCGCTTC-3'	5'-GAACCACTTTATATCAAGCTGCA-3'	XM_025278930.3	60	495
<i>ZNF503</i>	5'-TCTCCGCCAGGCTAACCGCCT-3'	5'-CTTATCCGAGCCCGGTTTGGAG-3'	XM_025285122.3	60	361
<i>KRRI</i>	5'-CCGACGTCGGAGAACCGAGATG-3'	5'-CTAATGATGATATATGGATCA-3'	XM_006048659.4	58	278
<i>MTPN</i>	5'-ATGTGGGACAAAGGAGTTCATG-3'	5'-GGCACCCAGTGACATCCAGAT-3'	XM_044946911.2	60	266
<i>IGFBP7</i>	5'-AGCTGGGGGAGACCCCGGATGC-3'	5'-GACAGAAACAATTCACGCAG-3'	XM_025290001.3	56	380
<i>LEP</i>	5'-CAGTCCGTCTCCTCCAAACAGA-3'	5'-TCAGCACCCAGGACTGAGGTC-3'	AF387814.1	58	360
<i>ABCC4</i>	5'-GCAGAAAGGCGAACTTCTGCTC-3'	5'-TATGGTTGAAC'TACTCTGGTG-3'	XM_045162483.1	60	355

MFSDI4A= Major facilitator superfamily domain containing 14A; *PALMD*= Palmdelphin; *VPSI3B*= Vacuolar protein sorting 13 homolog B; *BMP*= Bone morphogenetic protein; *MTNRI4*= Melatonin receptor 1B; *TSHR*= Thyroid stimulating hormone receptor; *CSGALNACT1*= Chondroitin sulfate N-acetylgalactosaminyltransferase 1; *CADM2*= Cell adhesion molecule 2; *ZNF503*= Zinc finger protein 503; *KRRI* = Small subunit processome component homolog; *MTPN*= Myotrophin; *IGFBP7*= insulin like growth factor binding protein 7; *LEP*= Leptin; and *ABCC4*= ATP-binding cassette sub-family C member 4.

RESULTS

SNP variations in amplified DNA nucleotides associated with repeat breeder were found in the results of PCR-DNA sequencing on both normal and repeat breeder buffaloes for the MFSD14A (457-bp), PALMD (343-bp), VPS13B (414-bp), BMP (480-bp), MTNR1A (465-bp), TSHR (442-bp), CS-GALNACT1 (385-bp), CADM2 (495-bp), ZNF503 (361-bp), KRR1 (278-bp), MTPN (266-bp), IGFBP7 (380-bp), LEP (360-bp), and ABCC4 (355-bp) genes. The DNA sequence differences between the sequences of reference genes obtained from GenBank and the reproductive indicators examined in the studied buffaloes were used to verify each SNP that has been discovered (Figures S1–S14).

Table 2 shows the dissemination of a single base variation as well as a type of inherited change for reproductive indicators in normal and repeat breeder buffaloes. The SNPs' Fisher's exact test analysis revealed considerably different occurrences of the investigated markers in the normal and repeat breeder buffaloes ($p < 0.01$). All of the reproductive markers under research had the exonic region mutations, which resulted in different coding DNA sequences in repeat breeder buffaloes versus nor-

mal ones. Thirty SNPs were discovered using DNA sequencing of reproductive genes; 16 of them are synonymous and 15 is non-synonymous. The amino acid variations between the reproductive genes in all the research buffaloes and the reference sequences retrieved in GenBank were used to validate all observed SNPs and the corresponding amino acids (Figures S15–S28).

For the MFSD14A gene (457-bp), two observed recurrent synonymous SNPs, 165CT and 322TC, resulted in 55C and 108G respectively. One recurrent SNP was found when the PALMD gene (343-bp) was sequenced. Synonymous mutation 36L occurred as a result of 108TC SNP. 175TC and 223TC involved a non-synonymous mutation, resulting in the substitution of the amino acid C59R, and Y75H respectively; were found in DNA sequences of the VPS13B gene (414-bp). The BMP gene (480-bp) contained three non-synonymous SNPs: 47TC, 80CA, and 184GA, which caused the amino acids V16A, P27Q, and V62I to be substituted, respectively. The nucleotide sequence of the MTNR1A gene (465-bp) revealed one recurrent synonymous SNP; 24N resulted from 72CT SNP. The 442-bp TSHR gene's DNA sequence revealed four identified

XM_055584984.1	GCGAACCGCAGTATCATGCTGGCCAAAAAGATCATCATTAAAGGACGGAGGCACGCCTCAA	60
H	GCGAACCGCAGTATCATGCTGGCCAAAAAGATCATCATTAAAGGACGGAGGCACGCCTCAA	60
RR	GCGAACCGCAGTATCATGCTGGCCAAAAAGATCATCATTAAAGGACGGAGGCACGCCTCAA	60

XM_055584984.1	GGAATAGGTTCTCCTAGTGTATATCATGCGAGTTATCGTCACTTTTTGGAGTTTTTCGCT	120
H	GGAATAGGTTCTCCTAGTGTATATCATGCGAGTTATCGTCACTTTTTGGAGTTTTTCGCT	120
RR	GGAATAGGTTCTCCTAGTGTATATCATGCGAGTTATCGTCACTTTTTGGAGTTTTTCGCT	120

XM_055584984.1	TGGGGATTATTGACAGCACCCACCTTGGTGGTATTACATGAAACCTTCCCTAAACATACA	180
H	TGGGGATTATTGACAGCACCCACCTTGGTGGTATTACATGAAACCTTCCCTAAACATACA	180
RR	TGGGGATTATTGACAGCACCCACCTTGGTGGTATTACATGAAACTTCCCTAAACATACA	180

XM_055584984.1	TTTCTGATGAATGGCCTAATTCAAGGAGTAAAGGGTTTGTGTGCATTCCCTCAGTGCCCT	240
H	TTTCTGATGAATGGCCTAATTCAAGGAGTAAAGGGTTTGTGTGCATTCCCTCAGTGCCCT	240
RR	TTTCTGATGAATGGCCTAATTCAAGGAGTAAAGGGTTTGTGTGCATTCCCTCAGTGCCCT	240

XM_055584984.1	CTTATTGGTGCTCTTTCTGATGTTTGGGGCCGAAATCCTTCTTGCTGCTAACAGTATTT	300
H	CTTATTGGTGCTCTTTCTGATGTTTGGGGCCGAAATCCTTCTTGCTGCTAACAGTATTT	300
RR	CTTATTGGTGCTCTTTCTGATGTTTGGGGCCGAAATCCTTCTTGCTGCTAACAGTATTT	300

XM_055584984.1	TTCACGTGTGCCCAATTCCTTTAATGAAATCAGCCCATGGTGGTACTTTGCTGTTATC	360
H	TTCACGTGTGCCCAATTCCTTTAATGAAATCAGCCCATGGTGGTACTTTGCTGTTATC	360
RR	TTCACGTGTGCCCAATTCCTTTAATGAAATCAGCCCATGGTGGTACTTTGCTGTTATC	360

XM_055584984.1	TCTGTTTCTGGGGTTTTTGCAGTGACTTCTCCGTGGTATTGCAATGTAGCAGATATA	420
H	TCTGTTTCTGGGGTTTTTGCAGTGACTTCTCCGTGGTATTGCAATGTAGCAGATATA	420
RR	TCTGTTTCTGGGGTTTTTGCAGTGACTTCTCCGTGGTATTGCAATGTAGCAGATATA	420

XM_055584984.1	ACCCAAGAACATGAAAGAAGTATGGCGTATGGCCTGG 457	
H	ACCCAAGAACATGAAAGAAGTATGGCGTATGGCCTGG 457	
RR	ACCCAAGAACATGAAAGAAGTATGGCGTATGGCCTGG 457	

Figure S1. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb|XM_055584984.1| and *MFSD14A* marker (457-bp) sequences.

XM_006045619.4	TGGCTTCTAGATGGAATCAGCAGTGGAAAAAGAGCAGGAAGAGATGAAGAAGCAAAATCAA	60
H	TGGCTTCTAGATGGAATCAGCAGTGGAAAAAGAGCAGGAAGAGATGAAGAAGCAAAATCAA	60
RR	TGGCTTCTAGATGGAATCAGCAGTGGAAAAAGAGCAGGAAGAGATGAAGAAGCAAAATCAA	60

XM_006045619.4	CAAGACCAGCACCAGATCCAGGTTCTAGAACAAAGTATCCTCAGACTTGAGAAAGAGATC	120
H	CAAGACCAGCACCAGATCCAGGTTCTAGAACAAAGTATCCTCAGACTTGAGAAAGAGATC	120
RR	CAAGACCAGCACCAGATCCAGGTTCTAGAACAAAGTATCCTCAGACTTGAGAAAGAGATC	120

XM_006045619.4	CAAGATCTTGAAAAGGCTGAACTGCRAATCTCAACCAATGAAGAAGCAATTTTAAAGAAA	180
H	CAAGATCTTGAAAAGGCTGAACTGCRAATCTCAACCAATGAAGAAGCAATTTTAAAGAAA	180
RR	CAAGATCTTGAAAAGGCTGAACTGCRAATCTCAACCAATGAAGAAGCAATTTTAAAGAAA	180

XM_006045619.4	CTGAAATCAGTTGAGAGGACAAACAGAGACATAATAAGGCTCTGTGAAAGTGGAAAAGGAA	240
H	CTGAAATCAGTTGAGAGGACAAACAGAGACATAATAAGGCTCTGTGAAAGTGGAAAAGGAA	240
RR	CTGAAATCAGTTGAGAGGACAAACAGAGACATAATAAGGCTCTGTGAAAGTGGAAAAGGAA	240

XM_006045619.4	GAAACATCAGGAGAGTCAGTTGAAGACATCTATGCTAATATCCCCGACCTTCCAAAATCC	300
H	GAAACATCAGGAGAGTCAGTTGAAGACATCTATGCTAATATCCCCGACCTTCCAAAATCC	300
RR	GAAACATCAGGAGAGTCAGTTGAAGACATCTATGCTAATATCCCCGACCTTCCAAAATCC	300

XM_006045619.4	TACATACCTTCCAGGTTAAGGAAGGAAAGAAATGAAGGAATAG	343
H	TACATACCTTCCAGGTTAAGGAAGGAAAGAAATGAAGGAATAG	343
RR	TACATACCTTCCAGGTTAAGGAAGGAAAGAAATGAAGGAATAG	343

Figure S2. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb| XM_006045619.4| and *PALMD* marker (343-bp) sequences.

XM_055546577.1	AATGAGCTATGTGAATCGCTACATCAAGAACTTAAAGCCGTCAGATCTACAGCTTTCGCT	60
H	AATGAGCTATGTGAATCGCTACATCAAGAACTTAAAGCCGTCAGATCTACAGCTTTCGCT	60
RR	AATGAGCTATGTGAATCGCTACATCAAGAACTTAAAGCCGTCAGATCTACAGCTTTCGCT	60

XM_055546577.1	ATGGGGTGGAGACGTTGGTTCTCAGCAAGCTGGAGTTAAAATTGGATGTGCTGGAGCAGGA	120
H	ATGGGGTGGAGACGTTGGTTCTCAGCAAGCTGGAGTTAAAATTGGATGTGCTGGAGCAGGA	120
RR	ATGGGGTGGAGACGTTGGTTCTCAGCAAGCTGGAGTTAAAATTGGATGTGCTGGAGCAGGA	120

XM_055546577.1	ATTGAAATTACCATTACATTTTTTAAGTGGACATATTCATGAACTGAGGATCCATGTACC	180
H	ATTGAAATTACCATTACATTTTTTAAGTGGACATATTCATGAACTGAGGATCCATGTACC	180
RR	ATTGAAATTACCATTACATTTTTTAAGTGGACATATTCATGAACTGAGGATCCATGTACC	180

XM_055546577.1	ATGGACAAAACCTGGGTTCCAGAACCAAGTGGTAATTACCATCAATACAATGGAATGCATTTT	240
H	ATGGACAAAACCTGGGTTCCAGAACCAAGTGGTAATTACCATCAATACAATGGAATGCATTTT	240
RR	ATGGACAAAACCTGGGTTCCAGAACCAAGTGGTAATTACCATCAATACAATGGAATGCATTTT	240

XM_055546577.1	GAACTTAAAGATGGAATACAGGATGATCATGAAAGCTGTGGTTCTAATTTCTACCAACCG	300
H	GAACTTAAAGATGGAATACAGGATGATCATGAAAGCTGTGGTTCTAATTTCTACCAACCG	300
RR	GAACTTAAAGATGGAATACAGGATGATCATGAAAGCTGTGGTTCTAATTTCTACCAACCG	300

XM_055546577.1	TAGTACTACTGAGAACACAAAACATCAGCAAAACCCCGAAGAAATTCACAGGCCACTCC	360
H	TAGTACTACTGAGAACACAAAACATCAGCAAAACCCCGAAGAAATTCACAGGCCACTCC	360
RR	TAGTACTACTGAGAACACAAAACATCAGCAAAACCCCGAAGAAATTCACAGGCCACTCC	360

XM_055546577.1	CACAGATCCTGATTTGCCACCAGGCTATGTACAGAGTCTGATCAGACGAGTTGT	414
H	CACAGATCCTGATTTGCCACCAGGCTATGTACAGAGTCTGATCAGACGAGTTGT	414
RR	CACAGATCCTGATTTGCCACCAGGCTATGTACAGAGTCTGATCAGACGAGTTGT	414

Figure S3. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb|XM_055546577.1| and *VPS13B* marker (414-bp) sequences.

frequent synonymous SNPs; 87CT, 126TC, 177CT, and 372CT, which were associated with the amino acids 29S, 42T, 59I, 42S and 124V.

For the *CSGALNACT1* gene (385-bp), one synonymous SNP was found; the results of 135GAs45T. Three recurrent synonymous SNPs; 105GA, 168TC,

and 354GT cleared during DNA sequencing of the *CADM2* (495-bp) gene, which resulted in the amino acids 50R, 56N, and 118T, respectively. One recurrent non-synonymous SNP, 217GC, was elaborated using DNA sequencing of the *ZNF503* gene (361-bp), which led to the substitution of the amino acid

```

EF375880.1   ATGGTCCTTCTGAGCATCCTTAGAATCCTTCTTCTTTGGGGACTGGTGCTTTTTATGGAA 60
H            ATGGTCCTTCTGAGCATCCTTAGAATCCTTCTTCTTTGGGGACTGGTGCTTTTTATGGAA 60
RR          ATGGTCCTTCTGAGCATCCTTAGAATCCTTCTTCTTTGGGGACTGGCGCTTTTTATGGAA 60
*****
EF375880.1   CATAGGGTCCAAATGACACCGGTAGGGCAGCCCTCTATTGCCCACTGCCTGAGGCCCT 120
H            CATAGGGTCCAAATGACACAGGTAGGGCAGCCCTCTATTGCCCACTGCCTGAGGCCCT 120
RR          CATAGGGTCCAAATGACACCGGTAGGGCAGCCCTCTATTGCCCACTGCCTGAGGCCCT 120
*****
EF375880.1   ACCTTGCCCTGATTGAGGAGCTGCTAGAGAAGCCCTGGCAAGCTGCAGAGGAAGCCG 180
H            ACCTTGCCCTGATTGAGGAGCTGCTAGAGAAGCCCTGGCAAGCTGCAGAGGAAGCCG 180
RR          ACCTTGCCCTGATTGAGGAGCTGCTAGAGAAGCCCTGGCAAGCTGCAGAGGAAGCCG 180
*****
EF375880.1   CGGGTCTTAGGGCATCCCTTACGGTATATGCTGGAGTTGTACCACCGTTGAGCTGACGCA 240
H            CGGATCTTAGGGCATCCCTTACGGTATATGCTGGAGTTGTACCACCGTTGAGCTGACGCA 240
RR          CGGGTCTTAGGGCATCCCTTACGGTATATGCTGGAGTTGTACCACCGTTGAGCTGACGCA 240
*****
EF375880.1   AGTGGACACCCTAGGGAARACCGCACCATTGGGGCCACCATGGTGAGGCTGGTGAGGCCA 300
H            AGTGGACACCCTAGGGAARACCGCACCATTGGGGCCACCATGGTGAGGCTGGTGAGGCCA 300
RR          AGTGGACACCCTAGGGAARACCGCACCATTGGGGCCACCATGGTGAGGCTGGTGAGGCCA 300
*****
EF375880.1   CTGGCTAGTGTAGCAAGGCCTCTCAGAGGCTCCTGGCACATACAGACCCTGGACTTTCC 360
H            CTGGCTAGTGTAGCAAGGCCTCTCAGAGGCTCCTGGCACATACAGACCCTGGACTTTCC 360
RR          CTGGCTAGTGTAGCAAGGCCTCTCAGAGGCTCCTGGCACATACAGACCCTGGACTTTCC 360
*****
EF375880.1   CTGAGACCAAACCGGGTAGCATAACCACTAGTCAGAGCCACTGTGGTTTACCGCCATCAA 420
H            CTGAGACCAAACCGGGTAGCATAACCACTAGTCAGAGCCACTGTGGTTTACCGCCATCAA 420
RR          CTGAGACCAAACCGGGTAGCATAACCACTAGTCAGAGCCACTGTGGTTTACCGCCATCAA 420
*****
EF375880.1   CTTACCTRACTCATTCCCACCTCTCCTGCCATGTGGAGCCCTGGGTCCAGAAAAGCCCA 480
H            CTTACCTRACTCATTCCCACCTCTCCTGCCATGTGGAGCCCTGGGTCCAGAAAAGCCCA 480
RR          CTTACCTRACTCATTCCCACCTCTCCTGCCATGTGGAGCCCTGGGTCCAGAAAAGCCCA 480
*****

```

Figure S4. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb|EF375880.1| and *BMP* marker (480-bp) sequences.

```

MF173058.1  GTGAGCCTGGCAGTTGCAGACCCTGCTGGTGGCCGTTGATCCGTACCCCTTGGCGCTGGCG 60
H            GTGAGCCTGGCAGTTGCAGACCCTGCTGGTGGCCGTTGATCCGTACCCCTTGGCGCTGGCG 60
RR          GTGAGCCTGGCAGTTGCAGACCCTGCTGGTGGCCGTTGATCCGTACCCCTTGGCGCTGGCG 60
*****
MF173058.1  TCTATAGTTAACGATGGGTGGAGCCTGAGCTCCCTGCATTGCCAAGTTAGTGGCTTCCTG 120
H            TCTATAGTTAATGATGGGTGGAGCCTGAGCTCCCTGCATTGCCAAGTTAGTGGCTTCCTG 120
RR          TCTATAGTTAACGATGGGTGGAGCCTGAGCTCCCTGCATTGCCAAGTTAGTGGCTTCCTG 120
*****
MF173058.1  ATGGGCTTGAGCGTCATCGGGTCCGTTTTCAATATCACGGGAATTGCCATCAACCGCTAT 180
H            ATGGGCTTGAGCGTCATCGGGTCCGTTTTCAATATCACGGGAATTGCCATCAACCGCTAT 180
RR          ATGGGCTTGAGCGTCATCGGGTCCGTTTTCAATATCACGGGAATTGCCATCAACCGCTAT 180
*****
MF173058.1  TGCTGCAICTGCCACAGCCTCAGATACGACAAGCTGTATAGCAGCACAAATCCCTCTGC 240
H            TGCTGCAICTGCCACAGCCTCAGATACGACAAGCTGTATAGCAGCACAAATCCCTCTGC 240
RR          TGCTGCAICTGCCACAGCCTCAGATACGACAAGCTGTATAGCAGCACAAATCCCTCTGC 240
*****
MF173058.1  TACGTGTTCCCTGATATGGCTGCTGACGTTTCGTGGCGATCGTGCCCAACCTGTGTGTGGGG 300
H            TACGTGTTCCCTGATATGGCTGCTGACGTTTCGTGGCGATCGTGCCCAACCTGTGTGTGGGG 300
RR          TACGTGTTCCCTGATATGGCTGCTGACGTTTCGTGGCGATCGTGCCCAACCTGTGTGTGGGG 300
*****
MF173058.1  ACCCTGCGGTACGACCCGAGGATCTATTCTTGTACCTTCACACAGTCCGTGAGCTCAGCC 360
H            ACCCTGCGGTACGACCCGAGGATCTATTCTTGTACCTTCACACAGTCCGTGAGCTCAGCC 360
RR          ACCCTGCGGTACGACCCGAGGATCTATTCTTGTACCTTCACACAGTCCGTGAGCTCAGCC 360
*****
MF173058.1  TACACGATCGCCGTGGTGGTGTCCATTTTCATAGTTCCAAATGCTCGTAGTCATCTTCTGT 420
H            TACACGATCGCCGTGGTGGTGTCCATTTTCATAGTTCCAAATGCTCGTAGTCATCTTCTGT 420
RR          TACACGATCGCCGTGGTGGTGTCCATTTTCATAGTTCCAAATGCTCGTAGTCATCTTCTGT 420
*****
MF173058.1  TACCTGAGAACTGCGCCCTGGTTCTTCAGGTCAGATGGAGAGTG 465
H            TACCTGAGAACTGCGCCCTGGTTCTTCAGGTCAGATGGAGAGTG 465
RR          TACCTGAGAACTGCGCCCTGGTTCTTCAGGTCAGATGGAGAGTG 465
*****

```

Figure S5. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb|MF173058.1| and *MTNR1A* marker (465-bp) sequences.

```

KC415275.1 ATCCGCCCTCTGGCAGCCCTACGTCATCATGCTGGGGGGCTGGGTTTGTCTGCTTCCCTGCTC 60
H ATCCGCCCTCTGGCAGCCCTACGTCATCATGCTGGGGGGCTGGGTTTGTCTGCTTCCCTGCTC 60
RR ATCCGCCCTCTGGCAGCCCTACGTCATCATGCTGGGGGGCTGGGTTTGTCTGCTTCCCTGCTC 60
*****
KC415275.1 GCCTTGTCCCTTTGGTGGGAATAGCAGCTATGCCAAGGTGAGCATCTGCCTGCCCATG 120
H GCCTTGTCCCTTTGGTGGGAATAGCAGCTATGCCAAGGTGAGCATCTGCCTGCCCATG 120
RR GCCTTGTCCCTTTGGTGGGAATAGTAGCTATGCCAAGGTGAGCATCTGCCTGCCCATG 120
*****
KC415275.1 GACACTGAGACTCCTCTTGCCTGGCGTACATTATCCTCGTTCTGTTACTCAACATCGTT 180
H GACACTGAGACTCCTCTTGCCTGGCGTACATTATCCTCGTTCTGTTACTCAACATCGTT 180
RR GACACCGAGACTCCTCTTGCCTGGCGTACATTATCCTCGTTCTGTTACTCAACATCGTT 180
*****
KC415275.1 GCCTTTATCATCGTCTGTGCTGTACGTGAAGATCTACATCACAGTCCGAAATCCCCAC 240
H GCCTTTATCATCGTCTGTGCTGTACGTGAAGATCTACATCACAGTCCGAAATCCCCAC 240
RR GCCTTTATCATCGTCTGTGCTGTACGTGAAGATCTACATCACAGTCCGAAATCCCCAC 240
*****
KC415275.1 TACAACCCGGGGGACAAAGATACTAGAATTGCCAAAAGGATGGCTGTGTGATCTTCACT 300
H TACAACCCGGGGGACAAAGATACTAGAATTGCCAAAAGGATGGCTGTGTGATCTTCACT 300
RR TACAACCCGGGGGACAAAGATACTAGAATTGCCAAAAGGATGGCTGTGTGATCTTCACT 300
*****
KC415275.1 GACTTCAATGTCATGGCCCAATCTCTTCTACGCTCTGTGCGCCCTTATGAACAAGCCT 360
H GACTTCAATGTCATGGCCCAATCTCTTCTACGCTCTGTGCGCCCTTATGAACAAGCCT 360
RR GACTTCAATGTCATGGCCCAATCTCTTCTACGCTCTGTGCGCCCTTATGAACAAGCCT 360
*****
KC415275.1 CTCATCACCGTACCCAATTCAAAATCTTGCTGGTCTCTTCTACCCACTTAACCTCTGT 420
H CTCATCACCGTACCCAATTCAAAATCTTGCTGGTCTCTTCTACCCACTTAACCTCTGT 420
RR CTCATCACCGTTACCCAATTCAAAATCTTGCTGGTCTCTTCTACCCACTTAACCTCTGT 420
*****
KC415275.1 GCCAATCCATTCTCTATGCCA 442
H GCCAATCCATTCTCTATGCCA 442
RR GCCAATCCATTCTCTATGCCA 442
*****

```

Figure S6. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb|KC415275.1| and *TSHR* marker (442-bp) sequences.

```

XM_025275645.3 CTGCTCGCCTGGGCTCCCGGGTGGTGGTCTTGTCTCGTCTTCTCTGCTGCGTGGCGTCT 60
H CTGCTCGCCTGGGCTCCCGGGTGGTGGTCTTGTCTCGTCTTCTCTGCTGCGTGGCGTCT 60
RR CTGCTCGCCTGGGCTCCCGGGTGGTGGTCTTGTCTCGTCTTCTCTGCTGCGTGGCGTCT 60
*****
XM_025275645.3 GTCCTCTACATGCTGGCCTGCACGCCGAGGGGCGATGATGAGCGCCAGGGGCTGCCCAGG 120
H GTCCTCTACATGCTGGCCTGCACGCCGAGGGGCGATGATGAGCGCCAGGGGCTGCCCAGG 120
RR GTCCTCTACATGCTGGCCTGCACGCCGAGGGGCGATGATGAGCGCCAGGGGCTGCCCAGG 120
*****
XM_025275645.3 GCCAACGGCCCCACGGGACGGGCGGGCTCCACGGCGGGTGTCTGGCTGGGAGGAGCGT 180
H GCCAACGGCCCCACGGGACGGGCGGGCTCCACGGCGGGTGTCTGGCTGGGAGGAGCGT 180
RR GCCAACGGCCCCACGGGACGGGCGGGCTCCACGGCGGGTGTCTGGCTGGGAGGAGCGT 180
*****
XM_025275645.3 CACCGGACCAAGTGGGAGCCTCAAGAGGCAATTCGCGAGCTGCAGGAGGAGCTGCAA 240
H CACCGGACCAAGTGGGAGCCTCAAGAGGCAATTCGCGAGCTGCAGGAGGAGCTGCAA 240
RR CACCGGACCAAGTGGGAGCCTCAAGAGGCAATTCGCGAGCTGCAGGAGGAGCTGCAA 240
*****
XM_025275645.3 GAGCGGAGCGAGCAGCTGAAGGCGCGCAGCAGCTGGCGGGGAGCCCGGGGTGCCCGG 300
H GAGCGGAGCGAGCAGCTGAAGGCGCGCAGCAGCTGGCGGGGAGCCCGGGGTGCCCGG 300
RR GAGCGGAGCGAGCAGCTGAAGGCGCGCAGCAGCTGGCGGGGAGCCCGGGGTGCCCGG 300
*****
XM_025275645.3 GGCAGGGCCCAGGCCGACCTGTGGCCTTCTGCGCTCGCAGGTGGACAGGGCGGAGGTG 360
H GGCAGGGCCCAGGCCGACCTGTGGCCTTCTGCGCTCGCAGGTGGACAGGGCGGAGGTG 360
RR GGCAGGGCCCAGGCCGACCTGTGGCCTTCTGCGCTCGCAGGTGGACAGGGCGGAGGTG 360
*****
XM_025275645.3 CACGCCGGCGTCAAGCAGGCCACCG 385
H CACGCCGGCGTCAAGCAGGCCACCG 385
RR CACGCCGGCGTCAAGCAGGCCACCG 385
*****

```

Figure S7. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb|XM_025275645.3| and *CSGALNACT1* marker (385-bp) sequences.

XM_025278930.3	CGCAGCGCCGTTCTCCGCTTCTACAGTGTCTGCGGGCTCCTGCTACAAGCGGCTGCTTCA	60
H	CGCAGCGCCGTTCTCCGCTTCTACAGTGTCTGCGGGCTCCTGCTACAAGCGGCTGCTTCA	60
RR	CGCAGCGCCGTTCTCCGCTTCTACAGTGTCTGCGGGCTCCTGCTACAAGCGGCTGCTTCA	60
*****	*****	*****
XM_025278930.3	ARGAATRAAGTTAAAGGCAGCCAGGGCAGTTTCCACTCACACAGAATGTAACGGTTGTT	120
H	ARGAATRAAGTTAAAGGCAGCCAGGGCAGTTTCCACTCACACAGAATGTAACGGTTGTT	120
RR	ARGAATRAAGTTAAAGGCAGCCAGGGCAGTTTCCACTCACACAGAATGTAACGGTTGTT	120
*****	*****	*****
XM_025278930.3	GAAGCGGAACCTGCAATTTTGACCTGCAGGGTTGATCAAAATGATAATACCTCCCTCCAG	180
H	GAAGCGGAACCTGCAATTTTGACCTGCAGGGTTGATCAAAATGATAATACCTCCCTCCAG	180
RR	GAAGCGGAACCTGCAATTTTGACCTGCAGGGTTGATCAAAATGATAATACCTCCCTCCAG	180
*****	*****	*****
XM_025278930.3	TGGTCARATCCAGCTCAACAGACTCTGTACTTTGATGACAAGAAAGCCTTAAGAGACAAT	240
H	TGGTCARATCCAGCTCAACAGACTCTGTACTTTGATGACAAGAAAGCCTTAAGAGACAAT	240
RR	TGGTCARATCCAGCTCAACAGACTCTGTACTTTGATGACAAGAAAGCCTTAAGAGACAAT	240
*****	*****	*****
XM_025278930.3	AGAATCGAGCTGGTTCGGGCTTCTGGCATGAAGTATCAGCGTCAGTGTGTCTCT	300
H	AGAATCGAGCTGGTTCGGGCTTCTGGCATGAAGTATCAGCGTCAGTGTGTCTCT	300
RR	AGAATCGAGCTGGTTCGGGCTTCTGGCATGAAGTATCAGCGTCAGTGTGTCTCT	300
*****	*****	*****
XM_025278930.3	CTGTCTGATGAAGGACAGTACACCTGTTCTTTATTTACAATGCCTGTCAAACGTCCTAAG	360
H	CTGTCTGATGAAGGACAGTACACCTGTTCTTTATTTACAATGCCTGTCAAACGTCCTAAG	360
RR	CTGTCTGATGAAGGACAGTACACCTGTTCTTTATTTACAATGCCTGTCAAACGTCCTAAG	360
*****	*****	*****
XM_025278930.3	GCCTATCTCACTGTCTGGGTGTTCTTGAGAAACCTCAAATTAAGTGGATTTTCATCCCCA	420
H	GCCTATCTCACTGTCTGGGTGTTCTTGAGAAACCTCAAATTAAGTGGATTTTCATCCCCA	420
RR	GCCTATCTCACTGTCTGGGTGTTCTTGAGAAACCTCAAATTAAGTGGATTTTCATCCCCA	420
*****	*****	*****
XM_025278930.3	GTTATGGAGGGAGACTTGATGCAGCTGACTTGTAACATCTGGTAGTAAACCTGCAGCT	480
H	GTTATGGAGGGAGACTTGATGCAGCTGACTTGTAACATCTGGTAGTAAACCTGCAGCT	480
RR	GTTATGGAGGGAGACTTGATGCAGCTGACTTGTAACATCTGGTAGTAAACCTGCAGCT	480
*****	*****	*****
XM_025278930.3	GATATAAGATGGTTC	495
H	GATATAAGATGGTTC	495
RR	GATATAAGATGGTTC	495
*****	*****	*****

Figure S8. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb|XM_025278930.3| and *CADM2* marker (495-bp) sequences.

XM_025285122.3	TCTCCGCCAGGCTAACCGCTGCCCATCAAGGTGCTGAAGATGCTGACGGCACGGACTGG	60
H	TCTCCGCCAGGCTAACCGCTGCCCATCAAGGTGCTGAAGATGCTGACGGCACGGACTGG	60
RR	TCTCCGCCAGGCTAACCGCTGCCCATCAAGGTGCTGAAGATGCTGACGGCACGGACTGG	60
*****	*****	*****
XM_025285122.3	CCATATTTTGCACCCCTGAGTACCTGCAGCCCCTGCCTTCCACTCCCGTCAGCCCCATAGA	120
H	CCATATTTTGCACCCCTGAGTACCTGCAGCCCCTGCCTTCCACTCCCGTCAGCCCCATAGA	120
RR	CCATATTTTGCACCCCTGAGTACCTGCAGCCCCTGCCTTCCACTCCCGTCAGCCCCATAGA	120
*****	*****	*****
XM_025285122.3	GCTCGATGCCAAGAAAGAGCCCGCTGGCGCTGTTGGCCCAACATGCTCGCAGATCGGGAA	180
H	GCTCGATGCCAAGAAAGAGCCCGCTGGCGCTGTTGGCCCAACATGCTCGCAGATCGGGAA	180
RR	GCTCGATGCCAAGAAAGAGCCCGCTGGCGCTGTTGGCCCAACATGCTCGCAGATCGGGAA	180
*****	*****	*****
XM_025285122.3	GCCCGACCCCTCGCCCTCGTCCAAACTCTCCTCAGTGGCCTCCAAACGGGGGTGGCACGGG	240
H	GCCCGACCCCTCGCCCTCGTCCAAACTCTCCTCAGTGGCCTCCAAACGGGGGTGGCACGGG	240
RR	GCCCGACCCCTCGCCCTCGTCCAAACTCTCCTCAGTGGCCTCCAAACGGGGGTGGCACGGG	240
*****	*****	*****
XM_025285122.3	CGGTGCCGCGCGCGCGCGGGGGCGACAGGACGCCAAGTGGGGCCCTCAAGCTGAG	300
H	CGGTGCCGCGCGCGCGCGGGGGCGACAGGACGCCAAGTGGGGCCCTCAAGCTGAG	300
RR	CGGTGCCGCGCGCGCGCGGGGGCGACAGGACGCCAAGTGGGGCCCTCAAGCTGAG	300
*****	*****	*****
XM_025285122.3	CGACATCGGCGTGAAGACAAGTCGAGTTTCAAGCCGTACTCCAAACCCGGCTCGGATAA	360
H	CGACATCGGCGTGAAGACAAGTCGAGTTTCAAGCCGTACTCCAAACCCGGCTCGGATAA	360
RR	CGACATCGGCGTGAAGACAAGTCGAGTTTCAAGCCGTACTCCAAACCCGGCTCGGATAA	360
*****	*****	*****
XM_025285122.3	G	361
H	G	361
RR	G	361
*****	*	*

Figure S9. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb|XM_025285122.3| and *ZNF503* marker (361-bp) sequences.

```

XM_006048659.4   CCGACGTCGGAGAACCAGAGATGAATCTGAACTCCTCACTGTTCCAGATGGTTGGAAGGAG   60
H_                CCGACGTCGGAGAACCAGAGATGAATCTGAACTCCTCACTGTTCCAGATGGTTGGAAGGAG   60
RR               CCGACGTCGGAGAACCAGAGATGAATCTGAACTCCTCACTGTTCCAGATGGTTGGAAGGAG   60
*****

XM_006048659.4   CCAGCTTTCTCCAAGAGGACAAATCCCAGAGGACTTCTGGAGGAGAGCAGTTTTGCAACT   120
H_                CCAGCTTTCTCCAAGAGGACAAATCCCAGAGGACTTCTGGAGGAGAGCAGTTTTGCAACT   120
RR               CCAGCTTTCTCCAAGAGGACAAATCCCAGAGGACTTCTGGAGGAGAGCAGTTTTGCAACT   120
*****

XM_006048659.4   TTGTTTCCGAARTACAGAGAGGCTTACTTGAAAGAGTGCTGGCCATTGGTACAGAAAGCC   180
H_                TTGTTTCCGAARTACAGAGAGGCTTACTTGAAAGAGTGCTGGCCATTGGTACAGAAAGCC   180
RR               TTGTTTCCGAARTACAGAGAGGCTTACTTGAAAGAGTGCTGGCCATTGGTACAGAAAGCC   180
*****

XM_006048659.4   TTGAATGAACATCATGTTAATGCCACCCTGGACCTGATTGAGGGCAGCATGACTGTCTGC   240
H_                TTGAATGAACATCATGTTAATGCCACCCTGGACCTGATTGAGGGCAGCATGACTGTCTGC   240
RR               TTGAATGAACATCATGTTAATGCCACCCTGGACCTGATTGAGGGCAGCATGACTGTCTGC   240
*****

XM_006048659.4   ACCACCAAGAAAGACATTTGATCCATATATCATCATTAG   278
H_                ACCACCAAGAAAGACATTTGATCCATATATCATCATTAG   278
RR               ACCACCAAGAAAGACATTTGATCCATATATCATCATTAG   278
*****

```

Figure S10. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb|XM_006048659.4| and *KRR1* marker (278-bp) sequences.

```

XM_044946911.2   ATGTGCGACRAGGAGTTCATGTGGGCCCTGAAAAACGGAGACCTAGATGAGGTGAAAGAC   60
H_                ATGTGCGACRAGGAGTTCATGTGGGCCCTGAAAAACGGAGACCTAGATGAGGTGAAAGAC   60
RR               ATGTGCGACRAGGAGTTCATGTGGGCCCTGAAAAACGGAGACCTAGATGAGGTGAAAGAC   60
*****

XM_044946911.2   TATGTGCCAAGGGAGAAGATGTCAACCGGACACTAGAAGGTGGAAGAAAGCCTCTTCAT   120
H_                TATGTGCCAAGGGAGAAGATGTCAACCGGACACTAGAAGGTGGAAGAAAGCCTCTTCAT   120
RR               TATGTGCCAAGGGAGAAGATGTCAACCGGACACTAGAAGGTGGAAGAAAGCCTCTTCAT   120
*****

XM_044946911.2   TATGCAGCAGATTGTGGACAGCTTGAAATCCTGGAATTTCTGCTGCTGAAAGGAGCAGAT   180
H_                TATGCAGCAGATTGTGGACAGCTTGAAATCCTGGAATTTCTGCTGCTGAAAGGAGCAGAT   180
RR               TATGCAGCAGATTGTGGACAGCTTGAAATCCTGGAATTTCTGCTGCTGAAAGGAGCAGAT   180
*****

XM_044946911.2   ATTAATGCTCCAGATAAACATCATATCACACCTCTTCTGTCTGCCGTCTATGAAGGTCAT   240
H_                ATTAATGCTCCAGATAAACATCATATCACACCTCTTCTGTCTGCCGTCTATGAAGGTCAT   240
RR               ATTAATGCTCCAGATAAACATCATATCACACCTCTTCTGTCTGCCGTCTATGAAGGTCAT   240
*****

XM_044946911.2   GTTTCCTGCGTGAATGCTTCTGTG   266
H_                GTTTCCTGCGTGAATGCTTCTGTG   266
RR               GTTTCCTGCGTGAATGCTTCTGTG   266
*****

```

Figure S11. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb|XM_044946911.2| and *MTPN* marker (266-bp) sequences.

G73R. The nucleotide sequence of the *KRR1* gene (278-bp) revealed one synonymous SNP; 56A resulted from 168GA SNP. Two recurrent SNPs in the *MTPN* gene (266-bp) were discovered using DNA sequencing; 45AG and 189TC, resulted in synonymous mutations, 15L and 63A, respectively. Three recurrent non-synonymous SNPs were found in the *IGFBP7* gene (380-bp); 38GC, 107TC, and 161AC changed the amino acids R13P, V36A and Q54P, respectively. Two recurrent synonymous SNPs, 93GA, and 216GA induced amino acids 31L, and 72P were found in the *LEP* gene (360-bp). Four recurrent SNPs in the *ABCC4* gene (355-bp) were discovered using DNA sequencing; one of these, 294CT resulted in

synonymous mutations, 98I. While the non-synonymous mutation brought about by the SNPs 65CT, 109AG and 306GT led to the substitution of the amino acids S22L, R37G, and L102F, respectively

DISCUSSION

Efficiency of investigated reproductive genes as candidates for repeat breeder incidence

For the effective exploitation of disease-resistant cattle or the complete eradication of ill animals, knowledge of the genes, it is necessary to understand the fundamental mutations and interactions that contribute to resistance (Pal & Chakravarty, 2020). The reproductive (*MFSD14A*, *PALMD*, *VPS13B*, *BMP*,

XM_025290001.3	AGCTGGGCGAGACCCGCGATGCAITGCGGCTGCTGCCCGGTGTGCGCCCGCGGCGAGGGCG	60
H	AGCTGGGCGAGACCCGCGATGCAITGCGGCTGCTGCCCGGTGTGCGCCCGCGGCGAGGGCG	60
RR	AGCTGGGCGAGACCCGCGATGCAITGCGGCTGCTGCCCGGTGTGCGCCCGCGGCGAGGGCG	60
XM_025290001.3	AGCCGTGCGGGGTGGCGGCGCCGCGAGGGGCACTGCCGCGCGGTATGGAGTGCCTGA	120
H	AGCCGTGCGGGGTGGCGGCGCCGCGAGGGGCACTGCCGCGCGGTATGGAGTGCCTGA	120
RR	AGCCGTGCGGGGTGGCGGCGCCGCGAGGGGCACTGCCGCGCGGTATGGAGTGCCTGA	120
XM_025290001.3	AGAGCCGCAAGAGGGCGGAAAGGTAAGCCGGGGCAGCAGCAGGCGGCCCGGTTGTGAGCG	180
H	AGAGCCGCAAGAGGGCGGAAAGGTAAGCCGGGGCAGCAGCAGGCGGCCCGGTTGTGAGCG	180
RR	AGAGCCGCAAGAGGGCGGAAAGGTAAGCCGGGGCAGCAGCAGGCGGCCCGGTTGTGAGCG	180
XM_025290001.3	GCGTGTGTGTGTGCAAGAGCCGCTACCCCGTGTGCGGCAGCGAGGTTGCACCTACTCCA	240
H	GCGTGTGTGTGTGCAAGAGCCGCTACCCCGTGTGCGGCAGCGAGGTTGCACCTACTCCA	240
RR	GCGTGTGTGTGTGCAAGAGCCGCTACCCCGTGTGCGGCAGCGAGGTTGCACCTACTCCA	240
XM_025290001.3	GCGGCTGCCAGCTGCCGCGCCGCGAGCCTCAGGGCCGAGAGCCGCGGGGAGAAGGCCATCA	300
H	GCGGCTGCCAGCTGCCGCGCCGCGAGCCTCAGGGCCGAGAGCCGCGGGGAGAAGGCCATCA	300
RR	GCGGCTGCCAGCTGCCGCGCCGCGAGCCTCAGGGCCGAGAGCCGCGGGGAGAAGGCCATCA	300
XM_025290001.3	CCCAGGTCAGCAAGGGCACTGCGAGCAAGGTCCTTCCATCGTGACACCCCCAAGGACA	360
H	CCCAGGTCAGCAAGGGCACTGCGAGCAAGGTCCTTCCATCGTGACACCCCCAAGGACA	360
RR	CCCAGGTCAGCAAGGGCACTGCGAGCAAGGTCCTTCCATCGTGACACCCCCAAGGACA	360
XM_025290001.3	TCTGGAATGTCACTGGTGCC	380
H	TCTGGAATGTCACTGGTGCC	380
RR	TCTGGAATGTCACTGGTGCC	380

Figure S12. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb|XM_025290001.3| and *IGFBP7* marker (380-bp) sequences.

AF387814.1	CAGTCCGTCCTCCAAACAGAGGGTCACTGGTTTGGACTTCATCCCTGGGCTCCACCCT	60
H	CAGTCCGTCCTCCAAACAGAGGGTCACTGGTTTGGACTTCATCCCTGGGCTCCACCCT	60
RR	CAGTCCGTCCTCCAAACAGAGGGTCACTGGTTTGGACTTCATCCCTGGGCTCCACCCT	60
AF387814.1	CTCCTGAGTTTGTCCAAGATGGACCAGACATTGGCGATCTACCAACAGATCCTCACCAGT	120
H	CTCCTGAGTTTGTCCAAGATGGACCAGACATTGGCGATCTACCAACAGATCCTCACCAGT	120
RR	CTCCTGAGTTTGTCCAAGATGGACCAGACATTGGCGATCTACCAACAGATCCTCACCAGT	120
AF387814.1	CTGCCTCCAGAAATGTGGTCCAAATATCTAATGACCTGGAGAACCTCCGGGACCTTCTC	180
H	CTGCCTCCAGAAATGTGGTCCAAATATCTAATGACCTGGAGAACCTCCGGGACCTTCTC	180
RR	CTGCCTCCAGAAATGTGGTCCAAATATCTAATGACCTGGAGAACCTCCGGGACCTTCTC	180
AF387814.1	CACCTGCTGGCCGCTCCAAAGAGCTGCCCCCTGCCGACAGTTCAGGGCCCTGGAGAGTTTG	240
H	CACCTGCTGGCCGCTCCAAAGAGCTGCCCCCTGCCGACAGTTCAGGGCCCTGGAGAGTTTG	240
RR	CACCTGCTGGCCGCTCCAAAGAGCTGCCCCCTGCCGACAGTTCAGGGCCCTGGAGAGTTTG	240
AF387814.1	GAGAGCTTGGGCGTGTCTGGAAGCCTCCCTCTACTCCACCGAGGTGGTGGCCCTGAGC	300
H	GAGAGCTTGGGCGTGTCTGGAAGCCTCCCTCTACTCCACCGAGGTGGTGGCCCTGAGC	300
RR	GAGAGCTTGGGCGTGTCTGGAAGCCTCCCTCTACTCCACCGAGGTGGTGGCCCTGAGC	300
AF387814.1	CGGCTGCAGGGGTCACTACAGGACATGTTGCGGCAGCTGGACCTCAGTCCCTGGGTGCTGA	360
H	CGGCTGCAGGGGTCACTACAGGACATGTTGCGGCAGCTGGACCTCAGTCCCTGGGTGCTGA	360
RR	CGGCTGCAGGGGTCACTACAGGACATGTTGCGGCAGCTGGACCTCAGTCCCTGGGTGCTGA	360

Figure S13. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb|AF387814.1| and *LEP* marker (360-bp) sequences.

MTNRI1A, *TSHR*, *CSGALNACT1*, *CADM2*, *ZNF503*, *KRR1*, *MTPN*, *IGFBP7*, *LEP*, and *ABCC4*) genes in repeat breeder and healthy Italian buffaloes were identified in this study utilizing sequenced amplified PCR products. The findings show that there are differences between the SNPs involving the two categories. The assessed buffaloes had a considerable

nucleotide polymorphism dispersion ($p < 0.01$), according to the Fisher's exact test. It is crucial to emphasize that, when compared to the relevant datasets obtained from GenBank, the variations discovered and materials readily accessible in this context offer fresh information on the markers under consideration.

```

XM_045162483.1 GCAGAAGGCGAAGCTTCTGCTCGCGCCTGTTTGTCTGGTGGCTAAACCCCTTGTTTAAAT 60
H GCAGAAGGCGAAGCTTCTGCTCGCGCCTGTTTGTCTGGTGGCTAAACCCCTTGTTTAAAT 60
RR GCAGAAGGCGAAGCTTCTGCTCGCGCCTGTTTGTCTGGTGGCTAAACCCCTTGTTTAAAT 60
*****
XM_045162483.1 TGATCACAAACCGGAAATTAGAACCAGATGACATGTACTCGGTGCTTCCAGAAGATCGCTC 120
H TGATCACAAACCGGAAATTAGAACCAGATGACATGTACTCGGTGCTTCCAGAAGATCGCTC 120
RR TGATCACAAACCGGAAATTAGAACCAGATGACATGTACTCGGTGCTTCCAGAAGATCGCTC 120
****
XM_045162483.1 CCAGTGCCTTGGAGAAGAGCTGCAAGGGTACTGGGATCARGAAGTTAAGAGAGCCAGAA 180
H CCAGTGCCTTGGAGAAGAGCTGCAAGGGTACTGGGATCARGAAGTTAAGAGAGCCAGAA 180
RR CCAGTGCCTTGGAGAAGAGCTGCAAGGGTACTGGGATCARGAAGTTAAGAGAGCCAGAA 180
*****
XM_045162483.1 GGATGGACAGGAACCCCTCTTTAGTGAAGCAATCGTAAAGTGTTACTGGAAATCCTATTT 240
H GGATGGACAGGAACCCCTCTTTAGTGAAGCAATCGTAAAGTGTTACTGGAAATCCTATTT 240
RR GGATGGACAGGAACCCCTCTTTAGTGAAGCAATCGTAAAGTGTTACTGGAAATCCTATTT 240
*****
XM_045162483.1 GGATGGACAGGAACCCCTCTTTAGTGAAGCAATCGTAAAGTGTTACTGGAAATCCTATTT 300
H GGATGGACAGGAACCCCTCTTTAGTGAAGCAATCGTAAAGTGTTACTGGAAATCCTATTT 300
RR GGATGGACAGGAACCCCTCTTTAGTGAAGCAATCGTAAAGTGTTACTGGAAATCCTATTT 300
*****
XM_045162483.1 AATTTGGGAATGTTTACATTTCTTGAGGAAGGCACCAGAGTAGTTCAACCCATA 355
H AATTTGGGAATGTTTACATTTCTTGAGGAAGGCACCAGAGTAGTTCAACCCATA 355
RR AATTTGGGAATGTTTACATTTCTTGAGGAAGGCACCAGAGTAGTTCAACCCATA 355
*****

```

Figure S14. Analysis of nitrogenous bases matching for DNA in healthy and repeat breeder buffaloes using GenBank gb|XM_045162483.1| and *ABCC4* marker (355-bp) sequences.

In recent studies, unique genes that are specifically related to the prevalence of bovine repeat breeders have been discovered using genome-wide association analysis for our investigated markers (Li et al., 2018; Mahrous et al., 2022), however, no research have yet looked into the relationship between repeat breeder risk and the SNPs in these genes. The gene sequences from the *Bubalus bubalis* employed in our study, which was published in PubMed, are the first to show this connection. The variance of the reproductive (*MFS14A*, *PALMD*, *VPS13B*, *BMP*, *MTNRIA*, *TSHR*, *CSGALNACT1*, *CADM2*, *ZNF503*, *KRR1*, *MTPN*, *IGFBP7*, *LEP*, and *ABCC4*) markers and how they relate to repeat breeder in Italian buffaloes have not, to our knowledge, been the subject of any prior research. However, the candidate gene technique has been used to monitor the validity of bovine fertility features. For instance, Egyptian buffaloes showed a connection between *IGF-I* genetic variations and issues with reproduction (Ramadan et al., 2018). Buffaloes with *MTNRIA* gene polymorphisms had seasonal suppression of fertility (Rangashamaiah et al., 2022). The frequency of repeat breeding in Egyptian buffaloes have been investigated in relation to the polymorphisms of the *LEP*, and *BMP* genes (Mahrous et al., 2022; El-Debaky et al., 2020). Buffalo genetic variations in the *FSHR* and *ESR* genes and were correlated with the frequency of repeat breeders (Fouda et al., 2021). *LHR* gene polymorphisms and Egyptian buffalo fertility was also investigated (Sosa et al., 2016). *CYP19A1*

gene polymorphisms were associated with anoestrus susceptibility and inactive ovaries in water buffaloes when a candidate gene approach to anoestrus was investigated (El-Bayomi et al., 2018; Abbaset al., 2014). Also discovered in Murrah buffaloes were polymorphisms in the *HSP70* gene, which may be related to post-partum anoestrus (Kumar et al., 2017).

The primary source of adaptation and selection is mutation (Chu & Wei, 2019). All of the reproductive markers that were being studied in this situation had exonic region mutations, which led to altered coding DNA sequences in repeat breeder buffaloes compared to healthy ones. Using DNA sequencing of reproductive genes, 30 SNPs were found; 16 of them are synonymous, and 15 are non-synonymous. Non-synonymous mutations alter protein sequences, and people who have these mutations are frequently the subject of natural selection (Chu, & Wei, 2019). The encoded amino acid at the mutant location is altered by genetic variation brought on by non-synonymous SNPs, which can result in structural and functional alterations in the mutated protein (Dakal et al., 2017). For a very long time, it was believed that selection on synonymous mutations was either nonexistent or very weak (Chu & Wei, 2019). In order to accurately characterize the investigated reproductive genes at the molecular level and to understand the physiological differences in reproductive performance between normal and repeat breeder buffaloes, our study discovered polymorphisms on the basis

Table 2. Single base differential dissemination as well as category of inherited alteration for reproductive markers in normal and repeat breeder buffaloes.

Gene	SNPs	Normal n = 120	Repeat breeder n = 120	Total n = 240	kind of inherited alteration	Amino acid order and sort
<i>MFSD14A</i>	C165T	-	72	72/240	Synonymous	55 C
	T322C	-	85	85/240	Synonymous	108 G
<i>PALMD</i>	T108C	91	-	91/240	Synonymous	36 L
<i>VPS13B</i>	T175C	48	-	48/240	Non-synonymous	59 C to R
	T223C	-	63	63/240	Non-synonymous	75 Y to H
	T47C	-	98	98/240	Non-synonymous	16 V to A
<i>BMP</i>	C80A	58	-	58/240	Non-synonymous	27 P to Q
	G184A	76	-	76/240	Non-synonymous	62 V to I
<i>MTNRI1A</i>	C72T	49	-	49/240	Synonymous	24 N
	C87T	-	83	83/240	Synonymous	29 S
<i>TSHR</i>	T126C	-	38	38/240	Synonymous	42 T
	C177T	69	-	69/240	Synonymous	59 I
	C372T	-	103	103/240	Synonymous	124 V
<i>CSGALNACT1</i>	G135A	47	-	47/240	Synonymous	45 T
	G150A	91	-	91/240	Synonymous	50 R
<i>CADM2</i>	T168C	63	-	63/240	Synonymous	56 N
	G354T	39	-	39/240	Synonymous	118 T
<i>ZNF503</i>	G217C	-	98	98/240	Non-synonymous	73 G to R
<i>KRR1</i>	G168A	107	-	107/240	Synonymous	56 A
<i>MTPN</i>	A45G	46	-	46/240	Synonymous	15 L
	T189C	-	104	104/240	Synonymous	63 A
	G38C	-	58	58/240	Non-synonymous	13 R to P
<i>IGFBP7</i>	T107C	-	112	112/240	Non-synonymous	36 V to A
	A161C	92	-	92/240	Non-synonymous	54 Q to P
<i>LEP</i>	G93A	71	-	71/240	Synonymous	31 L
	G216A	-	86	86/240	Synonymous	72 P
	C65T	-	89	89/240	Non-synonymous	22 S to L
<i>ABCC4</i>	A109G	-	109	109/240	Non-synonymous	37 R to G
	C294T	-	74	74/240	Synonymous	98 I
	G306T	93	-	93/240	Non-synonymous	102 L to F

Single base difference dispersal for reproductive markers in normal and repeat breeder buffaloes showed a highly significant variation ($p < 0.01$) according to Fisher's exact analysis.

MFSD14A= Major facilitator superfamily domain containing 14A; *PALMD*= Palmdelphin; *VPS13B*= Vacuolar protein sorting 13 homolog B; *BMP*= Bone morphogenic protein; *MTNRI1A*= Melatonin receptor 1B; *TSHR*= Thyroid stimulating hormone receptor; *CSGALNACT1*= Chondroitin sulfate N-acetylgalactosaminyltransferase 1; *CADM2*= Cell adhesion molecule 2; *ZNF503*= Zinc finger protein 503; *KRR1* = Small subunit processome component homolog; *MTPN*= Myotrophin; *IGFBP7*= insulin like growth factor binding protein 7; *LEP*= Leptin; and *ABCC4*= ATP-binding cassette sub-family C member 4. A = Alanine; C = Cysteine; F = Phenylalanine; G = Glycine; H = Histidine; I = Isoleucine; L = Leucine; N = Asparagine; P = Proline; Q = Glutamine; R = Arginine; S = Serine; T = Threonine; V = Valine; and Y = Tyrosine.

of translated DNA sequence to be of greater value than intronic parts.

Role of investigated genes in reproduction

A protein-coding gene called major facilitator superfamily domain containing 14A (*MFSD14A*) has product homology to the family of solute carrier proteins (Doran et al., 2016). Additionally, *MFSD14A* functions as a component of cellular fundamentals, a molecular controller of transporter action, and a biological tool involved in trans-membrane transport (Sreedharan et al., 2011). The nuclear transfer transcripts in pig blastocyst stage were discovered to be negatively controlled by *MFSD14A* (Whitworth, et al., 2011), and the gene may also play a role in mouse spermatogenesis (Doran, 2016). Two SNPs in the *MFSD14A* gene have been linked to characteristics of buffalo milk production (Liu et al., 2018). A cytosolic protein involved in the phosphorylation of p53 is encoded by the palmdelphin (*PALMD*) (Moioli et al., 2013). The variance in milk yield is likely to be caused by the *PALMD* gene (Seo et al., 2016; Moioli, et al., 2013). *PALMD* polymorphisms had a significant impact on back fat thickness in a research to identify probable causative mutations associated to pig production attributes (Martínez-Montes et al., 2017). These *PALMD* functional studies suggest that it may be a key target gene that regulates variables related to reproductive performance.

The eye, hematological system, and central nervous system all develop and function as a result of the potential trans-membrane protein that the vacuolar protein sorting 13 homolog B (*VPS13B*) gene encodes (Liu et al., 2018). Protein transport has been identified as the primary biological utility of *VPS13B*, and it may play a role in vesicle-mediated transport as well as protein sorting within cells. In a prior investigation, *VPS13B* was discovered inside a QTL linked to the shape of the legs in dairy cattle (Van den Berg et al., 2014). Additionally, it was hypothesized that attributes related to milk production in Holstein cattle as well as female fertility were linked to the genomic region on BTA14 encompassing *VPS13B* (Capitan et al., 2014).

The secreted factor known as transforming growth factor (TGF) was initially discovered to influence cell proliferation (Roberts & Sporn, 1990). The proteins can be divided into two main groups according to cell signaling pathways: activins/transforming growth factors and bone morphogenetic proteins/growth differentiation factors (Chang et al., 2002). Among the BMP family memberships expressed in-

side the uterus, ovary, granulosa cells, and oocyte are BMP 2, 3, 4, 5, 6, 7 and 15 (Tanwar& McFarlane, 2011; Sun et al., 2010; Otsuka et al., 2001). They provide significant biological tasks as controllers of ovarian follicle growth, female reproductive tract discrepancy, blastocyst embedding in the uterus, and organogenesis and morphogenesis during embryo expansion (Kishigami & Mishina, 2005). The *BMP* gene has been the subject of numerous investigations in mammals, including cattle (Baloza et al., 2017), sheep (Ibrahim, 2019), and goats (Latifah & Saa, 2021; Ortiz et al., 2015). The blastocyst rate was shown to be substantially correlated with *BMP* SNPs by Lari et al., (2012).

Photoperiod and melatonin have been identified as the hypothalamus level modulators of the reproductive axis in a number of ovine investigations (Karsch et al., 2013). The pineal gland produces melatonin hormone in direct proportion to the length of darkness (Malpaux et al., 2001). The nighttime plasma melatonin content was highest in winter and lowest in summer in the Mediterranean buffaloes that had a seasonal reproductive tendency (Parmeggianiet al., 1994). Subcutaneous insertion of sustained-release melatonin implants in buffaloes during summer anestrus lengthens the daily presence of melatonin, simulating the effect of short days, and has been extensively investigated for the activation of the reproductive axis (Kavita et al., 2018). Only the melatonin receptor 1A gene (*MTNR1A*) is connected to reproductive seasonality of the G-protein couple receptors melatonin receptor 1A gene (*MTNR1A*) and melatonin receptor 1B gene (*MTNR-RIB*), to which melatonin binds (Dubocovich et al., 2020).

Animal reproduction research have used hormones and antibiotics as therapies because hormones are crucial to animal reproduction (Refsdal, 2000). Seasonal cycles of body weight and reproduction are significantly influenced by thyroid hormones (Barrett et al., 2007). By chance, buffalo are a species that go into estrus seasonally. Thyroid hormone resistance (THR) has also been shown to be highly correlated with the THR mutation (Işik et al., 2013). The genes involved in the metabolic pathways that impact animal fertility have already been extensively studied (Morfeld & Brown, 2014). One of these genes, *CSGALNACT1*, is necessary for healthy cartilage development (Watanabe et al., 2010). The *CSGALNACT1* gene was hypothesized to affect bovine follicle growth by altering the degree of expression that controls glucose metabolism (Li et al., 2018). Buffalo reproductive efficiency was discovered to be

correlated with cell adhesion molecule 2 (*CADM2*) (Song et al., 2013). One SNP at the bovine *CADM2* gene was connected to the number of days between conception and the first calving as well as the first and second calving (Li et al., 2018).

The target genes for buffalo reproductive features include *ZNF503*, *KRR1*, and *MTPN* (Lappalainen et al., 2013). *ZNF503* enhances cell invasion and proliferation in mammary epithelial cells, which has a key role in embryogenesis (Shahi et al., 2015; Chang et al., 2013). *KRR1* has been linked to polycystic ovarian syndrome (PCOS) in European populations (Day et al., 2015). *MTPN* is crucial for the development of cells and skeletal muscles (Makinaet al., 2015) and is associated with the process of antigen recognition, a crucial step in the immune reaction (Wang & Wang, 2012). The elevated expression of these three genes in buffalo granulosa cells raises the possibility that they took part in dominant follicle selection (Li et al., 2018). *IGFBP7* and follistatin share sequence similarity (Kato, 2000), the latter of which was once thought to be an inhibitor of FSH secretion (Jorgezet et al., 2004) and is essential for ovarian and folliculogenesis (Muttukrishna et al., 2004). Additionally, *IGFBP7* was discovered to be expressed in the granulosa cells of the large antral follicles of the pig ovary (Wandji et al., 2002) and the bovine corpus luteum (Casey et al., 2004). Therefore, we assume that *IGFBP7* is in some way responsible for the regulation of follicle development and ovulation (Li et al., 2018).

The versatile peptide hormone leptin, which is mostly generated by adipocytes, plays a crucial role in reproduction alongside controlling energy expenditure and body weight (Wiles et al., 2014). Leptin and its receptors have been discovered to be expressed in buffalo ovaries. Contrary to the widely held idea that leptin primarily affects the neuroendocrine component of reproduction, this clearly demonstrates that leptin directly engages in ovarian activity (Reshma et al., 2016). According to Fu et al., (2016), the ovary has leptin receptors that can control steroidogenesis and improve the oocyte's ability to support later embryonic development. The *ABCC4* gene was linked to the number of services per conception (NSC). Since it affects prostaglandin efflux from cells, this gene appears to be crucial to support pregnancy in the endometrium of pregnant cows (Spencer et al., 2013) and pigs (Seo et al., 2014). Prostaglandin plays a number of roles in re-

production, including conceptus implantation and maternal recognition of pregnancy—processes that are intimately connected to NSC. The expression of *ABCC4* was substantially linked with residual feed intake (RFI) in Angus cattle and was up-regulated in animals with high RFI (Chen et al., 2012). Nucleotide sequence variants in *ABCC4* was linked to marbling score in Nelore cattle (Feitosa et al., 2014). The recently proposed theory is that *ABCC4*, which is highly polymorphic and acts in fundamental metabolic pathways, may have effects on reproductive and meat quality (De Camargo et al., 2015).

CONCLUSIONS

Using PCR-DNA sequencing, single nucleotide variations (SNPs) in the genes were found for reproductive (*MFSD14A*, *PALMD*, *VPS13B*, *BMP*, *MTNR1A*, *TSHR*, *CSGALNACT1*, *CADM2*, *ZNF503*, *KRR1*, *MTPN*, *IGFBP7*, *LEP*, and *ABCC4*) indicators in normal and repeat breeder Italian buffaloes. These special functional variations present a promising opportunity to lower the occurrence of repeat breeders by using genetic markers in conjunction with normal wellbeing during buffalo selection. Future approaches to dealing with repeat breeders may be facilitated based on the gene domains here.

FUNDING

Princess Nourah bint Abdulrahman University Researchers Supporting Project number (PNUR-SP2025R318), Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia, funds the publishing of this study. This research was also funded by the Researchers Supporting Project number (RSP-2024/R232), King Saud University, Riyadh, Saudi Arabia.

ACKNOWLEDGMENTS

Princess Nourah bint Abdulrahman University Researchers Supporting Project number (PNUR-SP2025R318), Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia acknowledged by the authors. The authors sincerely acknowledge the Researchers Supporting Project for funding this work number (RSP-2024/R232) at King Saud University, Riyadh, Saudi Arabia.

CONFLICT OF INTEREST

There are no competing interests, according to the authors.

REFERENCES

- Abbas, H. S., El-Magd, M. A., El-Bayomi, K. M. & Sosa, G. A. (2014). Detection of SNPs in exon7 locus of Cyp19 gene and their association with anestrus in Egyptian buffaloes (Bubalus bubalis). *Benha Veterinary Medical Journal*, 26, 151–160.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, 215, 403–410.
- Ashwell, M.S., Rexroad Jr, C. E., Miller, R. H., Van Raden, P.M. & Da, Y. (1997). Detection of loci affecting milk production and health traits in an elite US Holstein population using microsatellite markers. *Animal Genetics*, 28, 216–222.
- Baloza, S. H., Abo-Salem, M. E. S. & Hemeda, S. A. (2017). Effect of genetic variations in BMP 4 gene (exon 2 plus part of intron 2) on infertility in Egyptian Buffalo cows. *Benha Journal of Applied Sciences*, 21, 37–41.
- Barrett, P., Ebling, F. J. P., Schuhler, S., Wilson, D., Ross, A.W., Warner, A., Jethwa, P., Boelen, A., Visser, T. J., Ozanne, D. M. (2007). Hypothalamic thyroid hormone catabolism acts as a gatekeeper for the seasonal control of body weight and reproduction. *Endocrinology*, 148, 3608–17.
- Berry, D.P. & Evans, R. (2014). Genetics of reproductive performance in seasonal calving beef cows and its association with performance traits. *Journal of Animal Science*, 92, 1412–1422.
- Bishop, M. D., Hawkins, G.A. & Keeler, C.L. (1995). Use of DNA markers in animal selection. *Theriogenology*, 43, 61–70.
- Boesenberg-Smith, K. A., Pessaraki, M. M. & Wolk, D. M. (2012). Assessment of DNA Yield and Purity: An Overlooked Detail of PCR Troubleshooting. *Clinical Microbiology Newsletter*, 34, 1–6.
- Boichard, D. (1990). Estimation of the economic value of conception rate in dairy cattle. *Livestock Production Science Journal*, 24, 187–204.
- Boom, R., Sol, C. J., Salimans, M. M. & Jansen, C. L. (1990). Wertheim-van Dillen PM, Noordaa JVD. Rapid and simple method for purification of nucleic acids. *Journal of Clinical Microbiology*, 28, 495–503.
- Buch, L.H., Kargo, M., Berg, P., Lassen, J. & Sørensen A. C. (2012). The value of cows in reference populations for genomic selection of new functional traits. *Animals*, 6, 880–886.
- Cammack, K., Thomas, M. & Enns, R. (2009). Reproductive traits and their heritabilities in beef cattle. *The Professional Animal Scientist Journal*, 25, 517–528.
- Capitan, A., Michot, P., Baur, A., Saintilan, R., Hoze, C., Valour, D., Guillaume, F., Boichon, D., Barbat, A. & Boichard, D. (2014). Genetic tools to improve reproduction traits in dairy cattle. *Reproduction Fertility Development*, 27, 14–21.
- Casey, O. M., Fitzpatrick, R., Mcinerney, J. O., Morris, D. G., Powell, R., & Sreenan, J. M. (2004). Analysis of gene expression in the bovine corpus luteum through generation and characterisation of 960 ESTs. *Biochimica et Biophysica Acta - Gene Structure and Expression*, 1679, 10–7
- Chang, H., Brown, C.B. & Matzuk, M.M. (2002). Genetic analysis of the Mammalian transforming growth factor superfamily. *Endocrinology Reviews*, 23, 787–823.
- Chang, S. L., Liu, Y. C., Chen, S. Y., Huang, T. H., Liu, P. T. & Liu, F. C. (2013). Identification of two evolutionarily conserved 5' cis-elements involved in regulating spatiotemporal expression of Nolz-1 during mouse embryogenesis. *PLOS One*, 8, e54485.
- Chen, Y., Arthur, P. F., Barchia, I. M., Quinn, K., Parnell, P. F., Herd, R. M. (2012). Using gene expression information obtained by quantitative real-time PCR to evaluate Angus bulls divergently selected for feed efficiency. *Animal Production Science*, 52, 1058–67.
- Chu, D. & Wei, L. (2019). Nonsynonymous, synonymous and nonsense mutations in human cancer-related genes undergo stronger purifying selections than expectation. *BMC Cancer*, 19, 359.
- Dakal, T.C., Kala, D., Dhiman, G., Yadav, V., Krokhotin, A. & Dokholyan, N.V. (2017). Predicting the functional consequences of non-synonymous single nucleotide polymorphisms in IL8 gene. *Scientific Reports*, 7, 6525.
- Das, G.K. & Khan, F.A. (2010). Summer anestrus in buffalo-A review. *Reproduction in Domestic Animals*, 45:e483-e494.
- Day, F. R., Hinds, D. A., Tung, J. Y., Stolk, L., Styrkarsdottir, U., Saxena, R., Bjonnes, A., Broer, L., Dunger, D. B. & Halldorsson, B. V. et al. (2015). Causal mechanisms and balancing selection inferred from genetic associations with polycystic ovary syndrome. *Nature Communication*, 6, 8464.
- De Camargo, G. M., Aspilueta-Borquis, R. R., Fortes, M. R., Porto-Neto, R., Cardoso, D. F., Santos, D. J., Lehnert, S. A., Reverter, A., Moore, S. S. & Tonhati, H. (2015). Prospecting major genes in dairy buffaloes. *BMC Genomics*, 16, 872.
- Dekkers, J. (1991). Estimation of economic values for dairy cattle breeding goals: bias due to sub-optimal management policies. *Livestock Production Science Journal*, 29, 131–149.
- Devkota, B., Shah, S. & Gautam, G. (2022). Reproduction and Fertility of Buffaloes in Nepal. *Animals*, 13, 70.
- Doran, J., Walters, C., Kyle, V., Wooding, P., Hammett-Burke, R. & Colledge, W.H. (2016). Mfsd14a (Hiat1) gene disruption causes globozoospermia and infertility in male mice. *Reproduction*, 152, 91–99.
- Dubocovich, M. L., Cardinali, D. P., Delagrange, P. R. M., Krause, D. N., Strosberg, D. & Sugden, D. (2000). Melatonin Receptors. In Girdlestone, D. (ed.) *The IUPHAR Compendium of Receptor Characterization and Classification*, 2nd ed. London, UK. 270–277.
- El-Bayomi, K. M., Saleh, A. A., Awad, A., El-Tarabany, M. S., El-Qalioubi, H. S., Afifi, M., El-Komy, S., Essawi, W. M., Almadaly, E.A. & El-Magd, M. A. (2018). Association of CYP19A1 gene polymorphisms with anoestrus in water buffaloes. *Reproduction and Fertility Development*, 30, 487–497.
- El-Debaky, H., Mahmoud, K. M., Abd El-Razik, K. A., Sosa, A. S. A., Kandiel, M. M. M. & Ahmed, Y. F. (2020). PCR-SSCP and Sequencing Analysis for Studying Leptin Gene Polymorphism and Its Association with Reproductive Status of Egyptian Buffalo. *Egyptian Journal of Veterinary Science*, 51, 11–21.
- Feitosa, F. L. B., Pereira, A. S. C., Venturini, G. C., Tonussi, R. L., Espigolan, R., & Gordo, D. M. et al. (2014). Genome wide association study between copy number variation regions with marbling score in Nelore cattle. In: *Proceedings, 10th World Congress of Genetics Applied to Livestock Production*. Vancouver.
- Fouda, M., Hemeda, S., El-Bayomi, K., El-Araby, I., Hendam, B. & Ateya, A. (2021). Genetic polymorphisms in FSHR/ALUI and ESRα /BG11 loci and their association with repeat breeder incidence in buffalo. *Journal of the Hellenic Veterinary Medical Society*, 72, 2869–2878.
- Fu, Y.F., Li, L., Li, B., Fang, X. & Ren, S. (2016). Long form leptin receptor and SNP effect on reproductive traits during embryo attachment in Suzhong sows. *Animal Reproduction Science*, 168, 57–65.
- Heidari, M., Kafi, M., Mirzaei, A., Asaadi, A. & Mokhtari, A. (2019). Effects of follicular fluid of preovulatory follicles of repeat breeder dairy cows with subclinical endometritis on oocyte developmental competence. *Animal Reproduction Science*, 205, 62–69.
- Ibrahim, A. H. M. (2019). Association of growth performance and body conformational traits with BMP4 gene variation in Barki lambs. *Growth Factors*, 37, 153–163.
- Işık, E., Beck, P. P., Campi, I., Özon, A., Alikışıfoğlu, A., Gönc, N. & Kandemir N. (2013). Thyroid hormone resistance: a novel mutation in thyroid hormone receptor beta (THRB) gene - case report. *Turkish Journal of Pediatrics*, 55, 322–7
- Jorgez, C. J., Klysis, M., Jamin, S. P., Behringer, R. R., & Matzuk, M. M. (2004). Granulosa cell specific inactivation of follistatin causes female fertility defects. *Molecular Endocrinology*, 18, 953–67.
- Karsch, F. J., Bittman, E. L., Foster, D. L., Goodman, R. L., Legan, S. J. & Robinson, J. E. (2013). *Neuroendocrine basis of seasonal reproduction*. Recent Progress in Hormone Research, Academic Press, New York, USA.
- Kato, M. V. (2000). A secreted tumor-suppressor, mac25, with activin-binding activity. *Molecular Medicine*, 6, 126–35
- Kavita, K., Phogat, J. B., Pandey, A. K., Balhara, A. K., Ghuman, S. S. & Gunwant, P. (2018). Effects of melatonin supplementation prior to Ovsynch protocol on ovarian activity and conception rates in anestrus Murrah buffalo heifers during out of breeding season. *Reproductive Biology*, 18, 161–168.
- Kishigami, S. & Mishina, Y. (2005). BMP signaling and early embryonic patterning. *Cytokine Growth Factor Reviews*, 16, 265–278.

- Kukurba, K. R. & Montgomery, S. B. (2015). RNA Sequencing and Analysis. *Cold Spring Harbor Protocols*, 13, 951-69.
- Kumar, R., Ghosh, M., Kumar, N., Balhara, A. K., Gupta, M., Sharma, R. K. & Singh, I. (2017). Polymorphism in 5' untranslated region of heat-shock protein 70 gene as marker of post-partum anoestrus in Murrah buffaloes. *Reproduction Domestic Animal*, 52, 505-512.
- Lappalainen, T., Sammeth, M., Friedlander, M. R., Hoen, P. A., Monlong, J., Rivas, M. A., Gonzalez-Porta, M., Kurbatova, N., Griebel, T. & Ferreira, P. G. et al. (2013). Transcriptome and genome sequencing uncovers functional variation in humans. *Nature*, 501, 506-11.
- Lappalainen, T., Sammeth, M., Friedlander, M. R., Hoen, P. A., Monlong, J., Rivas, M. A., Gonzalez-Porta, M., Kurbatova, N., Griebel, T. & Ferreira, P. G. et al. (2013). Transcriptome and genome sequencing uncovers functional variation in humans. *Nature*, 501, 506-11.
- Lari, A. M., Mohebbi-Fani, M. & Rowshan-Ghasrodashti, A. (2012). Causes of culling in dairy cows and its relation to age at culling and interval from calving in Shiraz, Southern Iran. *Veterinary Research Forum*, 3, 233-237.
- Latifah, L. & Saa N.R. (2021). Single nucleotide Polymorphism gen BMP4 pada kambingberdasarakan data GenBank. *Journal of Tropical Animal Research*, 2, 1, 24-29
- Li, J., Liu, J., Campanile, G., Plastow, G., Zhang, C., Wang, Z., Cassandro, M., Gasparri, B., Salzano, A., Hua, G., Liang, A. & Yang, L. (2018). Novel insights into the genetic basis of buffalo reproductive performance. *BMC Genomics*, 9, 814.
- Li, J., Liu, X., Yang, J., Wang, H., Jiang, J., Liu, L., He, S., Ding, X., Liu, J. & Zhang, Q. (2014). Targeted resequencing of GWAS loci reveals novel genetic variants for milk production traits. *BMC Genomics*, 15, 1-9.
- Liu, J. J., Liang, A. X., Campanile, G., Plastow, G., Zhang, C., Wang, Z., Salzano, A., Gasparri, B., Cassandro, M. & Yang, L. G. (2018). Genome-wide association studies to identify quantitative trait loci affecting milk production traits in water buffalo. *Journal of Dairy Science*, 101, 433-444.
- Mahrous, K. F., El-Kader, H. A. M., Abdelhafez, M. A., Aboelenin, M. M., Balabel, E. A., Mabrouk, D. M., El Malky, O. M. & Hassanian, M. S. (2022). Genetic structure of some candidate genes of repeat breeder syndrome in Egyptian buffaloes. *Journal of genetic engineering and biotechnology*, 20, 110.
- Makina, S. O., Muchadeyi, F. C., Van Marle-Koster, E., Taylor, J. F., Makgahlela, M. L., & Maiwashe, A. (2015). Genome-wide scan for selection signatures in six cattle breeds in South Africa. *Genetics Selection Evolution*, 47, 92.
- Malpaux, B., Migaud, M., Tricoire, H. & Chemineau, P. (2001). Biology of mammalian photoperiodism and the critical role of the pineal gland and melatonin. *Journal of Biological Rhythms*, 16, 336-347.
- Martínez-Montes, A. M., Fernandez, A., Pérez-Montarelo, D., Alves, D., Benitez, R., Nunez, Y., Ovilo, C., Ibañez-Escriche, N., Folch, J. & Fernandez, A. (2017). Using RNA-Seq SNP data to reveal potential causal mutations related to pig production traits and RNA editing. *Animal Genetics*, 48, 151-165.
- Moioli, B., Scatà, M. C., Steri, R., Napolitano, F. & Catillo, G. (2013). Signatures of selection identify loci associated with milk yield in sheep. *BMC Genetics*, 14, 76.
- Moioli, B., Scatà, M., Catillo, G., Napolitano, F. & Pilla, F. (2013). Whole-genome genotyping for detecting mutations in the genes that are directly responsible for the trait variation. Page P0621 in Proc. Plant and Animal Genome XXI, San Diego, CA.
- Morfeld, K. A. & Brown, J. L. (2014). Ovarian acyclicity in zoo African elephants (*Loxodonta africana*) is associated with high body condition scores and elevated serum insulin and leptin. *Reproduction, Fertility and Development*, 28, 640-7.
- Musunuru, K., Strong, A., Frank-Kamenetsky, M., Lee, N. E., Ahfeldt, T., Sachs, K.V., Li, X., Li, H., Kuperwasser, N., & Ruda, V. M. (2010). From noncoding variant to phenotype via SORT1 at the 1p13 cholesterol locus. *Nature*, 466, 714-9.
- Muttukrishna, S., Tannetta, D., Groome, N., & Sargent I. (2004). Activin and follistatin in female reproduction. *Molecular and Cellular Endocrinology*, 225, 45-56.
- Ortiz, W. H., Quirino, C. R., Silva, A., Oliveira, C. S., Serapiao, R. V., Pacheco, A. & Bartholazzi, A. (2015). Association between BMP4 gene polymorphism and in vitro embryo production traits in Gyr cows. *Revista Colombiana de CienciasPecuarias*, 28, 156-164.
- Otsuka, F., Yamamoto, S., Erickson, G. F. & Shimasaki, S. (2001). Bonemorphogenetic protein-15 inhibits follicle stimulating hormone (FSH) action by suppressing FSH receptor expression. *Journal of Biological Chemistry*, 276, 11387-11392
- Pal, A. & Chakravarty, A. K. (2020). Disease resistance for different livestock species. *Genetics and Breeding for Disease Resistance of Livestock*, 19, 271-96.
- Parmeggiani, A., Di Palo, R., Zicarelli, L., Campanille, G., Esposito, L., Seren, E., Accorsi, P. A. & Sofai, S. M. (1994). Melatonin and reproductive seasonality in the buffalo cow. *Agricoltura Ricerca*, 153, 41-48
- Ramadan, S. I., Abo-Gamil, Z. H. & El-Qaliouby, H. S. (2018). Polymorphism of IGF-1 and Its Receptor (IGF-1R) Genes and Their Association with Fertility Problems of Egyptian Buffaloes. *Alexandria Journal of Veterinary Sciences*, 59, 119-124.
- Rangashamaiah, M. M., Ghuman, S. S. & Cheema, R. (2022). Association of melatonin receptor 1a (MTNR1A) gene polymorphism with seasonal suppression of fertility in buffaloes. *Buffalo Bulletin*, 41, 749-763.
- Refsdal, A.O. (2000). To treat or not to treat: a proper use of hormones and antibiotics. *Animal Reproduction Science*, 60, 109-19.
- Regmi, G. & Dhakal, I.P. (2020). Systemic levels of iron, phosphorus, and total protein in normocytic versus repeat breeder Holstein Friesian crossbred cows of Kesharbag, Chitwan. *Nepalese Veterinary Journal*, 13, 2353-2357.
- Reshma, R., Mishra, S. R., Thakur, N., Parmar, M. S., Somal, A., Bharti, M. K., Pandey, S., Chandr, V., Chouhan, V. S., Verma, M. R., Singh, G., Sharm, G. T., Maurya, V. P. & Sarkara, M. (2016). Modulatory role of leptin on ovarian functions in water buffalo (*Bubalus bubalis*). *Theriogenology*, 86, 1720-1739.
- Roberts, A.B. & Sporn, M. B. (1990). The transforming growth factors. In: Sporn, M.B., Roberts, A.B. (Eds.), *Peptide Growth Factors and Their Receptors*, Part I. *Springer-Verlag Berlin*, 419-472.
- Roxström, A. & Strandberg, E. (2002). Genetic analysis of functional, fertility-, mastitis-, and production-determined length of productive life in Swedish dairy cattle. *Livestock Production Science Journal*, 125-135.
- Sammad, A., Khan, M. Z., Abbas, Z., Hu, L., Ullah, Q., Wang, Y., Zhu, H. & Wang, Y. (2022). Major Nutritional Metabolic Alterations Influencing the Reproductive System of Postpartum Dairy Cows. *Metabolites*, 12, 60.
- Sanger, F., Nicklen, S. & Coulson, A. R. (1997). DNA sequencing with chain-terminating inhibitors. *The Proceedings of the National Academy of Sciences, USA*, 74, 5463-5467.
- Schaeffer, L. R. (2006). Strategy for applying genome-wide selection in dairy cattle. *Journal of Animal Breeding and Genetics*, 123, 218-223.
- Seo, H., Choi, Y., Shim, J., Yoo, I., & Ka, H. (2014). Prostaglandin Transporters ABCC4 and SLCO2A1 in the Uterine Endometrium and Conceptus During Pregnancy in Pigs. *Biology of Reproduction*, 90, 5.
- Seo, M., Lee, H. J., Kim, K., Caetano-Anolles, K., Jeong, J.Y., Park, S., Oh, Y. K., Cho, S. & Kim, H. (2016). Characterizing milk production related genes in Holstein using RNA-seq. *Asian-australas. Journal of Animal Science*, 29, 343.
- Shahi, P., Slorach E.M., Wang, C.Y., Chou, J., Lu, A., Ruderisch, A. & Werb, Z. (2015). The transcriptional repressor ZNF503/Zeppo2 promotes mammary epithelial cell proliferation and enhances cell invasion. *Journal Biological Chemistry*, 290, 3803-13.
- Shao, B., Sun, H., Ahmad, M. J., Ghanem, N., Abdel-Shafy, H., Du, C., Deng, T., Mansoor, S., Zhou, Y., Yang, Y., Zhang, S., Yang, L. & Hua, G. (2021). Genetic features of reproductive traits in bovine and Buffalo: lessons from bovine to Buffalo. *Frontiers in Genetics*, 12, 617128.
- Song, L. J., Wu, J. F., Yang, Z. B., Guo, L., Zhang S. J. & Wu, J.J. (2013). Investigation of transferability of BovineSNP50 BeadChip from cattle to water buffalo for genome wide association study. *Molecular Biology Reports*, 40, 743-50.
- Sosa, A. S. A., Mahmoud, K. M., Kandiel, M. M. M., Eldebaky, H. A. A., Nawito, M. F. & Abou El Roos, M. E. A. (2016). Genetic Polymorphism of Luteinizing Hormone Receptor Gene in Relation to Fertility of Egyptian Buffalo. *Biotechnology Journal*, 12, 1-11.
- Spencer, T. E., Forde, N., Dorniak, P., Hansen, T. R., Romero, J. J. & Lonergan, P. (2013). Conceptus-derived prostaglandins regulate gene expression in the endometrium prior to pregnancy recognition in ruminants. *Reproduction*, 146, 377-87.

- Sreedharan, S., Stephansson, O., Schiöth, H. B. & Fredriksson, R. (2011). Long evolutionary conservation and considerable tissue specificity of several atypical solute carrier transporters. *Gene*, 478, 11–18.
- Sun, R. Z., Lei, L., Cheng, L., Jin, Z. F., Zu, S. J. & Shan, Z. Y. (2010). Expression of GDF-9, BMP-15 and their receptors in mammalian ovary follicles. *Journal of Molecular Histology*, 41, 325–32.
- Tamura, K., Dudley, J., Nei, M. & Kumar, S. (2007). MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) Software Version 4.0. *Molecular Biology Evolution*, 24, 1596–1599.
- Tanwar, P.S. & McFarlane, J.R. (2011). Dynamic expression of bone morphogenetic protein 4 in reproductive organs of female mice. *Reproduction*, 142, 573–9.
- Van den Berg, I., Fritz, S., Rodriguez, S., Rocha, D., Boussaha, M., Lund, M. S. & Boichard, D. (2014). Concordance analysis for QTL detection in dairy cattle: A case study of leg morphology. *Genetics Selection Evolution*, 46, 31.
- Wandji, S. A., Gadsby, J. E., Barber, J. A., & Hammond, J. M. (2000). Messenger ribonucleic acids for IGFBP-7 and connective tissue growth factor (CTGF) are inversely regulated during folliculogenesis and early luteogenesis. *Journal of Endocrinology*, 141, 2648–57.
- Wang, L. & Wang, Y. (2012). Molecular characterization, expression patterns and subcellular localization of Myotrophin (MTPN) gene in porcine skeletal muscle. *Molecular Biology Reproduction*, 39, 2733–8.
- Warriach, H. M., McGill, D.M., Bush, R.D., Wynn, P.C. & Chohan, K.R. (2015). A review of recent developments in buffalo reproduction - a review. *Asian-Australasian Journal of Animal Sciences*, 28, 451–455.
- Watanabe, Y., Takeuchi, K., Onaga, S. H., Sato, M., Tsujita, M., Abe, M., Natsume, R., Li, M., Furuichi, T. & Saeki, M. (2010). Chondroitin sulfate N-acetylgalactosaminyltransferase-1 is required for normal cartilage development. *Biochemistry Journal*, 432, 47–55.
- Whitworth, K. M., Zhao, J., Spate, L. D., Li, R. & Prather, R. S. (2011). Scriptaid corrects gene expression of a few aberrantly reprogrammed transcripts in nuclear transfer pig blastocyst stage embryos. *Cellular Reprogramming*, 13, 191–204.
- Wiles, J.R., Katchko, R.A., Benavides, E.A., O’Gorman, C.W., Escudero, J.M., Keisler, D.H., Stanko, R.L., & Garcia, M.R. (2014). The effect of leptin on luteal angiogenic factors during the luteal phase of the estrous cycle in goats. *Animal Reproduction Science*, 148, 121–129.
- Yusuf, M., Toshihiko, N., Ranasinghe, B., Gautam, G., Long, S., Yoshida, C., Koike, K. & Hayashi, A. (2010). Reproductive performance of repeat breeders in dairy herds. *Theriogenology*, 73, 1220-1229.