



# Journal of the Hellenic Veterinary Medical Society

Vol 76, No 1 (2025)



## To cite this article:

Fallah Mehrabadi, M., Moluoki, A., Bashashati, M., Rabiee, M., Ghalyanchilangeroudi, A., Shoushtari, A., & Motamed, N. (2025). Prevalence and risk factors of viral diseases in broiler farms contracted to multifactorial respiratory syndrome. *Journal of the Hellenic Veterinary Medical Society*, *76*(1), 8637–8646. https://doi.org/10.12681/jhvms.36340

# Prevalence and risk factors of viral diseases in broiler farms contracted to multifactorial respiratory syndrome

# M.H. Fallah Mehrabadi<sup>1,2\*</sup>, A. Moluoki<sup>1</sup>, M. Bashashati<sup>1</sup>, M.H. Rabiee<sup>2</sup>, A. Ghalyanchilangeroudi<sup>3</sup>, A. Shoushtari<sup>1</sup>, N. Motamed<sup>1</sup>

<sup>1</sup>Department of Poultry Diseases, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Tehran. Iran

<sup>2</sup>Department of Epidemiology, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Tehran. Iran

<sup>3</sup>Department of Microbiology and Immunology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

**ABSTRACT:** *Newcastle Disease (ND), Avian Influenza (AI) H9N2* and *Infectious Bronchitis (IB)* are the major respiratory diseases affecting broiler poultry farming in Iran. This study aims to investigate the prevalence and risk factors associated with these viral diseases in commercial broiler chicken farms experiencing multi-factorial respiratory syndrome in the country. In this cross-sectional study, 157 affected commercial broiler chicken farms were sampled from 2017 to 2019. Samples were examined for *AIV H9N2, NDV*, and *IBV* by performing reverse transcription polymerase chain reaction. Possible risk factors for the occurrence of viral diseases were investigated by comparing the mean and frequency of the factors, as well as performing univariable and multivariable regression analysis. Highest frequency of investigated viral diseases was due to *ND* with 98 cases (62.43%, CI 95%= 54.35-70.01). Frequency of *AI H9N2* and *IB* were respectively 74 cases (47.13%, CI 95%= 39.13-55.25) and 22 cases (14.1%, CI 95%= 8.99-20.44). Flock age (Odd Ratio: 2.65, P value: 0.030) and first week mortality (Odd Ratio: 10.12, P value: 0.014) respectively had significant effect on occurrence of *AI H9N2* and co-infection of *AI H9N2*. In conclusion, it is essential to implement control strategies measures to mitigate the impact of multi-factorial respiratory syndrome in Iranian commercial broiler chicken farms. This should specially focus on prioritize viral diseases based on their prevalence and associated risk factors, determine the serotypes and genotypes of the circulating strains of the virus, and improve biosecurity measures.

Keywords: Newcastle Disease; Infectious Bronchitis; Avian Influenza H9N2, Broiler, Epidemiology

Corresponding Author:

Date of initial submission: 22-12-2023 Date of acceptance: 19-10-2024

Mohammad Hossein Fallah Mehrabadi, Department of Poultry Diseases, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Tehran. Iran, Department of Epidemiology, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Tehran. Iran E-mail address: Mhf2480@yahoo.com

## INTRODUCTION

In recent years, Iran's poultry farming, particularly broiler ones, has experienced significant growth in the country. This rapid expansion and development has led to an increase in the density of these farms in certain regions of the country. Poultry farming in a dense manner can lead to development of complex and multi-factorial infections caused by viruses and bacteria. (Ahmadi et al. 2014).

In Iran's broiler poultry farming, the occurrence of poultry respiratory diseases due to multiple agents called poultry multi-factorial respiratory syndrome, accounts for the highest number of mortality in these farms every year. Multi-factorial respiratory syndrome occurs either as individual infection due to viral agents or simultaneous infection due to presence of several viral and bacterial agents (Saif et al. 2020). Evidence shows that most of these agents have a synergistic effect on each other. Indeed, different mortality rates are observed depending on which of the pathogenic agents are present in the farm. Despite the use of various preventive and control measures as well as widespread use of numerous vaccines, the prevalence and occurrence of respiratory diseases in poultry broiler farms have remained significant in recent years. According to the statistics available from the Iran Veterinary Organization (IVO), the most significant economic losses in the country's broiler farms are associated with the occurrence of multi-factorial respiratory syndrome.

Among the causes of multi-factorial respiratory syndrome, *ND*, *IB* and, *AI H9N2* are endemic in Iran. Their occurrence results in significant losses to broiler farms every year. In recent years, despite national and domestic prevention and control measures against the three diseases, coupled with the widespread use of diverse vaccines, outbreaks of these viral diseases still lead to the appearance of respiratory syndromes in broiler flocks, resulting in high annual mortality (Nili and Asasi 2002, Fallah Mehrabadi et al. 2019a, Hosseini et al. 2015).

Considering the important role of *ND*, *IB* and *AI H9N2* in the occurrence of multi-factorial respiratory syndrome in Iran, it is important to prioritize them based on their prevalence and related factors to design and implement better preventive plans. This study determined the prevalence and risk factors for these diseases in commercial broiler chicken farms which contracted to multi-factorial respiratory syndrome in Iran from 2017 to 2019.

## MATERIALS AND METHODS

### Study Design and setting

In this cross-sectional study, conducted during 2017 to 2019, 157 commercial broiler chicken farms contracted to multi-factorial respiratory syndrome in Iran were sampled and evaluated.

According to the latest statistics of the Ministry of Jihad Agriculture of Iran, approximately 20,000 commercial broiler chicken farms have licenses in Iran (Ebadzadeh et al. 2015). All of these farms scattered in 31 provinces of the country with different densities. The largest numbers of farms are located in Mazandaran and Isfahan provinces. Likewise, the number of hatching times in a year also varies in different regions, so that it is up to 5 times a year in the Northern provinces of the country and up to 3 times a year in the Southern provinces of the country.

All broiler farms in Iran are under the supervision of a private vet as a farm vet or clinician, in groups of one to several, based on farm population and location. The farmers and their Vet supervisors must give online prompt report when a farm is involved in serious problems, including important diseases as defined by IVO as a National Assignment. So the study population including infected farms was self-reported by farmers or their private Vets supervisor, in addition to farms that were reported following the active surveillance programs of IVO by state Vets

Therefore the studied farm selected among farms contracted to multi-factorial respiratory syndrome reported by IVO during 2017 to 2019 in Iran. The number of required sample in each province was determined based on the proportion of active broiler farms in each province. Farms with multi-factorial respiratory syndrome defined as existence of respiratory symptoms including respiratory rales, chafing, and runny nose, with or without gastrointestinal symptoms including diarrhea and neurological symptoms (wristed neck, paralyzed wings or legs) along with increased losses during at least 3 days.

### **Data collection**

### **Detection of Viral Agents**

For this purpose, initially, 5-7 recently deceased broiler chickens or those with clinical symptoms were sampled in each selected infected farms. The samples included tracheal and cloacal swabs, tracheal tissue, lung, brain, spleen, tonsils and kidney. All collected samples were transported to the diagnostic center laboratory of the Iran Veterinary Organization and the department of poultry diseases at the *Razi vaccine and serum research Institute*. The samples were labeled with their respective characteristics. Furthermore, the samples were stored in a freezer at -70 degrees Celsius until the tests were conducted.

Finally, RT-PCR method was used in order to detect AIV H9N2, IBV and NDV respectively. RNA extraction was done by commercial kits as follow: RNA was isolated from tissue samples using the High Pure RNA Extraction Kit (Roche, Mannheim, Germany). The isolated RNA was subsequently converted into cDNA with the RevertAid<sup>™</sup> First Strand cDNA Synthesis Kit (Thermo Scientific, MA, USA), utilizing random primers. The PCR reaction was performed in a 25 µl reaction volume, consisting of 8.5 µl of nuclease-free water, 12.5 µl of Taq DNA Polymerase Master Mix RED (AMPLIQON, Odense M, Denmark), 1  $\mu$ l of each primer (10  $\mu$ M) (Table 1), and 2  $\mu$ l of cDNA. The thermal cycling conditions were as follows: an initial denaturation at 95°C for 2 minutes, followed by 35 cycles of 95°C for 20 seconds, 52-54°C for 30 seconds, and 72°C for 1 minute, with a final extension step at 72°C for 5 minutes. The PCR products were then separated on a 1% agarose gel and visualized under UV light using a gel documentation system. (Monne et al. 2008, Callison et al. 2006).

#### **Possible Risk Factors**

For this purpose, a questionnaire was designed that includes general information about the farm and information related to the occurrence of the disease. This questionnaire was completed during sampling and farms visits through observation and face-to-face interview. It should be noted that the relevant questionnaire only was completed in 2017 and 2018. Because of change in the policies of the Iran Veterinary Organization (IVO) in 2019, the data due to questionnaire wasn't collected.

#### **Statistical Analysis**

In order to analyze the data, the frequency and relative frequency with 95% CI were used to describe the occurrence of viral diseases.

To assess the factor related to occurrence of the disease (all combinations), we compared the mean, frequency and total indices of the investigated factors. For quantitative variables, t-test and Mann-Whitney test were used and for qualitative variables, chi square test was used.

To assess the factor related to occurrence of the disease (Simple Infection/No Infection) (Co-Infection/Simple Infection), we used univariable and multivariable binary logistic regression analysis. Factor with P value  $\leq 0.1$  in univariable regression analysis inserted to multivariable analysis. Eventually in multivariable regression analysis factor with P value  $\leq 0.05$  were considered as significant. All Statistical analysis was done using Stata software version 14.

### RESULTS

#### **Frequency of Viral Diseases**

In this study, samples were collected from 157 active commercial broiler chicken farms, comprising 55, 77 and, 25 farms in 2017, 2018 and, 2019 respectively. In this study, results indicated that the highest frequency of viral disease was due to *ND* with 98 cases (62.43%, CI 95%= 54.35-70.01). Frequency of *AI H9N2* and *IB* were respectively 74 cases (47.13%, CI 95%= 39.13-55.25) and 22 cases (14.1%, CI 95%= 8.99-20.44). (Table 2)

# Factor Related to Frequency of Viral Diseases (All Combinations)

Analysis of the data in order to investigate factor related to occurrence of the viral diseases (all combinations), indicated following results:

Table 1: List of j	Table 1: List of primers used in this study									
Name	Target gene	Sequence	Amplicon length	Detected of	Reference					
MB-NDV-F	Fusion	5'-GGTGAGTCTATCCGGARGATACAAG-3'	202	NDV	(Creelan					
MD-NDV-K					(1 2006)					
MB-N103F	Nucleocansid <u>5'-CCTGATGGTAATTTCCGTTGGG-3'</u>		- 357	IBV	(Loa 2006)					
MB-N102R	racicocapsia	5'-ACGCCCATCCTTAATACCTTCCTC-3'	557	ID (						
MB-H9-151f	Usmagalutinin	5'-CTYCACACAGARCACAATGG-3'	100	AIV subtype	(Lee et al.					
MB-H9-638r	nemaggiutiinii	5'-GTCACACTTGTTGTTGTRTC-3'	400	H9	2001)					
MB-NA2-1	Nourominidago	5'-TCCGTTTCATTTGGGAACC-3'	214	AIV subtype	(Qiu et al.					
MB-NA2-2	incurammuase	5'-CTGACAATGGRCTAATGTG-3'	514	N2	2009)					

# Table 1: List of primers used in this stud

J HELLENIC VET MED SOC 2025, 76 (1) ПЕКЕ 2025, 76 (1)

There was no statistically significant relationship between the vaccination administration status and the frequency of viral diseases (P>0.05). There was a significant relationship between the flock age and the frequency of viral diseases (P=0.027). Indeed, mean of flock age at syndrome occurrence onset time was higher in farms infected only with AI H9N2 than other farms. Abundance of ND vaccination also had no significant relationship with the frequency of viral diseases (P>0.05). Abundance of IB vaccination had significant relationship with the frequency of viral diseases (P=0.001). Indeed abundance of IB vaccination in farms infected to IB was lower than farms not infected to the disease. First week mortality, had significant relationship with the frequency of viral diseases (P=0.001). Indeed mean of first week mortality was higher in farms only infected to IB than other farms. Abundance of imported chick, had no significant relationship with the frequency of viral diseases

(P>0.05). Flock age at *AI H9N2* vaccination administration time and *ND* vaccination administration time had no significant relationship with the frequency of viral diseases (P>0.05). (Table 3)

# Factor Related to Frequency of Viral Diseases (Simple infection/No Infection)

Analysis of the data in order to investigate factor related to occurrence of the viral diseases (simple infection/no infection), indicated following results:

Investigation regarding relationship of the factor and frequency of AI H9N2 indicated that age flock has significant effect on AI H9N2 in the univariable and multivariable regression analysis. Indeed, in multivariable regression analysis we indicated that farms with age flock 30-60 days had the higher odds of AI H9N2 related to the farms with age flock less than 30 days (Odd Ratio: 2.65, P value: 0.030). There was

 Table 2: Frequency and Relative Frequency of Viral Disease in Commercial Broiler Chicken Farms contracted to Multi-Factorial

 Respiratory syndrome in 2017 - 2019

	2017		20	18	20	19	Total		
Viral Disease	Fraguanov	Relative	Fraguanau	Relative	Fraguanay	Relative	Fraguanay	Relative	
	requency	Frequency							
AI H9N2	26	47.3	16	20.8	5	20	47	29.94	
ND	20	36.4	43	55.8	5	20	68	43.31	
IB	1	1.8	2	2.6	2	8	5	3.18	
AI H9N2 + IB	3	5.5	0	0	4	16	7	4.46	
ND + AI H9N2	4	7.3	13	16.9	3	12	20	12.74	
ND + IB	1	1.8	3	3.9	6	24	10	6.37	
Total	55	100	77	100	25	100	157	100	

 Table 3: Factor Related to Occurrence of Viral Diseases (all combinations) in Commercial Broiler Chicken Farms contracted to

 Multi-Factorial Respiratory Syndrome in 2017 - 2018

Independent Factor		AI H9N2	ND	IB	<i>AI H9N2</i> + <i>IB</i>	ND + AI H9N2	ND + IB	P value	
	Mean	34.6	29.88	29.33	29.66	32.23	29.25 9.53		
Electron of Several several	SD	9.60	7.37	6.11	6.42	4.07	9.53		
Plock Age at Syndrome	Min	13	15	24	25	23	19	0.027	
Occurrence Onset Time (Day)	Max	59	42	36	37	39	42		
	Median	31	31	28	27	32	28		
	Mean	19479.64	19117.02	18400	30633.33	27153.53	14609.25		
Abundance of Immented Chieles	SD	13051.16	13453.16	9699.485	11623.4	27107.65	6669.488		
(Number)	Min	3000	5002	7200	22900	6000	10000	0.360	
(Indilider)	Max	74000	93300	24000	44000	121300	24437		
	Median	17750	17900	24000	25000	20500	12000		
	Mean	2675.35	7458.31	9095.33	9563	8166.35	4120		
	SD	3994.55	6705.55	4582.73	4153.54	8690.62	3048.47		
First Week Mortality (Number)	Min	40	220	4050	6626	1600	860	0.001	
	Max	19000	38000	13000	12500	39329	6900		
	Median	1200	5340	10236	9563	5250	4600		

J HELLENIC VET MED SOC 2025, 76 (1) ПЕКЕ 2025, 76 (1)

	AI H9N2	Yes, N=72	23	37	3	1	7	1	0.100			
	Vaccine Administration	No, N=60	19	26	0	2	10	3	0.120			
Vaccination	IB Vaccine	Yes , N=103	30	50	3	2	15	3	0.400			
	Administration	No, N=29	12	13	0	1	2	1	0.490			
Administration Status	ND Vaccine	Yes, N=126	40	61	3	2	17	3	0.055			
according to	Administration	No, N=6	2	2	0	1	0	1				
Viral Agents	ND Vaccine Administration	Yes, N=80	28	39	3	1	7	2	0.100			
	(Injectable Form)	No, N=52	14	24	0	2	10	2	0.100			
	AI H9N2+ <i>IB</i> +	Yes, N=57	18	29	3	1	6	0	0.110			
	Administration	No, N=75	24	34	0	2	11	4	0.110			
		Mean	5.52381	5.650794	8.666667	4.666667	7.058824	6.25				
Flock Age at Ne	ewcastle	SD	3.81421	3.208651	1.154701	4.163332	1.983387	4.5				
Vaccination adn	ninistration Time	Min	0	0	8	0	3	0	0.250			
(Day)		Max	11	11	10	8	11	10				
		Median	5.5	7	8	6	7	7.5				
		Mean	4	3.95	8.66	2	2.88	2.5				
Flock age at AI	H9N2	SD	4.29	4.01	1.15	3.46	3.58	5				
Vaccination adm	ninistration Time	Min	0	0	8	0	0	0	0.100			
(Day)		Max	11	11	10	6	8	10				
		Median	2	4	8	0	0	0				
		0	2	2	0	1	0	1				
		1	10	8	0	0	3	1				
		2	7	11	0	1	3	1				
Abundance of $\Lambda$	D Vaccination	3	6	22	1	0	6	0	0.260			
(Injectable Forn	1)	4	8	12	2	1	5	1	0.200			
		≥5	9	8	0	0	0	0				
		All	42	63	3	3	17	4				
		0	13	8	0	2	5	1				
		1	11	40	0	1	6	1				
Abundance of <i>I</i> .	B Vaccination	≥2	18	15	3	0	6	2	0.001			
		All	42	63	3	3	17	4				

no statistically significant relationship between the other possible factor and the frequency of *AI H9N2* diseases. Also there were no statistically significant relationship between the possible factor and the frequency of *ND* diseases. In addition, there were no statistically significant relationship between the possible factor and the frequency of *IB* disease. (Table 4)

## Factor Related to Frequency of Viral Diseases (Co-Infection /Simple Infection)

Analysis of the data in order to investigate factor related to occurrence of the viral diseases (Co-Infection/Simple Infection), indicated following results:

Investigation regarding relationship of the factor and frequency of co-infection of *AI H9N2* indicated that first week mortality has significant effect on co-infection of *AI H9N2* in the univariable and multivariable regression analysis. Indeed in multivariable regression analysis we indicated that first week mortality above 8000 can increase odds of occurrence of co-infection *AI H9N2* related to those first week mortality was less than 4000 (Odd Ratio: 10.12, P value: 0.014). There was no statistically significant relationship between the other possible factor and the frequency of co-infection of *AI H9N2* diseases.

Moreover, there were no statistically significant relationship between the possible factor and the fre-

quency of co-infection of *ND* diseases. As the same, there were no statistically significant relationship between the possible factor and the frequency of co-infection of *IB* diseases. (Table 5)

Table 4: Factor Related to Occurrence of Viral Diseases (Simple Infection/No Infection) in Commercial Broiler Chicken Far	ms con-
tracted to Multi-Factorial Respiratory Syndrome in 2017 - 2018	

N/inel		Univ	ariable Reg	Multivariable Regression Analysis						
Disease	Factor		Regression Coefficient	SD	OR	P-value	Regression Coefficient	SD	OR	P-value
	Flock Age at Syndrome	<30	-	-	-	-	-	-	-	-
	Occurrence Onset Time	30-60	1.31	0.431	3.70	0.002	0.97	0.45	2.65	0.030
	(Day)	>60	22.48	9748.2	5815709434	0.998	21.956	9669.182		0.998
	41 1 67 4 1	<10000	-	-	-	-	-	-	-	-
A AI H9N2	Abundance of Imported	10000-20000	0.50	0.47	1.66	0.282	-	-	-	-
	Chicks (Number)	>20000	0.58	0.47	1.79	0.214	-	-	-	-
		<4000	-	-	-	-	-	-	-	-
	First Week Mortality	4000-8000	-1.59	0.47	0.20	0.001	-1.028	0.51	0.35	0.055
	(Ivuiliber)	>8000	-1.17	0.46	0.31	0.011	-0.704	0.49	0.49	0.495
	AI H9N2	Yes	-0.346	0.35	0.70	0.324	-	-	-	-
	Vaccine Administration	No	-	-	-	-	-	-	-	-
	Flock Age at Syndrome	<30	-	-	-	-	-	-	-	-
	Occurrence Onset Time	30-60	-1.089	0.48	0.337	0.0.24	-0.68	0.50	0.50	0.178
	(Day)	>60	-22.92	9748.22	0	0.998	-22.342	9628.88	0	0.998
		<10000	-	-	-	-	-	-	-	-
	Abundance of Imported	10000-20000	-0.409	0.49	0.66	0.405	-	-	-	-
	Chicks (Number)	>20000	-0.326	0.49	0.72	0.508	-	-	-	-
	T' ( XV 1 M ( 1')	<4000	-	-	-	-	-	-	-	-
ND	(Number)	4000-8000	1.70	0.51	5.50	0.001	1.12	0.58	3.09	0.053
		>8000	1.59	0.52	4.91	0.002	1.11	0.58	3.04	0.056
	ND	Yes	0.588	0.83	1.80	0.483	-	-	-	-
	Vaccine Administration	No	-	-	-	-	-	-	-	-
	ND	Yes	-0.405	0.377	0.66	0.283	-	-	-	-
	Vaccine									
	Administration	No	-	-	-	-	-	-	-	-
	(Injectable Form)	0								
		0	-	-	-	-	-	-	-	-
		1	0.182	0.92	1.20	0.843	-	-	-	-
	Abundance of ND Veccination (Injectable	2	0.629	0.92	1.87	0.497	-	-	-	-
	Form)	3	1.38	0.91	4	0.132	-	-	-	-
	)	4	0.492	0.90	1.63	0.585	-	-	-	-
		$\geq_5$	-0.118	0.95	0.88	0.901	-	-	-	-
	Flock Age at Syndrome	<30	-	-	-	-	-	-	-	-
	Occurrence Onset Time	30-60	-1.37	0.71	0.25	0.056	-1.44	0.86	0.23	0.093
	(Day)	>60	-19.48	9748.22	0	0.998	-18.49	9559.031	0	0.998
	A 1 1	<10000	-	-	-	-	-	-	-	-
	Chicks (Number)	10000-20000	-0.56	1.02	0.57	0.586	-	-	-	-
	Chicks (Runiber)	>20000	0.62	0.85	1.86	0.463	-	-	-	-
	Einet Weste Mantality	<4000	-	-	-	-	-	-	-	-
<b>ID</b>	(Number)	4000-8000	2.079	1.14	8	0.068	1.49	1.16	4.43	0.200
IB	(ivuilioer)	>8000	1.861	1.17	6.42	0.114	1.46	1.19	4.33	0.220
	IB	Yes	-0.459	0.725	0.632	0.526	-	-	-	-
	Vaccine Administration	No	-	-	-	-	-	-	-	-
		0	-	-	-	-	-	-	-	-
	Abundance IR	1	-1.19	0.943	0.30	0.20	-	-	-	-
	Vaccination	$\geq_{2}$	0.10	0.773	1.11	0.89	-	-	-	-
		2								
		All	-	-	-	-	-	-	-	-

J HELLENIC VET MED SOC 2025, 76 (1) ПЕКЕ 2025, 76 (1)

Table 5: Factor Related to Occurrence of Viral Diseases (	Co-Infection/Simple infection) in Commercial Broiler Chicken Farms con-
tracted to Multi-Factorial Respiratory Syndrome in 2017 -	2018

				ariable Reg	Multivariable Regression Analysis					
Viral Disease	Facto	or	Regression Coefficient	SD	OR	P-value	Regression Coefficient	SD	OR	P-value
	Flock Age	<30								
Co-Infection AI H9N2	at Syndrome	30-60	-1.373	0.774	0.253	0.076	-0.660	0.909	0.517	0.468
	Onset Time	>60	-22.050	9748.22	0.	0.998	-21.168	9420.098	0	0.998
	Abundance	<10000	-	-	-	-	-	-	-	-
	of Imported Chicks	10000- 20000	-0.405	0.842	0.667	0.630	-	-	-	-
	(Number)	>20000	0.827	0.783	2.286	0.291	-	-	-	
	First Week	<4000	-	-	-	-	-	-	-	-
	Mortality	4000-8000	1.928	0.804	6.875	0.017	1.356	0.935	3.88	0.147
	(Number)	>8000	2.686	0.810	14.667	0.001	2.315	0.944	10.12	0.014
	AI H9N2+ IB+	Yes	-0.405	0.551	0.667	0.462	-	-	-	-
	ND Vaccine Administration	No	-	-	-	-	-	-	-	-
	Flock Age	<30	-	-	-	-	-	-	-	-
	at Syndrome	30-60	0.454	0.515	1.574	0.378	-	-	-	-
	Onset Time (Day)	>60	-	-	-	-	-	-	-	-
	Abundance	<10000	-	-	-	-	-	-	-	-
Co-Infection	of Imported Chicks	10000- 20000	0.215	0.702	1.240	0.760	-	-	-	-
ND	(Number)	>20000	0.658	0.674	1.932	0.329	-	-	-	-
	First Week	<4000	-	-	-	-	-	-	-	-
	Mortality	4000-8000	-0.201	0.635	0.818	0.752	-	-	-	-
	(Number)	>8000	0.154	0.624	1.167	0.805	-	-	-	-
	AI H9N2+ IB+	Yes	-0.709	0.516	0.492	0.170	-	-	-	-
	ND Vaccine	No								
	Administration	INO	-	-	-	-	-	-	-	-
	Flock Age	<30	-	-	-	-	-	-	-	-
	at Syndrome	30-60	-0.223	1.483	0.80	0.88	-	-	-	-
	Occurrence Onset Time	>60	-	-	-	-	-	-	-	-
	(Day) Abundance	<10000	_	-	_	-	-	-	-	-
Co-Infection	of Imported Chicks	10000- 20000	21.20	28420.72	1615474843	0.999	-	-	-	-
IB	(Number)	>20000	0.693	1.658	2	0.676	-	-	-	-
	First Week	<4000	-	-	-	-	-	-	-	-
	Mortality	4000-8000	-20.104	40192.96	0	1	-	-	-	-
	(Number)	>8000	-21.869	40192.96	0	1	-	-	-	-
	AI H9N2+ IB+	Yes	-21.608	17974.84	0	0.999	-	-	-	-
	ND Vaccine	No	-	-	-	-	-	-	-	-

### DISCUSSION

Respiratory diseases have been one of the most important problems in broiler farming in recent years in Iran. Among these diseases, *IB*, *ND* and *AI H9N2*, which are endemic in Iran, have played the greatest role. For example according to one investigation, the sum of costs and losses due to respiratory complex was 9.47 \$US Million, 2016-2017 among broiler farms in Iran. In addition to Iran, these three diseases are also the most important pathogens in the poultry farming, especially broiler one, in many other countries (Saif et al. 2020, Hosseini et al. 2015, Fallah Mehrabadi et al. 2019b, Hadipour and Golchin 2011).

In the present study, ND was the most frequent viral diseases in the farms contracted with multi-factorial respiratory syndrome. The disease, caused by a velogenic strain of the virus, is a significant problem in poultry farms in Iran, despite extensive vaccination efforts. The disease has been responsible for high mortality rates in recent years and continues to cause outbreaks across the country (Hosseini et al. 2014). Similar issues have been reported in other countries, such as Egypt and Pakistan, where velogenic strains of the virus have been detected in broiler chicken flocks and various bird species, respectively (Hassan et al. 2016) (Wajid et al. 2017) (Miller et al. 2015). Considering that backyard poultry are believed to be the main reservoirs of ND, and play a crucial role in transmitting the infection to commercial poultry farms (Awan et al. 1994), vaccinating backyard poultry could greatly reduce the reservoirs and limit the spread of the disease.

The study indicates that AI H9N2 is a significant viral contributor to multifactorial respiratory syndrome in broiler farms across Iran, affecting over 47% of farms. The disease is endemic in Iran since 1998 which for the first time reported in the country. Although AIV H9N2 is of low pathogenicity, its presence alongside other pathogens leads to increased mortality rates in poultry. This virus is also endemic in several neighboring countries, posing challenges to poultry farming. For instance, a study in Egypt revealed AI H9N2 in 53% of broiler flocks with respiratory issues, with a notable co-infection rate of 41.7% alongside IB. Between 2000 and 2003, AIV H9N2 was isolated from broiler chickens and quails in the UAE, where it caused mortality. In Pakistan, the virus has been endemic since 1996, resulting in ongoing economic losses (Hassan et al. 2016) (Cameron et al. 2000, Lee et al. 2016). Given the frequent mutations of AIV *H9N2* (Bashashati et al. 2013), continuous monitoring and appropriate vaccine strain selection based on the changes and the similarity between the circulating viruses are essential.

The current study found a notably low frequency of IB compared to the two other viral diseases examined, indicating that it had a lesser impact on the development of multi-factorial respiratory syndrome than in previous research. For example, Pourbakhsh et al in 2008 detected IBV in 73% of farms studied, with the 793/B genotype being the most prevalent (Pourbakhsh et al. 2008). Additionally, the Massachusetts genotype was also identified. Similarly, Seifi et al in 2010 found the virus in 40% of farms studied, with genotypes 4.91 and Massachusetts present. In neighboring countries, IB remains a significant issue for broiler poultry farming (Seger et al. 2016) (Hassan et al. 2016). It seems that geographic distance and control measures applied in various countries may play a role in preventing the introduction of IBV variants in Iran. Vaccination is crucial for the disease control, and using two different attenuated live vaccines can provide sufficient immunity against various IBV types as the current study supports the notion that increasing vaccination frequency can help reduce the incidence of IB.

The findings of this cross-sectional study indicate a significant relationship between flock age and the occurrence of viral diseases in broiler farms in Iran by comparing the mean age flock at syndrome occurrence time according to type of infection. Also the multivariable regression analysis further supports these observations, revealing that flocks aged at syndrome occurrence time between 30 to 60 days exhibited a higher likelihood of AI H9N2 compared to those less than 30 days. This age-related susceptibility may be attributed to several factors, including the immunological maturity of older birds, which could influence their response to viral infections. These results underscore the importance of age as a critical factor in managing viral diseases within poultry populations. Understanding the dynamics of disease occurrence in relation to flock age can inform biosecurity measures and vaccination strategies, ultimately aiding in the control of AI H9N2 and enhancing the overall health management practices in broiler farms. Further research is warranted to explore the underlying mechanisms of this relationship and to assess the effectiveness of interventions aimed at reducing disease prevalence in older flocks.

In this study, the role of different pathogens in various combinations in the occurrence of multi-factorial respiratory syndrome in Iranian commercial broiler chicken farms was shown. The most frequent disease was related to ND. In addition to NDV, the role of AIV H9N2 and IBV in the occurrence of multi-factorial respiratory syndrome in the farms was confirmed. The occurrence of these diseases may be related to poor hygiene and lack of prophylaxis methods. Also, poor biosecurity with poor conditions of buildings and equipment may lead to exposure of poultry in farms to pathogens circulating in wild bird populations. On the other hand, due to several reasons, in addition to the change of the dominant virus, the serotypes, genotypes and circulating strains of the same virus are constantly changing. Therefore, it is crucial to prioritize

viral diseases based on their prevalence and associated risk factors, determine the serotypes and genotypes of the circulating strains of the virus, and improve biosecurity measures. These strategies are essential control measures that must be implemented to mitigate the impact of multi-factorial respiratory syndrome in Iranian commercial broiler chicken farms.

#### **ACKNOWLEDGMENTS**

This study was approved by the ethics committee of *Razi Vaccine and Serum Research Institute*, Karaj, Iran (Number: 13-18-1851-067-97020-971046).

#### **CONFLICT OF INTERESTS**

The authors declare that they have no competing interests.

#### REFERENCES

- Ahmadi K, Ebadzadeh HR, Mohammadnia Afruzi S, Taghani RA, Saadat Akhtar A. (2014).Review the ptoduction process of protein products in the country in four decades (1974 to2013). Tehran, Iran: Center for Information and Communication Technology.
- Saif YM, Swayne DE, Pantin-Jackwood MJ, Spackman E, Johnson TJ, Day JM, et al. Emerging Diseases and Diseases of Complex or Unknown Etiology. Diseases of Poultry2020. p. 1383-410.
- Nili H, Asasi K. (2002). Natural cases and an experimental study of H9N2 avian influenza in commercial broiler chickens of Iran. Avian Pathol. 31(3):247-52.
- Fallah Mehrabadi MH, Ghalyanchilangeroudi A, Rabiee MH, Tehrani F. (2019a). Prevalence and risk factors of avian influenza H9N2 among backyard birds in Iran in 2015. Asian Pacific Journal of Tropical Medicine. 12(12).
- Hosseini H, Fard MH, Charkhkar S, Morshed R. (2015). Epidemiology of Avian Infectious Bronchitis Virus Genotypes in Iran (2010-2014). Avian Dis. 59(3):431-5.
- Ebadzadeh HR, Ahmadi K, Mohammadnia Afruzi S, Taghani RA, Moradi Eslami A, Yari SMA. (2015).Agricultural statistics. Tehran, Iran: Center for Information and Communication Technology.
- Monne I, Ormelli S, Salviato A, De Battisti C, Bettini F, Salomoni A, et al. (2008). Development and validation of a one-step real-time PCR assay for simultaneous detection of subtype H5, H7, and H9 avian influenza viruses. J Clin Microbiol. 46(5):1769-73.
- Callison SA, Hilt DA, Boynton TO, Sample BF, Robison R, Swayne DE, et al. (2006). Development and evaluation of a real-time Taqman RT-PCR assay for the detection of infectious bronchitis virus from infected chickens. J Virol Methods. 138(1-2):60-5.
- Creelan JL, Graham, D.A., and McCullough, S.J., . (2002). Detection and differentiation of pathogenicity of avian paramyxovirus serotype 1 from field cases using one-step reverse transcriptase-polymerase chain reaction,. Avian Pathology. 31:493-9.
- Loa CC, Lin, T.L., Wu, C.C., Bryan, T.A., Hooper, T.A., and Schrader,

D.L., . (2006). Differential detection of turkey coronavirus, infectious bronchitis virus, and bovine coronavirus by a multiplex polymerase chain reaction. Journal of Virological Methods. 131:86-91.

- Lee MS, Chang PC, Shien JH, Cheng MC, Shieh HK. (2001). Identification and subtyping of avian influenza viruses by reverse transcription-PCR. J Virol Methods. 97(1-2):13-22.
- Qiu BF, Liu WJ, Peng DX, Hu SL, Tang YH, Liu XF. (2009). A reverse transcription-PCR for subtyping of the neuraminidase of avian influenza viruses. J Virol Methods. 155(2):193-8.
- Fallah Mehrabadi MH, Ghalyanchi Langeroudi A, Bahonar A, Rabiee MH, Tehrani F, Amirhajloo S, et al. (2019b). Prevalence of Avian Influenza in Live Bird Markets, Bird Gardens, and Zoos in Iran in 2015: A Cross-sectional Study. Archives of Razi Institute. 74(3):243-50.
- Hadipour M, Golchin P. (2011). Serosurvey of H9N2 avian influenza virus during respiratory disease outbreaks in broiler flocks in Dezful, southern Iran. Bulg J Vet Med. 14:62-5.
- Hosseini H, Langeroudi AG, Torabi R. (2014). Molecular characterization and phylogenetic study of Newcastle disease viruses isolated in Iran, 2010-2012. Avian Dis. 58(3):373-6.
- Hassan KE, Shany SA, Ali A, Dahshan AH, El-Sawah AA, El-Kady MF. (2016). Prevalence of avian respiratory viruses in broiler flocks in Egypt. Poult Sci. 95(6):1271-80.
- Wajid A, Dimitrov KM, Wasim M, Rehmani SF, Basharat A, Bibi T, et al. (2017). Repeated isolation of virulent Newcastle disease viruses in poultry and captive non-poultry avian species in Pakistan from 2011 to 2016. Prev Vet Med. 142:1-6.
- Miller PJ, Haddas R, Simanov L, Lublin A, Rehmani SF, Wajid A, et al. (2015). Identification of new sub-genotypes of virulent Newcastle disease virus with potential panzootic features. Infect Genet Evol. 29:216-29.
- Awan MA, Otte MJ, James AD. (1994). The epidemiology of Newcastle disease in rural poultry: a review. Avian Pathol. 23(3):405-23.
- Cameron KR, Gregory V, Banks J, Brown IH, Alexander DJ, Hay AJ, et

8646

al. (2000). H9N2 subtype influenza A viruses in poultry in pakistan are closely related to the H9N2 viruses responsible for human infection in Hong Kong. Virology. 278(1):36-41.

- Lee DH, Swayne DE, Sharma P, Rehmani SF, Wajid A, Suarez DL, et al. (2016). H9N2 low pathogenic avian influenza in Pakistan (2012-2015). Vet Rec Open. 3(1):e000171.
- Bashashati M, Vasfi Marandi M, Sabouri F. (2013). Genetic diversity of early (1998) and recent (2010) avian influenza H9N2 virus strains isolated from poultry in Iran. Arch Virol. 158(10):2089-100.
- Pourbakhsh SA, Momayez R, Toroghi R R, Shoushtari AH. (2008). Ninetythree B type, the Predominant Circulating Type of Avian Infectious Bronchitis Viruses 1999 - 2004 in Iran: a retrospective study. Archives of Razi Institute. 63(1):1-5.
- Seger W, GhalyanchiLangeroudi A, Karimi V, Madadgar O, Marandi MV, Hashemzadeh M. (2016). Genotyping of infectious bronchitis viruses from broiler farms in Iraq during 2014-2015. Arch Virol. 161(5):1229-37.