


## Journal of the Hellenic Veterinary Medical Society

Vol 76, No 2 (2025)



### The first case of Co-infection of Sacbrood virus and Varroa destructor virus-1 in honey bees (*Apis mellifera* L., 1758)

A Zerek, M Şevik, I Erdem, M Yaman

doi: [10.12681/jhvms.36354](https://doi.org/10.12681/jhvms.36354)

Copyright © 2025, A Zerek, M Şevik, I Erdem, M Yaman



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

### To cite this article:

Zerek, A., Şevik, M., Erdem, I., & Yaman, M. (2025). The first case of Co-infection of Sacbrood virus and Varroa destructor virus-1 in honey bees (*Apis mellifera* L., 1758). *Journal of the Hellenic Veterinary Medical Society*, 76(2), 9385–9390. <https://doi.org/10.12681/jhvms.36354> (Original work published July 5, 2025)

## The first case of Co-infection of Sacbrood virus and Varroa destructor virus-1 in honey bees (*Apis mellifera* L., 1758)

A. Zerek,<sup>1a</sup> M. Şevik,<sup>2b(\*)</sup> İ. Erdem,<sup>1c</sup> M. Yaman<sup>1d</sup>

<sup>1</sup>Department of Parasitology, Veterinary Faculty, Hatay Mustafa Kemal University, Antakya, 31060 Hatay, Türkiye.

<sup>2</sup>Department of Virology, Veterinary Faculty, Necmettin Erbakan University, Ereğli, 42310 Konya, Türkiye.

**ABSTRACT:** Viral infections can occur at all developmental stages of honey bees and are of great concern for the beekeeping industry as they cause significant economic losses in beehives. The aim of this study is to investigate sudden bee mortalities and colony losses in an apiary. In 2022, high levels of bee mortality were observed in an apiary in Kırıkhan district of Hatay province. In order to determine the cause of bee mortality, adult worker bee samples (n = 50) were collected from hives within the apiary. Collected bee samples were processed for RNA extraction. One step real time RT-PCR methods were applied for the detection of Sacbrood virus (SBV) and Varroa destructor virus-1 (VDV-1). The one step real time RT-PCR analyses revealed the presence of mixed infection with SBV and VDV-1 viruses in the apiary. As far as we know, this study is the first report of a mixed infection of SBV and VDV-1 in honey bees. Additional comprehensive studies are needed to better understand the epidemiology of mixed infections caused by viruses in honey bees.

**Keyword:** Honey bee; SBV; VDV-1; mixed infection; Türkiye.

*Correspondence author:*

Murat ŞEVİK  
Department of Virology, Veterinary Faculty, Necmettin  
Erbakan University, Ereğli, 42310 Konya, Türkiye.  
E-mail address: dr\_muratank@hotmail.com  
Phone number: +90 332 777 0066

*Date of initial submission:* 26-12-2023

*Date of acceptance:* 7-3-2025

## INTRODUCTION

Honey bees are arguably the most significant living species on Earth in terms of maintaining ecological balance and agricultural food production due to their ability to pollinate flowering plants, ensuring biodiversity (Martin, 2001; Genersch, 2010; Parveen et al., 2022). However, the sudden decrease in the population of these highly vital species, along with the recent loss of colonies, are cause for concern (Aizen and Harder, 2009; Potts et al., 2010). Increasing numbers of global honey bee mortalities and colony losses have been reported in various parts of the world, particularly in the United States and Europe (Wyszkowska et al., 2019; Lu et al., 2020). Habitat loss, pesticides, bee pathogens and numerous stress factors are considered to be contributing factors to the decline in honey bee populations (Potts et al., 2010). Infections caused by bee pathogens, especially viruses, can be observed in all developmental stages of honey bees leading to significant losses in beehives and are of great concern for the beekeeping sector (Morawetz et al., 2019; Kalayci et al., 2020; Ullah et al., 2021 ).


The majority of bee viruses are known to be RNA viruses and are classified in the *Iflaviridae* and *Dicistroviridae* families (de Miranda, 2008). At least twenty-four different viruses have been reported to

date in honey bees (de Miranda et al., 2013; Remnant et al., 2017; McMenemy and Flenniken, 2018), several of these viruses have been reported to pose a substantial threat to beekeeping worldwide (Tantillo et al., 2015; McMenemy and Flenniken, 2018). Prominent viruses that cause severe diseases and economic losses in honey bee colonies; Deformed wing virus (DWV), Varroa destructor virus-1 (VDV-1), Sacbrood virus (SBV), Acute bee paralysis virus (ABPV), Israeli acute paralysis virus (IAPV), Black queen cell virus (BQCV), Slow bee paralysis virus (SBPV), Kashmir bee virus (KBV), Lake Sinai virus (LSV) and Chronic bee paralysis virus (CBPV) (Tantillo et al., 2015; McMenemy and Flenniken, 2018; Ullah et al., 2021). The present study aims to report on the findings from an apiary with high rates of bee mortality.

## CASE REPORT

In March 2022, an apiary conducting a fixed beekeeping practice in Hatay province was visited due to complaints of sudden bee mortalities and colony loss. The apiary was equipped with 50 hives.

Clinical symptoms, including flightless, crawling, paralysis, and wing abnormalities, were observed among the adult bees infected by VDV-1 (Figure 1). Samples of adult worker bees (n = 50) showing



**Figure 1.** Clinical signs detected in sampled bees; a) Paralysis b) Deformed wing c) Bee infected with Varroa mite and showing signs of deformed wing (marked with red arrows)

clinical symptoms were collected from 10 hives in the facility selected by random sampling method, with 5 samples from each hive. It has been reported that this sample size is sufficient for the molecular detection of infectious agents affecting 45% or more of the individuals within a colony (Daughenbaugh et al., 2015; Cavigli et al., 2016). Samples collected in sterile falcon tubes were immediately transferred to the laboratory under cold chain and stored at -80 °C until further analysis.

Bee samples were pooled in sterile falcon tubes containing phosphate-buffered saline (PBS) before being crushed utilizing a TissueRuptor device (Qia-gen, Hilden, Germany). The homogenates were then centrifuged at 5000 rpm for 30 minutes at 4 °C and 50 µl of supernatant was taken for RNA extraction. RNA extraction was conducted with the use of a commercial kit (High Pure Viral Nucleic Acid Kit, Roche, Germany) in accordance with the manufacturer's instructions. Obtained extracts were stored at -80°C until further analysis. The RNA extracts were analysed for the presence of SBV and VDV-1. These viruses were selected for analysis, based on the clinical symptoms of the afflicted bees. Probe and primer pairs as defined by Schurr et al. (2019) were employed for VDV-1 virus-specific RNA detection, and probe and primer pairs as defined by Blanchard et al. (2014) were employed for SBV detection (Table 1). One step real time RT-PCR master mix was prepared using a commercial kit (AgPath-ID One-Step RT-PCR, ThermoFisher Scientific, MA, USA). Amplification conditions: reverse transcription at 45°C for 10 minutes, initial PCR activation at 95°C for 10 minutes and 40 cycles at 95°C for 15 seconds and 60°C for 1 minute (binding/extension). Real time RT-PCR cycle threshold value (Ct) equal or lower than 35 was considered as positive (de Miranda et al., 2013). Throughout the analysis, sterile nuclease-free water was used as a negative control while

positive RNA samples of SBV and VDV-1 isolates previously confirmed by sequence analysis (obtained from Department of Parasitology, Veterinary Faculty, Hatay Mustafa Kemal University, Antakya, Türkiye) were used as positive controls.

## DISCUSSION

Viruses that affect honey bees threaten the beekeeping industry worldwide, causing significant economic losses (Kevill et al., 2017). SBV and DWV, the most widespread honey bee viruses globally, are the most prominent viruses associated with colony collapse disorder (Kevill et al., 2017; Li et al., 2019; Wei et al., 2022). According to reports, honey bees can become infected with multiple viruses, making it possible to identify mixed infections (Chen et al., 2006). In the present study, SBV and VDV-1 mixed infection was detected in honey bees. As far as we know, this study is the first report of a mixed infection of SBV and VDV-1 in honey bees.

SBV was the first virus identified in honey bees and remains one of the most prevalent honey bee viruses worldwide (Wei et al., 2022). SBV is a member of the *Iflavirus* genus of the *Iflaviridae* family and has positive-sense, single-stranded RNA genome (Valles et al., 2017). SBV infection can occur in both larval and adult stages (Ball and Bailey, 1997). Infected larvae are unable to pupate and take on a sac-like appearance. In adult bees, where there are no obvious clinical symptoms, it causes a latent/persistent infection with shortened lifespan (Berényi et al., 2006; de Miranda, 2008).

Studies carried out in honey bees throughout the world showed that the prevalence of SBV varies between 40.2% and 90.5% (Tentcheva et al., 2004; Choe et al., 2012; Tlak Gajger et al., 2014). In Türkiye, it has been reported that the prevalence of SBV in apiaries varies between 2.7% and 22.3% (Kalayci et al., 2020; Cagiran and Yazici, 2021). The differ-

**Table 1.** Primers and probes used in one step real time RT-PCR analyses

Primer/ Probe	Sequence (5'-3')	Concentration	Reference
VDV-1-F	GGTCTGAAGCGAAAATAG	1200 nM	Schurr et al. (2019)
VDV-1-R	CTAGCATATCCATGATTATAAAC	1200 nM	
VDV-1-P	Fam-CCTTGTCAGTAGATACAGCATCACA-Tamra	400 nM	
SBV-F	AACGTCCACTACACCGAAATGTC	320 nM	Blanchard et al. (2014)
SBV-R	AACTGCGCGTCTAACATTCC	320 nM	
SBV-P	Fam-TGATGAGAGTGGACGAAGA-MGB	200 nM	



ence in SBV rates detected in honey bees in various countries and regions may be associated with the number of sampled bees, the method of analysis used and the awareness of beekeepers against diseases.

VDV-1 is classified as the DWV-B genotype of DWV and has positive-sense, single-stranded RNA genome within the genus *Iflavirus* of the family *Iflaviridae* (Ongus et al., 2004; Lanzi et al., 2006). VDV-1 causes wing deformities in developing honey bees and a shortened life span in infected adult bees (Benaets et al., 2017; Brettell et al., 2017).

VDV-1 was detected at very high levels of 66% in the United States (Ryabov et al., 2017), 47% in Argentina (Brascesco et al., 2021), in 82% and 98% of adult honey bees and hives in Türkiye, respectively (Oz et al., 2023). In addition, it was reported that the province of Hatay had a high rate of virus infection during the colony losses in 2010-2011 (Tozkar et al., 2015). The current research also detected VDV-1 in an apiary in Hatay province. This result shows that VDV-1 is still circulating in honey bees located in Hatay province.

Mixed infections in bees can lead to severe colony collapse disorders. Studies conducted throughout the globe have reported mixed infections including SBV and DWV-A viruses in honey bees in Argentina (Sguazza et al., 2013), Croatia (Tlak Gajger et al.,

2014), Austria (Berényi et al., 2006), France (Tentcheva et al., 2004), Iran (Moharrami and Modirrousta, 2018), Denmark (Nielsen et al., 2008), USA (Chen et al., 2004), Jordan (Haddad et al., 2008). Although there are studies in Türkiye in which SBV and DWV-A viruses have been detected together (Rüstemoğlu and Sipahioğlu, 2019; Kalayci et al., 2020; Kızıltepe et al., 2023; Muz and Muz, 2023a; Muz and Muz, 2023b), the present study was the first to detect SBV and VDV-1 in honey bees.

In conclusion, the results of this study showed that SBV and VDV-1 mixed infection can cause high rates of bee mortality in honey bees. Recent studies have shown that virus interactions play a role in colony collapse disorder (Li et al., 2019; Wei et al., 2022). The way of transmission plays a crucial role in the dynamics of viral infections and in the evolutions of host-pathogen interactions. The spread of the infection can occur through the robbing and drifting of virus-infected colonies between hives. Further comprehensive research is necessary to fully understand the epidemiology of mixed viral infections in honey bees.

## ACKNOWLEDGMENTS

This study has been supported by Necmettin Erbakan University Scientific Research Projects Coordination Unit under project number 221229001.

## REFERENCES

- Aizen MA, Harder LD (2009) The global stock of domesticated honey bees is growing slower than agricultural demand for pollination. *Curr biol*, 19 (11): 915–918. DOI: 10.1016/j.cub.2009.03.071
- Ball BV, Bailey L (1997) Viruses. In: Morse RA, Flottum K. (Eds.), *Honey Bee Pest, Predators, & Diseases*. The A. I. Root Co., Medina, OH, pp. 11–31.
- Benaets K, Van Geystelen A, Cardoen D, De Smet L, de Graaf DC, Schoofs L, Larmuseau MH, Brettell LE, Martin SJ, Wenseleers T (2017) Covert deformed wing virus infections have long-term deleterious effects on honeybee foraging and survival. *Proc Biol Sci*. 284: 20162149. DOI: 10.1098/rspb.2016.2149.
- Berényi O, Bakonyi T, Derakhshifar I, Köglberger H, Nowotny N (2006) Occurrence of six honeybee virus in diseased Austrian apiaries. *Appl Environ Microbiol*, 72 (4):2414–2420. DOI: 10.1128/AEM.72.4.2414–2420.2006
- Blanchard P, Guillot S, Antúnez K, Köglberger H, Kryger P, de Miranda JR, Franco S, Chauzat MP, Thiéry R, Ribiére M (2014) Development and validation of a real-time two-step RT-qPCR TaqMan® assay for quantitation of Sacbrood virus (SBV) and its application to a field survey of symptomatic honey bee colonies. *J Virol Methods*, 197: 7–13. DOI: 10.1016/j.jviromet.2013.09.012
- Brascesco C, Quintana S, Di Gerónimo V, Genchi García ML, Sguazza G, Bravi ME, Fagnoli L, Reynaldi FJ, Eguaras M, Maggi M (2021) Deformed wing virus type a and b in managed honeybee colonies of Argentina. *Bull Entomol Res*, 1–11. DOI: 10.1017/S000748532000036X.
- Brettell LE, Mordecai GJ, Schroeder DC, Jones IM, da Silva JR, Vicente-Rubiano M, Martin SJ (2017) A Comparison of Deformed Wing Virus in Deformed and Asymptomatic Honey Bees. *Insects*. 8 (1): 28. DOI: 10.3390/insects8010028.
- Cagirgan AA, Yazici Z (2021) The prevalence of seven crucial honeybee viruses using multiplex RT-PCR and their phylogenetic analysis, *Turk J Vet Anim Sci*, 45: 44–55. DOI: 10.3906/vet-2004-139
- Cavigli I, Daughenbaugh KF, Martin M, Lerch M, Banner K, Garcia E, Brutscher LM, Flenniken ML (2016) Pathogen prevalence and abundance in honey bee colonies involved in almond pollination. *Apidologie*. 47(2):251–266. DOI: 10.1007/s13592-015-0395-5.
- Chen Y, Evans J, Feldlaufer M (2006) Horizontal and vertical transmission of viruses in the honey bee, *Apis mellifera*. *J Invertebr Pathol*, 92: 152–159. DOI: 10.1016/j.jip.2006.03.010
- Chen Y, Zhao Y, Hammond J, Hsu HT, Evans J, Feldlaufer M (2004) Multiple virus infections in the honey bee and genome diver-

- gence of honey bee viruses. *J Invertebr Pathol*, 87: 84-93. DOI: 10.1016/j.jip.2004.07.005.
- Choe SE, Nguyen TTD, Hyun BH, Noh JH, Lee HS, Lee CH, Kang SW (2012) Genetic and phylogenetic analysis of South Korean sacbrood virus isolates from infected honey bees (*Apis cerana*). *Vet Microbiol*, 157: 32-40. DOI: 10.1016/j.vetmic.2011.12.007
- Daughenbaugh KF, Martin M, Brutscher LM, Cavigli I, Garcia E, Lavin M, Flenniken ML (2015) Honey Bee Infecting Lake Sinai Viruses. *Viruses*, 7(6): 3285-309. DOI: 10.3390/v7062772.
- de Miranda JR (2008) Diagnostic techniques for virus detection in honey bees. In: Aubert MFA, Ball BV, Fries I, Morritz RFA, Milani N, Bernardinelli I (Eds.): *Virology and the Honey Bee*. Luxembourg: EEC Publications, pp. 121-232.
- de Miranda JR, Bailey L, Ball BV, Blanchard P, Budge GE, Chejanovsky N, Chen YP, Gauthier L, Genersch E, de Graaf DC, Ribière M, Ryabov E, De Smet J, Van Der Steen JJM (2013) Standard methods for virus research in *Apis mellifera*. *J Apic Res*, 52(4): 1-56. DOI: 10.3896/IBRA.1.52.4.22.
- Genersch E (2010) Honey bee pathology: current threats to honey bees and beekeeping. *Appl Microbiol Biotechnol*, 87: 87-97. DOI: 10.1007/s00253-010-2573-8
- Haddad N, Brake M, Migdadi H, de Miranda JR (2008) First detection of honey bee viruses in Jordan by RT-PCR. *Jordan Journal of Agricultural Sciences*, 4 (3): 242-247.
- Kalayci G, Cagiran AA, Kaplan M, Pekmez K, Beyazit A, Ozkan B, Yesiloz B, Arslan F (2020) The role of viral and parasitic pathogens affected by colony losses in Turkish apiaries. *Kafkas Univ Vet Fak Derg*. DOI: 10.9775/kvfd.2020. 24154.
- Kevill JL, Highfield A, Mordecai GJ, Martin SJ, Schroeder DC (2017) ABC Assay: Method Development and Application to Quantify the Role of Three DWV Master Variants in Overwinter Colony Losses of European Honey Bees. *Viruses*, 9 (11): 314. DOI: 10.3390/v9110314.
- Kızıltepe Ş, Akpınar R, Akman A, Kaya S, Çelik S (2023) İğdir İlinde Görülen Arı Kolonisi Kayıplarında Viral ve Paraziter Etkenlerin Rolü. *Journal of the Institute of Science and Technology*, 13 (2): 858-871. DOI: 10.21597/jist.1184308.
- Lanzi G, de Miranda JR, Boniotti MB, Cameron CE, Lavazza A, Capucci L, Camazine SM, Rossi C (2006) Molecular and biological characterization of deformed wing virus of honeybees (*Apis mellifera* L.). *J Virol*, 80(10): 4998-5009. DOI: 10.1128/jvi.80.10.4998-5009.2006.
- Li J, Wang T, Evans JD, Rose R, Zhao Y, Li Z, Li J, Huang S, Heerman M, Rodríguez-García C, Banmekea O, Brister JR, Hatcher EL, Cao L, Hamilton M, Chen Y (2019) The Phylogeny and Pathogenesis of Sacbrood Virus (SBV) Infection in European Honey Bees, *Apis mellifera*. *Viruses*, 11(1):61. DOI: 10.3390/v11010061.
- Lu C, Hung YT, Cheng Q (2020) A review of sub-lethal neonicotinoid insecticides exposure and effects on pollinators. *Curr Pollution Rep*, 6(2): 137-151. DOI: 10.1007/s40726-020-00142-8
- Martin SJ (2001) The role of Varroa and viral pathogens in the collapse of honeybee colonies: a modeling approach. *J Appl Ecol*, 38: 1082-1093. DOI: 10.1046/j.1365-2664.2001.00662.x
- McMenamin AJ, Flenniken ML (2018) Recently identified bee viruses and their impact on bee pollinators. *Curr Opin Insect Sci*, 26: 120-129. DOI: 10.1016/j.cois.2018.02.009
- Moharrami M, Modirrousta H (2018) Molecular identification of six honeybee viruses in Iranian apiaries. *Arch Razi Inst*, 73 (4): 311-318. DOI: 10.22092/ARI.2017.107481.1068
- Morawetz L, Kogelberger H, Griesbacher A, Derakhshifa I, Crailsheim K, Brodschneide R, Moosbeckhofe R (2019) Health status of honey bee colonies (*Apis mellifera*) and disease-related risk factors for colony losses in Austria. *PLoS ONE* 14 (7): e0219293. DOI: 10.1371/journal.pone.0219293
- Muz D, Muz MN (2023a) Israel acute bee paralysis virus and its prevalence in apiaries with colony loss in Türkiye. *U. Bee J*, 23(1): 105-114. DOI: 10.31467/uluaricilik.1265816.
- Muz D, Muz MN (2023b) Acute bee paralysis virus field isolates from apiaries suffering colony losses in Türkiye. *J Apic Res*, 62 (5): 1169-1175. DOI: 10.1080/00218839.2023.2242123.
- Nielsen SL, Nicolaisen M, Kryger P (2008) Incidence of acute bee paralysis virus, black queen cell virus, chronic bee paralysis virus, deformed wing virus, Kashmir bee virus and sacbrood virus in honey bees (*Apis mellifera*) in Denmark. *Apidologie*, 39: 310-314. DOI: 10.1051/apido:2008007.
- Ongus JR, Peters D, Bonmatin JM, Bengsch E, Vlak JM, van Oers MM (2004) Complete sequence of a picorna-like virus of the genus Iflavirus replicating in the mite Varroa destructor. *J Gen Virol*, 85 (12):3747-3755. DOI: 10.1099/vir.0.80470-0.
- Oz ME, Avci O, Dogan M (2023) Impact of deformed wing virus master variants (DWV-A, DWV-B, and DWV-C) in managed honey bee colonies of Türkiye. *Eurasian J Vet Sci*, 39 (3): 124-131. DOI: 10.15312/EurasianJVetSci.2023.407.
- Parveen N, Miglani R, Kumar A, Dewali S, Kumar K, Sharma N, Bisht SS (2022) Honey bee pathogenesis posing threat to its global population: A short review. *Proc Indian Natl Sci Acad*, 88(1): 11-32. DOI: 10.1007/s43538-022-00062-9
- Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE (2010) Global pollinator declines: Trends, impacts and drivers. *Trends Ecol Evol*, 25: 345-353. DOI: 10.1016/j.tree.2010.01.007
- Remnant EJ, Shi M, Buchmann G, Blacquière T, Holmes EC, Beekman M, Ashe A (2017) A diverse range of novel RNA viruses in geographically distinct honey bee populations. *J Virol*, 91:e00158-17. DOI: 10.1128/jvi.00158-17.
- Rüstemoğlu M, Sipahioğlu HM (2019) Occurrence and prevalence of six honey bee viruses in Hakkari (Türkiye) and their genomic divergence. *Mun Ent Zool*, 14(2): 574-583.
- Ryabov EV, Childers AK, Chen Y, Madella S, Nessa A, vanEngelsdorp D, Evans JD (2017) Recent spread of Varroa destructor virus-1, a honey bee pathogen, in the United States. *Sci Rep*, 7: 17447. DOI:10.1038/s41598-017-17802-3.
- Schurr F, Tison A, Militano L, Cheviron N, Sircoulomb F, Riviere MP, Chabert MR, Thierry R, Dubois E (2019) Validation of Quantitative Real-Time RT-PCR Assays for the Detection of Six Honeybee Viruses. *J Virol Methods*, 270: 70-78. DOI: 10.1016/j.jviromet.2019.04.020
- Sguazza GH, Reynaldi FJ, Galosi CM, Pecoraro MR (2013) Simultaneous detection of bee viruses by multiplex PCR. *J Virol Methods*, 194:102-106. DOI: 10.1016/j.jviromet.2013.08.003.
- Tantillo G, Bottaro M, Di Pinto A, Martella V, Di Pinto P, Terio V (2015) Virus infections of honeybees *Apis mellifera*. *Ital J Food Saf*, 30: 4 (3): 5364. DOI: 10.4081/ijfs.2015.5364
- Tentcheva D, Gauthier L, Zappulla N, Dainat B, Cousserans F, Colin ME, Bergoin M (2004) Prevalence and seasonal variations of six bee viruses in *Apis mellifera* L. and Varroa destructor mite populations in France. *Appl Environ Microbiol*, 70 (12):7185-7191. DOI: 10.1128/AEM.70.12.7185-7191.2004.
- Tlak Gajger I, Kolodziejek J, Bakonyi T, Nowotny N (2014) Prevalence and distribution patterns of seven different honeybee viruses in diseased colonies: a case study from Croatia. *Apidologie*, 45:701-706. DOI: 10.1007/s13592-014-0287-0.
- Tozkar CÖ, Kence M, Kence A, Huang Q, Evans JD (2015) Metatranscriptomic analyses of honey bee colonies. *Front Genet*, 6:100. DOI: 10.3389/fgene.2015.00100.
- Ullah A, Gajger IT, Majoros A, Dar SA, Khan S, Shah AH, Khabir MN, Hussain R, Khan HU, Hameed M, Anjum SI (2021) Viral impacts on honey bee populations: A review. *Saudi J Biol Sci*, 28(1): 523-530. DOI: 10.1016/j.sjbs.2020.10.037

- Valles SM, Chen Y, Firth AE, Guérin DMA, Hashimoto Y, Herrero S, de Miranda JR, Ryabov E (2017) ICTV Report Consortium. ICTV Virus Taxonomy Profile: Iflaviridae. *J Gen Virol*, 98(4):527-528. DOI: 10.1099/jgv.0.000757.
- Wei R, Cao L, Feng Y, Chen Y, Chen G, Zheng H (2022) Sacbrood virus: a growing threat to honeybees and wild pollinators. *Viruses*, 14(9): 1871. DOI: 10.3390/v14091871
- Wyszkowska J, Grodzicki P, Szczygieł M (2019) Electromagnetic fields and colony collapse disorder of the Honey bee. *Przegląd Elektrotechniczny*, 95(1): 137-40.