

Journal of the Hellenic Veterinary Medical Society

Vol 71, No 3 (2020)



In vitro anticoccidial activity of *Vitis vinifera* extract on oocysts of different *Eimeria* species of Broiler Chicken

R.Z. ABBAS, A. ABBAS, Z. IQBAL, M.A. RAZA, K. HUSSAIN, T. AHMED, M.U. SHAFI

doi: [10.12681/jhvms.25071](https://doi.org/10.12681/jhvms.25071)

Copyright © 2020, R.Z. ABBAS, A. ABBAS, Z. IQBAL, M.A. RAZA, K. HUSSAIN, T. AHMED, M.U. SHAFI



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/).

To cite this article:

ABBAS, R., ABBAS, A., IQBAL, Z., RAZA, M., HUSSAIN, K., AHMED, T., & SHAFI, M. (2020). In vitro anticoccidial activity of *Vitis vinifera* extract on oocysts of different *Eimeria* species of Broiler Chicken. *Journal of the Hellenic Veterinary Medical Society*, 71(3), 2267–2272. <https://doi.org/10.12681/jhvms.25071>

In vitro anticoccidial activity of *Vitis vinifera* extract on oocysts of different *Eimeria* species of Broiler Chicken

R.Z. Abbas¹, A. Abbas*², Z. Iqbal², M.A. Raza², K. Hussain², T. Ahmed³, M.U. Shafi³

¹Department of Parasitology, University of Agriculture, Faisalabad, Pakistan

²Department of Veterinary and Animal Sciences, Muhammad Nawaz Sharif University of Agriculture Multan, Pakistan

³Department of Clinical Medicine, Bahauddin Zakriya University, Multan

⁴Department of Animal Breeding and Genetics, University of Agriculture, Faisalabad, Pakistan

ABSTRACT: In the current experiment, the *in vitro* anticoccidial effect of *Vitis vinifera* (grape seed) extract was evaluated. For this purpose, an *in vitro* sporulation inhibition assay was used. Collected oocysts of four *Eimeria* species (*E. tenella*, *E. necatrix*, *E. brunetti* and *E. mitis*) were exposed to six different concentrations (w/v) of *Vitis vinifera* extract (VVE) in 10% Dimethylsulphoxide solution (DMSO), while Dimethylsulphoxide (DMSO) and Potassium dichromate solution ($K_2Cr_2O_7$) served as control groups. The results of the present study revealed that *V. vinifera* extract showed inhibitory effect on sporulation (%) and damage (%) of *Eimeria* oocysts in a dose dependent manner as compared to both control groups. *V. vinifera* extract also damaged the morphology of oocysts in terms of shape, size and number of sporocysts.

Keywords: *Vitis vinifera*, *in vitro*, *Eimeria*, oocysts

Corresponding Author:

A. Abbas, Department of Veterinary and Animal Sciences, Muhammad Nawaz Sharif University of Agriculture Multan, Pakistan
E-mail address: abbasasghar255@gmail.com

Date of initial submission: 14-12-2018
Date of revised submission: 14-12-2018
Date of acceptance: 24-04-2020

INTRODUCTION

Coccidiosis is an important disease infecting the intestine of chicken, which is caused by *Eimeria* (protozoa) species (Alzahrani *et al.*, 2016; Abbas *et al.*, 2019a). Coccidiosis causes heavy economic losses to poultry farming in different parts of world (Bachaya *et al.*, 2015; Abbas *et al.*, 2017). Disease has various clinical features such as poor weight gain, high mortality and bloody feces (Masood *et al.*, 2013). Oocysts of *Eimeria* sporulate rapidly in soil having high multiplication rate due to which its prevention is difficult once its outbreak has occurred at poultry farm (Zaman *et al.*, 2012). Poultry coccidiosis has been controlled by using synthetic anticoccidial drugs but, due to their frequent and irrationale use resistance has been developed due to which this method is ineffective in controlling of disease (Alzahrani *et al.*, 2016; Abbas *et al.*, 2017).

So, in matter of achieving success in controlling this severe disease other options and protocols are effectively used in different countries of world (Zhang *et al.*, 2018; Bakr *et al.*, 2019; Sarwar *et al.*, 2019). Among other options plant-derived compounds and their products have shown better anticoccidial effects (Gadelhaq *et al.*, 2018; Fariha *et al.*, 2019; Zhang *et al.*, 2020). Botanicals such as *Camellia sinensis* (Abbas *et al.*, 2017), *Ageratum conyzoides* (Nweze and Obiwulu, 2009), *Vitis vinifera* (Wang *et al.*, 2008), *Saccharum officinarum* (Abbas *et al.*, 2015) are reported to have excellent anticoccidial and immunomodulatory activity against coccidiosis.

Vitis vinifera commonly known as grape is one of the most abundant cultivated plants all over the world and is rich with various usefull antioxidant compounds including flavanoids, anthocyanins, catechin and epicatechins. These antioxidant compounds are well known for their therapeutic, health beneficial and immunomodulatory effects in poultry and other livestock (Kara *et al.*, 2016; Kasapidou *et al.*, 2016). Due to its availibility in large quantity and its content in bioactive compounds contained in it is a suitable candidate for improving poultry production.

Based on the various therapeutic and health beneficial effects of *V. vinifera* and very limited published research on its effect against *Eimeria*, the current experiment was conducted to evalaute *in vitro* anticoccidial potential of *V. vinifera* extract a on oocysts of four *Eimeria* species of broiler chicken.

MATERIALS AND METHODS

Preparation of *Vitis vinifera* extract

Seeds of *V. vinifera* were obtained from local market in Faisalabad, Pakistan. The plant material were authenticated by botanist of University of Agriculture Faisalabad and were dried and converted in powder form using an electric grinder. Aqueous methanolic extract of *V. vinifera* (seeds) was prepared using using Soxhlet's apparatus (Velp Italy) following method described by Abbas *et al.* (2015). Prepared *V. vinifera* extract (VVE), was stored at 4°C untill further use.

Collection of *Eimeria* oocysts

Oocysts of four *Eimeria* species were collected from the caeca of infected broilers from different reported cases in Faisalabad. Collected oocysts were preserved in potassium dichromate solution (2.5%) following the procedure as described by Ryley *et al.* (1976).

Experimental design

The experiment was reviewed and approved by Research and Ethics Committee of Department of Parasitology, University of Agriculture Faisalabad, Pakistan. *In vitro* efficacy of *V. vinifera* extract (VVE) was evaluated by a sporulation inhibition assay. For this purpose, unsporulated oocysts (100 oocysts/5ml) of four *Eimeria* species (*E. tenella*, *E. brunetti*, *E. necatrix* and *E. mitis*) were exposed to different concentrations (w/v; 10, 5, 2.5, 1.25, 0.625 and 0.31%) of VVE in 10% Dimethylsulfoxide (DMSO) solution in 5cm petri dishes by making two fold serial dilutions. DMSO and potassium dichromate solution ($K_2Cr_2O_7$) served as control groups. Incubation of *Eimeria* oocysts was done for 48 hours at 27-29°C and 60% humidity. Three replications were made for each concentration. The sporulation process of *Eimeria* oocysts was checked under light microscope at 40x. A total of 40 oocysts for each *Eimeria* species were counted in all treatment and control groups. The oocysts with 4 sporocysts was considered sporulated regardless the shape and size of the sporocysts. *Eimeria* oocysts having damaged wall and misshapen were considered damaged. The Percentage of sporulation and damaged of each *Eimeria* species was determined out of total counted oocysts. Sporulation inhibition (SI) and damage of *Eimeria* oocysts was determined in percentage by following the method by You (2014).

Statistical analysis

Data were analyzed by Analysis of Variance

(ANOVA) and significance among groups was determined at $P < 0.05$. For comparisons of means Duncan's multiple range test was used.

RESULTS

The statistical analysis showed that all dilutions of *V. vinifera* extract (VVE) significantly affected the sporulation process of *Eimeria* oocysts of all the four species (*E. tenella*, *E. necatrix*, *E. brunetti* and *E. mitis*) as compared to both control groups (Control 1: DMSO, Control 2: Potassium dichromate solution ($K_2Cr_2O_7$)) Figure 1. Effect of *Vitis vinifera* on percent damage of *Eimeria* oocysts is shown in Figure 2. *V. vinifera* extract damaged internal and external the morphology of of *Eimeria* oocysts in concentration-de-

pendent manner as compared to both control groups (Control 1: DMSO, Control 2: Potassium dichromate solution ($K_2Cr_2O_7$)). Among all tested *Eimeria* species *V. vinifera* extract at higher dose significantly reduced sporulation process and damaged oocysts of *E. tenella* and *E. necatrix*. Oocysts of four *Eimeria* species (*E. tenella*, *E. necatrix*, *E. brunetti* and *E. mitis*) were differentiated on the basis of their morphology following method as described Abbas *et al.* (2019b). Photomicrographs of normal *Eimeria* oocysts and damaged *Eimeria* oocysts by *V. vinifera* extract in terms shape and wall are shown in Figure 3. It can be clearly seen in Figure 3 that *V. vinifera* extract inhibited sporulation and also damaged the morphology of *Eimeria* oocysts.




Figure 1: Effect of *V. vinifera* extract on % sporulation of oocysts of four *Eimeria* species Control-1 (DMSO) and Control-2 ($K_2Cr_2O_7$). Asterisks (*) indicate a difference from the control groups




Figure 2: Effect of *Vitis vinifera* extract on percent damage of oocysts of four *Eimeria* species. Control-1 (DMSO) and Control-2 ($K_2Cr_2O_7$). Asterisks (*) indicate a difference from the control groups




Figure 3: Photomicrographs of *Eimeria* oocysts

A: Normal sporulated oocysts of *Eimeria*.

B & C: Damaged *Eimeria* oocysts by *V. Vinifera* extract in terms shape and wall

DISCUSSION

Many botanicals and their products are reported to have excellent anticoccidial activity as proven by different *in vitro* and *in vivo* studies (Abbas *et al.*, 2015, 2017). In the present, as well as in previous studies (Zaman *et al.*, 2012; Gadelhaq *et al.*, 2018; Mujahid *et al.*, 2019), the *in vitro* anticoccidial effect of *V. vinifera* extract was measured in terms of percent sporulation inhibition and damage of *Eimeria* oocysts. The results showed an inhibitory effect on sporulation and damage of *Eimeria* oocysts in dose dependent manner.

In addition, *V. vinifera extract* also affected the morphology of *Eimeria* oocysts in terms of abnormal shape of oocysts and sporocysts. Likewise, an aqueous extract of pine bark also showed a similar effect on sporulation of *Eimeria* oocysts (Molan *et al.*, 2009). In a recent study, Gadelhaq *et al.* (2018) have reported the *in vitro* anticoccidial effects of chemicals and natural products. They concluded that commonly used disinfectants such as formalin and ethanol (70%) are the most effective in inhibition of sporulation process of different *Eimeria* species.

Abbas *et al.* (2019b) has reported *in vitro* anticoccidial effects of *Trachyspermum ammi* extract on oocysts of four *Eimeria* species of chickens. *T. ammi* inhibited sporulation of *Eimeria* oocysts and also damaged them. *T. ammi* effect was in dose dependent manner against *Eimeria* oocysts.

Abbas *et al.* (2015) have reported a similar, dose-dependent, *in vitro* anticoccidial effect of *S. officinarum* (sugar cane) extract on inhibition sporulation of *Eimeria* oocysts in dose dependent manner. Such high *in vitro* anticoccidial potential of *V. vinifera* extract might be due to action of its antioxidant compounds against *Eimeria*.

Somewhat similar *in vitro* effects of *Camellia sinensis* on the sporulation of various *Eimeria* species

has been reported previously and a significant reduction in sporulation rate of *Eimeria* oocysts was observed after exposure to *C. sinensis* extract (Molan and Thomas, 2007).

In another study, *in vitro* destruction of *Eimeria* oocysts by essential oils has been reported (Remmal *et al.*, 2013).

In the present study *V. vinifera* showed an *in vitro* anticoccidial potential against *Eimeria* oocysts which might be due to the action of its various flavonoids so-called antioxidant compounds including anthocyanins, catechin and epicatechins. However further studies and *in vivo* trials are needed to understand its anticoccidial effect in poultry.

CONCLUSION

It was concluded from the results of the present study that *V. vinifera* (grape seed) extract have *in vitro* anticoccidial potential against four *Eimeria* species. *V. vinifera* extract damaged the morphology and inhibited the sporulation process of *Eimeria* oocysts of all tested species. *In vitro* results of this study can help in future to explore and develop effective herbal remedy based on antioxidant compounds of *V. vinifera* by conducting *in vivo* trials for treating poultry coccidiosis.

ACKNOWLEDGEMENTS

All Authors acknowledge the financial support for this work from Punjab Agricultural research board (PARB), Pakistan.

CONFLICT OF INTEREST

The authors have no conflict of interest with this publication.

REFERENCES

- Abbas A, Z Iqbal, Abbas RZ, Khan MK, Khan JA (2017) Immunomodulatory activity of *Pinus radiata* extract against coccidiosis in broiler chicken. Pak Vet J 37: 145-149.
- Abbas A, Abbas RZ, Khan MK, Raza MA, Mahmood MS, Saleemi MK, Hussain T, Khan JA, Sindhu ZD (2019a) Anticoccidial Effects of *Trachyspermum ammi* (Ajwain) in broiler chickens. Pak Vet J 39: 301-304.
- Abbas A, Abbas RZ, Raza MA, Khan MK, Saleemi MK, Saeed Z (2019b) *In vitro* anticoccidial activity of *Trachyspermum ammi* (Ajwain) extract on oocysts of *Eimeria* species of Chicken. Adv Life Sci 7: 44-47.
- Alzahrani F, Al-Shaebi EM, Dkhil MA, Al-Quraishy S (2016) *In vivo* anti-*Eimeria* and *in vitro* anthelmintic activity of *Ziziphusspina-christi* leaf extracts fares, Pak J Zool 48: 409-413.
- Abbas A, Iqbal Z, Abbas RZ, Khan MK, Khan JA (2015) *In-vitro* anticoccidial potential of *Sacharrumofficinarum* extract against *Eimeria* Oocysts. Bollatinoam. Caribe Plantas Med Aromát14: 456-46.
- Bachaya HA, Abbas RZ, Raza MAM, Iqbal Z, Rehman TU, Baber W, Hussain R (2015). Existence of coccidiosis and associated risk factors in broiler chickens in Southern Punjab, Pakistan. Pak Vet J 35: 81-84.
- Bakr AF, Abdelgayed SS, EL-Tawil OS, Bakeer AM (2019) Assessment of ginger extract and ginger nanoparticles protective activity against acetaminophen-induced hepatotoxicity and nephrotoxicity in rats. Pak Vet J 39: 479-486.
- Farieha, Jahan N, Rehman KU and Ali S (2019) Effect of herbal mixture as Angiotensin converting enzyme inhibitor in Angiotensin-II dependent hypertension. Pak Vet J 39: 25-30.
- Gadelhaq SM, Arafa WM, Abolhadid SM (2018) *In vitro* activity of natural and chemical products on sporulation of *Eimeria* species oocysts of chickens. Vet Parasitol 251:12-6.
- Kara K, Guclu BK, Baytok E, Senturk M (2016) Effects of grape pomace supplementation to laying hen diet on performance, egg quality, egg lipid peroxidation and some biochemical parameters. J Appl Animal Res 44: 303-10.
- Kasapidou E, Soddidou EN, Zdragas A, Papadaki C, Vafeas G, Mtianga P (2016) Effect of grape pomace supplementation on broiler meat quality characteristics. Eur Poult Sci 80:135-42
- Mujahid H, Hashmi AS, Khan MZ, Tayyab M, Shezad W (2019) Protective effect of yeast sludge and whey powder against ochratoxigenin in broiler chicks. Pak Vet J 39: 588-592.
- Masood S, Abbas RZ, Iqbal Z, Mansoor MK, Sindhu ZD, Zia MA, Khan JA (2013) Role of natural antioxidants for the control of coccidiosis in Poultry. Pak Vet J 33: 401-407.
- Molan AL, Thomas DV (2007) Green tea water extracts inhibit the sporulation of *Eimeria* oocysts *in vitro*. Advancing Poultry Production – Proceedings of the Massey Technical Update Conference. 9: 89–97.
- Molan AL, Zhuojian L, De S (2009) Effect of pine bark (*Pinusradiata*) extracts on sporulation of coccidian oocysts. Folia Parasitol 56: 1–5.
- Nweze NE, Obiwulu IS (2009) Anticoccidial effects of *Ageratum conyzoides*. J Ethnopharmacol 122: 6–9
- Remmal A, Sanaa A, Latifa B, Fouzia C, Najat C (2013) Oocysticidal Effect of Essential Oil Components against Chicken *Eimeria* Oocysts. Int J Vet Med 2: 133-139.
- Ryley JF, Meade R, burst JH, Robinson TE (1976) Methods in coccidiosis research: Separation of oocysts from faeces. J Parasitol 73: 311-326.
- Sarwar N, Mehmood A, Sheraz A, Noman M (2019) Protective effect of probiotics in combination with vaccination on antibody response, biochemical and hematological indices in broilers. Pak Vet J 39: 451-454.
- You MJ (2014) Suppression of *Eimeria tenella* sporulation by disinfectants. Korean J Parasitol 52: 435–438.
- Wang MI, Suo X, Gu JH, Zhang WW, Fang Q, Wang X (2008) Influence of grape seed proanthocyanidin extract in broiler chickens: effect on chicken coccidiosis and antioxidant status. Poult Sci 87:2273-2280.
- Zaman MA, Iqbal Z, Abbas RZ, Khan MN (2012) Anticoccidial activity of herbal complex in broiler chickens challenged with *Eimeria tenella*. Parasitology 139: 237-243.
- Zhang QH, Nie FH, Wang XN, Lin HY, Wang HCR, Gooneratne R, Chen JJ (2018) Effect of flavonoids from *Eupatorium odoratum* L. on immunoglobulins and cecal microflora in broilers. Pak Vet J 38: 296-300.
- Zhang K, Li X, Na C, Abbas A, Abbas RZ, Zaman MA (2020) Anticoccidial effects of *Camellia sinensis* (green tea) extract and its effect on Blood and Serum chemistry of broiler chickens. PakVet J 40: 77-80